

Appendix 3.2. Breakdown of river channel length by water quality category within hydrometric areas.

NWIRBD: HYDROMETRIC AREA NO. 01

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|----------|-------|------|-------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Bunadaowen | 2007 | 01B01 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 |
| Burn Daurnett | 2007 | 01B02 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Carrigans | 2007 | 01C01 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Clogher (Finn) | 2008 | 01C06 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 7.0 |
| Cloghroe | 2007 | 01C05 | 0.0 | 6.0 | 0.0 | 1.0 | 0.0 | 7.0 |
| Cronamuck ¹ | 2008 | 01C02 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 1.0 |
| Cross Roads Stream | 2007 | 01C04 | 0.0 | 3.0 | 0.0 | 1.0 | 0.0 | 4.0 |
| Cummirk | 2007 | 01C03 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 9.0 |
| Deele (Donegal) | 2007 | 01D01 | 11.0 | 24.5 | 2.0 | 0.5 | 0.0 | 38.0 |
| Elatagh | 2007 | 01E02 | 0.0 | 0.0 | 0.0 | 7.0 | 0.0 | 7.0 |
| Finn | 2007 | 01F01 | 0.0 | 20.5 | 3.0 | 28.0 | 0.0 | 51.5 |
| Greenhill Stream ¹ | 2008 | 01G02 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Mourne Beg | 2007 | 01M01 | 0.0 | 0.0 | 0.0 | 16.5 | 0.0 | 16.5 |
| Reelan | 2008 | 01R01 | 0.0 | 0.0 | 15.0 | 0.0 | 0.0 | 15.0 |
| Rough Burn | 2007 | 01R02 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| St Johnston ¹ | 2008 | 01S01 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| Stranagappoge | 2007 | 01S02 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Swilly Burn | 2007/08 | 01S03 | 0.0 | 4.0 | 8.5 | 2.0 | 1.5 | 16.0 |
| Total Length (km) surveyed this cycle | 2007/'08 | | 11.0 | 69.5 | 42.5 | 82.0 | 1.5 | 206.5 |
| Adjustments (See below)* | | | 0.0 | 0.0 | 7.0 | 1.0 | 0.0 | 8.0 |
| Percentages | | | 5.5 | 35.0 | 17.9 | 40.8 | 0.8 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline: Current adjusted Status (km) | 2007/'08 | | | 80.5 | 35.5 | 81.0 | 1.5 | 198.5 |
| Percentages | | | | 40.6 | 17.9 | 40.8 | 0.8 | 100.0 |
| Baseline: Previous Unadjusted Status (km) | 2004-'06 | | | 92.5 | 55.5 | 50.5 | 0.0 | 198.5 |
| Percentages | | | | 46.6 | 28.0 | 25.4 | 0.0 | |
| Changes since Previous Survey (Km) | | | | -12.0 | -20.0 | 30.5 | 1.5 | 0.0 |

* Adjustments : Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics

Hydrometric Area 01 : Trends % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 01 : 2007 - '09 % Surveyed Channel Length in Five WFD Quality Classes



NBIRBD: HYDROMETRIC AREA NO. 03

The baseline rivers, showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|----------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Blackwater (Monaghan) | 2007 | 03B01 | 0.0 | 6.0 | 10.0 | 10.5 | 0.0 | 26.5 |
| Clontibret Stream | 2007 | 03C01 | 0.0 | 0.0 | 0.0 | 13.0 | 0.0 | 13.0 |
| Conawary (Lower) | 2007 | 03C02 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Mountain Water | 2007 | 03M01 | 0.0 | 13.5 | 2.5 | 8.0 | 0.0 | 24.0 |
| Scotstown | 2007 | 03S02 | 6.5 | 5.0 | 0.0 | 0.0 | 0.0 | 11.5 |
| Shambles ¹ (no distances, all C) | 2007 | 03S01 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total Length (km) surveyed this cycle | 2007 | | 6.5 | 24.5 | 12.5 | 34.5 | 0.0 | 78.0 |
| Adjustments (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 8.3 | 31.4 | 16.0 | 44.2 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | | | | | Total |
| Baseline: Current adjusted Status (km) | 2007-'09 | | | A | B | C | D | I |
| | | | | 31.0 | 12.5 | 34.5 | 0.0 | 78.0 |
| Percentages | | | | 39.7 | 16.0 | 44.2 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km) | 2004-'06 | | | 31.0 | 9.5 | 37.5 | 0.0 | 78.0 |
| | | | | 39.7 | 12.2 | 48.1 | 0.0 | 100.0 |
| Percentages | | | | 39.7 | 12.2 | 48.1 | 0.0 | 0 |
| Changes since Previous Survey (km) | | | | 0.0 | 3.0 | -3.0 | 0.0 | 0.0 |

* Adjustments : Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics

Hydrometric Area 03 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 03 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



NBIRBD: HYDROMETRIC AREA NO. 06

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---------------------------------------|-------|-------|----------|----------|----------|----------|--------------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballymascanlan | 2009 | 06B02 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Big (Louth) | 2009 | 06B01 | 0.0 | 8.0 | 0.0 | 2.0 | 0.0 | 10.0 |
| Carrickaslane Lough Stream | 2009 | 06C04 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Castletown | 2009 | 06C01 | 0.0 | 4.3 | 5.1 | 4.2 | 0.0 | 13.5 |
| County Water | 2009 | 06C03 | 0.0 | 5.3 | 4.8 | 2.4 | 0.0 | 12.5 |
| Cully Water | 2009 | 06C02 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Dee | 2009 | 06D01 | 0.0 | 24.1 | 16.0 | 9.4 | 0.0 | 49.5 |
| Drumsallagh Stream | 2009 | 06D07 | 3.3 | 3.7 | 0.0 | 0.0 | 0.0 | 7.0 |
| Fane | 2009 | 06F01 | 0.0 | 17.5 | 7.2 | 13.9 | 0.0 | 38.5 |
| Flurry | 2009 | 06F02 | 0.0 | 0.0 | 8.5 | 2.0 | 0.0 | 10.5 |
| Gentle Owen's Lake Stream | 2009 | 06G04 | 0.0 | 6.5 | 1.0 | 0.0 | 0.0 | 7.5 |
| Glyde | 2009 | 06G02 | 7.8 | 32.2 | 0.0 | 0.0 | 0.0 | 40.0 |
| Kilcurry | 2009 | 06K02 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Killary Water | 2009 | 06K01 | 0.0 | 0.0 | 5.5 | 1.0 | 0.0 | 6.5 |
| Kilmainham | 2009 | 06K04 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Magheracloone Stream | 2009 | 06M01 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Proules | 2009 | 06P01 | 0.0 | 3.3 | 3.7 | 3.0 | 0.0 | 10.0 |
| Termonfeckin | 2009 | 06T01 | 0.0 | 0.0 | 3.5 | 1.5 | 0.0 | 5.0 |
| White (Louth) | 2009 | 06W01 | 0.0 | 12.4 | 2.6 | 0.0 | 0.0 | 15.0 |
| Total Length (km) surveyed this cycle | 2009 | | 11.1 | 143.2 | 62.9 | 40.3 | 0.0 | 257.5 |
| Adjustments (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 4.3 | 55.6 | 24.4 | 15.6 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline: Current adjusted Status | 2007- | | | | | | | |
| (km) | '09 | | 154.3 | 62.9 | 40.3 | 0.0 | 257.5 | |
| Percentages | | | 59.9 | 24.4 | 15.6 | 0.0 | 100.0 | |
| Baseline: Previous unadjusted Status | 2004- | | | | | | | |
| (km) | '06 | | 137.0 | 92.5 | 26.0 | 2.0 | 257.5 | |
| Percentages | | | 53.2 | 35.9 | 10.1 | 0.8 | 100.0 | |
| Changes since Previous Survey (km) | | | 17.3 | -29.6 | 14.3 | -2.0 | 0.0 | |


Raskeagh (06R02) & Ballykelly (06B03) discontinued in 2001- '03 period.

* Adjustments : Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics

Hydrometric Area 06 : Trends % Surveyed Channel in Four Quality Classes



Hydrometric Area 06 : 2007 - '09 % Surveyed Channel Length in Five WFD Quality Classes



ERBD: HYDROMETRIC AREA NO. 07

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class WFD Quality Class | | | A High | A Good | B Moderate | C Poor | D Bad | Total |
|---|----------|-------|-----------|-----------|---------------|-----------|----------|--------------|
| River Name | Year | Code | | | | | | |
| Athboy | 2009 | 07A01 | 0.0 | 19.1 | 10.9 | 0.0 | 0.0 | 30.0 |
| Blackwater (Kells) | 2009 | 07B01 | 0.0 | 30.9 | 25.1 | 8.0 | 0.0 | 64.0 |
| Blackwater (Longwood) | 2009 | 07B02 | 0.0 | 11.2 | 9.3 | 2.6 | 0.0 | 23.0 |
| Boycetown | 2009 | 07B03 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 9.0 |
| Boyne | 2009 | 07B04 | 15.8 | 40.0 | 38.2 | 0.0 | 0.0 | 94.0 |
| Castlejordan | 2009 | 07C04 | 0.0 | 3.5 | 7.4 | 6.2 | 0.0 | 17.0 |
| Clady (Meath) ¹ | 2009 | 07C01 | 0.0 | 0.0 | 0.0 | 13.0 | 0.0 | 13.0 |
| Crosskeys Stream | 2009 | 07C07 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| D'Arcy's Crossroads Stream | 2009 | 07D06 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 7.0 |
| Deel (Raharney) | 2009 | 07D01 | 0.0 | 13.5 | 20.5 | 0.0 | 0.0 | 34.0 |
| Devlins | 2009 | 07D02 | 0.0 | 0.0 | 3.0 | 4.6 | 0.0 | 7.5 |
| Glash | 2009 | 07G02 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Kinnegad | 2009 | 07K01 | 0.0 | 15.1 | 1.9 | 0.0 | 0.0 | 17.0 |
| Knightsbrook | 2009 | 07K02 | 0.0 | 0.0 | 5.1 | 6.9 | 0.0 | 12.0 |
| Lislea | 2009 | 07L01 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Mattock | 2009 | 07M01 | 0.0 | 12.6 | 3.5 | 0.0 | 0.0 | 16.0 |
| Moynalty | 2009 | 07M03 | 0.0 | 10.7 | 23.8 | 0.0 | 0.0 | 34.5 |
| Mullagh Lough Stream | 2009 | 07M06 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 1.0 |
| Nadreegeel Lough Stream | 2009 | 07N01 | 0.0 | 9.5 | 2.0 | 0.0 | 0.0 | 11.5 |
| Riverstown | 2009 | 07R01 | 0.0 | 0.0 | 15.0 | 0.0 | 0.0 | 15.0 |
| Skane | 2009 | 07S01 | 0.0 | 0.0 | 0.0 | 16.5 | 0.0 | 16.5 |
| Stonyford | 2009 | 07S02 | 0.0 | 6.5 | 15.0 | 0.0 | 0.0 | 21.5 |
| Yellow (Blackwater) | 2009 | 07Y01 | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 2.5 |
| Yellow (Castlejordan) | 2009 | 07Y02 | 0.0 | 15.0 | 3.5 | 0.0 | 0.0 | 18.5 |
| Total Length (km) surveyed this cycle | 2009 | | 15.8 | 188.5 | 207.0 | 65.3 | 0.0 | 476.5 |
| Adjustments (See below)* | | | 0.0 | -6.0 | -4.0 | 7.5 | 0.0 | -2.5 |
| Percentages | | | 3.3 | 40.6 | 44.1 | 12.1 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline: Current Adjusted Status (km) | 2007-'09 | | | 210.3 | 211.0 | 57.8 | 0.0 | 479.0 |
| Percentages | | | | 43.9 | 44.1 | 12.1 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status. (km)** | 2004-'06 | | | 217.0 | 205.5 | 56.5 | 0.0 | 479.0 |
| Percentages | | | | 45.3 | 42.9 | 11.8 | 0.0 | |
| Changes since Previous Survey (km) | | | | -6.8 | 5.5 | 1.3 | 0.0 | 0.0 |


* Adjustments : Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics


Hydrometric Area 07 : Trends

% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 07 : 2007 - '09

% Surveyed Channel Length in Five WFD Quality Classes



ERBD: HYDROMETRIC AREA NO. 08

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|-------|------|----------|------|-------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballough Stream | 2008 | 08B03 | 0.0 | 0.0 | 0.0 | 8.0 | 0.0 | 8.0 |
| Ballyboghil | 2008 | 08B01 | 0.0 | 0.0 | 0.0 | 12.0 | 0.0 | 12.0 |
| Broadmeadow | 2008 | 08B02 | 0.0 | 0.0 | 0.0 | 17.0 | 0.0 | 17.0 |
| Delvin | 2008 | 08D01 | 0.0 | 0.0 | 8.0 | 5.5 | 0.0 | 13.5 |
| Dunshaughlin Stream | 2008/9 | 08D03 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Fairyhouse Stream | 2008 | 08F01 | 0.0 | 0.0 | 0.0 | 3.5 | 0.0 | 3.5 |
| Hurley | 2008 | 08H01 | 0.0 | 2.0 | 12.0 | 3.0 | 0.0 | 17.0 |
| Mosney | 2008 | 08M02 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Nanny | 2008 | 08N01 | 0.0 | 6.5 | 6.5 | 15.0 | 0.0 | 28.0 |
| Ratoath Stream | 2008 | 08R01 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Ward | 2007/8 | 08W01 | 0.0 | 0.0 | 0.0 | 15.0 | 0.0 | 15.0 |
| Total Length (km) surveyed this cycle | 2007-'09 | | 0.0 | 8.5 | 26.5 | 87.0 | 0.0 | 122.0 |
| Adjustments (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 0.0 | 7.0 | 21.7 | 71.3 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline: Current Adjusted Status (km) | 2007 - '09 | | 8.5 | 26.5 | 87.0 | 0.0 | 122.0 | |
| Percentages | | | 7.0 | 21.7 | 71.3 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 22.5 | 23.0 | 70.0 | 6.5 | 122.0 | |
| Percentages | | | 18.4 | 18.9 | 57.4 | 5.3 | | |
| Changes since Previous Survey (km) | | | -14.0 | 3.5 | 17.0 | -6.5 | 0.0 | |

* Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics

Hydrometric Area 08 : Trends % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 08 : 2007 - '09 % Surveyed Channel Length in Five WFD Quality Classes



ERBD: HYDROMETRIC AREA NO. 09

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|--------|-------|------|------|----------|------|------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Annalecka Brook | 2007 | 09A01 | 4.0 | 0.0 | 0.0 | 0.0 | 0.00 | 4.0 |
| Ballinagee | 2007 | 09B08 | 0.0 | 3.5 | 0.0 | 0.0 | 0.00 | 3.5 |
| Ballydonnel Brook | 2007 | 09B03 | 5.0 | 0.0 | 0.0 | 0.0 | 0.00 | 5.0 |
| Ballylow Brook | 2007 | 09B04 | 0.0 | 3.5 | 0.0 | 0.0 | 0.00 | 3.5 |
| Brittas | 2007 | 09B02 | 0.0 | 6.0 | 4.5 | 0.0 | 0.00 | 10.5 |
| Camac | 2007/8 | 09C02 | 4.0 | 0.0 | 0.0 | 15.5 | 0.00 | 19.5 |
| Clonshanbo | 2007 | 09C03 | 0.0 | 0.0 | 0.0 | 4.0 | 0.00 | 4.0 |
| Cock Brook | 2007 | 09C04 | 2.5 | 0.0 | 0.0 | 0.0 | 0.00 | 2.5 |
| Dodder | 2007 | 09D01 | 0.0 | 13.5 | 5.5 | 1.5 | 0.00 | 20.5 |
| Douglas (Liffey) | 2007 | 09D02 | 3.5 | 0.0 | 0.0 | 0.0 | 0.00 | 3.5 |
| Dunboyne Stream | 2007 | 09D04 | 0.0 | 0.0 | 0.0 | 2.5 | 0.00 | 2.5 |
| Glashaboy Brook | 2007 | 09G02 | 0.0 | 2.0 | 0.0 | 0.0 | 0.00 | 2.0 |
| Kilcullen Stream | 2007 | 09K02 | 0.0 | 6.0 | 0.0 | 0.0 | 0.00 | 6.0 |
| Kings (Liffey) | 2007 | 09K01 | 7.3 | 8.7 | 0.0 | 0.0 | 0.00 | 16.0 |
| Lemonstown Stream | 2007 | 09L03 | 9.0 | 0.0 | 0.0 | 0.0 | 0.00 | 9.0 |
| Liffey | 2007 | 09L01 | 4.5 | 38.5 | 55.5 | 1.5 | 0.00 | 100.0 |
| Lyreen | 2007 | 09L02 | 0.0 | 0.0 | 6.5 | 1.5 | 0.00 | 8.0 |
| Mayne | 2007 | 09M03 | 0.0 | 0.0 | 0.0 | 2.0 | 0.00 | 2.0 |
| Morell ¹ | 2007 | 09M01 | 3.5 | 3.0 | 4.5 | 0.0 | 0.00 | 11.0 |
| Owenadoher | 2007 | 09O01 | 4.5 | 0.0 | 1.5 | 1.0 | 0.00 | 7.0 |
| Painestown ¹ | 2007 | 09P01 | 0.0 | 0.0 | 6.0 | 0.0 | 0.00 | 6.0 |
| Poddle ¹ | 2007 | 09P03 | 0.0 | 0.0 | 0.0 | 2.0 | 0.00 | 2.0 |
| Pinkeen | 2007 | 09P03 | 0.0 | 0.0 | 2.5 | 0.0 | 0.00 | 2.5 |
| Rathmore Stream ¹ | 2007 | 09R02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.00 | 7.0 |
| Rye Water | 2007 | 09R01 | 0.0 | 11.0 | 9.5 | 0.5 | 0.00 | 21.0 |


| | | | | | | | | |
|---|------------|-------|----------|----------|----------|----------|--------------|-------|
| Santry | 2007 | 09S01 | 0.0 | 0.0 | 0.0 | 5.0 | 0.00 | 5.0 |
| Tolka | 2007/8 | 09T01 | 0.0 | 0.0 | 15.0 | 13.0 | 0.00 | 28.0 |
| Total Length (km) surveyed this cycle | 2007/'08 | | 47.8 | 102.7 | 111.0 | 50.0 | 0.0 | 311.5 |
| Adjustments (See below)* | | | 3.5 | 10.0 | 10.5 | 2.0 | 0.0 | 26.0 |
| <i>Percentages</i> | | | 15.5 | 32.5 | 35.2 | 16.8 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline: Current Adjusted Status (km) | 2007 - '09 | | 137.0 | 100.5 | 48.0 | 0.0 | 285.5 | |
| <i>Percentages</i> | | | 48.0 | 35.2 | 16.8 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 148.5 | 44.0 | 84.5 | 8.5 | 285.5 | |
| <i>Percentages</i> | | | 52.0 | 15.4 | 29.6 | 3.0 | | |
| Changes since Previous Survey (km) | | | -11.5 | 56.5 | -36.5 | -8.5 | 0.0 | |

* Adjustments : Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 09 : Trends % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 09 : 2007 - '09 % Surveyed Channel Length in Five WFD Quality Classes



ERBD: HYDROMETRIC AREA NO. 10

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Askanagap Stream | 2009 | 10A06 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Aughrim | 2009 | 10A02 | 0.0 | 9.5 | 0.0 | 0.0 | 0.0 | 9.5 |
| Avoca | 2009 | 10A03 | 0.0 | 0.0 | 0.0 | 0.0 | 11.5 | 11.5 |
| Avonbeg | 2009 | 10A04 | 7.0 | 6.5 | 7.5 | 0.0 | 0.0 | 21.0 |
| Avonmore | 2009 | 10A05 | 17.1 | 16.5 | 0.0 | 0.0 | 0.0 | 33.5 |
| Ballycreen Brook | 2009 | 10B02 | 0.0 | 3.5 | 1.5 | 0.0 | 0.0 | 5.0 |
| Ballyduff Stream | 2009 | 10B01 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 |
| Cloghoge Brook | 2009 | 10C01 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Coolalug Stream | 2009 | 10C02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Dargle | 2009 | 10D01 | 10.5 | 2.0 | 1.5 | 0.0 | 0.0 | 14.0 |
| Derry Water | 2009 | 10D02 | 13.5 | 0.0 | 1.5 | 0.0 | 0.0 | 15.0 |
| Glencree | 2009 | 10G01 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 9.0 |
| Glencullen | 2009 | 10G02 | 6.5 | 2.6 | 0.0 | 0.0 | 0.0 | 9.0 |
| Glendasan | 2009 | 10G06 | 4.8 | 0.2 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glenealo | 2009 | 10G05 | 5.5 | 0.0 | 0.5 | 0.0 | 0.0 | 6.0 |
| Glenmacnass | 2009 | 10G03 | 3.1 | 8.9 | 0.0 | 0.0 | 0.0 | 12.0 |
| Goldmine | 2009 | 10G04 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Kill O' the Grange Stream | 2009 | 10K02 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Kilmacanoge | 2009 | 10K03 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Newcastle | 2009 | 10N01 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 4.0 |
| Newtownmountkennedy | 2009 | 10N02 | 0.0 | 4.5 | 0.0 | 4.5 | 0.0 | 9.0 |
| Ow | 2009 | 10O01 | 0.0 | 2.0 | 13.5 | 0.0 | 0.0 | 15.5 |
| Potters | 2009 | 10P01 | 0.0 | 0.0 | 6.1 | 8.0 | 0.0 | 14.0 |


| | | | | | | | | |
|--|------------|-------|-------|----------|----------|----------|----------|--------------|
| Redcross | 2009 | 10R01 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Rathnew Stream ¹ (no distances yet) | 2009 | 10R02 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Shanganagh | 2009 | 10S01 | 0.0 | 6.0 | 7.0 | 0.0 | 0.0 | 13.0 |
| Templeraing Stream | 2009 | 10T04 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Three Mile Water | 2009 | 10T01 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Vartry | 2009 | 10V01 | 0.0 | 13.5 | 3.5 | 0.0 | 0.0 | 17.0 |
| <hr/> | | | | | | | | |
| Total Length (km) surveyed this cycle | 2007 - '09 | | 75.9 | 103.1 | 60.6 | 17.5 | 11.5 | 268.5 |
| Adjustments (See below)* | | | 0.0 | -1.5 | 0.0 | 0.0 | 0.0 | -1.5 |
| <i>Percentages</i> | | | 28.1 | 38.7 | 22.4 | 6.5 | 4.3 | 100.0 |
| <hr/> | | | | | | | | |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline: Current Adjusted Status (km) | 2007 - '09 | | 180.5 | 60.6 | 17.5 | 11.5 | | 270.0 |
| <i>Percentages</i> | | | 66.9 | 22.4 | 6.5 | 4.3 | | 100.0 |
| <hr/> | | | | | | | | |
| Baseline : Previous Unadjusted Status (km)** | 2004 - '06 | | 216.0 | 48.5 | 5.5 | 0.0 | | 270.0 |
| <i>Percentages</i> | | | 80.9 | 17.0 | 2.1 | 0.0 | | |
| Changes since Previous Survey (km) | | | -35.5 | 12.1 | 12.0 | 11.5 | | 0.0 |

* Adjustments : Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics

Hydrometric Area 10 : Trends
 % Surveyed Channel Length in Four Water Quality Classes



Hydrometric Area 10 : 2007 - '09
 % Surveyed Channel Length in Five WFD Water Quality Classes



SERBD: HYDROMETRIC AREA NO. 11

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|-------|------|----------|------|-------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Askinch Upper Stream | 2007 | 11A03 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 1.5 |
| Aughboy | 2007/9 | 11A02 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 2.0 |
| Ballyedmond | 2007 | 11B01 | 0.0 | 0.0 | 0.5 | 3.0 | 0.0 | 3.5 |
| Banoge | 2007 | 11B02 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 5.5 |
| Blackwater (Wexford) | 2007 | 11B03 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 5.5 |
| Brackan | 2007 | 11B04 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Clonough | 2007 | 11C01 | 0.0 | 2.5 | 0.0 | 4.0 | 0.0 | 6.5 |
| Gorteen Upper Stream | 2007 | 11G01 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Inch (Wexford) | 2007 | 11I01 | 0.0 | 6.5 | 3.0 | 0.0 | 0.0 | 9.5 |
| Owenavorrach | 2007 | 11O01 | 0.0 | 9.0 | 15.0 | 4.0 | 0.0 | 28.0 |
| Total Length (km) surveyed this cycle | 2007 - '09 | | 0.0 | 28.0 | 24.0 | 20.0 | 0.0 | 72.0 |
| Adjustments (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 0.0 | 38.9 | 33.3 | 27.8 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline: Current adjusted Status (km) | 2007 - '09 | | 28.0 | 24.0 | 20.0 | 0.0 | 72.0 | |
| Percentages | | | 38.9 | 33.3 | 27.8 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 40.5 | 21.0 | 10.5 | 0.0 | 72.0 | |
| Percentages | | | 56.3 | 29.2 | 14.6 | 0.0 | 100.0 | |
| Changes since Previous Survey (km) | | | -12.5 | 3.0 | 9.5 | 0.0 | 0.0 | |

* Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics

Hydrometric Area 11 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 11 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



SERBD: HYDROMETRIC AREA NO. 12

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|-------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Askinvillar Stream | 2007 | 12A03 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Assaly ¹ | 2007 | 12A02 | 0.0 | 1.0 | 7.0 | 0.0 | 0.0 | 8.0 |
| Ballingale Stream | 2007 | 12B06 | 0.0 | 9.0 | 0.0 | 0.5 | 0.0 | 9.5 |
| Ballycarney Stream | 2007 | 12B07 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Bann | 2007 | 12B01 | 31.3 | 8.4 | 0.4 | 0.0 | 0.0 | 40.0 |
| Blacklion Stream | 2007 | 12B04 | 0.0 | 3.5 | 3.5 | 0.0 | 0.0 | 7.0 |
| Boro | 2007 | 12B02 | 0.0 | 26.0 | 0.0 | 0.0 | 0.0 | 26.0 |
| Borris Stream | 2007 | 12B05 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Brown's Beck Brook | 2007 | 12B03 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Camolin Stream | 2007 | 12C08 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Carrigower | 2007 | 12C06 | 0.0 | 3.0 | 5.0 | 0.0 | 0.0 | 8.0 |
| Chapel Stream ¹ | 2007 | 12C01 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Clody | 2007 | 12C03 | 6.4 | 1.7 | 0.0 | 0.0 | 0.0 | 8.0 |
| Coolboy Stream | 2007 | 12C07 | 3.0 | 3.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Corbally Stream | 2007 | 12C04 | 0.0 | 8.5 | 7.0 | 0.0 | 0.0 | 15.5 |
| Derreen | 2007 | 12D01 | 13.9 | 18.7 | 7.5 | 0.0 | 0.0 | 40.0 |
| Derry | 2007 | 12D02 | 0.0 | 31.0 | 0.0 | 0.0 | 0.0 | 31.0 |
| Douglas (Ballon) ¹ | 2007 | 12D03 | 0.0 | 8.5 | 4.0 | 1.0 | 0.0 | 13.5 |
| Douglas (Kiltegan) | 2007 | 12D04 | 0.0 | 4.0 | 5.5 | 0.0 | 0.0 | 9.5 |
| Glasha (Slaney) | 2007 | 12G01 | 0.0 | 8.5 | 0.0 | 0.0 | 0.0 | 8.5 |
| Killeen Stream (Boro) | 2007 | 12K03 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Knicken | 2007 | 12K01 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Lask | 2007 | 12L01 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Little Slaney | 2007 | 12L02 | 8.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Mine | 2007 | 12M01 | 0.0 | 9.5 | 0.0 | 0.0 | 0.0 | 9.5 |
| Rosnastraw Stream | 2007 | 12R01 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |


| | | | | | | | | |
|---|------------|-------|----------|----------|----------|----------|--------------|-------|
| Shillelagh | 2007 | 12S01 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 |
| Slaney | 2007 | 12S02 | 15.6 | 70.0 | 11.5 | 0.0 | 0.0 | 97.0 |
| Sow | 2007 | 12S03 | 0.0 | 10.0 | 6.0 | 0.0 | 0.0 | 16.0 |
| Tinnacross Stream | 2007 | 12T01 | 0.0 | 5.0 | 8.5 | 0.0 | 0.0 | 13.5 |
| Tinnokilla Stream | 2007 | 12T02 | 0.0 | 6.5 | 2.0 | 0.0 | 0.0 | 8.5 |
| Urrin | 2007 | 12U01 | 8.0 | 14.0 | 0.0 | 0.0 | 0.0 | 22.0 |
| Total Length (km) surveyed this cycle | 2007 | | 94.6 | 291.7 | 67.8 | 2.5 | 0.0 | 456.5 |
| Adjustments (km) (See below)* | | | -2.0 | 1.0 | 7.0 | 2.0 | 0.0 | 8.0 |
| <i>Percentages</i> | | | 21.5 | 64.8 | 13.6 | 0.1 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 387.2 | 60.8 | 0.5 | 0.0 | 448.5 | |
| <i>Percentages</i> | | | 86.3 | 14 | 0.1 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 359.5 | 77.0 | 11.5 | 0.5 | 448.5 | |
| <i>Percentages</i> | | | 80.2 | 17.2 | 2.6 | 0.1 | | |
| Changes since Previous Survey (km) | | | 27.7 | -16.2 | -11.0 | -0.5 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 12 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 12 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



SERBD: HYDROMETRIC AREA NO. 13

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|------|------|----------|------|-------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Battlestown | 2007 | 13B04 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Begerin Stream | 2007 | 13B05 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Bridgetown (Wexford) | 2007 | 13B01 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Cleristown Stream | 2007 | 13C04 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Corock | 2007 | 13C01 | 0.0 | 14.0 | 4.0 | 0.0 | 0.0 | 18.0 |
| Duncormick | 2007 | 13D01 | 0.0 | 7.5 | 3.0 | 1.5 | 0.0 | 12.0 |
| Heathpark Stream | 2007 | 13H01 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Mulmontry | 2007 | 13M01 | 0.0 | 0.0 | 9.0 | 6.5 | 0.0 | 15.5 |
| Owenduff (Wexford) | 2007 | 13O01 | 8.5 | 2.5 | 0.0 | 0.0 | 0.0 | 11.0 |
| Tintern Abbey Stream | 2007 | 13T01 | 0.0 | 0.0 | 0.0 | 7.0 | 0.0 | 7.0 |
| Total Length (km) surveyed this cycle | 2007 | | 8.5 | 41.5 | 21.0 | 15.0 | 0.0 | 86.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 9.9 | 48.3 | 24.4 | 17.4 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 50.0 | 21.0 | 15.0 | 0.0 | 86.0 | |
| <i>Percentages</i> | | | 58.1 | 24.4 | 17.4 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 57.5 | 19.0 | 9.5 | 0.0 | 86.0 | |
| <i>Percentages</i> | | | 66.9 | 22.1 | 11.0 | 0.0 | 100.0 | |
| Changes since Previous Survey (km) | | | -7.5 | 2.0 | 5.5 | 0.0 | 0.0 | |

* Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 13 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 13 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



SERBD: HYDROMETRIC AREA NO. 14

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aghalona | 2009 | 14A02 | 0.0 | 0.0 | 10.0 | 0.0 | 0.0 | 10.0 |
| Athy Stream | 2009 | 14A06 | 0.0 | 0.0 | 10.5 | 0.0 | 0.0 | 10.5 |
| Aughavaud | 2009 | 14A04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Aughnabriskey | 2009 | 14A05 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Aughnacrew | 2009 | 14A07 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Barrow | 2009 | 14B01 | 6.0 | 90.5 | 41.2 | 2.4 | 0.0 | 140.0 |
| Black Borris | 2009 | 14B06 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Blackwater (Laois) | 2009 | 14B03 | 0.0 | 2.7 | 5.9 | 0.0 | 0.0 | 8.5 |
| Bothoge | 2009 | 14B04 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Burren | 2009 | 14B05 | 0.0 | 4.7 | 26.8 | 2.5 | 0.0 | 34.0 |
| Cloncumber Stream | 2009 | 14C17 | 0.0 | 0.0 | 8.5 | 0.0 | 0.0 | 8.5 |
| Crooked (Stradbally) | 2009 | 14C02 | 0.0 | 10.0 | 0.0 | 0.5 | 0.0 | 10.5 |
| Cushina | 2009 | 14C04 | 0.0 | 0.0 | 11.9 | 6.2 | 0.0 | 18.0 |
| Daingean | 2009 | 14D06 | 0.0 | 0.0 | 9.4 | 13.2 | 0.0 | 22.5 |
| Douglas (Laois) | 2009 | 14D03 | 9.0 | 3.0 | 4.0 | 0.0 | 0.0 | 16.0 |
| Duiske | 2009 | 14D04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Dunrally Stream | 2009 | 14D05 | 0.0 | 0.7 | 10.3 | 0.0 | 0.0 | 11.0 |
| Esker Stream | 2009 | 14E01 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| Figile | 2009 | 14F01 | 0.0 | 3.0 | 31.5 | 5.5 | 0.0 | 40.0 |
| Fushoge | 2009 | 14F03 | 8.5 | 2.5 | 0.0 | 0.0 | 0.0 | 11.0 |
| Glenlahan | 2009 | 14G02 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Gowran | 2009 | 14G03 | 5.5 | 1.5 | 11.0 | 0.5 | 0.0 | 18.5 |
| Graney (Lerr) | 2009 | 14G07 | 0.0 | 0.0 | 0.0 | 8.0 | 0.0 | 8.0 |
| Grangecon Stream | 2009 | 14G06 | 0.0 | 4.3 | 1.2 | 0.0 | 0.0 | 5.5 |
| Greese | 2009 | 14G04 | 0.0 | 0.0 | 13.5 | 17.5 | 0.0 | 31.0 |
| Guillie ¹ | 2009 | 14G11 | 0.0 | 8.5 | 0.0 | 0.0 | 0.0 | 8.5 |

| | | | | | | | | |
|---|------------|-------|------|----------|----------|----------|----------|--------------|
| Lerr | 2009 | 14L01 | 0.0 | 0.0 | 11.9 | 2.1 | 0.0 | 14.0 |
| Monefelim | 2009 | 14M03 | 0.0 | 11.9 | 0.1 | 0.0 | 0.0 | 12.0 |
| Mountain (Carlow) | 2009 | 14M01 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Old Leighlin Stream | 2009 | 14O02 | 0.0 | 0.0 | 3.5 | 0.0 | 0.0 | 3.5 |
| Owenass | 2009 | 14O01 | 0.0 | 8.0 | 8.0 | 0.0 | 0.0 | 16.0 |
| Palatine Stream | 2009 | 14P04 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Pollmounty | 2009 | 14P03 | 0.0 | 5.0 | 1.0 | 0.0 | 0.0 | 6.0 |
| Powerstown | 2009 | 14P02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Ratheven Stream ¹ no distances yet | 2009 | 14R05 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Red Hill Stream ¹ | 2009 | 14R06 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Slate | 2009 | 14S01 | 0.0 | 0.0 | 5.6 | 19.5 | 0.0 | 25.0 |
| Stradbally | 2009 | 14S02 | 0.0 | 5.5 | 12.5 | 0.0 | 0.0 | 18.0 |
| Triogue | 2009 | 14T01 | 0.0 | 0.0 | 3.3 | 17.7 | 0.0 | 21.0 |
| Tully Stream | 2009 | 14T02 | 0.0 | 0.0 | 5.3 | 14.3 | 0.0 | 19.5 |
| Total Length (km) surveyed this cycle | 2009 | | 34.0 | 218.8 | 252.6 | 112.7 | 0.0 | 618.0 |
| Adjustments (km) (See below)* | | | 0.0 | 7.5 | 0.0 | -3.0 | 0.0 | 4.5 |
| <i>Percentages</i> | | | 5.5 | 34.4 | 41.2 | 18.9 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 245.3 | 252.6 | 115.7 | 0.0 | 613.5 |
| <i>Percentages</i> | | | | 40.0 | 41.2 | 18.9 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 319.5 | 178.5 | 109.0 | 6.5 | 613.5 |
| <i>Percentages</i> | | | | 52.1 | 29.1 | 17.8 | 1.1 | 100.0 |
| Changes since Previous Survey (km) | | | | -74.3 | 74.1 | 6.7 | -6.5 | 0.0 |


Ballynaboley (14B08) & Enaghan streams (14E02) not surveyed in the 2007 - '09 period.

* Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 14 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 14 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



SERBD: HYDROMETRIC AREA NO. 15

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Arigna | 2007 | 15A01 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Arrigle | 2007 | 15A02 | 0.0 | 9.0 | 3.5 | 0.0 | 0.0 | 12.5 |
| Ballyroan | 2007 | 15B01 | 0.0 | 6.0 | 0.0 | 1.5 | 0.0 | 7.5 |
| Baunballinlough Stream | 2007 | 15B12 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Bregagh | 2007 | 15B02 | 0.0 | 0.0 | 0.0 | 10.0 | 0.0 | 10.0 |
| Brownstown Stream | 2007 | 15B04 | 0.0 | 1.5 | 3.0 | 0.0 | 0.0 | 4.5 |
| Caherlesk Stream | 2007 | 15C12 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Cappanacloghy | 2007 | 15C06 | 0.0 | 4.5 | 5.0 | 3.5 | 0.0 | 13.0 |
| Castlecomer Stream | 2007 | 15C01 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Clogh | 2007 | 15C03 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Cloghnagh | 2007 | 15C04 | 0.0 | 2.5 | 4.5 | 0.0 | 0.0 | 7.0 |
| Cloonawoolan Stream | 2007 | 15C19 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Coalbrook Stream | 2007 | 15C20 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Delour | 2007 | 15D01 | 11.0 | 5.0 | 0.0 | 0.0 | 0.0 | 16.0 |
| Desart Stream | 2007 | 15D04 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Dinin | 2007 | 15D02 | 0.0 | 14.5 | 0.0 | 0.0 | 0.0 | 14.5 |
| Dinin North | 2007 | 15D07 | 0.0 | 18.5 | 0.0 | 0.0 | 0.0 | 18.5 |
| Dinin South | 2007 | 15D08 | 0.0 | 15.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Donaghmore Stream | 2007 | 15D03 | 0.0 | 7.5 | 1.5 | 0.0 | 0.0 | 9.0 |
| Ennisnag Stream | 2007 | 15E02 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 1.0 |
| Erkina | 2007 | 15E01 | 0.0 | 0.0 | 28.0 | 0.0 | 0.0 | 28.0 |
| Errill | 2007 | 15E03 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 1.0 |
| Garranacool | 2007 | 15G10 | 0.0 | 0.0 | 3.5 | 0.0 | 0.0 | 3.5 |
| Glory | 2007 | 15G01 | 0.0 | 3.0 | 4.0 | 5.0 | 0.0 | 12.0 |
| Gorteenahilla | 2007 | 15G08 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Goul | 2007 | 15G02 | 0.0 | 13.0 | 12.0 | 0.0 | 0.0 | 25.0 |
| Gully | 2007 | 15G03 | 0.0 | 12.0 | 3.5 | 1.5 | 0.0 | 17.0 |


| | | | | | | | | |
|---|------------|-------|------|----------|----------|----------|----------|--------------|
| Killeen | 2007 | 15K01 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Kings | 2007 | 15K02 | 0.0 | 27.5 | 11.0 | 0.0 | 0.0 | 38.5 |
| Lisdowney | 2007 | 15L02 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Mountrath | 2007 | 15M01 | 0.0 | 13.0 | 1.0 | 0.0 | 0.0 | 14.0 |
| Muckalee | 2007 | 15M02 | 6.5 | 1.0 | 0.0 | 0.0 | 0.0 | 7.5 |
| Munster | 2007 | 15M03 | 9.2 | 5.4 | 1.0 | 0.0 | 0.0 | 15.5 |
| Needleford Stream | 2007 | 15N04 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Nore | 2007 | 15N01 | 0.0 | 73.5 | 36.5 | 8.0 | 0.0 | 118.0 |
| Nuenna | 2007 | 15N02 | 0.0 | 6.0 | 0.0 | 3.5 | 0.0 | 9.5 |
| Owveg (Nore) | 2007 | 15O01 | 0.0 | 20.0 | 4.0 | 0.0 | 0.0 | 24.0 |
| Rathdowney Stream | 2007 | 15R03 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Tonet | 2007 | 15T01 | 8.0 | 2.0 | 2.0 | 0.0 | 0.0 | 12.0 |
| Tullaroan Stream | 2007 | 15T02 | 0.0 | 13.0 | 0.0 | 0.0 | 0.0 | 13.0 |
| Total Length (km) surveyed this cycle | 2007 | | 39.7 | 299.4 | 137.0 | 43.0 | 0.0 | 519.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 7.6 | 57.7 | 26.4 | 8.3 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 339.0 | 137.0 | 43.0 | 0.0 | 519.0 |
| <i>Percentages</i> | | | | 65.3 | 26.4 | 8.3 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 254.0 | 201.5 | 61.0 | 2.5 | 519.0 |
| <i>Percentages</i> | | | | 49.0 | 38.7 | 11.8 | 0.5 | |
| Changes since Previous Survey (km) | | | | 85.0 | -64.5 | -18.0 | -2.5 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 15 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 15 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



SERBD: HYDROMETRIC AREA NO. 16

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------------|--------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aherlow | 2008 | 16A01 | 0.0 | 43.0 | 0.0 | 0.0 | 0.0 | 43.0 |
| Anner | 2008 | 16A02 | 0.0 | 19.5 | 4.5 | 10.0 | 0.0 | 34.0 |
| Ara | 2008 | 16A03 | 0.0 | 8.0 | 5.0 | 11.0 | 0.0 | 24.0 |
| Arglo | 2008 | 16A04 | 0.0 | 0.0 | 12.0 | 0.0 | 0.0 | 12.0 |
| Aughnaglanny | 2008 | 16A05 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Ballintemple Stream | 2008 | 16B07 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Black (Twomileborris) | 2008 | 16B01 | 0.0 | 0.0 | 10.0 | 0.0 | 0.0 | 10.0 |
| Black Stream (Cashel) ¹ | 2008 | 16B05 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Blackwater (Kilmacow) | 2008 | 16B02 | 0.0 | 16.5 | 3.5 | 0.0 | 0.0 | 20.0 |
| Borrisoleigh Stream | 2008/9 | 16B06 | 0.0 | 6.0 | 3.0 | 0.0 | 0.0 | 9.0 |
| Burncourt | 2008 | 16B04 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Clashawley | 2008 | 16C01 | 0.0 | 17.5 | 3.5 | 0.0 | 0.0 | 21.0 |
| Clodiagh (Portlaw) | 2008 | 16C03 | 0.0 | 18.0 | 6.0 | 1.0 | 0.0 | 25.0 |
| Clodiagh (Tipperary) | 2008 | 16C02 | 13.0 | 15.0 | 3.5 | 0.0 | 0.0 | 31.5 |
| Clonmore Stream | 2008 | 16C11 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Clover | 2008 | 16C04 | 0.0 | 0.5 | 2.0 | 1.0 | 0.0 | 3.5 |
| Drish | 2008 | 16D02 | 0.0 | 0.0 | 0.0 | 26.0 | 0.0 | 26.0 |
| Duag | 2008 | 16D03 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Farneybridge | 2008 | 16F02 | 0.0 | 2.0 | 4.0 | 7.0 | 0.0 | 13.0 |
| Fidaghta | 2008 | 16F01 | 0.0 | 0.0 | 8.0 | 0.0 | 0.0 | 8.0 |
| Fishmoyne | 2008 | 16F03 | 0.0 | 0.0 | 0.0 | 12.5 | 0.0 | 12.5 |
| Glasha (Waterford) | 2008 | 16G01 | 7.5 | 0.0 | 0.0 | 0.0 | 0.0 | 7.5 |
| Glenboy | 2008 | 16G03 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Glengala | 2008 | 16G05 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Half-Way House Stream | 2008 | 16H02 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 |
| Killenaule Stream | 2008 | 16K05 | 0.0 | 4.5 | 5.0 | 1.5 | 0.0 | 11.0 |
| Lingaun | 2008 | 16L01 | 0.0 | 23.0 | 0.0 | 0.0 | 0.0 | 23.0 |


| | | | | | | | | |
|---|------------|-------|----------|----------|----------|----------|--------------|-------|
| Moyle | 2008 | 16M01 | 0.0 | 0.0 | 0.0 | 18.0 | 0.0 | 18.0 |
| Multeen East | 2008 | 16M08 | 0.0 | 25.5 | 0.0 | 0.0 | 0.0 | 25.5 |
| Multeen | 2008 | 16M02 | 24.5 | 1.0 | 0.0 | 0.0 | 0.0 | 25.5 |
| Nier | 2008 | 16N01 | 18.5 | 0.5 | 0.0 | 0.0 | 0.0 | 19.0 |
| Outeragh Stream | 2008 | 16O01 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 4.0 |
| Owenbeg (Tipperary) | 2008 | 16O02 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Pollanassa | 2008 | 16P02 | 0.0 | 9.5 | 4.5 | 0.0 | 0.0 | 14.0 |
| Rossadrehid Stream | 2008 | 16R06 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Rossestown | 2008 | 16R01 | 0.0 | 0.0 | 0.0 | 13.0 | 0.0 | 13.0 |
| Shanbally | 2008 | 16S01 | 8.5 | 0.0 | 0.0 | 0.0 | 0.0 | 8.5 |
| Smartscastle Stream | 2008 | 16S07 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Suir | 2008 | 16S02 | 0.0 | 107.5 | 11.0 | 15.5 | 0.0 | 134.0 |
| Tar | 2008 | 16T01 | 12.5 | 14.5 | 0.0 | 0.0 | 0.0 | 27.0 |
| Thonoge | 2008 | 16T02 | 0.0 | 15.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Total Length (km) surveyed this cycle | 2008 - '09 | | 100.5 | 391.5 | 94.5 | 118.5 | 0.0 | 705.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 13.5 | 0.0 | 13.5 |
| <i>Percentages</i> | | | 14.5 | 56.6 | 13.7 | 15.2 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 492.0 | 94.5 | 105.0 | 0.0 | 691.5 | |
| <i>Percentages</i> | | | 71.1 | 13.7 | 15.2 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 457.5 | 142.5 | 90.0 | 1.5 | 691.5 | |
| <i>Percentages</i> | | | 66.2 | 20.6 | 13.0 | 0.2 | 100.0 | |
| Changes since Previous Survey (km) | | | 34.5 | -48.0 | 15.0 | -1.5 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 16 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 16 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



SERBD: HYDROMETRIC AREA NO. 17



The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|------|------|----------|------|-------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Araglin (Colligan) | 2007 | 17A01 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Brickey | 2007 | 17B01 | 0.0 | 0.0 | 3.5 | 3.5 | 0.0 | 7.0 |
| Colligan | 2007 | 17C01 | 14.0 | 7.5 | 0.0 | 0.0 | 0.0 | 21.5 |
| Dalligan | 2007 | 17D01 | 3.3 | 6.8 | 0.0 | 0.0 | 0.0 | 10.0 |
| Dunhill | 2007 | 17D02 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 5.5 |
| Leperstown Stream | 2007 | 17L01 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 2.0 |
| Mahon | 2007 | 17M01 | 6.5 | 9.0 | 2.0 | 0.0 | 0.0 | 17.5 |
| Morrigen | 2007 | 17M02 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Tay | 2007 | 17T01 | 11.5 | 5.0 | 0.0 | 0.0 | 0.0 | 16.5 |
| Total Length (km) surveyed this cycle | 2007 | | 35.3 | 36.8 | 11.0 | 5.5 | 0.0 | 88.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 39.8 | 41.5 | 12.4 | 6.2 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 72.0 | 11.0 | 5.5 | 0.0 | 88.5 | |
| Percentages | | | 81.4 | 12.4 | 6.2 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 74.0 | 12.5 | 2.0 | 0.0 | 88.5 | |
| Percentages | | | 83.6 | 14.1 | 2.3 | 0.0 | 100.0 | |
| Changes since Previous Survey (km) | | | -2.0 | -1.5 | 3.5 | 0.0 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



SWRBD: HYDROMETRIC AREA NO. 18

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|--------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Allow | 2009 | 18A02 | 24.3 | 9.8 | 0.0 | 0.0 | 0.0 | 34.0 |
| Araglin (Blackwater) | 2009 | 18A03 | 13.0 | 9.0 | 0.0 | 0.0 | 0.0 | 22.0 |
| Awbeg (Buttevant) East | 2009 | 18A08 | 0.0 | 13.0 | 0.0 | 0.0 | 0.0 | 13.0 |
| Awbeg (Buttevant) West | 2009 | 18A09 | 0.0 | 8.3 | 3.2 | 0.0 | 0.0 | 11.5 |
| Awbeg (Buttevant) Main Channel | 2009 | 18A05 | 0.0 | 29.5 | 4.6 | 3.4 | 0.0 | 37.5 |
| Awnaskirtaun | 2009 | 18A07 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Ballard Stream | 2009 | 18B10 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Ballyclogh Stream | 2009 | 18B08 | 0.0 | 0.0 | 2.5 | 3.5 | 0.0 | 6.0 |
| Behanagh | 2009 | 18B01 | 5.0 | 0.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Blackwater (Munster) | 2009 | 18B02 | 65.2 | 73.0 | 2.8 | 0.0 | 0.0 | 141.0 |
| Bregoge | 2009 | 18B04 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Bride (Blackwater) | 2009 | 18B05 | 0.0 | 55.0 | 0.0 | 0.0 | 0.0 | 55.0 |
| Brogeen | 2009 | 18B06 | 0.0 | 8.2 | 2.1 | 4.2 | 1.5 | 16.0 |
| Clyda | 2009 | 18C02 | 21.5 | 1.0 | 0.0 | 0.0 | 0.0 | 22.5 |
| Coom | 2009 | 18C03 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Crinnaghtane | 2009 | 18C07 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Cullavaw Stream | 2009 | 18C04 | 0.0 | 0.1 | 4.4 | 0.0 | 0.0 | 4.5 |
| Curraheen (Cork) | 2009 | 18C06 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Dalua | 2009 | 18D01 | 8.5 | 14.5 | 0.0 | 0.0 | 0.0 | 23.0 |
| Douglas (Araglin) | 2009 | 18D03 | 5.0 | 0.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Douglas (Bride) | 2009 | 18D02 | 0.0 | 0.0 | 11.0 | 0.0 | 0.0 | 11.0 |
| Farahy | 2009 | 18F01 | 4.0 | 6.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Farnane | 2009 | 18F06 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |

| | | | | | | | | |
|-----------------------|------|-------|------|------|-----|-----|-----|------|
| Finisk | 2009 | 18F02 | 0.0 | 26.0 | 0.0 | 0.0 | 0.0 | 26.0 |
| Finnow (Blackwater) | 2009 | 18F03 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Flesk (Bride) | 2009 | 18F04 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Funshion | 2009 | 18F05 | 44.8 | 8.3 | 0.0 | 0.0 | 0.0 | 53.0 |
| Glashanabrack | 2009 | 18G02 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Glashawee (Allow) | 2009 | 18G03 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glen (Banteer) | 2009 | 18G04 | 17.0 | 0.0 | 0.0 | 0.0 | 0.0 | 17.0 |
| Glenaboy | 2009 | 18G05 | 0.0 | 7.0 | 0.5 | 0.0 | 0.0 | 7.5 |
| Glenakeefe | 2009 | 18G06 | 5.5 | 0.0 | 0.0 | 0.0 | 0.0 | 5.5 |
| Glencorra Stream | 2009 | 18G15 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 1.5 |
| Glendine (Blackwater) | 2009 | 18G07 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Glenlara | 2009 | 18G08 | 0.0 | 6.5 | 2.5 | 0.0 | 0.0 | 9.0 |
| Glennafallia | 2009 | 18G10 | 8.5 | 0.5 | 0.0 | 0.0 | 0.0 | 9.0 |
| Glenshelane | 2009 | 16G11 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Goish | 2009 | 18G12 | 0.0 | 6.5 | 5.5 | 0.0 | 0.0 | 12.0 |
| Gradoge | 2009 | 18G13 | 0.0 | 0.0 | 9.5 | 0.0 | 0.0 | 9.5 |
| Knoppoge | 2009 | 18K02 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Licky | 2009 | 18L01 | 0.0 | 13.1 | 2.0 | 0.0 | 0.0 | 15.0 |
| Lyre | 2009 | 18L02 | 0.0 | 8.5 | 0.0 | 0.5 | 0.0 | 9.0 |
| Monavugga | 2009 | 18M01 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Nad | 2009 | 18N01 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Ogeen | 2009 | 18O01 | 11.0 | 0.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Owbeg (Waterford) | 2009 | 18O02 | 0.0 | 3.0 | 1.6 | 3.5 | 0.0 | 8.0 |
| Owenageeragh | 2009 | 18O03 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Owenanare | 2009 | 18O04 | 0.5 | 9.5 | 0.0 | 0.0 | 0.0 | 10.0 |
| Owenbaun (Rathcool) | 2009 | 18O05 | 3.5 | 6.5 | 0.0 | 0.0 | 0.0 | 10.0 |
| Owenkeale | 2009 | 18O06 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Owennagloo | 2009 | 18O07 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owennashad | 2009 | 18O08 | 10.5 | 0.5 | 0.0 | 0.0 | 0.0 | 11.0 |


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|-------|-------|----------|------|-------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Owentaraglin | 2009 | 18O09 | 13.3 | 17.8 | 0.0 | 0.0 | 0.0 | 31.0 |
| Rathcool | 2009 | 18R01 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Ross (Killavullen) | 2009 | 18R02 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Sheep | 2009 | 18S03 | 8.0 | 4.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Tourig | 2009 | 18T03 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Watergrasshill | 2009 | 18W01 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 1.5 |
| Total Length (km) surveyed this cycle | 2009 | | 280.0 | 465.4 | 53.6 | 16.5 | 1.5 | 817.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 9.5 | 0.0 | 0.0 | 9.5 |
| Percentages | | | 34.7 | 57.6 | 5.5 | 2.0 | 0.2 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 745.4 | 44.1 | 16.5 | 1.5 | 807.5 | |
| Percentages | | | 92.3 | 5.5 | 2.0 | 0.2 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 696.0 | 85.0 | 26.5 | 0.0 | 807.5 | |
| Percentages | | | 86.2 | 10.5 | 3.3 | 0.0 | 100.0 | |
| Changes since Previous Survey (km) | | | 49.4 | -40.9 | -10.0 | 1.5 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 18 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 18 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



SWRBD: HYDROMETRIC AREA NO. 19

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aughnaboy (Cork) | 2008 | 19A02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Awboy | 2008 | 19A03 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Blarney | 2008 | 19B02 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Bride (Lee) | 2008 | 19B04 | 19.0 | 13.0 | 0.0 | 0.0 | 0.0 | 32.0 |
| Butlerstown | 2008 | 19B06 | 0.0 | 8.0 | 3.0 | 0.0 | 0.0 | 11.0 |
| Cosloura | 2008 | 19C01 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Cummer | 2008 | 19C02 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Dissour | 2008 | 19D03 | 0.0 | 0.0 | 12.0 | 0.0 | 0.0 | 12.0 |
| Douglas (Sullane) | 2008 | 19D04 | 3.3 | 3.3 | 0.0 | 0.0 | 0.0 | 6.5 |
| Dripsey | 2008 | 19D06 | 10.3 | 4.0 | 4.8 | 0.0 | 0.0 | 19.0 |
| Dungourney | 2008 | 19D07 | 0.0 | 15.5 | 0.0 | 0.5 | 0.0 | 16.0 |
| Finnow (Foherish) | 2008 | 19F01 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Foherish | 2008 | 19F02 | 18.0 | 0.0 | 0.0 | 0.0 | 0.0 | 18.0 |
| Garrane (Lee) | 2008 | 19G03 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Glashaboy | 2008 | 19G01 | 0.0 | 23.0 | 0.0 | 0.0 | 0.0 | 23.0 |
| Keel | 2008 | 19K02 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Laney | 2008 | 19L01 | 14.0 | 10.0 | 0.0 | 0.0 | 0.0 | 24.0 |
| Lee (Cork) | 2008 | 19L03 | 8.5 | 26.0 | 2.5 | 1.0 | 0.0 | 38.0 |
| Martin | 2008 | 19M01 | 0.0 | 14.0 | 1.5 | 0.0 | 0.0 | 15.5 |
| Owenboy (Cork) | 2008 | 19O01 | 0.0 | 21.0 | 2.0 | 0.0 | 0.0 | 23.0 |
| Owennacurra | 2008 | 19O03 | 0.0 | 17.5 | 0.0 | 0.0 | 0.0 | 17.5 |
| Shournagh | 2008 | 19S01 | 14.8 | 11.3 | 0.0 | 0.0 | 0.0 | 26.0 |
| Sullane | 2008 | 19S02 | 18.0 | 13.0 | 0.0 | 0.0 | 0.0 | 31.0 |
| Templebodan | 2008 | 19T01 | 0.0 | 0.0 | 4.5 | 0.0 | 0.0 | 4.5 |


| | | | | | | | | |
|---|------------|-------|-------|----------|----------|----------|----------|--------------|
| Toon | 2008 | 19T02 | 15.0 | 0.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Womanagh | 2008 | 19W01 | 0.0 | 12.0 | 9.0 | 0.0 | 0.0 | 21.0 |
| Total Length (km) surveyed this cycle | 2008 | | 127.8 | 220.0 | 39.3 | 1.5 | 0.0 | 388.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 32.9 | 56.6 | 10.1 | 0.4 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 347.8 | 39.3 | 1.5 | 0.0 | 388.5 |
| <i>Percentages</i> | | | | 89.5 | 10.1 | 0.4 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 344.5 | 30.5 | 13.5 | 0.0 | 388.5 |
| <i>Percentages</i> | | | | 88.7 | 7.9 | 3.5 | 0.0 | 100.0 |
| Changes since Previous Survey (km) | | | | 3.3 | 8.8 | -12.0 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 19 : Trends % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 19 : 2007 - '09 % Surveyed Channel Length in Five WFD Quality Classes



SWRBD: HYDROMETRIC AREA NO. 19

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aughnaboy (Cork) | 2008 | 19A02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Awboy | 2008 | 19A03 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Blarney | 2008 | 19B02 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Bride (Lee) | 2008 | 19B04 | 19.0 | 13.0 | 0.0 | 0.0 | 0.0 | 32.0 |
| Butlerstown | 2008 | 19B06 | 0.0 | 8.0 | 3.0 | 0.0 | 0.0 | 11.0 |
| Cosloura | 2008 | 19C01 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Cummer | 2008 | 19C02 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Dissour | 2008 | 19D03 | 0.0 | 0.0 | 12.0 | 0.0 | 0.0 | 12.0 |
| Douglas (Sullane) | 2008 | 19D04 | 3.3 | 3.3 | 0.0 | 0.0 | 0.0 | 6.5 |
| Dripsey | 2008 | 19D06 | 10.3 | 4.0 | 4.8 | 0.0 | 0.0 | 19.0 |
| Dungourney | 2008 | 19D07 | 0.0 | 15.5 | 0.0 | 0.5 | 0.0 | 16.0 |
| Finnow (Foherish) | 2008 | 19F01 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Foherish | 2008 | 19F02 | 18.0 | 0.0 | 0.0 | 0.0 | 0.0 | 18.0 |
| Garrane (Lee) | 2008 | 19G03 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Glashaboy | 2008 | 19G01 | 0.0 | 23.0 | 0.0 | 0.0 | 0.0 | 23.0 |
| Keel | 2008 | 19K02 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Laney | 2008 | 19L01 | 14.0 | 10.0 | 0.0 | 0.0 | 0.0 | 24.0 |
| Lee (Cork) | 2008 | 19L03 | 8.5 | 26.0 | 2.5 | 1.0 | 0.0 | 38.0 |
| Martin | 2008 | 19M01 | 0.0 | 14.0 | 1.5 | 0.0 | 0.0 | 15.5 |
| Owenboy (Cork) | 2008 | 19O01 | 0.0 | 21.0 | 2.0 | 0.0 | 0.0 | 23.0 |
| Owennacurra | 2008 | 19O03 | 0.0 | 17.5 | 0.0 | 0.0 | 0.0 | 17.5 |
| Shournagh | 2008 | 19S01 | 14.8 | 11.3 | 0.0 | 0.0 | 0.0 | 26.0 |
| Sullane | 2008 | 19S02 | 18.0 | 13.0 | 0.0 | 0.0 | 0.0 | 31.0 |
| Templebodan | 2008 | 19T01 | 0.0 | 0.0 | 4.5 | 0.0 | 0.0 | 4.5 |


| | | | | | | | | |
|---|------------|-------|-------|----------|----------|----------|----------|--------------|
| Toon | 2008 | 19T02 | 15.0 | 0.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Womanagh | 2008 | 19W01 | 0.0 | 12.0 | 9.0 | 0.0 | 0.0 | 21.0 |
| Total Length (km) surveyed this cycle | 2008 | | 127.8 | 220.0 | 39.3 | 1.5 | 0.0 | 388.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 32.9 | 56.6 | 10.1 | 0.4 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline: Current Adjusted Status (km) | 2007 - '09 | | | 347.8 | 39.3 | 1.5 | 0.0 | 388.5 |
| <i>Percentages</i> | | | | 89.5 | 10.1 | 0.4 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 344.5 | 30.5 | 13.5 | 0.0 | 388.5 |
| <i>Percentages</i> | | | | 88.7 | 7.9 | 3.5 | 0.0 | 100.0 |
| Changes since Previous Survey (km) | | | | 3.3 | 8.8 | -12.0 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 19 : Trends % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 19 : 2007 - '09 % Surveyed Channel Length in Five WFD Quality Classes



SWRBD: HYDROMETRIC AREA NO. 20

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Argideen | 2009 | 20A02 | 0.0 | 23.0 | 0.0 | 0.0 | 0.0 | 23.0 |
| Ballinspittle | 2009 | 20B09 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Ballymahane | 2009 | 20B01 | 6.5 | 2.5 | 0.0 | 0.0 | 0.0 | 9.0 |
| Bandon | 2009 | 20B02 | 0.0 | 48.0 | 9.1 | 0.0 | 0.0 | 57.0 |
| Bawnaknockane | 2009 | 20B03 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Bealanascartane | 2009 | 20B08 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Blackwater (Bandon) | 2009 | 20B04 | 4.3 | 6.6 | 1.2 | 0.0 | 0.0 | 12.0 |
| Brinny | 2009 | 20B07 | 4.3 | 7.8 | 0.0 | 0.0 | 0.0 | 12.0 |
| Burrane | 2009 | 20B10 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Caha | 2009 | 20C01 | 3.5 | 10.0 | 0.0 | 0.0 | 0.0 | 13.5 |
| Carhoo | 2009 | 20C06 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 1.5 |
| Cashel (Cork) | 2009 | 20C02 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Clodagh | 2009 | 20C03 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Clonakilty Stream | 2009 | 20C05 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Cullenagh lake Stream | 2009 | 20C04 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Dirty | 2009 | 20D01 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Glashagloragh | 2009 | 20G02 | 2.8 | 2.3 | 0.0 | 0.0 | 0.0 | 5.0 |
| Ilen | 2009 | 20I01 | 10.5 | 14.0 | 0.0 | 0.0 | 0.0 | 24.5 |
| Kilbrittain | 2009 | 20K01 | 3.0 | 1.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Leamawaddra | 2009 | 20L01 | 8.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Leap Stream | 2009 | 20L04 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Minane | 2009 | 20M01 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Owenkeagh | 2009 | 20O01 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owennashingaun | 2009 | 20O02 | 13.5 | 0.0 | 0.0 | 0.0 | 0.0 | 13.5 |

| | | | | | | | | | | |
|---|------|-------|------------|------|-------|-------|------|------|-------|-------|
| Ownahinchy | 2009 | 20O03 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 | | |
| Rathruane | 2009 | 20R01 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 | | |
| Roury | 2009 | 20R02 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 | | |
| Saivnose | 2009 | 20S01 | 0.0 | 13.5 | 0.0 | 0.0 | 0.0 | 13.5 | | |
| Sall | 2009 | 20S02 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 | | |
| Stick | 2009 | 20S03 | 0.0 | 10.5 | 0.0 | 0.0 | 0.0 | 10.5 | | |
| Tinneel Stream | 2009 | 20T02 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 | | |
| Total Length (km) surveyed this cycle | | | 2009 | | 56.3 | 220.6 | 12.2 | 5.0 | 0.0 | 294.0 |
| Adjustments (km) (See below)* | | | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | | | 19.1 | 75.0 | 4.1 | 1.7 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | | A | B | C | D | Total | |
| Baseline: Current Adjusted Status (km) | | | 2007 - '09 | | 276.8 | 12.2 | 5.0 | 0.0 | 294.0 | |
| Percentages | | | | | 94.1 | 4.1 | 1.7 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | | | 2004 - '06 | | 281.0 | 9.0 | 3.0 | 1.0 | 294.0 | |
| Percentages | | | | | 95.9 | 3.1 | 0.7 | 0.3 | | |
| Changes since Previous Survey (km) | | | | | -4.2 | 3.2 | 2.0 | -1.0 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



SWRBD: HYDROMETRIC AREA NO. 21

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Adrigole | 2009 | 21A01 | 6.5 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 |
| Ardsheelhane | 2009 | 21A02 | 8.5 | 0.0 | 0.0 | 0.0 | 0.0 | 8.5 |
| Blackwater (Kerry) | 2009 | 21B03 | 13.0 | 0.0 | 0.0 | 0.0 | 0.0 | 13.0 |
| Cleady | 2009 | 21C02 | 6.5 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 |
| Cloghane (Allihies) | 2009 | 21C08 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Cloonee (Kerry) | 2009 | 21C06 | 0.0 | 3.0 | 5.0 | 0.0 | 0.0 | 8.0 |
| Coomelian Stream | 2009 | 21C14 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Coomhola | 2009 | 21C03 | 15.0 | 0.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Croanshagh | 2009 | 21C05 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Cummeragh | 2009 | 21C04 | 4.0 | 7.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Derreendaragh | 2009 | 21D03 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Drimminboy | 2009 | 21D01 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Drumoghty | 2009 | 21D04 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Emlaghmore | 2009 | 21E01 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Finnihy | 2009 | 21F01 | 0.0 | 10.5 | 0.0 | 0.0 | 0.0 | 10.5 |
| Four Mile Water | 2009 | 21F02 | 0.0 | 10.5 | 0.0 | 0.0 | 0.0 | 10.5 |
| Glan Stream | 2009 | 21G13 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Glengarriff | 2009 | 21G03 | 8.5 | 0.0 | 0.0 | 0.0 | 0.0 | 8.5 |
| Inny (Kerry) | 2009 | 21I01 | 0.0 | 24.0 | 0.0 | 0.0 | 0.0 | 24.0 |
| Isknagahiny Lough Stream | 2009 | 21I03 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Kealduff | 2009 | 21K01 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Kealinya | 2009 | 21K02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Lough Fadda Stream | 2009 | 21L03 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Magannagan Stream | 2009 | 21M02 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |

| | | | | | | | | |
|---|------------|-------|-------|----------|----------|----------|----------|--------------|
| Mealagh | 2009 | 21M01 | 16.5 | 0.5 | 0.0 | 0.0 | 0.0 | 17.0 |
| Owbeg (Roughy) | 2009 | 21O02 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Owenbeg (Owvane) | 2009 | 21O03 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Owengar (Cork) | 2009 | 21O04 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owenshagh | 2009 | 21O08 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Ownagappul | 2009 | 21O09 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Owreagh | 2009 | 21O05 | 3.0 | 4.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owroe | 2009 | 21O06 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Owvane (Cork) | 2009 | 21O07 | 10.5 | 6.5 | 0.0 | 0.0 | 0.0 | 17.0 |
| Roughy | 2009 | 21R01 | 27.0 | 0.0 | 0.0 | 0.0 | 0.0 | 27.0 |
| Sheen | 2009 | 21S01 | 20.0 | 0.0 | 0.0 | 0.0 | 0.0 | 20.0 |
| Slaheny | 2009 | 21S02 | 11.5 | 0.5 | 0.0 | 0.0 | 0.0 | 12.0 |
| Sneem | 2009 | 21S03 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Tahilla | 2009 | 21T01 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 |
| Trafrask Stream | 2009 | 21T03 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Total Length (km) surveyed this cycle | 2009 | | 223.5 | 109.5 | 5.0 | 0.0 | 0.0 | 338.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 66.1 | 32.4 | 1.5 | 0.0 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 333.0 | 5.0 | 0.0 | 0.0 | 338.0 |
| <i>Percentages</i> | | | | 98.5 | 1.5 | 0.0 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 327.0 | 11.0 | 0.0 | 0.0 | 338.0 |
| <i>Percentages</i> | | | | 96.7 | 3.3 | 0.0 | 0.0 | 100.0 |
| Changes since Previous Survey (km) | | | | 6.0 | -6.0 | 0.0 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



SWRBD: HYDROMETRIC AREA NO. 22

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Beheenagh | 2007 | 22B01 | 0.0 | 8.5 | 0.0 | 0.0 | 0.0 | 8.5 |
| Behy (Kerry) | 2007 | 22B02 | 5.0 | 2.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Brown Flesk | 2007 | 22B03 | 0.0 | 33.0 | 0.0 | 0.0 | 0.0 | 33.0 |
| Caherlehillan Stream | 2007 | 22C20 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 |
| Caragh | 2007 | 22C02 | 5.0 | 10.5 | 3.0 | 0.0 | 0.0 | 18.5 |
| Carhan | 2007 | 22C03 | 0.0 | 0.0 | 7.5 | 0.0 | 0.0 | 7.5 |
| Coomaglaslaw Lough Stream | 2007 | 22C19 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Coomnacarrig | 2007 | 22C06 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Coomnacronia Lough Stream | 2007 | 22C18 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Cottoner's (Laune) | 2007 | 22C05 | 13.0 | 0.0 | 0.5 | 0.0 | 0.0 | 13.5 |
| Crinnagh | 2007 | 22C07 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Croaghane | 2007 | 22C09 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Deenagh | 2007 | 22D01 | 0.0 | 10.0 | 3.0 | 0.0 | 0.0 | 13.0 |
| Derreen (Kerry) | 2007 | 22D02 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Dogue | 2007 | 22D03 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Emlagh | 2007 | 22E01 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Fahaduff | 2007 | 22F09 | 0.0 | 4.0 | 1.0 | 0.0 | 0.0 | 5.0 |
| Ferta | 2007 | 22F01 | 0.0 | 6.5 | 0.0 | 0.5 | 0.0 | 7.0 |
| Finglas (Laune) | 2007 | 22F03 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Finow | 2007 | 22F04 | 6.5 | 0.0 | 0.5 | 0.0 | 0.0 | 7.0 |
| Flesk (Kerry) | 2007 | 22F02 | 32.3 | 11.8 | 1.0 | 0.0 | 0.0 | 45.0 |
| Gaddagh | 2007 | 22G01 | 8.0 | 4.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Gearhameen | 2007 | 22G03 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Glantane | 2007 | 22G07 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |


| | | | | | | | | |
|---|------------|-------|-------|----------|----------|----------|----------|--------------|
| Groin | 2007 | 22G08 | 4.5 | 0.0 | 0.0 | 0.0 | 0.0 | 4.5 |
| Gweestin | 2007 | 22G06 | 0.0 | 19.0 | 0.0 | 1.0 | 0.0 | 20.0 |
| Laune | 2007 | 22L01 | 0.0 | 10.0 | 6.0 | 0.0 | 0.0 | 16.0 |
| Little Maine | 2007 | 22L02 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Loe | 2007 | 22L03 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Loo | 2007 | 22L04 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Maine | 2007 | 22M01 | 0.0 | 12.0 | 7.0 | 0.0 | 0.0 | 19.0 |
| Meelagh | 2007 | 22M02 | 6.5 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 |
| Milltown (Kerry) | 2007 | 22M03 | 0.0 | 1.5 | 1.0 | 3.0 | 0.0 | 5.5 |
| Owenalondrig | 2007 | 22O01 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owenascaul | 2007 | 22O02 | 0.0 | 5.0 | 2.0 | 0.0 | 0.0 | 7.0 |
| Owenreagh | 2007 | 22O03 | 13.5 | 0.0 | 0.0 | 0.0 | 0.0 | 13.5 |
| Owenroe (Caragh) | 2007 | 22O04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Owgarriff (Finow) | 2007 | 22O06 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Owneykeagh | 2007 | 22O05 | 4.0 | 4.5 | 0.0 | 0.0 | 0.0 | 8.5 |
| Quagmire | 2007 | 22Q01 | 10.3 | 2.8 | 0.0 | 0.0 | 0.0 | 13.0 |
| Shanowen (Maine) | 2007 | 22S01 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Teermoyle Stream | 2007 | 22T04 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Total Length (km) surveyed this cycle | 2007 | | 144.0 | 213.0 | 32.5 | 4.5 | 0.0 | 394.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 36.5 | 54.1 | 8.2 | 1.1 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 357.0 | 32.5 | 4.5 | 0.0 | 394.0 |
| <i>Percentages</i> | | | | 90.6 | 8.2 | 1.1 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 352.0 | 34.0 | 5.5 | 2.5 | 394.0 |
| <i>Percentages</i> | | | | 89.3 | 8.6 | 1.4 | 0.6 | |
| Changes since Previous Survey (km) | | | | 5.0 | -1.5 | -1.0 | -2.5 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 22 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 22 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



SHIRBD: HYDROMETRIC AREA NO. 23

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Allaghaun | 2007 | 23A01 | 0.0 | 23.0 | 0.0 | 0.0 | 0.0 | 23.0 |
| Breanagh | 2007 | 23B02 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Brick | 2007 | 23B03 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Caher (Limerick) | 2007 | 23C01 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Cloghane (Owenmore) | 2007 | 23C08 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Clydagh (Feale) | 2007 | 23C03 | 0.0 | 9.0 | 0.5 | 0.0 | 0.0 | 9.5 |
| Cruttia Lough Stream | 2007 | 23C11 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Feale | 2007 | 23F01 | 7.5 | 44.5 | 8.0 | 0.0 | 0.0 | 60.0 |
| Feohanagh | 2007 | 23F02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Finglas | 2007 | 23F03 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Galey | 2007 | 23G01 | 0.0 | 27.5 | 10.1 | 4.4 | 0.0 | 42.0 |
| Glashacooncore | 2007 | 23G02 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Glashoreag | 2007 | 23G03 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 5.5 |
| Glenacarney | 2007 | 23G06 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Glennahoo | 2007 | 23G05 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Knockfinnisk | 2007 | 23K02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Lee (Tralee) | 2007 | 23L01 | 0.0 | 7.5 | 2.5 | 1.0 | 0.0 | 11.0 |
| Lyracrumpane | 2007 | 23L02 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Milltown House Stream | 2007 | 23M04 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Oolagh | 2007 | 23O01 | 0.0 | 12.5 | 0.0 | 0.0 | 0.0 | 12.5 |
| Owencashla | 2007 | 23O02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owenmore (Kerry) | 2007 | 23O03 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Owennafeana | 2007 | 23O04 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Owveg (Kerry) | 2007 | 23O05 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |


| | | | | | | | | |
|---|------------|-------|-------|-------|------|------|-------|-------|
| Scorid | 2007 | 23S01 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Smearlagh | 2007 | 23S02 | 0.0 | 22.5 | 0.0 | 0.0 | 0.0 | 22.5 |
| Tarmon Stream | 2007 | 23T03 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 5.5 |
| Tullaleague | 2007 | 23T01 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| Tyshe | 2007 | 23T02 | 0.0 | 0.0 | 7.5 | 0.0 | 0.0 | 7.5 |
| Total Length (km) surveyed this cycle | 2007 | | 22.0 | 227.5 | 40.1 | 10.9 | 0.0 | 300.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 7.3 | 75.7 | 13.3 | 3.6 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 249.5 | 40.1 | 10.9 | 0.0 | 300.5 | |
| Percentages | | | 83.0 | 13.3 | 3.6 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 249.5 | 31.5 | 19.5 | 0.0 | 300.5 | |
| Percentages | | | 83.0 | 10.5 | 6.5 | 0.0 | | |
| Changes since Previous Survey (km) | | | 0.0 | 8.6 | -8.6 | 0.0 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 23 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 23 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



SHIRBD: HYDROMETRIC AREA NO. 24

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ahacronane | 2008 | 24A01 | 0.0 | 0.0 | 0.0 | 8.5 | 0.0 | 8.5 |
| Ahavarraga Stream | 2008 | 24A02 | 0.0 | 0.5 | 0.0 | 4.0 | 1.0 | 5.5 |
| Arra | 2008 | 24A04 | 0.0 | 4.5 | 0.0 | 1.5 | 0.0 | 6.0 |
| Ballylongford | 2008 | 24B03 | 0.0 | 6.5 | 2.0 | 0.0 | 0.0 | 8.5 |
| Barnakyle | 2008 | 24B05 | 0.0 | 0.0 | 0.0 | 14.0 | 0.0 | 14.0 |
| Broadford Stream | 2008 | 24B07 | 0.0 | 3.5 | 1.0 | 0.0 | 0.0 | 4.5 |
| Bunoke | 2008 | 24B06 | 0.0 | 7.5 | 3.5 | 2.0 | 0.0 | 13.0 |
| Camoge | 2008 | 23C01 | 0.0 | 14.5 | 13.5 | 0.0 | 0.0 | 28.0 |
| Charleville Stream | 2008 | 24C02 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Clonshire | 2008 | 24C03 | 0.0 | 0.0 | 0.0 | 13.0 | 0.0 | 13.0 |
| Cloonlahard Stream | 2008 | 24C06 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Deel (Newcastlewest) | 2008 | 2D02 | 0.0 | 21.5 | 23.0 | 7.0 | 0.0 | 51.5 |
| Dooncaha Stream | 2008 | 24D07 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 5.5 |
| Flemingstown Stream | 2008 | 24F03 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Glencorbry | 2008 | 24G03 | 0.0 | 0.0 | 4.5 | 0.0 | 0.0 | 4.5 |
| Greanagh | 2008 | 24G05 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 |
| Lismakeery Stream | 2008 | 24L03 | 0.0 | 0.0 | 0.0 | 7.0 | 0.0 | 7.0 |
| Loobagh | 2008 | 24L01 | 0.0 | 18.0 | 2.0 | 0.0 | 0.0 | 20.0 |
| Mahore | 2008 | 24M04 | 0.0 | 0.0 | 12.0 | 0.0 | 0.0 | 12.0 |
| Maigue | 2008 | 24M01 | 0.0 | 16.0 | 8.5 | 11.0 | 0.0 | 35.5 |
| Morningstar | 2008 | 24M02 | 0.0 | 27.5 | 5.5 | 0.0 | 0.0 | 33.0 |
| Owvane (Limerick) | 2008 | 24O02 | 0.0 | 13.0 | 0.0 | 0.0 | 0.0 | 13.0 |
| Shanagolden Stream | 2008 | 24S02 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Tarbert Stream | 2008 | 24T01 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 1.5 |


| | | | | | | | |
|---|------------|-----|----------|----------|----------|----------|--------------|
| Total Length (km) surveyed this cycle | 2008 | 0.0 | 154.0 | 75.5 | 83.5 | 1.0 | 314.0 |
| Adjustments (km) (See below)* | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | 0.0 | 49.0 | 24.0 | 26.6 | 0.3 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 154.0 | 75.5 | 83.5 | 1.0 | 314.0 |
| <i>Percentages</i> | | | 49.0 | 24.0 | 26.6 | 0.3 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 122.0 | 94.5 | 94.5 | 3.0 | 314.0 |
| <i>Percentages</i> | | | 38.9 | 30.1 | 30.1 | 1.0 | |
| Changes since Previous Survey (km) | | | 32.0 | -19.0 | -11.0 | -2.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 24 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 24 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



SHIRBD: HYDROMETRIC AREA NO. 26

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|-------------------------------|---------|-------|------|------|----------|------|------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aghnashannagh Stream | 2008 | 26A11 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Ahascragh | 2008 | 26A01 | 0.0 | 22.0 | 3.5 | 1.5 | 0.0 | 27.0 |
| Anaderryboy | 2008 | 26A03 | 5.3 | 11.8 | 0.0 | 0.0 | 0.0 | 17.0 |
| Arigna (Roscommon) | 2008 | 26A02 | 0.0 | 0.0 | 0.0 | 0.0 | 18.0 | 18.0 |
| Ballinure | 2009 | 26B01 | 0.0 | 20.0 | 0.0 | 0.0 | 0.0 | 20.0 |
| Ballymore Stream ¹ | 2008 | 26B33 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Ballydangan | 2009 | 26B14 | 0.0 | 4.2 | 0.8 | 0.0 | 0.0 | 5.0 |
| Bellavally Stream | 2008 | 26B32 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Black (South Leitrim) | 2008 | 26B04 | 3.3 | 13.8 | 0.0 | 0.0 | 0.0 | 17.0 |
| Black (Westmeath) | 2008 | 26B05 | 0.0 | 3.5 | 4.5 | 7.0 | 0.0 | 15.0 |
| Boleybaun Stream | 2008 | 26B30 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Boor | 2008 | 26B07 | 0.0 | 0.0 | 13.0 | 0.0 | 0.0 | 13.0 |
| Boyle | 2008/09 | 26B08 | 0.0 | 16.5 | 0.0 | 0.0 | 0.0 | 16.5 |
| Breedoge | 2008 | 26B09 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Breensford | 2008 | 26B10 | 0.0 | 0.0 | 3.0 | 4.0 | 0.0 | 7.0 |
| Camlin | 2008 | 26C01 | 0.0 | 10.8 | 20.2 | 4.1 | 0.0 | 35.0 |
| Carricknabraher | 2008 | 26C02 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Castlegar | 2008 | 26C03 | 0.0 | 0.0 | 9.0 | 4.0 | 0.0 | 13.0 |
| Clogher (Roscommon) | 2008 | 26C18 | 0.0 | 0.0 | 3.5 | 3.5 | 0.0 | 7.0 |
| Clooncoose Stream | 2008 | 26C20 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Cloone | 2008 | 26C05 | 8.5 | 5.0 | 3.0 | 0.0 | 0.0 | 16.5 |
| Clooneigh | 2008/09 | 26C06 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Cloonfower Stream | 2008 | 26C21 | 0.0 | 0.0 | 4.5 | 0.0 | 0.0 | 4.5 |
| Comoge | 2008 | 26C12 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |

| | | | | | | | | |
|--------------------------------|---------|-------|------|------|------|------|-----|------|
| Creelaghta | 2008 | 26C22 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Cross (Roscommon) | 2009 | 26C10 | 0.0 | 2.1 | 6.3 | 7.6 | 0.0 | 16.0 |
| Cuilleen Stream | 2008 | 26C17 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Curraghroe Stream | 2008 | 26C15 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 5.5 |
| Derryhippoo | 2009 | 26D01 | 0.0 | 0.0 | 0.0 | 10.5 | 0.0 | 10.5 |
| Derrymullan Stream | 2008 | 26D07 | 0.0 | 11.5 | 0.0 | 0.0 | 0.0 | 11.5 |
| Derrynananta Stream | 2008 | 26D23 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 4.0 |
| Diffagher | 2008 | 26D02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Drumshanbo Stream ¹ | 2008 | 26D05 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.5 |
| Dungolman | 2008 | 26D06 | 0.0 | 0.0 | 14.0 | 0.0 | 0.0 | 14.0 |
| Eslin | 2008 | 26E01 | 0.0 | 0.0 | 18.5 | 2.5 | 0.0 | 21.0 |
| Fallan | 2008 | 26F01 | 0.0 | 0.0 | 16.5 | 0.0 | 0.0 | 16.5 |
| Fardrumman Stream | 2008 | 26F06 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 |
| Feorish (Ballyfarnan) | 2008 | 26F02 | 20.0 | 6.5 | 0.0 | 0.0 | 0.0 | 26.5 |
| Feorish (Tarmonbarry) | 2008 | 26F03 | 0.0 | 0.0 | 0.0 | 10.0 | 0.0 | 10.0 |
| Francis | 2008 | 26F05 | 13.5 | 0.5 | 0.0 | 0.0 | 0.0 | 14.0 |
| Gaine | 2008 | 26G01 | 0.0 | 0.0 | 8.7 | 0.9 | 0.0 | 9.5 |
| Glore (Westmeath) | 2008 | 26G02 | 0.0 | 0.0 | 8.5 | 3.5 | 0.0 | 12.0 |
| Hind | 2008 | 26H01 | 0.0 | 0.0 | 3.5 | 7.0 | 0.0 | 10.5 |
| Inny | 2008/09 | 26I01 | 0.0 | 34.5 | 44.0 | 3.1 | 0.0 | 81.5 |
| Island | 2008 | 26I03 | 9.5 | 9.5 | 0.0 | 0.0 | 0.0 | 19.0 |
| Jiggy (Hind) | 2008/09 | 26J01 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 3.0 |
| Kiltaclare Stream | 2008 | 26K06 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Killadiskert Stream | 2008 | 26K16 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Killeglan | 2008 | 26K04 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Killian | 2008 | 26K01 | 0.0 | 15.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Killukin | 2008 | 26K02 | 0.0 | 2.7 | 1.9 | 4.5 | 0.0 | 9.0 |
| Kilmacricard | 2008 | 26K17 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|---------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Kilmactranny | 2008 | 26K03 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Kinard | 2008 | 26K07 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Laurencetown Stream | 2009 | 26L07 | 0.0 | 0.0 | 0.0 | 2.6 | 3.9 | 6.5 |
| Lenamore Stream | 2008 | 26L06 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Lisdaly Stream | 2008 | 26L02 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 9.0 |
| Lissaphobble | 2008 | 26L04 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Lissydaly Stream | 2008 | 26L10 | 8.3 | 6.8 | 0.0 | 0.0 | 0.0 | 15.0 |
| Lung | 2008 | 26L03 | 8.0 | 13.0 | 0.0 | 0.0 | 0.0 | 21.0 |
| Mantua Stream | 2008 | 26M13 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Mountain (Roscommon) | 2008 | 26M03 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 |
| Mountnugent | 2008 | 26M02 | 0.0 | 0.0 | 16.0 | 1.0 | 0.0 | 17.0 |
| Mullenmeehan Stream | 2008 | 26M12 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 7.0 |
| Owengar (Leitrim) | 2008/09 | 26O02 | 0.0 | 1.5 | 1.0 | 2.5 | 3.0 | 8.0 |
| Owenmore (Glengevin) | 2008 | 26O03 | 0.0 | 0.0 | 11.0 | 0.0 | 0.0 | 11.0 |
| Owennaforeesha | 2008 | 26O04 | 0.0 | 2.5 | 4.5 | 0.0 | 0.0 | 7.0 |
| Owennayle | 2008 | 26O05 | 0.0 | 3.5 | 5.5 | 0.0 | 0.0 | 9.0 |
| Owenur | 2008 | 26O06 | 0.0 | 16.5 | 0.0 | 0.0 | 0.0 | 16.5 |
| Rath | 2008 | 26R01 | 0.0 | 0.0 | 14.0 | 0.0 | 0.0 | 14.0 |
| Relagh | 2008 | 26R05 | 8.5 | 0.5 | 0.0 | 0.0 | 0.0 | 9.0 |
| Rhine | 2008 | 26R04 | 0.0 | 5.0 | 2.0 | 1.5 | 0.0 | 8.5 |
| Rinn | 2008 | 26R02 | 0.0 | 0.0 | 10.0 | 3.5 | 0.0 | 13.5 |
| Rocksavage | 2008 | 26R07 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 1.5 |
| Scramoge | 2008 | 26S01 | 0.0 | 19.0 | 0.0 | 1.5 | 0.0 | 20.5 |
| Shad Lough Stream | 2009 | 26S13 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Shannon (Upper) | 2008 | 26S02 | 25.8 | 28.5 | 35.8 | 13.5 | 0.0 | 103.5 |
| Shiven (South) | 2008 | 26S03 | 6.8 | 12.8 | 6.5 | 2.5 | 0.0 | 28.5 |


| | | | | | | | | |
|---|------------|-------|-------|----------|----------|----------|----------|--------------|
| Smaghraan | 2008 | 26S04 | 0.0 | 0.0 | 0.0 | 12.0 | 0.0 | 12.0 |
| Springfield | 2008 | 26S05 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Strokestown | 2008 | 26S08 | 0.0 | 2.0 | 1.5 | 2.5 | 0.0 | 6.0 |
| Suck | 2008 | 26S07 | 37.3 | 62.8 | 9.5 | 6.5 | 0.0 | 116.0 |
| Tang | 2008 | 26T03 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Termon Stream | 2008 | 26T03 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 1.5 |
| Yellow (Ballinaglera) | 2008 | 26Y01 | 0.0 | 4.5 | 7.5 | 0.0 | 0.0 | 12.0 |
| Yellow (Castlepollard) | 2008 | 26Y02 | 0.0 | 0.0 | 11.0 | 0.0 | 0.0 | 11.0 |
| Total Length (km) surveyed this cycle | 2008 - '09 | | 158.5 | 471.0 | 351.5 | 131.6 | 27.9 | 1140.5 |
| Adjustments (See below)* | | | 0.0 | -11.0 | 1.0 | 2.0 | 0.0 | -8.0 |
| Percentages | | | 13.8 | 42.0 | 30.5 | 11.3 | 2.4 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline: Current Adjusted Status (km) | 2007 - '09 | | | 640.5 | 350.5 | 129.6 | 27.9 | 1148.5 |
| <i>Percentages</i> | | | | 55.8 | 30.5 | 11.3 | 2.4 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 795.5 | 229.5 | 114.0 | 9.5 | 1148.5 |
| <i>Percentages</i> | | | | 69.3 | 20.0 | 9.9 | 0.8 | 100.0 |
| Baseline : Current Adjusted Status (km) | | | | -155.0 | 121.0 | 15.6 | 18.4 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 26 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 26 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes



SHIRBD: HYDROMETRIC AREA NO. 27

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|--|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballymacooda Lough Stream ¹ | 2007 | 27B01 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 1.5 |
| Ballynacally | 2007 | 27B04 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 1.0 |
| Broadford | 2007 | 27B02 | 9.5 | 0.0 | 1.5 | 0.0 | 0.0 | 11.0 |
| Carrownanelly | 2007 | 27C07 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 5.5 |
| Castlelodge | 2007 | 27C01 | 0.0 | 0.0 | 0.0 | 7.0 | 0.0 | 7.0 |
| Clareen (Fergus) | 2004 | 27C06 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 5.5 |
| Cloon (Clare) | 2005 | 27C02 | 0.0 | 13.5 | 0.0 | 0.0 | 0.0 | 13.5 |
| Cloondanagh Lough Stream | 2007 | 27C13 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Clooneen (Clare) | 2007 | 27C03 | 0.0 | 2.0 | 1.0 | 2.0 | 0.0 | 5.0 |
| Cloverhill Stream | 2007 | 27C10 | 0.0 | 0.0 | 0.0 | 7.0 | 0.0 | 7.0 |
| Craggaunboy | 2007 | 27C04 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 | 6.5 |
| Cratloe | 2007 | 27C08 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Crompaum (East) | 2007 | 27C09 | 4.0 | 1.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Crompaum (West) | 2007 | 27C05 | 0.0 | 2.5 | 5.0 | 0.0 | 0.0 | 7.5 |
| Cullaun | 2007 | 27C14 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 1.0 |
| Doonaha | 2007 | 27D01 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Druminshin | 2007 | 27D02 | 0.0 | 0.0 | 2.5 | 0.0 | 0.0 | 2.5 |
| Fergus | 2007 | 27F01 | 0.0 | 29.0 | 0.0 | 6.0 | 0.0 | 35.0 |
| Gourna | 2007 | 27G02 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Hell | 2007 | 27H01 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Inch (Clare) | 2007 | 27I01 | 6.5 | 13.0 | 0.0 | 0.5 | 0.0 | 20.0 |
| Kilcarrol Stream | 2007 | 27K06 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 2.0 |
| Kilmore North Stream | 2007 | 27K04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Laghtyshaughnessy Lough Stream | 2007 | 27L03 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 1.5 |


| | | | | | | | | | | |
|---|------------|-------|------|------|-------|-------|------|------|-------|-------|
| Liskenny | 2007 | 27L01 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 | | |
| Lissycasey Stream | 2007 | 27L04 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 2.0 | | |
| Mill Brook | 2007 | 27M03 | 7.4 | 2.1 | 0.0 | 0.0 | 0.0 | 9.5 | | |
| Moyarta | 2007 | 27M01 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 | | |
| Moyree | 2007 | 27M02 | 5.4 | 10.7 | 0.0 | 0.0 | 0.0 | 16.0 | | |
| Owenogarney | 2007 | 27O01 | 19.5 | 5.5 | 0.0 | 0.0 | 0.0 | 25.0 | | |
| Owenslieve | 2007 | 27O02 | 9.5 | 2.0 | 0.0 | 0.0 | 0.0 | 11.5 | | |
| Rine | 2007 | 27R01 | 9.0 | 19.0 | 2.5 | 0.0 | 0.0 | 30.5 | | |
| Shallee | 2007 | 27S01 | 0.0 | 0.0 | 0.0 | 11.0 | 0.0 | 11.0 | | |
| Spancelhill | 2007 | 27S03 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 | | |
| Tarmon Lough Stream | 2007 | 27T02 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 | | |
| Wood | 2007 | 27W01 | 0.0 | 0.0 | 0.0 | 3.5 | 0.0 | 3.5 | | |
| Total Length (km) surveyed this cycle | | | 2007 | | 70.8 | 128.3 | 24.5 | 70.5 | 0.0 | 294.0 |
| Adjustments (km) (See below)* | | | | | 0.0 | -4.0 | 1.5 | 0.0 | 0.0 | -2.5 |
| Percentages | | | | | 23.9 | 44.6 | 7.8 | 23.8 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | | 203.0 | 23.0 | 70.5 | 0.0 | 296.5 | |
| Percentages | | | | | 68.5 | 7.8 | 23.8 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | | 202.5 | 48.5 | 44.0 | 1.5 | 296.5 | |
| Percentages | | | | | 68.3 | 16.4 | 14.8 | 0.5 | | |
| Changes since Previous Survey (km) | | | | | 0.5 | -25.5 | 26.5 | -1.5 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 27 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 27 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



SHIRBD: HYDROMETRIC AREA NO. 28

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|------|-------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aille (Clare) | 2009 | 28A01 | 0.0 | 5.8 | 5.2 | 0.0 | 0.0 | 11.0 |
| Annageeragh | 2009 | 28A02 | 4.4 | 8.7 | 2.5 | 0.0 | 0.0 | 15.5 |
| Annagh (Clare) | 2009 | 28A03 | 0.0 | 7.1 | 2.5 | 1.4 | 0.0 | 11.0 |
| Aughaglanna | 2009 | 28A06 | 2.9 | 0.0 | 1.7 | 0.9 | 0.0 | 5.5 |
| Aughaveema | 2009 | 28A04 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Ballinphonta | 2009 | 28B03 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Ballymacravan | 2009 | 28B02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Caher (Clare) | 2009 | 28C01 | 4.5 | 0.0 | 0.0 | 0.0 | 0.0 | 4.5 |
| Castlepark Stream | 2009 | 28C09 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Cloonenagh Stream | 2009 | 28C08 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Cooleen | 2009 | 28C05 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Creegh | 2009 | 28C02 | 0.0 | 14.0 | 0.0 | 0.0 | 0.0 | 14.0 |
| Dealagh | 2009 | 28D01 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Doonbeg | 2009 | 28D02 | 0.0 | 35.0 | 0.0 | 0.0 | 0.0 | 35.0 |
| Glendine (Clare) | 2009 | 28G02 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Inagh (Ennistymon) | 2009 | 28I01 | 0.0 | 24.0 | 0.0 | 0.0 | 0.0 | 24.0 |
| Kildeema | 2009 | 28K01 | 0.0 | 5.0 | 2.5 | 0.0 | 0.0 | 7.5 |
| Kilmihil Stream ¹ | 2007 | 28K02 | 0.0 | 0.0 | 0.0 | 8.5 | 0.0 | 8.5 |
| Sruhaunakit Stream | 2009 | 28S04 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Total Length (km) surveyed this cycle | 2007/09 | | 14.8 | 144.1 | 14.3 | 10.8 | 0.0 | 184.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | -3.5 | 5.5 | 0.0 | 2.0 |
| Percentages | | | 8.1 | 79.2 | 9.8 | 2.9 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 158.9 | 17.8 | 5.3 | 0.0 | 182.0 |

| | | | | | | |
|---|------------|-------|------|------|-----|-------|
| <i>Percentages</i> | | 87.3 | 9.8 | 2.9 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | 156.0 | 11.5 | 14.5 | 0.0 | 182.0 |
| <i>Percentages</i> | | 85.7 | 6.3 | 8.0 | 0.0 | |
| Changes since Previous Survey (km) | | 2.9 | 6.3 | -9.2 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



WRBD: HYDROMETRIC AREA NO. 29



The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|-------|------|-------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballymabilla | 2009 | 29B03 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Beagh | 2009 | 29B02 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Boleyneendorish | 2009 | 29B04 | 16.5 | 1.0 | 0.0 | 0.0 | 0.0 | 17.5 |
| Cannahowna | 2009 | 29C01 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Carra Stream | 2009 | 29C03 | 0.0 | 10.9 | 8.2 | 0.0 | 0.0 | 19.0 |
| Clarinbridge | 2009 | 29C02 | 0.0 | 13.5 | 0.0 | 11.0 | 1.5 | 26.0 |
| Kilcolgan | 2009 | 29K01 | 0.0 | 9.8 | 11.7 | 0.0 | 0.0 | 21.5 |
| Knocknarebana | 2009 | 29K04 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Lecarrow Stream | 2009 | 29L01 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Owendalulleegh | 2009 | 29O01 | 22.5 | 0.0 | 0.0 | 0.0 | 0.0 | 22.5 |
| Raford | 2009 | 29R01 | 0.0 | 35.0 | 0.0 | 0.0 | 0.0 | 35.0 |
| Toberdoney | 2009 | 29T01 | 0.0 | 2.1 | 4.0 | 5.9 | 0.0 | 12.0 |
| Total Length (km) surveyed this cycle | 2009 | | 46.0 | 86.8 | 23.9 | 16.9 | 1.5 | 175.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 26.3 | 49.6 | 13.6 | 9.7 | 0.9 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 132.8 | 23.9 | 16.9 | 1.5 | 175.0 |
| Percentages | | | | 75.9 | 13.6 | 9.7 | 0.9 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 134.0 | 23.0 | 16.5 | 1.5 | 175.0 |
| Percentages | | | | 76.6 | 13.1 | 9.4 | 0.9 | 100.0 |
| Changes since Previous Survey (km) | | | | -1.3 | 0.9 | 0.4 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



WRBD: HYDROMETRIC AREA NO. 30

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Abbert | 2009 | 30A01 | 0.0 | 17.5 | 17.5 | 0.0 | 0.0 | 35.0 |
| Aghinish | 2009 | 30A03 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 1.5 |
| Aille (Mayo) | 2009 | 30A02 | 23.7 | 2.3 | 5.6 | 0.0 | 0.0 | 31.5 |
| Ballindine | 2009 | 30B03 | 0.0 | 0.0 | 1.5 | 2.5 | 0.0 | 4.0 |
| Bealanabrack | 2009 | 30B01 | 0.0 | 17.0 | 0.0 | 0.0 | 0.0 | 17.0 |
| Black (Shrule) | 2009 | 30B02 | 9.0 | 12.0 | 0.0 | 0.0 | 0.0 | 21.0 |
| Bunowen (Oughterard) | 2009 | 30B08 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Ballycuike | 2009 | 30B14 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Cammanagh | 2009 | 30C04 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Clare (Galway) | 2009 | 30C01 | 5.5 | 28.9 | 24.6 | 0.0 | 0.0 | 59.0 |
| Claureen (Mayo) | 2009 | 30C12 | 0.0 | 2.0 | 2.0 | 4.5 | 0.0 | 8.5 |
| Cloondaver Stream North | 2009 | 30C09 | 0.0 | 3.0 | 3.0 | 0.0 | 0.0 | 6.0 |
| Cloonfad | 2009 | 30C11 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 7.0 |
| Cong Canal ¹ | 2009 | 30C06 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Corrib | 2009 | 30C02 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Cregg | 2009 | 30C03 | 0.0 | 0.0 | 11.0 | 0.0 | 0.0 | 11.0 |
| Dalgan | 2009 | 30D01 | 0.0 | 14.0 | 4.3 | 5.8 | 0.0 | 24.0 |
| Dooghta | 2009 | 30D02 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Drimneen | 2009 | 30D03 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Failmore | 2009 | 30F01 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Finny | 2009 | 30F03 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Foey | 2009 | 30F02 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Glennamucka Stream ¹ | 2009 | 30G04 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Glensaul | 2009 | 30G01 | 3.6 | 5.4 | 0.0 | 0.0 | 0.0 | 9.0 |

| | | | | | | | | |
|------------------------------|------|-------|-----|------|------|-----|-----|------|
| Gortgarrow Stream | 2009 | 30G05 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| Grange (Galway) | 2009 | 30G02 | 0.0 | 20.0 | 4.6 | 1.5 | 0.0 | 26.0 |
| Glengawbeg ¹ | 2009 | 30G06 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Headford Stream ¹ | 2009 | 30H01 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Joyces | 2009 | 30J01 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Knockaunranny Stream | 2009 | 30K02 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Levally Stream | 2009 | 30L07 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Loughkip | 2009 | 30L01 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Lough Nacorralea Stream | 2009 | 30L03 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Nanny (Tuam) | 2009 | 30N01 | 6.5 | 0.0 | 0.0 | 2.0 | 0.0 | 8.5 |
| Owenbrin | 2009 | 30O01 | 0.0 | 3.0 | 6.0 | 0.0 | 0.0 | 9.0 |
| Owenriff (Corrib) | 2009 | 30O02 | 7.0 | 1.5 | 0.0 | 0.0 | 0.0 | 8.5 |
| Owenwee (Corrib) | 2006 | 30O03 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Robe | 2009 | 30R01 | 0.0 | 43.0 | 11.0 | 2.6 | 0.0 | 56.5 |
| Scardaun | 2009 | 30S04 | 0.0 | 2.5 | 0.0 | 1.0 | 0.0 | 3.5 |
| Sinking | 2009 | 30S01 | 0.0 | 10.6 | 7.4 | 0.5 | 0.0 | 18.5 |
| Srahnalong | 2009 | 30S03 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Srah Stream ¹ | 2009 | 30S02 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 1.5 |
| Terryland | 2009 | 30T01 | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 2.5 |
| Tullaghaun | 2009 | 30T03 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Yellow (Sinking) | 2009 | 30Y01 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |


| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|------|------|-------|----------|-------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Total Length (km) surveyed this cycle | 2009 | | 81.8 | 257.6 | 114.3 | 22.8 | 0.0 | 476.5 |
| Adjustments (km) (See below)* | | | 0.0 | 8.5 | 3.0 | 0.0 | 0.0 | 11.5 |
| <i>Percentages</i> | | | 17.6 | 53.6 | 23.9 | 4.9 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 330.9 | 111.3 | 22.8 | 0.0 | 465.0 |
| <i>Percentages</i> | | | | 71.2 | 23.9 | 4.9 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 351.5 | 77.5 | 36.0 | 0.0 | 465.0 |
| <i>Percentages</i> | | | | 75.7 | 16.6 | 7.7 | 0.0 | |
| Changes since Previous Survey (km) | | | | -20.6 | 33.8 | -13.2 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 30 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 30 : 2007- '09
% Surveyed Channel Length in Five WFD Quality Classes



WRBD: HYDROMETRIC AREA NO. 31



The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|-------------|-------------|-------------|-------------|-----------------|-------------|--------------|--------------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Cashla | 2009 | 31C01 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Crumlin (Galway Bay) | 2009 | 31C02 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Glencoaghan | 2009 | 31G01 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Gowlabeg | 2009 | 31G03 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Invermore | 2009 | 31I01 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Knock (Furbo) | 2009 | 31K01 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Knockadoagh | 2009 | 31K02 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Loughinch | 2009 | 31L01 | 0.0 | 2.1 | 2.5 | 0.0 | 0.0 | 4.5 |
| Lough Nabrocky Stream | 2009 | 31N01 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Owenboliska | 2009 | 31O01 | 2.3 | 4.8 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owengowla | 2009 | 31O02 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Owenriff (South Galway) | 2009 | 31O04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Owentooey | 2009 | 31O03 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Polleen | 2009 | 31P01 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Recess | 2009 | 31R01 | 6.6 | 9.0 | 0.0 | 0.0 | 0.0 | 15.5 |
| Screeb | 2009 | 31S01 | 0.0 | 7.5 | 1.5 | 0.0 | 0.0 | 9.0 |
| Tooreenacoona | 2009 | 31T01 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Total Length (km) surveyed this cycle | 2009 | | 18.8 | 63.3 | 9.0 | 0.0 | 0.0 | 91.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 20.7 | 69.5 | 9.8 | 0.0 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 07 - '09 | | 82.1 | 9.0 | 0.0 | 0.0 | 91.0 | |
| <i>Percentages</i> | | | 90.2 | 9.8 | 0.0 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 79.5 | 11.5 | 0.0 | 0.0 | 91.0 | |
| <i>Percentages</i> | | | 87.4 | 12.6 | 0.0 | 0.0 | 100.0 | |
| Total Length (km) surveyed this cycle | | | 2.6 | -2.6 | 0.0 | 0.0 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



WRBD: HYDROMETRIC AREA NO. 32

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Altaconey | 2009 | 32A02 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Ballinaboy | 2009 | 32B07 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 |
| Bundorragha | 2009 | 32B01 | 9.5 | 0.0 | 0.0 | 0.0 | 0.0 | 9.5 |
| Bunnahowna | 2009 | 32B02 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Bunowen (Louisburg) | 2009 | 32B03 | 15.0 | 0.0 | 2.0 | 0.0 | 0.0 | 17.0 |
| Carrowbeg (Westport) | 2009 | 32C05 | 7.7 | 3.8 | 1.0 | 0.0 | 0.0 | 12.5 |
| Carrowniskey | 2009 | 32C01 | 6.5 | 8.3 | 2.2 | 0.0 | 0.0 | 17.0 |
| Cross (Mayo) ¹ | 2009 | 32C02 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Crumpaun | 2009 | 32C03 | 12.0 | 0.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Culfin | 2009 | 32C04 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Clogher (Moyour) | 2009 | 32C07 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Dawros | 2009 | 32D01 | 6.0 | 0.0 | 6.5 | 0.0 | 0.0 | 12.5 |
| Derrycraff | 2009 | 32D02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Derryehorraun | 2009 | 32D04 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Erriff | 2009 | 32E01 | 0.0 | 28.0 | 0.0 | 0.0 | 0.0 | 28.0 |
| Glendavock | 2009 | 32G01 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Glenisland | 2009 | 32G07 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glenlaur | 2009 | 32G02 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glenamong | 2009 | 32G03 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Goulaun | 2009 | 32G06 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Glaishwy | 2009 | 32G12 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Moyour | 2009 | 32M01 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Newport (Mayo) | 2009 | 32N01 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Owenduff (Erriff) | 2009 | 32O08 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |

| | | | | | | | | |
|---|------------|-------|-------|----------|----------|----------|----------|--------------|
| Owengarve (Mayo) | 2009 | 32O02 | 0.0 | 0.0 | 7.5 | 0.0 | 0.0 | 7.5 |
| Owenglin | 2009 | 32O03 | 0.0 | 15.0 | 0.0 | 0.0 | 0.0 | 15.0 |
| Owennabrockagh | 2009 | 32O04 | 0.0 | 14.0 | 0.0 | 0.0 | 0.0 | 14.0 |
| Owennadornaun | 2009 | 32O07 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Owenwee (Mayo) | 2009 | 32O06 | 0.0 | 13.0 | 0.0 | 0.0 | 0.0 | 13.0 |
| Skerdagh | 2009 | 32S01 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Srahmore | 2009 | 32S02 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Streamstown (Clifden) | 2009 | 32S04 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 |
| Traheen | 2009 | 32T01 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Total Length (km) surveyed this cycle | 2009 | | 83.2 | 151.1 | 24.2 | 4.0 | 0.0 | 262.5 |
| Adjustments (km) (See below)* | | | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| <i>Percentages</i> | | | 32.4 | 56.6 | 9.4 | 1.6 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 228.3 | 24.2 | 4.0 | 0.0 | | 256.5 |
| <i>Percentages</i> | | | 89.0 | 9.4 | 1.6 | 0.0 | | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 220.5 | 11.5 | 24.5 | 0.0 | | 256.5 |
| <i>Percentages</i> | | | 86.0 | 4.5 | 9.6 | 0.0 | | 100.0 |
| Changes since Previous Survey (km) | | | 7.8 | 12.7 | -20.5 | 0.0 | | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



WRBD: HYDROMETRIC AREA NO. 33

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Altnabrocky | 2008 | 33A02 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Ballingen | 2008 | 33B01 | 9.5 | 1.5 | 0.0 | 0.0 | 0.0 | 11.0 |
| Barroosky | 2008 | 33B08 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Belderg | 2008 | 33B02 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Bellagarvaun | 2008 | 33B04 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Bellanaboy | 2008 | 33B07 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Bunanioo | 2008 | 33B09 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Cartron | 2008 | 33C02 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Cloonalaghan | 2008 | 33C01 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Clooneen (Mayo) | 2008 | 33C03 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Doega | 2008 | 33D01 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Doolough Stream | 2008 | 33D02 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 |
| Glenamoy | 2008 | 33G01 | 13.5 | 2.5 | 0.0 | 0.0 | 0.0 | 16.0 |
| Glencastle | 2008 | 33G08 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Glencullin (North Mayo) | 2008 | 33G02 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glencullin (West Mayo) | 2008 | 33G03 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Glenglassera | 2008 | 33G05 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Gortmore Stream (Mayo) | 2008 | 33G04 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Gweedaney | 2008 | 33G06 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Keerglen | 2008 | 33K01 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Muing | 2008 | 33M01 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 4.0 |
| Muingnabo | 2008 | 33M02 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Munhin | 2008 | 33M03 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 3.5 |
| Owenduff (Blacksod) | 2008 | 33O01 | 0.0 | 25.5 | 0.0 | 0.0 | 0.0 | 25.5 |

| | | | | | | | | |
|---|------------|-------|------|----------|----------|----------|----------|--------------|
| Owenmore (Mayo) | 2008 | 33O04 | 33.0 | 9.0 | 0.0 | 0.0 | 0.0 | 42.0 |
| Sheskin Stream | 2008 | 33S03 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Srahduggan Stream | 2008 | 33S05 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Tarsaghaunmore | 2008 | 33T01 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Total Length (km) surveyed this cycle | 2008 | | 74.0 | 129.5 | 7.0 | 5.0 | 0.0 | 215.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 34.3 | 60.1 | 3.2 | 2.3 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 203.5 | 7.0 | 5.0 | 0.0 | 215.5 |
| <i>Percentages</i> | | | | 94.4 | 3.2 | 2.3 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 190.5 | 18.0 | 7.0 | 0.0 | 215.5 |
| <i>Percentages</i> | | | | 89.2 | 8.4 | 2.3 | 0.0 | |
| Changes since Previous Survey (km) | | | | 13.0 | -11.0 | -2.0 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



WRBD: HYDROMETRIC AREA NO. 34

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Addergoole | 2007 | 34A01 | 8.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Bar Deela | 2007 | 34B02 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Behy (North Mayo) | 2007 | 34B08 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Bellanamean | 2007 | 34B04 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Bellawaddy | 2007 | 34B05 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Breaghwy | 2007 | 34B06 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Brusna (North Mayo) | 2007 | 34B07 | 0.0 | 12.0 | 0.0 | 0.0 | 0.0 | 12.0 |
| Burren Stream (Clydagh) | 2007 | 34B13 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Callow Loughs Stream | 2007 | 34C08 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 |
| Carroward | 2007 | 34C09 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Carrowkerribly Lough Stream | 2007 | 34C07 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Castlebar | 2007 | 34C01 | 5.0 | 4.2 | 1.5 | 11.3 | 0.0 | 22.0 |
| Charlestown Stream ¹ | 2007 | 34C28 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Cloonaghmore | 2007 | 34C03 | 23.0 | 0.0 | 0.0 | 0.0 | 0.0 | 23.0 |
| Cloonlavis | 2007 | 34C10 | 2.3 | 0.0 | 5.7 | 0.0 | 0.0 | 8.0 |
| Clydagh (Castlebar) | 2007 | 34C05 | 28.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.0 |
| Crumlin (Lough Cullin) | 2007 | 34C11 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Deel (Crossmolina) | 2007 | 34D01 | 13.7 | 28.4 | 0.0 | 0.0 | 0.0 | 42.0 |
| Duvowen (Cloonaghmore) | 2007 | 34D03 | 3.3 | 5.8 | 0.0 | 0.0 | 0.0 | 9.0 |
| Eignagh | 2007 | 34E01 | 12.5 | 0.0 | 0.0 | 0.0 | 0.0 | 12.5 |
| Fiddaunatooghaun Stream | 2007 | 34F06 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glenree | 2007 | 34G01 | 15.5 | 0.5 | 0.0 | 0.0 | 0.0 | 16.0 |
| Glore (Mayo) | 2007 | 34G02 | 4.2 | 5.6 | 5.9 | 4.4 | 0.0 | 20.0 |
| Gweestion | 2007 | 34G03 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |



| | | | | | | | | |
|---------------------------------|---------|-------|------|------|-----|------|-----|------|
| Leaffony | 2007 | 34L01 | 0.0 | 12.5 | 0.0 | 0.0 | 0.0 | 12.5 |
| Lenyvee | 2007 | 34L06 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Little (Strade) | 2007 | 34L02 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Lough Muck Stream | 2007 | 34L05 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Loughnaminoe Stream | 2007 | 34L04 | 2.0 | 2.2 | 4.8 | 0.0 | 0.0 | 9.0 |
| Mad | 2007 | 34M04 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 |
| Manulla | 2007 | 34M01 | 3.7 | 8.4 | 7.0 | 0.0 | 0.0 | 19.0 |
| Meander | 2007 | 34M05 | 0.0 | 3.0 | 2.0 | 0.0 | 0.0 | 5.0 |
| Moy | 2007 | 34M02 | 21.3 | 60.4 | 3.4 | 0.0 | 0.0 | 85.0 |
| Mullaghanoe | 2007 | 34M03 | 0.0 | 5.0 | 5.5 | 3.0 | 0.0 | 13.5 |
| Oughtagh | 2007 | 34O05 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Owenaher | 2007 | 34O01 | 11.0 | 3.0 | 0.0 | 0.0 | 0.0 | 14.0 |
| Owengarve (Sligo) | 2007 | 34O03 | 22.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.0 |
| Owenlobnaglaur | 2007 | 34O04 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Pollagh | 2007 | 34P01 | 8.0 | 2.0 | 1.5 | 1.5 | 0.0 | 13.0 |
| Shanvolahan | 2007 | 34S01 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Slieveclaur | 2007 | 34S06 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Sonnagh (Moy) | 2007 | 34S02 | 0.0 | 0.0 | 0.0 | 11.0 | 0.0 | 11.0 |
| Spaddagh | 2007 | 34S03 | 5.0 | 6.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Swinford ¹ | 2007 | 34S05 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Strade | 2007 | 34S04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Tubbercurry ¹ | 2007/09 | 34T02 | 0.0 | 0.0 | 0.0 | 2.5 | 2.5 | 5.0 |
| Tubbercurry Stream ¹ | 2007 | 34T03 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Trimoge | 2007 | 34T01 | 0.0 | 16.5 | 3.5 | 0.0 | 0.0 | 20.0 |
| Yellow (Foxford) | 2007 | 34Y01 | 16.0 | 0.0 | 0.0 | 0.0 | 0.0 | 16.0 |
| Yellow (Knock) | 2007 | 34Y02 | 3.8 | 11.0 | 2.2 | 0.0 | 0.0 | 17.0 |

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|------|-------|-------|----------|------|-------|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Total Length (km) surveyed this cycle | 2007 | | 247.6 | 260.8 | 46.0 | 46.7 | 2.5 | 603.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 5.5 | 2.5 | 8.0 |
| <i>Percentages</i> | | | 41.6 | 43.8 | 7.7 | 6.9 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 508.4 | 46.0 | 41.2 | 0.0 | 595.5 | |
| <i>Percentages</i> | | | 85.4 | 7.7 | 6.9 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 534.0 | 45.0 | 16.5 | 0.0 | 595.5 | |
| <i>Percentages</i> | | | 89.7 | 7.6 | 2.8 | 0.0 | | |
| Changes since Previous Survey (km) | | | -25.6 | 1.0 | 24.7 | 0.0 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



NWIRBD: HYDROMETRIC AREA NO. 35

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballymote Stream | 2009 | 35B04 | 0.0 | 2.5 | 0.5 | 2.0 | 0.0 | 5.0 |
| Ballysodare | 2009 | 35B05 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Bonet | 2009 | 35B06 | 10.2 | 20.4 | 4.5 | 0.0 | 0.0 | 35.0 |
| Brackary | 2009 | 35B10 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Bradoge | 2009 | 35B07 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Buncrowey | 2009 | 35B09 | 0.0 | 0.0 | 11.5 | 0.0 | 0.0 | 11.5 |
| Bunнанaddan Stream | 2009 | 35B08 | 0.0 | 0.5 | 0.0 | 3.5 | 0.0 | 4.0 |
| Cashel Stream (Bonet) | 2009 | 35C08 | 5.5 | 5.0 | 0.0 | 0.0 | 0.0 | 10.5 |
| Clooneen (Sligo) | 2009 | 35C01 | 0.0 | 14.0 | 0.0 | 0.0 | 0.0 | 14.0 |
| Diffreen | 2009 | 35D01 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Doonbeakin | 2009 | 35D09 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| Doonflin | 2009 | 35D10 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Douglas (Sligo) | 2009 | 35D02 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 4.0 |
| Drowes | 2009 | 35D03 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Drumcliffe | 2009 | 35D04 | 0.0 | 4.0 | 2.0 | 1.0 | 0.0 | 7.0 |
| Drumfin | 2009 | 35D11 | 0.0 | 5.5 | 2.0 | 0.0 | 0.0 | 7.5 |
| Duff | 2009 | 35D05 | 18.0 | 1.0 | 0.0 | 0.0 | 0.0 | 19.0 |
| Dunmorán | 2009 | 35D16 | 6.5 | 3.5 | 0.0 | 0.0 | 0.0 | 10.0 |
| Dunneill | 2009 | 35D06 | 13.0 | 1.0 | 0.0 | 0.0 | 0.0 | 14.0 |
| Easky | 2009 | 35E01 | 3.6 | 22.4 | 0.0 | 0.0 | 0.0 | 26.0 |
| Finned | 2009 | 35F01 | 7.2 | 5.8 | 0.0 | 0.0 | 0.0 | 13.0 |
| Garavoge | 2009 | 35G01 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Glenaniff | 2009 | 35G02 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Gowlan (Sligo) | 2009 | 35G03 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |

| | | | | | | | | |
|---|------------|-------|----------|----------|----------|----------|--------------|-------|
| Grange (Sligo) | 2009 | 35G04 | 7.5 | 0.5 | 0.0 | 0.0 | 0.0 | 8.0 |
| Gurteen Stream ¹ | 2009 | 35G05 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 4.0 |
| Killanummery | 2009 | 35K03 | 2.8 | 9.3 | 0.0 | 0.0 | 0.0 | 12.0 |
| Killoran Lough Stream | 2009 | 35K02 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Liskeagh | 2009 | 35L02 | 9.5 | 0.0 | 0.0 | 0.0 | 0.0 | 9.5 |
| Lugdoon Stream | 2009 | 35L01 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 5.5 |
| Owenbeg (Coolaney) | 2009 | 35O01 | 0.0 | 23.5 | 0.0 | 0.0 | 0.0 | 23.5 |
| Owenmore (Manorhamilton) | 2009 | 35O08 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Owenmore (Sligo) | 2009 | 35O06 | 0.0 | 38.5 | 8.5 | 3.0 | 0.0 | 50.0 |
| Shanvaus | 2009 | 35S01 | 8.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Tullinwillin Stream | 2009 | 35T03 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Tullynascreen | 2009 | 35T06 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Unshin | 2009 | 35U01 | 6.4 | 14.7 | 1.9 | 0.0 | 0.0 | 23.0 |
| Willsborough Stream | 2009 | 35W01 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Total Length (km) surveyed this cycle | 2009 | | 111.7 | 245.5 | 53.4 | 9.5 | 0.0 | 420.0 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 |
| <i>Percentages</i> | | | 26.7 | 58.7 | 12.3 | 2.3 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 357.1 | 51.4 | 9.5 | 0.0 | 418.0 | |
| <i>Percentages</i> | | | 85.4 | 12.3 | 2.3 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | 362.5 | 35.5 | 20.0 | 0.0 | 418.0 | |
| <i>Percentages</i> | | | 86.7 | 8.5 | 4.8 | 0.0 | 100.0 | |
| Changes since Previous Survey (km) | | | -5.4 | 15.9 | -10.5 | 0.0 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



NWIRBD: HYDROMETRIC AREA NO. 36

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aghacashlaun | 2007 | 36A03 | 7.5 | 8.5 | 0.0 | 0.0 | 0.0 | 16.0 |
| Aghnacliffe Stream | 2007 | 36A06 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Annadale Stream | 2007 | 36A05 | 0.0 | 8.5 | 1.5 | 0.0 | 0.0 | 10.0 |
| Annalee | 2007 | 36A02 | 0.0 | 29.5 | 17.2 | 2.4 | 0.0 | 49.0 |
| Avaghon lake Stream | 2007 | 36A07 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Bawnboy | 2007 | 36B07 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Blackwater (Newtowngore) | 2007 | 36B04 | 0.0 | 15.5 | 3.0 | 0.0 | 0.0 | 18.5 |
| Blackwater (Swanlinbar) | 2007 | 36B03 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Bunnoe | 2007 | 36B05 | 0.0 | 15.5 | 5.5 | 0.0 | 0.0 | 21.0 |
| Cavan | 2007 | 36C02 | 0.0 | 14.0 | 0.0 | 7.0 | 0.0 | 21.0 |
| Conawary (Upper) | 2007 | 36C11 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Cornavannoge | 2007 | 36C04 | 0.0 | 16.0 | 0.0 | 0.0 | 0.0 | 16.0 |
| Cullies | 2007 | 36C03 | 0.0 | 2.5 | 10.0 | 3.5 | 0.0 | 16.0 |
| Derradda Stream | 2007 | 36D07 | 0.0 | 7.5 | 0.0 | 2.0 | 0.0 | 9.5 |
| Dromore | 2007 | 36D02 | 0.0 | 9.0 | 16.0 | 11.0 | 0.0 | 36.0 |
| Drumane Stream ¹ | 2007 | 36D04 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Erne | 2007 | 36E01 | 0.0 | 0.0 | 57.0 | 3.0 | 0.0 | 60.0 |
| Finn (Monaghan) | 2007 | 36F01 | 0.0 | 10.5 | 14.0 | 0.0 | 0.0 | 24.5 |
| Knappagh | 2007 | 36K01 | 0.0 | 0.0 | 3.0 | 4.5 | 0.0 | 7.5 |
| Laheen Stream | 2007 | 36L02 | 0.0 | 6.5 | 4.5 | 0.0 | 0.0 | 11.0 |
| Laragh | 2007 | 36L01 | 0.0 | 11.0 | 10.5 | 0.0 | 0.0 | 21.5 |
| Legga Stream | 2007 | 36L03 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Madabawn Stream | 2007 | 36M02 | 0.0 | 4.0 | 6.5 | 0.0 | 0.0 | 10.5 |
| Magherarney | 2007 | 36M01 | 0.0 | 0.0 | 0.0 | 8.0 | 0.0 | 8.0 |

| | | | | | | | | | | |
|---|------|------------|------|-----|-------|-------|-------|------|-------|-------|
| Maghery | 2007 | 36M03 | 0.0 | 0.0 | 14.5 | 1.0 | 0.0 | 15.5 | | |
| Owensallagh | 2007 | 36O01 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 9.0 | | |
| Rag (Cavan) | 2007 | 36R01 | 0.0 | 7.5 | 0.0 | 0.0 | 0.0 | 7.5 | | |
| Roo | 2007 | 36R02 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 | | |
| Stradone | 2007 | 36S02 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 | | |
| Swanlinbar | 2007 | 36S01 | 10.0 | 2.0 | 0.0 | 0.0 | 0.0 | 12.0 | | |
| Waterfoot ¹ | 2007 | 36W03 | 6.5 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 | | |
| Yellow (Ballinamore) | 2007 | 36Y01 | 0.0 | 8.5 | 3.0 | 1.5 | 0.0 | 13.0 | | |
| Total Length (km) surveyed this cycle | | | 2007 | | 35.0 | 207.0 | 171.2 | 51.9 | 0.0 | 465.0 |
| Adjustments (km) (See below)* | | | | | 6.5 | 5.0 | 0.0 | -2.5 | 0.0 | 9.0 |
| Percentages | | | | | 6.3 | 44.3 | 37.5 | 11.9 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | | A | B | C | D | Total | |
| Baseline : Current Adjusted Status (km) | | 2007 - '09 | | | 230.5 | 171.2 | 54.4 | 0.0 | 456.0 | |
| Percentages | | | | | 50.5 | 37.5 | 11.9 | 0.0 | 100.0 | |
| Baseline: Previous Unadjusted Status (km)** | | 2004 - '06 | | | 296.0 | 108.0 | 50.5 | 1.5 | 456.0 | |
| Percentages | | | | | 64.9 | 23.7 | 11.1 | 0.3 | | |
| Changes since Previous Survey (km) | | | | | -65.6 | 63.2 | 3.9 | -1.5 | 0.0 | |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



NWIRBD: HYDROMETRIC AREA NO. 37

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.



| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballaghdoe | 2008 | 37B01 | 0.0 | 0.0 | 1.2 | 3.3 | 1.6 | 6.0 |
| Ballintra | 2008 | 37B02 | 0.0 | 6.3 | 5.1 | 4.1 | 0.0 | 15.5 |
| Bridgetown (Donegal) | 2008 | 37B03 | 12.4 | 1.6 | 0.0 | 0.0 | 0.0 | 14.0 |
| Cloghanmore | 2008 | 37C05 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 2.0 |
| Clogher (Donegal) | 2005 | 37C06 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Corabber | 2008 | 37C01 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 |
| Crow | 2008 | 37C03 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 7.0 |
| Cunlin Lough Stream | 2008 | 37C08 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Crowanrudda (Ballaghdoe) | 2008 | 37C09 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Drummenny | 2008 | 37D01 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Eanybeg Water | 2008 | 37E01 | 0.0 | 13.0 | 1.0 | 0.0 | 0.0 | 14.0 |
| Eanymore Water | 2008 | 37E02 | 0.0 | 13.3 | 2.5 | 1.3 | 0.0 | 17.0 |
| Eglis | 2008 | 37E04 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Eske | 2008 | 37E05 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Fintragh | 2008 | 37F01 | 4.5 | 0.0 | 0.0 | 0.0 | 0.0 | 4.5 |
| Glen (Carrick) | 2008 | 37G01 | 0.0 | 13.0 | 0.0 | 0.0 | 0.0 | 13.0 |
| Glenaddragh | 2008 | 37G02 | 0.0 | 0.0 | 7.0 | 2.0 | 0.0 | 9.0 |
| Laghy Stream | 2008 | 37L02 | 0.0 | 8.5 | 1.0 | 0.0 | 0.0 | 9.5 |
| Loughaderry | 2008 | 37L03 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Lowerymore | 2008 | 37L01 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 11.0 |
| Oily | 2008 | 37O01 | 15.0 | 1.0 | 0.0 | 0.0 | 0.0 | 16.0 |
| Owenteskiny | 2008 | 37O02 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Owenwee (Carrick) | 2008 | 37O03 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Roechrow | 2008 | 37R01 | 0.0 | 0.0 | 0.0 | 1.0 | 2.0 | 3.0 |

| | | | | | | | | |
|---|------------|-------|------|----------|----------|----------|----------|--------------|
| Stragar | 2008 | 37S02 | 7.4 | 1.6 | 0.0 | 0.0 | 0.0 | 9.0 |
| Tullinteane | 2008 | 37T01 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Total Length (km) surveyed this cycle | 2008 | | 59.8 | 104.3 | 20.8 | 13.6 | 3.6 | 202.0 |
| Adjustments (km) (See below)* | | | -2.3 | -9.8 | 0.0 | 0.0 | 0.0 | -12.0 |
| <i>Percentages</i> | | | 29.0 | 53.3 | 9.7 | 6.4 | 1.7 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 176.1 | 20.8 | 13.6 | 3.6 | 214.0 |
| <i>Percentages</i> | | | | 82.3 | 9.7 | 6.4 | 1.7 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 175.5 | 13.5 | 22.0 | 3.0 | 214.0 |
| <i>Percentages</i> | | | | 82.0 | 6.3 | 10.3 | 1.4 | 100.0 |
| Changes since Previous Survey (km) | | | | 0.6 | 7.3 | -8.4 | 0.6 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



NWIRBD: HYDROMETRIC AREA NO. 38

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Abberachrin | 2009 | 38A01 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Aighe | 2009 | 38A03 | 0.0 | 2.0 | 1.0 | 0.0 | 0.0 | 3.0 |
| Aspick | 2009 | 38A02 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Big Burn | 2009 | 38B03 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Bracky | 2009 | 38B02 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 1.5 |
| Bunlin | 2009 | 38B04 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Burnside | 2009 | 38B05 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Calabber | 2009 | 38C01 | 4.0 | 3.5 | 0.0 | 0.0 | 0.0 | 7.5 |
| Carrownamaddy | 2009 | 38C02 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Clady (Donegal) | 2009 | 38C04 | 0.0 | 3.5 | 4.0 | 0.0 | 0.0 | 7.5 |
| Cloghernagore | 2009 | 38C07 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Corveen | 2009 | 38C05 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Cronaniv Burn | 2009 | 38C06 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Devlin (Donegal) | 2009 | 38D01 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Dungloe | 2009 | 38D02 | 0.0 | 3.0 | 7.0 | 0.0 | 0.0 | 10.0 |
| Duntally | 2009 | 38D03 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 |
| Duvoge | 2009 | 38D05 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Duvowen (Ray) | 2009 | 38D04 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Faymore | 2009 | 38F01 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 7.0 |
| Glen (Lackagh) | 2009 | 38G04 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Glen (Meenclady) | 2009 | 38G05 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Glenleheen Stream | 2009 | 38G07 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 5.5 |
| Glenna | 2009 | 38G01 | 2.5 | 6.5 | 0.0 | 0.0 | 0.0 | 9.0 |
| Glentornan | 2009 | 38G06 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |



| | | | | | | | | |
|----------------------|------|-------|-----|------|-----|-----|-----|------|
| Gweebarra | 2009 | 38G02 | 9.0 | 4.0 | 1.0 | 0.0 | 0.0 | 14.0 |
| Gweedore | 2009 | 38G03 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 5.5 |
| Keel Lough Stream | 2009 | 38K01 | 0.0 | 2.0 | 1.0 | 2.0 | 0.0 | 5.0 |
| Lough Agher Stream | 2009 | 38L02 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Loughkeel Burn | 2009 | 38L03 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Murlin | 2009 | 38M03 | 0.0 | 4.0 | 0.0 | 1.0 | 0.0 | 5.0 |
| Owenator | 2009 | 38O01 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| Owenawillin | 2009 | 38O10 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 3.0 |
| Owencarrow | 2009 | 38O03 | 4.6 | 3.4 | 0.0 | 0.0 | 0.0 | 8.0 |
| Owencronahulla | 2009 | 38O09 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Owenea | 2009 | 38O04 | 0.0 | 18.0 | 4.0 | 0.0 | 0.0 | 22.0 |
| Owennamarve | 2009 | 38O05 | 0.0 | 9.5 | 0.5 | 0.0 | 0.0 | 10.0 |
| Owentocker | 2009 | 38O06 | 0.0 | 17.0 | 0.0 | 0.0 | 0.0 | 17.0 |
| Owenveagh | 2009 | 38O14 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Owenwee (Doocharry) | 2009 | 38O07 | 0.0 | 6.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Owenwee (Dunlewey) | 2009 | 38O12 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Owenwee (Glen Lough) | 2009 | 38O13 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Owenwee (Loughros) | 2009 | 38O08 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Port Stream | 2009 | 38P01 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 2.0 |
| Ray | 2009 | 38R01 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 |
| Shallogan | 2009 | 38S03 | 0.0 | 0.0 | 6.0 | 0.0 | 0.0 | 6.0 |
| Sruhanameel | 2009 | 38S02 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Stracashel | 2009 | 38S01 | 0.0 | 3.5 | 0.0 | 0.5 | 0.0 | 4.0 |
| Tullaghobegley | 2009 | 38T01 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 9.0 |

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---|------------|------|------|-------|----------|-------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Total Length (km) surveyed this cycle | 2009 | | 44.1 | 170.4 | 54.0 | 5.0 | 0.0 | 273.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| <i>Percentages</i> | | | 16.1 | 62.3 | 19.7 | 1.8 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 214.5 | 54.0 | 5.0 | 0.0 | 273.5 |
| <i>Percentages</i> | | | | 78.4 | 19.7 | 1.8 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 225.0 | 18.5 | 30.0 | 0.0 | 273.5 |
| <i>Percentages</i> | | | | 82.3 | 6.8 | 11.0 | 0.0 | |
| Changes since Previous Survey (km) | | | | -10.5 | 35.5 | -25.0 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.



NWIRBD: HYDROMETRIC AREA NO. 39

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Aghaweel | 2007 | 39A01 | 0.0 | 1.5 | 0.5 | 0.0 | 0.0 | 2.0 |
| Bullaba | 2007 | 39B01 | 6.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.0 |
| Burnfoot | 2007 | 39B02 | 0.0 | 0.0 | 10.0 | 0.0 | 0.0 | 10.0 |
| Camowen (Crana) ¹ | 2007 | 39C08 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Cashelnacor | 2007 | 39C01 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Corravaddy Burn | 2007 | 39C03 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Crana | 2007 | 39C02 | 0.0 | 2.5 | 3.0 | 11.5 | 0.0 | 17.0 |
| Dooballagh Burn | 2007 | 39D02 | 0.0 | 0.0 | 0.0 | 6.0 | 0.0 | 6.0 |
| Drumbarnet Stream | 2007 | 39D03 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 5.0 |
| Drumhallagh | 2007 | 39D01 | 3.5 | 0.0 | 0.0 | 0.5 | 0.0 | 4.0 |
| Glashagh (Upper) | 2007 | 39G01 | 0.0 | 8.0 | 1.5 | 0.0 | 0.0 | 9.5 |
| Gliskeelan | 2007 | 39G05 | 0.0 | 0.0 | 4.5 | 0.0 | 0.0 | 4.5 |
| Glenalla | 2007 | 39G03 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 5.0 |
| Glenvar | 2007 | 39G04 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Leannan | 2007 | 39L01 | 12.2 | 7.5 | 3.3 | 1.6 | 0.0 | 24.5 |
| Leslie Hill Stream | 2007 | 39L05 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |
| Lownagh | 2007 | 39L04 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Lurgy | 2007 | 39L02 | 0.0 | 7.0 | 0.0 | 0.5 | 0.0 | 7.5 |
| Mill (Donegal) | 2007 | 39M02 | 5.5 | 4.5 | 0.0 | 0.5 | 0.0 | 10.5 |
| Maggy's Burn ¹ | 2007 | 39M01 | 0.0 | 1.0 | 0.0 | 2.0 | 0.0 | 3.0 |
| Owenboy (Crana) | 2007 | 39O04 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 5.0 |
| Owenerk | 2007 | 39O02 | 0.0 | 0.0 | 8.0 | 0.0 | 0.0 | 8.0 |
| Owennasop | 2007 | 39O05 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 |
| Owenwee (Lough Gartan) | 2007 | 39O03 | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 |


| | | | | | | | | |
|---|------------|-------|------|----------|----------|----------|----------|--------------|
| Skeoge | 2007 | 39S01 | 0.0 | 0.0 | 0.0 | 7.0 | 0.0 | 7.0 |
| Swilly | 2007 | 39S02 | 5.3 | 5.8 | 0.0 | 6.0 | 0.0 | 17.0 |
| Total Length (km) surveyed this cycle | 2007 | | 35.4 | 58.2 | 35.8 | 52.6 | 0.0 | 182.0 |
| Adjustments (km) (See below)* | | | 0.0 | -1.0 | -4.5 | 5.0 | 0.0 | -0.5 |
| <i>Percentages</i> | | | 19.4 | 32.5 | 22.1 | 26.1 | 0.0 | 100.0 |
| Channel Length (km) in Class | | | | A | B | C | D | Total |
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | | 94.6 | 40.3 | 47.6 | 0.0 | 182.5 |
| <i>Percentages</i> | | | | 51.8 | 22.1 | 26.1 | 0.0 | 100.0 |
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | | | 133.0 | 21.0 | 27.5 | 1.0 | 182.5 |
| <i>Percentages</i> | | | | 72.9 | 11.5 | 15.1 | 0.5 | |
| Changes since Previous Survey (km) | | | | -38.4 | 19.3 | 20.1 | -1.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).


** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 39 : Trends
 % Surveyed Channel Length in Four Quality Classes



Hydrometric Area 39 : 2007 - '09
 % Surveyed Channel Length in Five WFD Quality Classes



NWIRBD: HYDROMETRIC AREA NO. 40

The baseline rivers showing the channel length surveyed (km) and the estimated channel length in four biological quality classes:- A - Unpolluted, B - Slightly polluted/eutrophic, C - Moderately polluted and D - Seriously polluted. Data from biological surveys in period 2007-2009.

| Channel Length (km) in Class | | | A | A | B | C | D | Total |
|---------------------------------------|------|-------|------|------|----------|------|-----|-------|
| WFD Quality Class | | | High | Good | Moderate | Poor | Bad | |
| River Name | Year | Code | | | | | | |
| Ballyboe | 2007 | 40B03 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Ballyhallan | 2007 | 40B01 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 4.0 |
| Bredagh | 2007 | 40B02 | 8.0 | 0.0 | 0.0 | 0.0 | 0.5 | 8.5 |
| Cabry | 2007 | 40C03 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 4.0 |
| Clonmany | 2007 | 40C01 | 0.0 | 0.0 | 0.0 | 9.5 | 0.0 | 9.5 |
| Cloontagh | 2004 | 40C04 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 3.0 |
| Culdaff | 2004 | 40C02 | 0.0 | 0.0 | 0.0 | 17.0 | 0.0 | 17.0 |
| Donagh | 2007 | 40D01 | 0.0 | 9.0 | 0.0 | 3.0 | 0.0 | 12.0 |
| Drung | 2007 | 40D02 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 3.0 |
| Glennagannon | 2007 | 40G01 | 11.0 | 0.0 | 1.0 | 0.0 | 0.0 | 12.0 |
| Keenagh | 2007 | 40K01 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Long Glen | 2007 | 40L01 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Lough Nastackan Stream | 2007 | 40L03 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 |
| Malin Stream | 2007 | 40M01 | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 2.5 |
| Portaleen Stream | 2007 | 40P02 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 1.5 |
| Roosky | 2007 | 40R01 | 0.0 | 0.0 | 0.0 | 0.5 | 1.0 | 1.5 |
| Shivnagh | 2007 | 40S05 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Straid | 2007 | 40S01 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 |
| Total Length (km) surveyed this cycle | 2007 | | 33.0 | 30.5 | 1.0 | 39.5 | 1.5 | 105.5 |
| Adjustments (km) (See below)* | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Percentages | | | 31.3 | 28.9 | 0.9 | 37.4 | 1.4 | 100.0 |

| Channel Length (km) in Class | | | A | B | C | D | Total |
|---|------------|--|------|-----|------|-----|-------|
| Baseline : Current Adjusted Status (km) | 2007 - '09 | | 63.5 | 1.0 | 39.5 | 1.5 | 105.5 |
| Percentages | | | 60.2 | 0.9 | 37.4 | 1.4 | 100.0 |


| | | | | | | |
|---|------------|-------|------|------|-----|-------|
| Baseline: Previous Unadjusted Status (km)** | 2004 - '06 | 87.5 | 4.0 | 12.5 | 1.5 | 105.5 |
| <i>Percentages</i> | | 82.9 | 3.8 | 11.8 | 1.4 | |
| Changes since Previous Survey (km) | | -24.0 | -3.0 | 27.0 | 0.0 | 0.0 |

*Adjustments: Deduct the 'extras' (+), add the 'shortages' (-).

** Table II.4 Clabby *et al.*, 2008

¹ Rivers not previously included in the National Statistics.

Hydrometric Area 40 : Trends
% Surveyed Channel Length in Four Quality Classes



Hydrometric Area 40 : 2007 - '09
% Surveyed Channel Length in Five WFD Quality Classes

