



## **IRELAND**

# **INFORMATIVE INVENTORY REPORT 2025**

### **AIR POLLUTANT EMISSIONS IN IRELAND 1990–2023 REPORTED TO THE SECRETARIAT OF THE UNECE CONVENTION ON LONG-RANGE TRANSBOUNDARY AIR POLLUTION AND TO THE EUROPEAN UNION**

N. Rahman, D. Rossiter, B. Hyde, P. Duffy, S. Monaghan, N. Streethran  
F. Martini and A. Brandonisio

Environmental Protection Agency  
An Ghníomhaireacht um Chaomhnú Comhshaoil  
PO Box 3000, Johnstown Castle, Co. Wexford. Ireland

Telephone: +353 53 9160600 Fax: +353 53 9160699  
Email: [info@epa.ie](mailto:info@epa.ie) Web site: [www.epa.ie](http://www.epa.ie)  
LoCall: 0818 335 599

ISBN 978-1-80009-238-9

March 2025

## Contents

<b>Contents</b> .....	<b>iii</b>
<b>List of Tables</b> .....	<b>ix</b>
<b>List of Figures</b> .....	<b>xi</b>
<b>EXECUTIVE SUMMARY</b> .....	<b>1</b>
<b>Chapter One - Introduction</b> .....	<b>3</b>
1.1 Convention on Long-Range Transboundary Air Pollution .....	3
1.2 Inventory Reporting and Review under the Convention on Long-Range Transboundary Air Pollution (CLRTAP) .....	3
1.3 Inventory Reporting and Review under Directive (EU) 2016/2284 .....	4
1.4 Inventory Reporting by the Environmental Protection Agency .....	5
1.4.1 National Atmospheric Inventory System .....	5
1.4.2 Scope of Inventories under the LRTAP Convention and Directive (EU) 2016/2284 .....	6
1.4.3 Inventory Preparation .....	9
1.4.4 Data Acquisition .....	10
1.4.5 Quality Assurance and Quality Control (QA/QC) .....	11
1.4.6 Inventory Compilation .....	13
1.5 Key Category Analysis .....	14
1.6 Uncertainty Assessment .....	16
<b>Chapter Two - Analysis of Key Emission Trends</b> .....	<b>18</b>
2.1 Introduction .....	18
2.2 Main Pollutants .....	18
2.2.1 Sulphur Dioxide (SO <sub>2</sub> ) .....	18
2.2.2 Nitrogen Oxides (NO <sub>x</sub> ) .....	19
2.2.3 Ammonia (NH <sub>3</sub> ) .....	20
2.2.4 Non-Methane Volatile Organic Compounds (NMVOCs) .....	22
2.2.5 Carbon Monoxide (CO) .....	23
2.3 Particulate Matter .....	24
2.3.1 Particulate Matter <10 µm Diameter (PM <sub>10</sub> ) .....	24
2.3.2 Particulate Matter <2.5 µm Diameter (PM <sub>2.5</sub> ) .....	25
2.3.3 Total Suspended Particulates (TSP) .....	27
2.3.4 Black Carbon (BC) .....	28
2.4 Priority Metals .....	29
2.4.1 Lead (Pb) .....	29
2.4.2 Cadmium (Cd) .....	29
2.4.3 Mercury (Hg) .....	30
2.5 Other Metals .....	31
2.5.1 Arsenic (As) .....	31
2.5.2 Chromium (Cr) .....	32
2.5.3 Copper (Cu) .....	33
2.5.4 Nickel (Ni) .....	34
2.5.5 Selenium (Se) .....	35

2.5.6 Zinc (Zn) .....	36
2.6 Persistent Organic Pollutants (POPs).....	37
2.6.1 Dioxins and Furans (PCDD/F).....	38
2.6.2 Hexachlorobenzene (HCB).....	39
2.6.3 Polychlorinated Biphenyls (PCBs) .....	39
2.6.4 Polycyclic Aromatic Hydrocarbons (PAHs) .....	40
<b>Chapter - Three Energy .....</b>	<b>42</b>
3.1 Overview of the Energy (NFR 1) Sector .....	42
3.2 Public Electricity and Heat Production (NFR 1A1a) .....	43
3.2.1 Emissions of Sulphur Dioxide and Nitrogen Oxides.....	43
3.2.2 Emissions of Other Substances.....	45
3.3 Petroleum Refining (NFR 1A1b).....	45
3.4 Manufacture of Solid Fuels and Other Energy Industries (NFR 1A1c) .....	46
3.5 Manufacturing Industries and Construction (NFR 1A2) .....	46
3.6 Transport (NFR 1A3).....	48
3.6.1 Domestic and International LTOs (NFRs 1A3ai(i) and 1A3ai(ii)) .....	49
3.6.2 Road Transport (NFR 1A3b).....	51
3.6.3 Railways (NFR 1A3c) .....	54
3.6.4 National Navigation (NFR 1A3dii) .....	54
3.6.5 Other Transportation (NFR 1A3e) .....	55
3.7 Other Sectors (NFR 1A4).....	55
3.7.1 Commercial / Institutional (NFR 1A4a) and Residential (1A4b) .....	55
3.7.2 Agriculture/Forestry/Fishing (NFR 1A4ci and ii) .....	55
3.7.3 Agriculture/Forestry/Fishing: National Fishing (NFR 1A4ciii) .....	56
3.8 Other Stationary and Mobile (including military) (NFR 1A5).....	56
3.9 Fugitive emissions from Solid Fuels (NFR 1B1).....	56
3.9.1 Coal mining and handling (1B1a).....	56
3.10 Fugitive emissions from Oil and Natural Gas (NFR 1B2).....	58
3.10.1 Oil (NFR 1B2a) .....	58
3.10.2 Natural Gas (NFR 1B2b).....	58
3.10.3 Venting and flaring (oil, gas, combined oil and gas) (NFR1B2c) .....	60
3.11 Recalculations in the Energy Sector .....	60
3.12 Quality Assurance/Quality Control.....	71
3.13 Planned Improvements .....	71
3.14 Memo Items .....	71
<b>Chapter Four - Industrial Processes and Product Use .....</b>	<b>72</b>
4.1 Overview of the Industrial Processes and Product Use (NFR 2) Sector .....	72
4.2 Mineral Products (NFR 2A) .....	72
4.2.1 Cement Production (NFR 2A1) .....	72
4.2.2 Lime Production (NFR 2A2) .....	73
4.2.3 Glass Production (NFR 2A3).....	73
4.2.4 Quarrying and Mining of Minerals Other than Coal (NFR 2A5a) .....	75

4.2.5 Construction and Demolition (NFR 2A5b) .....	75
4.2.6 Storage, Handling and Transport of Mineral Products (NFR 2A5c) .....	75
4.2.7 Other (NFR 2A6) .....	75
4.3 Chemical Industry (NFR 2B) .....	76
4.3.1 Nitric Acid Production (NFR 2B2) .....	76
4.3.2 Storage, handling and transport of chemical products (NFR 2B10b) .....	77
4.4 Metal Production (NFR 2C) .....	77
4.4.1 Iron and Steel Production (NFR 2C1) .....	77
4.4.2 Ferroalloys Production (NFR 2C2) .....	78
4.4.3 Aluminium Production (NFR 2C3) .....	79
4.4.4 Lead Production (NFR 2C5) .....	79
4.4.5 Other Metal Production (NFR 2C7) .....	79
4.5 Overview of NMVOC emissions from Solvent and Other Product Use (NFR 2D-2L) Sector .....	80
4.5.1 Domestic Solvent Use including fungicides (NFR 2D3a) .....	84
4.5.2 Road Paving with asphalt (NFR 2D3b) .....	84
4.5.3 Coating Application (NFR 2D3d) .....	85
4.5.4 Degreasing and Dry Cleaning (NFR 2D3e and 2D3f) .....	88
4.5.5 Chemical Products, Manufacture and Processing (NFR 2D3g) .....	89
4.5.6 Printing (2D3h) .....	92
4.5.7 Other Use of Solvents and Related Activities (2D3i-2G) .....	92
4.5.8 Food and Beverage Industry (NFR 2H2) .....	95
4.6 Consumption of POPs and heavy metals 2K (e.g. electrical and scientific equipment) .....	96
4.7 Other production, consumption, storage, transportation or handling of bulk product (NFR 2L) .....	97
4.7.1 Leakage from Electrical Equipment (NFR 2L) .....	97
4.7.2 Fragmentisers and Shredders (NFR 2L) .....	98
4.8 Recalculations in the Industrial Processes Sector .....	99
4.8.1 Heavy Metals recalculations .....	99
4.8.2 Particulate matter recalculations .....	99
4.8.3 Nitrogen Oxides (NO <sub>x</sub> ) recalculations .....	99
4.8.4 Non- Methane Volatile Organic Compounds (NMVOC) recalculations .....	100
4.9 Quality Assurance/Quality Control .....	100
4.10 Overview of Notation Keys .....	100
4.11 Planned Improvements .....	100
<b>Chapter Five - Agriculture .....</b>	<b>110</b>
5.1 Overview of the Agriculture (NFR 3) Sector .....	110
5.2 Manure Management (NFR 3B) .....	113
5.2.1 Cattle (NFR 3B1) .....	114
5.2.2 Sheep (3B2) .....	115
5.2.3 Swine (NFR 3B3) .....	116
5.2.4 Poultry (NFR 3B4g) .....	117
5.2.5 Other livestock – Goats (NFR 3B4d), Horses (3B4e), Mules and Asses (NFR 3B4f) and Other Animals (3B4h) .....	118

5.2.6 Uncertainties .....	119
5.3 Agricultural Soils (NFR 3D).....	119
5.3.1 Direct Soil Emissions - Inorganic N-fertilizers (NFR 3Da1) .....	119
5.3.2 Direct Soil Emissions – Livestock manure applied to soils (NFR 3Da2a).....	120
5.3.3 Direct Soil Emissions – Sewage sludge applied to soils (NFR 3Da2b) .....	120
5.3.4 Direct Soil Emissions – Other organic fertilizers applied to soils (NFR 3Da2c) .....	120
5.3.5 Direct Soil Emissions – Urine and dung deposited during grazing (NFR 3Da3) .....	120
5.3.6 Farm-level agricultural operations (NFR 3Dc) .....	121
5.3.7 Off-farm storage, handling and transport (NFR 3Dd).....	121
5.3.8 Cultivated crops (NFR 3De) .....	121
5.3.9 Use of Pesticides (NFR 3Df) .....	121
5.4 Field burning of agricultural residues (NFR 3F) .....	121
5.5 Uncertainties associated with Synthetic Fertilizer (3Da1), Organic fertilizers (3Da2c) and urine and dung deposited by grazing animals (3Da3). .....	122
5.6 Recalculations in the Agriculture Sector .....	122
5.7 Quality Assurance/Quality Control.....	122
5.8 Planned Improvements .....	122
<b>Chapter Six - Waste .....</b>	<b>133</b>
6.1 Overview of the Waste (NFR 5) Sector .....	133
6.2 Biological treatment of waste - Solid waste disposal on land (NFR 5A) .....	133
6.2.1 Main Pollutants .....	133
6.2.2 Heavy Metals.....	134
6.3 Biological treatment of waste - Composting (NFR 5B1) and Anaerobic Digestion Biogas (NFR 5B2).....	134
6.4 Waste Incineration (NFR 5C) .....	135
6.4.1 Clinical Waste (5C1biii).....	135
6.4.2 Industrial Waste (5C1bi), Hazardous Waste (5C1bii) and Sewage Sludge (5C1biv) .....	137
6.4.3 Crematoria (5C1bv) .....	138
6.4.4 Open Burning of Agricultural Wastes (5C2).....	139
6.5 Wastewater handling (NFR 5D) .....	140
6.6 Other Waste (NFR 5E) .....	141
6.6.1 Accidental Fires .....	141
6.6.2 Other Burning.....	142
6.7 Recalculations in the Waste Sector .....	144
6.8 Quality Assurance/Quality Control.....	144
6.9 Planned Improvements .....	144
<b>Chapter Seven - Gridded and LPS data .....</b>	<b>150</b>
7.1 Overview of Gridded and LPS data reporting .....	150
7.2 Mapping methodology .....	154
7.3 Mapping methods for each GNFR .....	157
7.3.1 Public Power (GNFR A_PublicPower) .....	158
7.3.2 Industry (GNFR B_Industry).....	158
7.3.3 Other stationary combustion (GNFR C_OtherStationaryComb).....	158

7.3.4 Fugitive Emissions (GNFR D_Fugitive) .....	158
7.3.5 Solvents (GNFR E_Solvents) .....	159
7.3.6 Road transport (GNFR F_RoadTransport) .....	159
7.3.7 National navigation (GNFR G_Shipping) .....	159
7.3.8 Aviation (LTO) (GNFR H_Aviation) .....	159
7.3.9 Off road mobile sources (GNFR I_Offroad) .....	159
7.3.10 Waste handling and treatment (GNFR J_Waste) .....	160
7.3.11 Agricultural livestock (GNFR K_AgriLivestock) .....	160
7.3.12 Agricultural soils (Other emissions) (GNFR L_AgriOther) .....	160
7.3.13 Aviation (Cruise) (GNFR O_AviCruise) .....	160
7.3.14 International navigation (GNFR P_IntShipping) .....	161
7.4 National total emission mapped by the EMEP 0.1 degree x 0.1 degree grid .....	161
<b>Chapter Eight - Projections .....</b>	<b>167</b>
8.1 Overview of Emissions Projections .....	167
8.2 Emission Reduction Commitments for 2020 and 2030 .....	168
8.3 With Existing Measures and With Additional Measures Scenarios - Approach .....	168
8.4 Key Trends .....	170
8.4.1 Sulphur Dioxide (SO <sub>2</sub> ) .....	170
8.4.2 Nitrogen Oxides (NO <sub>x</sub> ) .....	172
8.4.3 Ammonia (NH <sub>3</sub> ) .....	175
8.4.4 Non-Methane Volatile Organic Compounds (NMVOCs) .....	177
8.4.5 Particulate Matter < 2.5 µm in diameter (PM <sub>2.5</sub> ) .....	180
8.5 Energy Industries (NFR 1A1) .....	182
8.5.1 Oil and Gas Refining and Solid Fuel Manufacturing (NFR 1A1b and 1A1c) .....	182
8.6 Manufacturing Industries and Construction (NFR 1A2) .....	183
8.7 Transport (NFR 1A3) .....	183
8.7.1 Domestic aviation (NFR 1A3a) .....	183
8.7.2 Road transportation (NFR 1A3b) .....	183
8.7.3 Rail (NFR 1A3c) .....	184
8.7.4 Navigation (NFR 1A3d) .....	184
8.7.5 Gas Transmission (NFR 1A3e) .....	184
8.8 Residential and Commercial/Institutional (NFR 1A4) .....	184
8.8.1 Commercial/Institutional (NFR 1A4a) .....	185
8.8.2 Residential (1A4b) .....	185
8.9 Combustion in Agriculture and Fishing (NFR 1A4c) .....	186
8.10 Cross cutting measures impacting key energy sectors .....	186
8.11 Fugitive emissions (NFR 1B) .....	186
8.12 Mineral Products (NFR 2A), Chemical industry (NFR 2B), Metal Industry (NFR 2C) .....	187
8.13 Non-Energy Products from Fuels and Non-Energy Products from Fuels and Solvent Use (NFR 2D) and NFR 2H, I, J, K, L .....	188
8.14 Agriculture (NFR 3) .....	188
8.15 Waste (NFR 5) .....	190

<b>Chapter Nine - Adjusted annual national emission inventories .....</b>	<b>191</b>
9.1 Meeting the Requirements for an Adjustment in the period 2020 to 2030 .....	191
9.1.1 Emission Reduction Commitments .....	191
9.1.2 NMVOC Adjustment Application .....	192
<b>Glossary.....</b>	<b>195</b>
<b>References.....</b>	<b>197</b>
<b>Annex A to H .....</b>	<b>203</b>
<b>Annex H - 2023 NECD Review findings and responses .....</b>	<b>204</b>



## List of Tables

Table 1.1 Key Data Providers and Information Covered by Memoranda of Understanding	13
Table 1.2. Key Category Analysis of Ireland's Air Pollutant Inventory 2023	15
Table 1.3 Emissions Uncertainties	16
Table 3.1. Fugitive Emissions from Coal mining and handling	57
Table 3.2. Fugitive Non-Methane Volatile Organic Compound Emissions	59
Table 3.3. Recalculations in Energy NFR 1A1	64
Table 3.3. Recalculations in Energy NFR 1A1 (continued)	65
Table 3.4. Recalculations in Energy NFR 1A2	66
Table 3.5. Recalculations in Energy NFR 1.A.3 (a & b)*	67
Table 3.6. Recalculations in Energy NFR 1A3 (c, d & e)	68
Table 3.7. Recalculations in Energy NFR 1A4	69
Table 4.12 Emission Time Series for Lead Production	79
Table 4.49 Notation keys used in 2023 NMVOC inventory	100
Table 4.50 Recalculations for Industrial Processes 1990–2023 (Heavy Metals)	102
Table 4.51 Recalculations for Industrial Processes 1990–2023 (Particulates and NO <sub>x</sub> )	103
Table 4.52 Recalculations for Industrial Processes 1990–2023 (NMVOC)	108
Table 5.1 Emission Estimates for Hexachlorobenzene Emissions from Pesticide Use	121
Table 5.2. NH <sub>3</sub> Recalculations for Agriculture 1990–2023	124
Table 5.3. NMVOC Recalculations for Agriculture 1990–2023	126
Table 5.4. TSP Recalculations for Agriculture 1990–2023	127
Table 5.5. PM <sub>10</sub> Recalculations for Agriculture 1990–2023	129
Table 5.6. PM <sub>2.5</sub> Recalculations for Agriculture 1990–2023	131
Table 6.1 Pollutant Emissions by Waste Source Category	133
Table 6.2 Emission Time Series for NMVOC TSP, PM <sub>10</sub> and PM <sub>2.5</sub> from Solid Waste Disposal on Land	134
Table 6.3 Emission Time Series for Mercury from Solid Waste Disposal on Land	134
Table 6.4 Time Series of Activity Data and Emissions of NH <sub>3</sub> and CO from Composting	135
Table 6.5 Time Series of Emissions from the Incineration of Clinical Waste	136
Table 6.6 Time Series of Emissions from the Incineration of Industrial (incl. Hazardous & Sludge) Waste	137
Table 6.7 Time Series of Emissions from Crematoria	139
Table 6.8 Time Series of Emissions from the Open Burning of Farm Plastics	140
Table 6.9 Time Series of activity data from Accidental Fires from Vehicles and Buildings	141
Table 6.10 Time Series of Emissions from Accidental Fires from Vehicles and Buildings	142
Table 6.11 Time Series of Emissions from Other Waste Burning	144
Table 6.12 Recalculations for Waste 1990–2023	146
Table 7.1. List of GNFR categories	151
Table 7.2. Correspondence list for GNFR and NFR categories	151
Table 8.1. Emission Reduction Commitments for 2020 and 2030 (expressed as a percentage reduction on 2005 levels)	168
Table 8.2. Projected Emissions of SO <sub>2</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)	172
Table 8.3. Projected Emissions of NO <sub>x</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)	174
Table 8.4. Projected Emissions of NH <sub>3</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)	176
Table 8.5. Projected Emissions of NMVOC under the With Existing Measures and With Additional Measures Scenarios (kt)	179
Table 8.6. Projected Emissions of PM <sub>2.5</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)	181
Table 9.1 Summary of National Emissions in 2005 and 2023, the ERCs for 2020-2030 and actual emission reduction	191

Table 9.2. Adjustments for emissions from the Food and Beverage Industry (NMVOC) -----	193
--	-----

## List of Figures

Figure 1.1. National Atmospheric Inventory System Overview -----	8
Figure 1.1. National Atmospheric Inventory System Overview -----	8
Figure 2.1 Emission Trend for Sulphur Dioxide 1990–2023 -----	19
Figure 2.2 Emission Trend for Nitrogen Oxides 1990–2023 -----	20
Figure 2.3 Emission Trend for Ammonia from agriculture 1990–2023 -----	22
Figure 2.4 Emission Trend for Non-Methane Volatile Organic Compounds 1990–2023 -----	23
Figure 2.5 Emission Trend for Carbon Monoxide 1990–2023 -----	24
Figure 2.6 Emission Trend for Particulate Matter <10 µm in Diameter 1990–2023 -----	25
Figure 2.7 Emission Trend for Particulate Matter <2.5 µm in Diameter 1990–2023 -----	26
Figure 2.8 Emission Trend for Total Suspended Particulates 1990–2023 -----	27
Figure 2.9 Emission Trend for Black Carbon 1990–2023 -----	28
Figure 2.10 Emission Trend for Lead 1990–2023 -----	29
Figure 2.11 Emission Trend for Cadmium 1990–2023 -----	30
Figure 2.12 Emission Trend for Mercury 1990–2023 -----	31
Figure 2.13 Emission Trend for Arsenic 1990–2023 -----	32
Figure 2.14 Emission Trend for Chromium 1990–2023 -----	33
Figure 2.15 Emission Trend for Copper 1990–2023 -----	34
Figure 2.16 Emission Trend for Nickel 1990–2023 -----	35
Figure 2.17 Emission Trend for Selenium 1990–2023 -----	36
Figure 2.18 Emission Trend for Zinc 1990–2023 -----	37
Figure 2.19 Emission Trend for Dioxins and Furans 1990–2023 -----	38
Figure 2.20 Emission Trend for Hexachlorobenzene 1990–2023 -----	39
Figure 2.21 Emission Trend for Polychlorinated Biphenyls 1990–2023 -----	40
Figure 2.22 Emission Trend for Polycyclic Aromatic Hydrocarbons 1990–2023 -----	41
Figure 3.1 Nitrogen Oxide Implied Emission Factors (IEF) for Category 1A1a -----	44
Figure 3.2 Sulphur Dioxide Implied Emission Factors (IEF) for Category 1A1a -----	45
Figure 3.3 Number of Domestic LTOs from Irish airports 1990–2023 -----	50
Figure 3.4 Number of International LTOs from Irish airports 1990–2023 -----	51
Figure 4.1 NMVOC Emission Trend for 2D-2H Other Solvent and Product 1990–2023 -----	82
Figure 5.1. Emission Trend for Ammonia from Agriculture 1990–2023 -----	111
Figure 5.2. Emission Trend for NMVOC from Agriculture 1990–2023 -----	112
Figure 5.3. Emission Trend for NO <sub>x</sub> from Agriculture 1990–2023 -----	112
Figure 5.4. Emission Trend for PM <sub>2.5</sub> from Agriculture 1990–2023 -----	113
Figure 7.1 General decision tree for emissions mapping (EMEP/EEA 2019) -----	155
Figure 7.2 Example of GeoKey preparation -----	157
Figure 7.3 National Total Emissions in 2015 for a) NO <sub>x</sub> , b) SO <sub>2</sub> , c) NMVOC, d) NH <sub>3</sub> , and e) PM <sub>2.5</sub> -----	166
Figure 8.1 SO <sub>2</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios -----	171
Figure 8.2 SO <sub>2</sub> Emission Projections for the With Additional Measures Scenario by Source Sector -----	171
Figure 8.3 NO <sub>x</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios -----	173
Figure 8.4 NO <sub>x</sub> Emission Projections for the With Additional Measures Scenario by Source Sector -----	173
Figure 8.5 NH <sub>3</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios -----	175
Figure 8.6 NH <sub>3</sub> Emission Projections for the With Additional Measures Scenario by Source Sector -----	176
Figure 8.7 NMVOC Emission Projections for the With Existing Measures and With Additional Measures Scenarios -----	177
Figure 8.8 NMVOC Emission Projections for the With Existing Measures and With Additional Measures Scenarios where an adjustment is applied for category 2H2 Spirit production -----	178
Figure 8.9 NMVOC Emission Projections for the With Additional Measures Scenario by Source Sector -----	179
Figure 8.10 PM <sub>2.5</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios --	180
Figure 8.11 PM <sub>2.5</sub> Emission Projections for the With Additional Measures Scenario by Source Sector -----	181

## **Acknowledgements**

*The authors wish to express their appreciation to the various Government Departments, the Sustainable Energy Authority of Ireland, the Central Statistics Office, the Electricity Supply Board and the many other data suppliers and individuals who have contributed to the subject matter of this inventory report.*

## EXECUTIVE SUMMARY

As a Party to the United Nations Economic Commission for Europe (UNECE) Convention on Long-Range Transboundary Air Pollution (CLRTAP), Ireland is required to annually report emission data for a wide range of air pollutants and other substances released into the atmosphere. The data are needed to support the work of the Convention in addressing well-known environmental problems such as urban pollution, acidification and tropospheric ozone formation arising from classic pollutants, such as nitrogen oxides (NO<sub>x</sub>), sulphur dioxide (SO<sub>2</sub>), non-methane volatile organic compounds (NMVOCs), ammonia (NH<sub>3</sub>), carbon monoxide (CO) and particulate matter (PM), and for the implementation of its Protocols on Heavy Metals and Persistent Organic Pollutants.

The UNECE revised 2014 Reporting Guidelines, Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution<sup>1</sup>, describe the scope and reporting of the emission inventories and projections under the Convention. They specify the methodologies and procedures to be followed for submitting consistent and comparable data on an annual basis in a timely, efficient, and transparent manner to meet the needs of the Convention. Under the Guidelines, Parties are strongly encouraged to submit an Informative Inventory Report (IIR) to support the evaluation of their up-to-date annual inventories and projections. The objective of the IIR is to describe the methodologies, input data, background information and the entire process of inventory compilation for transboundary air pollutant emissions and to give explanations for any improvements and recalculations of the inventories reported in previous submissions. The report is needed by expert review teams to assess the transparency, completeness and overall quality of the inventories as part of the review process of submissions from Parties to the Convention.

Member States of the European Union are required to report an Informative Inventory Report annually under Article 8(3) of Directive [\(EU\) 2016/2284](#). This Directive sets out emission reduction commitments of certain atmospheric pollutants and repeals Directive [2001/81/EC](#). This report to the European Union fulfils this reporting obligation.

The Environmental Protection Agency (EPA) in Ireland has overall responsibility for national air emission inventories and projections pursuant to the establishment of the National Atmospheric Inventory System (NAIS) in 2007. The EPA Office of Evidence and Assessment (OEA) performs the role of inventory agency in Ireland and undertakes all aspects of inventory preparation and management and is responsible for the submission of the results to CLRTAP. The present report constitutes Ireland's sixteenth IIR submitted under the Convention, covering annual inventories for the period 1990–2023. The report aims to provide a comprehensive description of the procedures, methodologies and activity data used for the compilation of Ireland's air emission inventories and projections as presented in Ireland's 2025 submission under CLRTAP and to the European Union under Directive (EU) 2016/2284. The report shows how Ireland follows the guidelines for estimating and reporting of emission data to ensure the transparency, accuracy, consistency, comparability and completeness (TACCC) of the reported emissions. In addition to complying with reporting requirements in this

---

<sup>1</sup> [Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution](#)

regard, the 2025 IIR is intended to inform the Government Departments and institutions involved, as well as other stakeholders in Ireland, of the level of emissions and the state of the art of Ireland's emission inventories and projections as they address the challenges to comply with commitments already established for air pollutants and to control emissions in general. An attempt has been made to give adequate descriptions of all methodological approaches and to provide pertinent information to facilitate the assessment of the emission estimates and the understanding of emission trends. The IIR is published on the web site of the EPA (<https://www.epa.ie/publications/monitoring--assessment/climate-change/air-emissions/>). It will be further developed for future submissions and updated annually in accordance with the UNECE Reporting Guidelines and Directive (EU) 2016/2284.

# Chapter One - Introduction

## 1.1 Convention on Long-Range Transboundary Air Pollution

The Convention on Long-Range Transboundary Air Pollution (CLRTAP) came into being in 1979 following the recognition that co-operation at international level was necessary to address environmental problems such as acidification associated with the transboundary transport and deposition of acidifying gases emitted into the atmosphere. The Convention was the first international legally binding instrument to deal with problems of air pollution on a broad regional basis. Besides laying down the general principles of international co-operation for air pollution abatement, the Convention sets up an institutional framework bringing together research and policy. The Executive Secretary of the United Nations Economic Commission for Europe (UNECE) acts as Secretariat to CLRTAP, and the Convention entered into force in 1983.

The aim of the Convention is that Parties shall endeavour to limit and, as far as possible, gradually reduce and prevent air pollution, including long-range transboundary air pollution. This objective is pursued under eight protocols that identify specific measures to be taken by Parties to cut their emissions of a wide range of air pollutants. The extent to which Parties to the Convention have ratified the various protocols varies. Of the eight protocols to date, Ireland has ratified the 1994 Oslo Protocol on Further Reduction of Sulphur Emissions, the 1988 Sofia Protocol concerning the Control of Emissions of Nitrogen Oxides, the 1998 Aarhus Protocol on Persistent Organic Pollutants (POPs) and the 1998 Aarhus Protocol on Heavy Metals and the 1999 Gothenburg Protocol.

## 1.2 Inventory Reporting and Review under the Convention on Long-Range Transboundary Air Pollution (CLRTAP)

The Guidelines for Reporting Emissions and Projections Data under the Convention on Long-Range Transboundary Air Pollution (ECE/EB.AIR/125)<sup>2</sup> specify the scope, methodologies, formats and deadlines for annual inventory submissions by Parties to the Convention. These Guidelines were adopted by the Executive Body in December 2013 (ECE/EB.AIR/122/Add.1, decisions 2013/3 and 2013/4) and published in 2014. They are a revised version of the 2009 Guidelines for Reporting Emission data under the Convention (ECE/EB.AIR/97), which were approved by the Executive Body in 2008 (ECE/EB.AIR/96, para. 83 (b)). While the Guidelines make it clear that Parties are required to report only on the substances and for the years set forth in the protocols that they have ratified and that have entered into force, Ireland endeavours to estimate and report emissions for the full range of substances set down in Annex I of the Guidelines. These substances are nitrogen oxides (NO<sub>x</sub>), sulphur oxides (SO<sub>x</sub>), non-methane volatile organic compounds (NMVOCs), carbon monoxide (CO), ammonia (NH<sub>3</sub>), particulate matter (PM), black carbon (BC), heavy metals (HM) and persistent organic pollutants (POPs).

The Guidelines state that an Informative Inventory Report (IIR) should be prepared for inclusion in the annual submission, and it is also required under Article 8(3) of Directive (EU) 2016/2284. The objective of the IIR is to describe the methodologies, input data, background information and the entire process

---

<sup>2</sup> The Guidelines for Reporting Emissions and Projections Data under the Convention on Long-Range Transboundary Air Pollution ([ECE/EB.AIR/125](#))

of inventory compilation for air pollutants, as well as any improvements and recalculations of the inventories reported in previous submissions. The report is needed to support the evaluation of emission trends and is used by expert review teams to assess the transparency, completeness and overall quality of the inventories as part of the review process established for submissions by the Parties to CLRTAP and the review process established under the NECD.

The present report constitutes Ireland's IIR for 2025, the sixteenth such report. It contains specific information on the national inventory for the years 1990–2023 as submitted to CLRTAP in 2025, including descriptions of methods, data sources, quality assurance/quality control (QA/QC) activities and trend analysis. The structure of the report follows the structure proposed in Annex II to the reporting guidelines.

The IIR focuses on the year 2023 and the status of the inventories achieved for the time series up to 2023 and is the basis for methodological description for the purposes of facilitating technical review and general assessment of Ireland's emission inventories. The IIR is designed to capture the cyclical nature of the reporting process and to clarify the chronology of changes and revisions that are part of normal inventory development. In this way, the report provides the basis for technical assessment and expert review of Ireland's air pollutant inventories. An attempt has been made to give adequate descriptions of all methodological approaches and to provide all the pertinent inventory information to facilitate the assessment of the emission estimates and the understanding of emission trends.

The IIR will be further developed and updated annually in accordance with the UNECE guidelines and is published on the web site of the EPA (<https://www.epa.ie/publications/monitoring--assessment/climate-change/air-emissions/>) Such updating is necessary to keep the UNECE Secretariat and other interested parties informed of the status of Ireland's air pollutant inventories and to document on-going improvements, recalculations and other developments affecting the estimates of emissions. Ireland is contributing to the Stage 3 in-depth review process for transboundary emission inventories by not only providing this IIR, but also providing expert reviewers to evaluate the submissions from other Parties to the Convention.

### 1.3 Inventory Reporting and Review under Directive (EU) 2016/2284

Directive (EU) 2016/2284<sup>3</sup> came into force on the 31<sup>st</sup> of December 2016. This inventory report and data submission fulfils Ireland's reporting requirements as set out in Article 10(2) of the directive. The inventory submission is fully consistent with the submission under the LRTAP Convention and will be subject to review in accordance with Article 10(3) of the directive.

For 2023 Ireland's national total emissions of Sulphur Dioxide (SO<sub>2</sub>), Nitrogen Oxides (NO<sub>x</sub>), Ammonia (NH<sub>3</sub>), Particulate matter <2.5µm (PM<sub>2.5</sub>) are in compliance, however non-methane volatile organic compounds (NMVOCs) are not in compliance with the emission reduction commitments shown in Annex II of Directive 2016/2284. Ireland has established adjusted annual national emission inventories for NMVOCs as allowed for under the flexibilities outlined in 5(1) of the Directive 2016/2284. The

---

<sup>3</sup> [DIRECTIVE \(EU\) 2016/2284 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC](#)



adjusted inventory is established in accordance with Part 4 of Annex IV of the directive and is explained in detail in chapter 9 of this report.

## 1.4 Inventory Reporting by the Environmental Protection Agency

Under Section 52 of the Environmental Protection Agency Act, 1992 (DOE, 1992), the Agency is required to establish and maintain databases of information on the environment and to disseminate such information to interested parties. Section 55 of the Act states that the Agency must provide, of its own volition or upon request, information and advice to Ministers of the Government in the performance of their duties. This includes making available such data and materials as are necessary to comply with Ireland's reporting obligations and commitments within the framework of international agreements. These requirements are the regulatory basis on which the EPA prepares annual inventories of air pollutants in Ireland. The activities related to the compilation and reporting of air pollutant emissions constitute one specific on-going project in the Agency's work programme.

The Department Environment, Climate and Communications (DECC) has designated the EPA as the agency with responsibility for the submission of emission data to international bodies, including the Secretariat for the CLRTAP and the European Union. The Agency's Office of Evidence and Assessment (OEA) currently compiles the national air pollutant emission inventories on behalf of the DECC for submission under the CLRTAP and the NECD.

### 1.4.1 National Atmospheric Inventory System

In 2005, UK consultants from the National Environmental Technology Centre (NETCEN) carried out a scoping study to identify the essential elements and structure of a national inventory system for Ireland to meet the needs of Decision 280/2004/EC (EP and CEU, 2004a) and to comply with obligations under Articles 5 and 7 of the Kyoto Protocol. The report (Thistlethwaite et al., 2005) describes how institutional arrangements among the EPA, its parent government department and other stakeholders may be reorganised, extended and legally consolidated across all participating institutions to strengthen inventory capacity within the EPA and to ensure that more formal and comprehensive mechanisms of data collection and processing are established for long-term implementation. The report sets out the extent of institutional participation, resource requirements and the form of legal arrangements necessary to perform the functions prescribed in the guidelines for national systems and enable Ireland to meet the objectives specified in those guidelines. The scoping study developed a QA/QC system as an integral part of the national system and the report made recommendations on internal inventory review and proposed a system to facilitate more efficient data management and reporting. Whilst developed to meet the needs of Decision 280/2004/EC and the Kyoto Protocol, Ireland's national system is also implemented to achieve emission inventories for transboundary gases for submission under the CLRTAP and the NECD.

The National Atmospheric Inventory System (NAIS) for Ireland was adopted by Government decision in April 2007. It establishes the necessary institutional, legal and procedural arrangements for the compilation of robust inventories of emissions of greenhouse gases (GHGs) and air pollutants to the atmosphere. It sets out formal procedures for the planning, preparation and management of the national atmospheric inventory and identifies clearly the roles and responsibilities of all the organisations involved in inventory compilation, reporting and review. A schematic overview of the national system is presented in Figure 1.1.

The principal objective of the NAIS is to ensure that Ireland can compile robust and verifiable annual inventories of emissions and report its emission estimates in accordance with relevant international obligations. The NAIS also facilitates the formal review of information submitted under international obligations, including the Paris Agreement, protocols under the CLRTAP and the NECD. Implementation of the national system ensures the transparency, consistency, comparability, completeness and accuracy of the national inventory in accordance with the established reporting guidelines, which incorporate methodological guidance and good practice.

Within the NAIS, the EPA's Office of Evidence and Assessment (OEA) is designated as the single national entity with overall responsibility for the national emission inventory in Ireland. The OEA also performs the role of inventory agency, i.e. it compiles the annual inventory and delivers Ireland's submissions to the various international organisations (European Commission (EC), European Environment Agency (EEA), United Nations Framework Convention on Climate Change (UNFCCC) and UNECE) in accordance with agreed deadlines and reporting formats. In addition to the primary data received from the Key Data Providers (KDPs), the inventory team obtains considerable supplementary information from other teams in the OEA, the Office of Environmental Sustainability and the Office of Environmental Enforcement within the EPA. These sources include Annual Environmental Reports (AERs) submitted by licensed facilities and the National Waste Database. The inventory team also draws on national research related to air pollutant emissions and special studies undertaken from time to time to acquire the information needed to improve the estimates for categories and gases. The approval of the completed annual inventory involves sign-off by the QA/QC manager and the inventory manager before it is transmitted to the Board of Directors of the EPA via the Programme Manager of OEA. Any issues arising from the Board of Directors' examination of the estimates are communicated to the inventory experts for resolution before final adoption of the inventory for submission and publication.

#### 1.4.2 Scope of Inventories under the LRTAP Convention and Directive (EU) 2016/2284


The scope of Ireland's emission inventories under the LRTAP Convention and Directive (EU) 2016/2284 is provided in this report. It covers a wide range of air pollutants and other substances, which are reported in a standard electronic format for a predefined nomenclature of source categories set down in the UNECE Reporting Guidelines. The air pollutants are referred to in seven groups as follows:

1. Main pollutants (NO<sub>x</sub>, NMVOCs, SO<sub>x</sub>, NH<sub>3</sub>, CO);
2. Particulate matter (PM<sub>2.5</sub>, PM<sub>10</sub>, total suspended particulates (TSP), black carbon (BC));
3. Priority heavy metals (lead (Pb), cadmium (Cd), mercury (Hg));
4. Other heavy metals (arsenic (As), chromium (Cr), copper (Cu), nickel (Ni), selenium (Se), zinc (Zn));
5. Annex I Persistent Organic Pollutants: the POPs listed in Annex I to the Protocol on POPs which are substances scheduled for elimination;
6. Annex II Persistent Organic Pollutants: the POPs listed in Annex II to the Protocol on POPs which are substances scheduled for restrictions on use;

7. Annex III Persistent Organic Pollutants: the POPs listed in Annex III to the Protocol on POPs which are substances referred to in Article 3, Para. 5 (a), of the Protocol. Polycyclic aromatic hydrocarbons (PAHs): for the purpose of the emission inventories, the following four indicator compounds should be used: benzo[b]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene and indeno [1,2,3-cd] pyrene. Hexachlorobenzene (HCB) is also included in Annex I to the Protocol as a substance for elimination.

The list of source categories for inventory purposes is known as the NFR (Nomenclature for Reporting). It comprises coded activities across all socio-economic sectors identified as sources of one or more of the substances listed above and provides for the inclusion of other activities that may be specific to individual countries. Many of the NFR categories are split into a varying number of subcategories, which are designed to reflect their importance as sources of one or more pollutants and to provide an adequate level of transparency. In the compilation of annual inventories, significant subdivision of the given NFR categories is normally applied for the process of calculating the relevant emissions. The NFR facilitates the comparison of emissions among reporting countries and the synthesis and assessment of submissions at the UNECE level. The current version of the NFR, NFR 19 is a revision of NFR14, which was included in Annex I of the Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution (ECE/EB.AIR/125), which were adopted by the Executive Body in December 2013. This revised NFR was approved by the Steering Body of the European Monitoring and Evaluation Programme (EMEP) during its 5<sup>th</sup> Joint Session in September 2019 and should be used for reporting under the CLRTAP for 2020 onwards.

The reporting format also includes a number of *Memo Item* entries. These items refer to sources of emissions whose contributions are not included in a Party's national total but which are to be reported because of their importance in relation to the overall assessment of emissions and for comparisons among Parties. The notable emission sources excluded from the reported national total for transboundary gases and included as Memo Items are emissions from international and domestic aviation during the cruise phase of a flight, and international shipping.



*Figure 1.1. National Atmospheric Inventory System Overview*

A set of notation keys has been adopted for use in completing the NFR templates to provide explanation and transparency where a numerical value does not appear for a particular pollutant and/or category combination. The notation keys are as follows:

- (a) NO (not occurring) for activities that do not occur in the country;
- (b) NE (not estimated) where emissions do occur but are not estimated, usually because they are considered negligible or the necessary data cannot be obtained;
- (c) NA (not applicable) for activities that do not generate emissions of a particular pollutant;
- (d) IE (included elsewhere) for emissions relating to a subcategory that are reported in another subcategory, usually at the next highest level;
- (e) C (confidential) for emissions that could lead to the disclosure of confidential information;
- (f) NR (not relevant) for emissions that are not required by the ratified protocols.

At four year intervals from 2017 onwards, the inventory submissions under the CLRTAP should include compilations of emissions for a list of defined large point sources and aggregated sectoral gridded data for the European Monitoring and Evaluation Programme (EMEP) grid cells overlying the national territory. This information is used in EMEP models for evaluating long-range transport of air pollutants and for assessing emission deposition relationships in Europe. Ireland's 2025 submission contains data for large point sources and sectoral gridded data of emissions on the EMEP grid that were submitted as required in 2025. This information is provided in chapter 7 of this report.

Parties to the Gothenburg Protocol shall report national projections every four years from 2015 onwards, for the years 2020, 2025 and 2030 and where available, also for 2040 and 2050, by 15th March for the pollutants: SO<sub>x</sub> (as SO<sub>2</sub>), NO<sub>x</sub>, NMVOCs, NH<sub>3</sub> and PM<sub>2.5</sub>, with voluntary reporting for black carbon. Other Parties are encouraged to provide projections for these pollutants. The Directive (EU) 2016/2284 requires emission projections every two years from 2017 onwards from European Member States. Ireland produces national air pollutant emission projections on an annual basis. Information on emission projections is provided in chapter 8 of this report.

### 1.4.3 Inventory Preparation

The air pollutant emission inventory database normally contains information on measured emission quantities, activity statistics (populations, fuel consumption, vehicle/kilometres of travel, industrial production and land areas), emission factors and the associated emission estimates for the NFR list of source categories. In practice, very few measured emission data are available for the range of gases covered and, consequently, the emissions from most activities are estimated by applying emission factors for each source/gas combination to appropriate activity data for the activity concerned. Virtually all emissions may be ultimately derived on the basis of the product of activity data and emission factor. Even in the case where emission estimates for particular categories are reported directly to the inventory agency they will normally have been derived in this manner.

The reporting guidelines provide the general guidance for the preparation and reporting of annual inventories by Parties. They incorporate the methodologies given in the EMEP/EEA (European Environment Agency) Emission Inventory Guidebook, hereafter referred to as the Inventory Guidebook. The inventory preparation process involves the acquisition of the required statistical data for the inventory year concerned and the application of emission factors that characterise the rate of

emission of the gases concerned. Some data analysis and preparatory calculations are generally needed to make available suitable combinations of activity data and emission factors at the level of disaggregation that gives the best estimate of emissions in the individual emission source categories. In the case of some source/gas combinations, it may be necessary to apply sophisticated models to generate the activity data, the emission factors or the emissions. The methods recommended by the Inventory Guidebook use a tiered system. This provides methodologies at different levels of detail and sophistication, which take account of these issues and other factors, such as data availability, technical expertise, inventory capacity and other circumstances, which may vary considerably across countries.

#### 1.4.4 Data Acquisition

In its capacity as the inventory agency, the OEA of the EPA acquires the principal items of activity data from identified Key Data Providers (KDP) relevant to each of the NFR sectors. Most KDPs provide data directly to the OEA, but some secondary KDPs provide their input to one of the primary KDPs for processing and incorporation into the information subsequently transmitted to the OEA. Some KDPs may also deliver estimates of emissions for their particular area of coverage or expertise. Table 1.1 lists the KDPs and the data they supply for use in transboundary air pollutant emission inventories.

The NAIS provides for a formal Memorandum of Understanding (MoU) between each KDP and the inventory agency regarding the scope, quality and submission date of the data to be provided for the purposes of the national emission inventory compilation. In the majority of cases, the data concerned are already routinely collected and published by the KDPs under existing mandates and established reporting programmes. Additional MoUs may be developed under the NAIS in cases where new or supplementary data sources need to be targeted. Under Section 69 of the EPA Act 1992, formal legal powers are assigned to the EPA, whereby the Agency may require any public body to provide information related to environmental quality and may make arrangements with other bodies for the provision of similar information. This provision can also be invoked by the OEA to acquire specific information for inventory purposes as the need arises.

The Emissions Trading Unit was established under the EPA Office of Licensing and Guidance (OLG) in late 2003 to implement Directive 2003/87/EC (EP and CEU, 2003) in Ireland. The Emissions Trading Unit currently forms part of the OES and is another key component of the national system. Information compiled for participants in the Emissions Trading Scheme (ETS) under Directive 2003/87/EC is an important source of activity-specific and company-specific data on GHG emissions for over 110 installations in Ireland. The inventories for transboundary air pollutants draw on relevant information regarding fuel quantities and fuel properties available under the ETS for these installations and fuel data are used for reconciliation with the national energy balance for major categories and matching of activity data for GHG emission inventories. The inventory agency in the OEA obtains useful support and activity data from other EPA offices and programme areas, including the Environmental Licensing Programme, Office of Environmental Enforcement, and the Environmental Research Programme. These programmes and offices make various contributions that are used to determine or substantiate the activity data or emission factors for particular categories or individual activities, which ensures that country-specific information is exploited to the maximum extent possible. In all cases, consistency is maintained with data application for GHG emission inventories and vice versa.

### 1.4.5 Quality Assurance and Quality Control (QA/QC)

**Quality Assurance (QA)** activities include a planned system of review procedures conducted by personnel not directly involved in the inventory compilation and development process. Reviews, preferably by independent third parties, should be performed upon a finalised inventory following the implementation of QC procedures. Reviews verify that data quality objectives were met, ensure that the inventory represents the best possible estimates of emissions, given the current state of scientific knowledge and data available, and support the effectiveness of the QC programme.

A QA/QC spreadsheet tool has been developed to manage and maintain Ireland's QA/QC system. This tool consists of several spreadsheets that provide procedures, guidance, forms and templates as required for the general QA/QC functions. The supporting manual (Thistlethwaite et al., 2005) provides a general overview to the QA/QC system and guidance on the application of the plan and procedures. The QA/QC plan identifies the specific data quality objectives related to the principles of transparency, accuracy, consistency, completeness, and comparability (TACCC) required for Ireland's national inventory and provides specific guidance and documentation forms and templates for the practical implementation of QA/QC procedures. The spreadsheets include a brief introduction and a statement of the data quality objectives (DQOs) and how they will be met through the QA/QC system with reference to the relevant spreadsheet tool template sheets and forms. The Introduction sheet links to the QA/QC plan which provides the schedules and procedures for the QA/QC system and lists all of the QA/QC activities that exist or are planned to make up Ireland's QA/QC system. The Plan sheet consists of tables that contain three different categories of QA/QC activity:

1. General activities covering the planning and management practices and procedures;
2. Activities that should be undertaken on an annual basis for management and preparation of the inventory;
3. Periodic activities that should be undertaken in response to specific events in the inventory and for periodic peer review or verification.

The inventory agency has implemented this approach to QA/QC for nineteen annual reporting cycles. This involved the allocation of responsibilities linked to the national system and the use of the template spreadsheet system to record the establishment and maintenance of general inventory checking and management activities covering the overall compilation process, as well as the undertaking of specific annual activities and any necessary periodic activities in response to specific events or outcomes in inventory reporting and review. The system facilitates record keeping related to the chain of activities from data capture, through emission calculations and checking, to archiving and the identification of improvements.

Ireland's calculation spreadsheets in all sectors are structured on a time-series basis. This organisation is designed to facilitate the QA/QC process as well as more efficient trend analysis and to ensure ease of transfer of the outputs to the NFR tables. The inventory compilation is directly linked to the primary statistical inputs, which facilitates rapid year-on-year extension of the time series and efficient updating and recalculation, where appropriate, in the annual reporting cycle. Internal aggregation to various levels corresponding to the NFR tables provides immediate and complete checks of the results.



**Quality Control (QC)** is a system of routine technical activities to measure and control the quality of the inventory as it is being developed. The QC system is designed to:

- (i) Provide routine and consistent checks to ensure data integrity, correctness, and completeness;
- (ii) Identify and address errors and omissions;
- (iii) Document and archive inventory material and record all QC activities.

Quality control activities include general methods such as accuracy checks on data acquisition and calculations and the use of approved standardised procedures for emission calculations, measurements, estimating uncertainties, archiving information and reporting. Higher-tier QC activities include technical reviews of source categories, activity and emission factor data, and methods.

The inventory agency has created and implemented a number of QA\QC tools. These spreadsheet-based tools are used to better inform the QA\QC process whilst providing transparent descriptions of the outcomes of checks. In each spreadsheet-based tool comments are added to explain anything highlighted by the checking process. The text from these tools (such as the recalculations assessment) then helps inform the update of the IIR. The tools used include:

1. Recalculations assessment – this spreadsheet tool calculates the percentage change of emission estimates between the current and previous inventory submissions using conditional formatting to highlight the significant changes. There is a separate table for each pollutant in which the entire time series is evaluated for all NFR codes. This check highlights the recalculations that have been made in the inventory so that they can be verified and justified by the inventory agency.
2. Trend assessment – this spreadsheet tool calculates the percentage change between the most recent year and the preceding year of the current inventory submission using conditional formatting to highlight the significant changes. There is a separate table for each pollutant in which the entire time series is evaluated for all NFR codes. This check helps identify any time series inconsistency with the newly reported data of the most recent year. Provided this is run annually and alongside the recalculations check, time series consistency should be maintained.
3. Pollutant specific assessments – there are two tools that check the following rules that should be maintained in an inventory:  $TSP \geq PM_{10} \geq PM_{2.5}$ ; Total PAHs (1-4) =  $B(a)p + B(b)F + B(k)P + I(123)-cd$ . These simple checks help maintain the accuracy of the inventory.
4. Data value assessment – two tools check the entire time series for all pollutants to ensure that none of the following are reported: zero values, errors, negative values. These simple checks help maintain the accuracy and transparency of the inventory.
5. Annex I reporting template assessment – this tool evaluates whether all compiled Annex I reporting template files are comparable in structure and content to the template. This helps maintain the comparability of the inventory.
6. Notation keys assessment – this tool summarises the use of the different notation keys within the inventory. This tool has been implemented for all NEC pollutants across the entire time series. The tool helps the inventory agency evaluate, justify and document the use of notation



keys in the inventory. By increasing the accuracy of the notation keys, the transparency of the inventory is improved and this allows the inventory agency to clearly identify areas where potential improvements could be made (e.g. the use of IE or NE).

The online tool provided by the Centre on Emissions Inventories and Projections (CEIP), RepDab, is used by the inventory agency to check the format, completeness and internal consistency of the Annex I reporting template submission files. Further details regarding the checks that are carried out can be found on the [CEIP](#) website.

In the 2025 reporting cycle, the inventory agency updated some of the default emission factors in accordance with the EMEP/EEA Air Pollutant Emission Inventory Guidebook (2023) and implemented suggested changes/requests for further information as presented in the Final NECD Review Report of the 2024 Comprehensive Technical Review of National Emission Inventories. These are outlined in the sectoral chapters of this report.

**Table 1.1 Key Data Providers and Information Covered by Memoranda of Understanding**

Key Data Provider	Data Supplied	Deadline	Sector in which Data are Used
Sustainable Energy Authority of Ireland (SEAI)	National Energy Balance; Detailed national energy consumption disaggregated by economic sector and fuel	30 September	Energy, Waste
Department of Agriculture Food and the Marine (DAFM)	Nitrogen fertiliser sales, cattle populations from the AIM (Animal Identification and Movement) database, sheep statistics, poultry statistics	30 September	Agriculture
Central Statistics Office (CSO)	Annual population, livestock populations, crop statistics, housing survey data	30 September	Agriculture, Industrial Processes, Waste
Gas Networks Ireland (GNI)	Analysis results for indigenous and imported natural gas	30 September	Energy
Marine Institute	Annual report on discharges, spills and emissions from offshore gas production installations	30 October	Energy
Emissions Trading Unit (OES, EPA)	Verified CO <sub>2</sub> estimates and related fuel and production data for installations covered by the EU ETS	30 April	Energy, Industrial Processes
Department of Environment, Climate and Communications (DECC)	National Oil Balance (as a component of the energy balance)	30 September	Energy
Road Safety Authority (RSA)*	Road transport statistics from the National Car Testing (NCT) Service	30 April	Energy

\*These bodies have MoUs with the SEAI rather than with the OES.

## 1.4.6 Inventory Compilation

The source data, calculation workbooks and outputs for all emissions to air are held on EPA servers. The annual inventory compilation for transboundary gases is undertaken in separate *Data Processing* folders for each sector, which are linked to the *Source Data* folders for the respective sectors at the same level. The *Outputs* folder and the *QA/QC* folder are also at this level. The *Outputs* folder contains the files used for the official submissions to the EU and the UNECE and for preparing summary reports

and relevant media statements at national level. All calculation workbooks for the individual sectors contain a QA/QC worksheet, which are compiled collectively in the QA/QC folder. Data processing to compute the emission estimates is carried out at the most detailed level of aggregation possible, consistent with data availability and the outputs needed to populate the reporting template format for the category concerned. These outputs are primarily the estimates of emissions and the corresponding activity data for each category.

Quality control procedures are an integral part of the inventory preparation and reporting cycle. Within the inventory team, quality control for each sector is undertaken by an inventory compiler who has not produced the emission estimates for that sector. Quality control involves a series of checks covering the data inputs and any necessary pre-processing, the calculation of emissions, and the generation of the output records that are subsequently compiled in the NFR templates. The checks cover such items such as the comparison of inputs with those of previous years, the identification of errors and omissions, validating internal linking and calculation algorithms, replicating the aggregation of subcategories and ensuring an adequate level of completeness in NFR files to achieve transparency for external review purposes. A colour code system is used to distinguish between such elements as data taken from another spread sheet, calculated values, extrapolated or interpolated values, outputs for the NFR, and checks and annotations.

## 1.5 Key Category Analysis

Key category analysis for transboundary air emissions is explained in the Inventory Guidebook (EMEP/EEA, 2023), and is the same concept as that presented in the Intergovernmental Panel on Climate Change (IPCC) 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006). This defines a key emission category as one that is prioritised within the national inventory system because its emission estimate has a significant influence on the Party's total inventory in terms of the absolute level of emissions, the trend in emissions, or both. The Inventory Guidebook provides several methods for undertaking the analysis of key categories that can be applied at any appropriate level of source aggregation, depending on the information available. The simplest approach (Approach 1) – identification based on contribution to emission level – is used here for the inventories of all substances to highlight which sources of emissions are the most important in Ireland.

In level assessment, key categories are those categories whose combined contribution to the total emission level, determined from the ranking of all categories on the basis of their individual contributions to the level, is 80 per cent. Information about key categories is considered to be crucial to the choice of methodology for individual sources and to the management and reduction of overall inventory uncertainty. The identification of such categories is recommended in order that inventory agencies can give them priority in the preparation of annual inventories, especially in cases where resources may be limited. Information on key categories is clearly also vital for the development of policies and measures for emissions reduction.

It is well established that fuel combustion in a small number of economic sectors is the major source of most air pollutants. This is true for classic pollutants such as SO<sub>x</sub> and NO<sub>x</sub>, which are reasonably well quantified in emission inventories and for which emissions have decreased considerably, and also for other substances (PM, POPs) for which inventories have much higher levels of uncertainty. The relative

contributions of key categories are clearly shown by the results of the simple key category analysis shown in Table 1.2 and Annex A.2, Tables 1-15, which summarises NFR Level 2 key categories by pollutant. The dominance of categories under 1A (Energy: Combustion) highlights the importance of combustion sources. The four key categories (1A1a, 1A2f, 1A3b and 1A4b) dominate the results of the key category analysis presented in Table 1.2. Agriculture sources (3B, 3D) are the main sources of emissions for NH<sub>3</sub>, NMVOC's, particulate matters, HCB and the additional categories 2D and 2H for NMVOCs. Waste sector sources are main driver of emissions for As and PCBs and an important source for Hg, Cr and Dioxins.

**Table 1.2. Key Category Analysis of Ireland's Air Pollutant Inventory 2023**

Pollutant	Key categories										Total (%)
NOx	3Da3 12.81%	3Da1 12.57%	1A3bi 11.27%	1A3bii 9.26%	1A3biii 8.52%	3Da2a 7.84%	1A3dii 5.76%	1A4bi 5.40%	1A2f 5.04%	1A1a 5.04%	83.50%
CO	1A4bi 58.88%	1A3bi 14.96%	1A1a 8.89%								82.73%
NMVOC	2H2 27.31%	3B1b 20.28%	2D3a 11.13%	3B1a 10.63%	1A4bi 5.84%	3De 4.12%	2D3i 2.11%				81.43%
SOx	1A4bi 64.90%	1A2f 11.92%	1A1a 9.65%								86.47%
NH3	3Da2a 26.97%	3B1b 25.97%	3B1a 13.44%	3Da1 13.25%	3Da3 10.84%						90.46%
TSP	2D3b 60.43%	1A4bi 9.43%	2A5b 6.06%	1A3bvi 3.40%	3B3 2.55%						81.87%
PM10	1A4bi 24.15%	2D3b 22.37%	1A3bvi 7.37%	3Dc 5.93%	2A5b 5.04%	2A1 4.41%	1A2f 4.15%	3B1b 3.45%	1A3bvii 3.31%		80.17%
PM2.5	1A4bi 45.95%	1A3bvi 7.35%	1A2f 7.29%	2A1 4.68%	1A2gviii 4.36%	3B1b 4.31%	1A3bvii 3.41%	3B1a 2.99%			80.34%
Pb	1A3bvi 69.42%	1A4bi 14.31%									83.73%
Cd	1A2gviii 26.27%	2D3i 16.65%	1A1a 12.48%	1A3bvi 10.24%	1A4bi 9.21%	2G 7.65%					82.51%
Hg	1A4bi 20.59%	1A1a 17.38%	1A2f 16.44%	5A 9.29%	5C1bv 6.54%	1A3bi 5.39%	1A2b 4.88%				80.50%
As	5C1bi 66.10%	1A1a 15.80%									81.90%
Cr	1A3bvi 53.72%	5C1bi 19.89%	1A2f 4.83%	2D3i 4.38%							82.82%
Cu	1A3bvi 83.82%										83.82%
Ni	1A4ai 28.20%	1A2gviii 15.27%	1A1a 13.21%	1A2f 13.04%	1A2e 6.53%	1A3bvi 5.87%					82.13%
Se	1A4bi 67.53%	1A1a 22.02%									89.55%
Zn	1A3bvi 53.06%	2D3i 13.70%	1A4bi 9.67%	1A2gviii 8.59%							85.02%
PCDD/F	1A4bi 68.69%	5E 19.83%									88.52%
PCBs	5E 53.46%	1A4bi 33.29%									86.74%
HCB	3Df 86.13%										86.13%
PAHs	1A4bi 90.24%										90.24%
	1 Energy		2 IPPU		3 Agriculture		5 Waste				

## 1.6 Uncertainty Assessment

Undertaking a quantitative estimate of emissions uncertainty requires a substantial amount of detailed data on the uncertainty of both activity data and emissions factors for a diverse range of source types. It has not been possible to collect these data in full. However, it has been possible to characterise the uncertainties associated with sources in a more approximate way.

A semi-quantitative uncertainty analysis has been used to determine the overall emissions uncertainty for a number of pollutants for 2023 data. This uses a Tier 1 propagation of errors to obtain an uncertainty for the total emission. However, the uncertainty assigned to the activity data and emission factor for each individual source is obtained from a combination of expert judgement and ranges of uncertainty obtained from the EMEP/EEA emission inventory guidebook. The results provide a good indication as to which sources are contributing the most to the overall uncertainty, and therefore where improvement effort should be targeted.

The methodology and results of the Tier 1 uncertainty analysis are presented in detail in Annex G, tables G.1 to G.6. The results can be summarised as follows:

**Table 1.3 Emissions Uncertainties**

Pollutant	Emission (kilotonnes, 2023)	Uncertainty in 2023 (%)	Trend Uncertainty 1990-2023 (%)
NO <sub>x</sub>	85.87	49.0	14.6
SO <sub>2</sub>	7.36	14.2	0.5
NM VOC	105.69	72.9	18.0
NH <sub>3</sub>	116.36	82.3	8.0
CO	88.97	41.8	1.4
PM <sub>2.5</sub>	9.31	85.3	15.4

The total uncertainty in the NO<sub>x</sub> emission in 2023 is dominated by the contribution of inorganic N-fertilisers (3.D.a.1), animal manure applied to soils (3.D.a.2.a) and urine and dung deposited by grazing animals (3.D.a.3), ±200 per cent for each subcategory of 3.D. These three categories combined account for 98.4 per cent of the total NO<sub>x</sub> emissions uncertainty. The next largest contributor to uncertainty for NO<sub>x</sub> emissions is landfill gas used as a fuel in Public electricity and Heat Production with an emission factor uncertainty of over 300 per cent.

Emissions of SO<sub>2</sub> are well characterised when compared to NO<sub>x</sub> for most emissions because they are combustion related, with emission factors for SO<sub>2</sub> (i.e. the sulphur content of the fuel) more readily determined than for NO<sub>x</sub>. Solid fuel combustion, coal, biomass and peat in the residential sector combined account for 94.5 per cent of the overall uncertainty.

Emissions of NM VOC from non-combustion sources are typically high in uncertainty because they are difficult to characterise by measurement. Manure management from cattle (3.B.1.a & 3.B.1.b) contributes to 88.8 per cent of the overall uncertainty because of the magnitude of the emission of these categories (30.9 per cent in 2023) and, as with all manure management categories, has poorly characterised emission factors (±300 per cent). Fugitive NM VOC losses from fuel extraction and

distribution (1.B.2) and domestic solvent use (2.D.3.a) are large contributors because they have poorly characterised emission factors and activity data, respectively.

The uncertainty associated in  $\text{NH}_3$  emissions are driven by the emission factors, with the activity data; number of livestock and nitrogen amounts, typically being well characterised by comparison. The sources making the largest contributions to the overall uncertainty are ammonia losses from inorganic N-fertiliser use (3.D.a.1), animal manure applied to soils (3.D.a.2.a) and urine and dung deposited by grazing animals (3.D.a.3). The emission factors for these sources are currently assigned an uncertainty of  $\pm 200$  per cent, and they contribute to 96.3 per cent of the overall uncertainty.

Emissions of  $\text{PM}_{2.5}$  are generally high in uncertainty because many combustion sources are either not well characterised, or are variable in emission, with small changes to combustion conditions having very large impacts on  $\text{PM}_{2.5}$  emissions. Residential coal, biomass and peat combustion (1.A.4.b) and road paving with asphalt (2.D.3.b) are the largest contributors to the overall uncertainty, contributing 97.3 per cent and 1.6 per cent respectively. These are relatively large sources with very uncertain emission factors.

It is interesting to note that electricity generation and industrial combustion do not feature as major contributors to the overall uncertainty for any pollutants. This is because the use of point specific data allows the emission estimates to be particularly well characterised.

## Chapter Two - Analysis of Key Emission Trends

### 2.1 Introduction

Ireland's 2025 submission under the CLRTAP and the Directive (EU) 2016/2284 includes emission estimates for the period 1990–2023 in respect of all substances listed in Section 1.4.2 above. The primary emission time series is prepared on the basis of Ireland's published national energy balances, which record the amounts of fuels sold in the country. In recognition of the significant cross-border movement that occurs with respect to automotive fuels in some parts of Europe, the reporting guidelines allow for the reporting of emissions from road transport on the basis of fuels used within the country. This may result in a substantial decrease in the national total emissions for some pollutants and the adjusted total is considered more appropriate for the assessment of performance in relation to certain protocols. This issue is relevant to Ireland in the case of the Sofia Protocol on NO<sub>x</sub> emissions and, to facilitate the assessment, Ireland has also submitted inventories in which the estimates for road transport are based on fuels used in the country. Emission inventories based on fuel sold and fuel used are provided in this submission for all pollutants for the period 1990–2023 and for the year 1987, the base year for the Sofia Protocol.

This chapter provides an overview of the emission trends for the period 1990–2023 for all substances included in Ireland's 2025 submission under the CLRTAP and the NECD. The general analysis of trends is performed only in respect of emissions estimated on the basis of fuels used in Ireland.

### 2.2 Main Pollutants

#### 2.2.1 Sulphur Dioxide (SO<sub>2</sub>)


Total sulphur dioxide emissions decreased by 96.0 per cent, from 184.78 kt in 1990 to 7.36 kt in 2023 (Figure 2.1). The Commercial/Institutional and Residential (1A4a and 1A4b) sectors combined account for 66.7 per cent of the total in 2023 and decreased by 87.1 per cent between 1990 (38.15 kt) and 2023 (4.91 kt). The Public Electricity and Heat Production (1A1a) sector contributed 9.6 per cent of the total in 2023 and decreased by 99.3 per cent between 1990 (103.04 kt) and 2023 (0.71 kt). Consumption of coal, peat and oil in this sector has generally declined to date. Emissions from Public Electricity and Heat Production decreased in 2023 by 61.7 per cent compared to 2022 due to reduced emissions from coal (64.8 per cent), oil (84.9 per cent) and peat (27.7 per cent).

Combustion sources in the Manufacturing Industries and Construction (1A2) sector largely account for the remainder of the emissions, with contribution of 18.7 per cent in 2023. Emissions in this sector peaked in 1994 due to an increase in SO<sub>2</sub> from oil combustion in an installation which is the main contributor to emissions in Non-ferrous metals (1A2b). Emissions in Manufacturing Industries and Construction (1A2) have decreased in the 1990–2023 time series by 96.0 per cent and are the lowest ever recorded since 1990. Combustion in Agriculture/Forestry/Fishing (1A4c) sector accounts for 0.4 per cent and Transport (1A3) combustion sources account for 3.3 per cent of national total emissions of SO<sub>2</sub> in 2023. The remainder of the SO<sub>2</sub> emissions arise from combustion sources in the Petroleum Refining (1A1b) and Manufacture of Solid Fuels and Other Energy Industries (1A1c) sectors, Other Product Use (2G) Waste Incineration (5C1), which combined account for 1.3 per cent of the total in 2023 and are presented in Other NFR (Nomenclature for Reporting) sectors in Figure 2.1. Overall SO<sub>2</sub> emissions decreased by 21.3 per cent between 2022 and 2023 as a result of reductions in the

combustion of fossil fuels in the energy generation, residential and commercial sectors. In 2023, the combustion of coal and oil for public electricity and heat generation reduced by 44.2 per cent and 78.2 per cent, respectively, compared to 2022. Reduction in the use of coal and home heating oil in 2023 were driven by fewer heating degree days in 2023 than in 2022 as well as increased retrofit activity and heat pump installation in households.

*Figure 2.1 Emission Trend for Sulphur Dioxide 1990–2023*

## 2.2.2 Nitrogen Oxides (NO<sub>x</sub>)




Total nitrogen oxide emissions have decreased by 51.6 per cent, from 174.28 kt in 1990 to 84.42 kt in 2023 (Figure 2.2). Road Transport (1A3b) is a major source of NO<sub>x</sub> emissions, contributing 27.8 per cent (and 23.50 kt) of the total in 2023, with the transport sector as a whole accounting for 38.1 per cent (and 32.18 kt) of the national total. The Manufacturing Industries and Construction (1A2) sector accounts for an increasing percentage of the national total. The contribution of the sector in 1990 to the national total was 5.2 per cent (9.08 kt), which increased to 11.2 per cent share in 2007 (and 17.33 kt) as a result of the increases in cement production for construction during the economic boom in Ireland over the previous decade, then reduced to 7.3 per cent share (7.53 kt) of the national total in 2011 due to the economic crisis impacting upon the sector. In 2023 the sector contribution increased to an 9.7 per cent share (and 8.15 kt) of the national total largely driven by an increase in cement production.

The Public Electricity and Heat Production (1A1a) sector is another key source of NO<sub>x</sub> emissions, accounting for 5.1 per cent of emissions in 2023. The application of low-NO<sub>x</sub> burner technology in several major power stations, in addition to increasing use of natural gas, have reduced NO<sub>x</sub> emissions from electricity generation by 90.7 per cent between 1990 (46.37 kt) and 2023 (4.33 kt), even though electricity total final consumption has increased by 166.7 per cent over the same period. Emissions from Public Electricity and Heat Production decreased by 42.1 per cent in 2023 from 2022. This reflects the considerable reduction of coal, oil and peat use in this sector due to the improvement in efficiency of modern gas fired power plants replacing older peat and oil-fired plants and the increased share of renewables, primarily wind power, and increased interconnection. Commercial/Institutional and Residential (1A4a and 1A4b) sectors combined account for 7.8 per cent of the total and combustion sources in Agriculture/Forestry/Fishing (1A4c) sector account for 4.0 per cent in 2023. The remainder

of the combustion sources of NO<sub>x</sub> arise in the Petroleum Refining (1A1b) and Manufacture of Solid Fuels and Other Energy Industries (1A1c) sectors, as well as combustion sources in Other product Use (2G) and the Waste sector (5C1), which are presented in Other NFR sectors that together account for 0.4 per cent of the total in 2023. Agricultural sources of NO<sub>x</sub>, accounted for 34.9 per cent of emissions in 2023 (29.44 kt), having decreased by 11.8 per cent since 1990 (33.38 kt).

The largest sources of NO<sub>x</sub> emissions within agriculture are associated with Inorganic nitrogen fertilizer application (3Da1) and Urine and Dung Deposited by grazing animals (3Da3).

The reductions in NO<sub>x</sub> emissions arising from the use of catalytic converters in cars and heavy-duty vehicles have only become apparent in recent years, as the technology has been offset by large increases in vehicle numbers in the past decade. This effect is exaggerated in later years by ‘fuel tourism’, whereby a proportion of the automotive fuel sold in Ireland, the basis for the emission time series given in Figure 2.2, is used by vehicles in other countries. The estimated level of fuel tourism is provided in Annex A.3, together with the adjusted annual NO<sub>x</sub> emissions based on fuels used in Ireland and facilitates assessment of obligations in relation to the Sofia Protocol on NO<sub>x</sub> emissions.



**Figure 2.2 Emission Trend for Nitrogen Oxides 1990–2023**

### 2.2.3 Ammonia (NH<sub>3</sub>)

National total emissions have decreased by 9.8 per cent, from 128.98 kt in 1990 to 116.36 kt in 2023. Agriculture accounted for 99.3 per cent of total ammonia emissions in 2023. Livestock production has historically accounted for the bulk of national total ammonia emissions in Ireland and, in 2023, Manure Management (3B) and Animal manure applied to soil (3Da2a) and nitrogen from urine and dung deposition by grazing animals combined accounted for 86.1 per cent of the national total. In 2023, Manure Management at 55.78 kt and 47.9 per cent share of the total in 2023 showed a decrease of 1.9 per cent from 56.84 kt in 1990. Organic fertiliser use covers emissions from two sectors: Animal Manure applied to soils (3Da2a) and Sewage Sludge applied to soils (3Da2b), which combined represent 31.74 kt and account for 27.3 per cent of the total ammonia emissions in 2023 (indicating




an 19.8 per cent decrease from 39.55 kt in 1990). Urine and Dung deposited by grazing animals (3Da3) accounted for 10.8 per cent of total emissions in 2023 (12.61 kt), having increased by 4.0 per cent since 1990. Inorganic/synthetic N-fertilizers applied to soils (sector 3Da1) decreased by 22.6 per cent from 1990 (19.91 kt) and at 15.41 kt in 2023 accounted for 13.2 per cent of the national total. The small contribution by Transport (1A3) sources peaked in 2005, the main driver of which was the increased use of cars with early generation three-way exhaust catalysts in Road Transport (1A3b). Transport emissions have increased from 0.04 kt in 1990 to 0.52 kt (and 0.4 per cent share of the total) in 2023. The remainder of the ammonia emissions arise from Commercial/Institutional and Residential (1A4a and 1A4b) sectors combined (0.1 per cent share) and Other NFR sectors (Combustion in Manufacturing Industries and Construction (1A2), Combustion in off road Agricultural machinery (1A4cii), Other Product Use (2G), Biological Treatment of Waste (5B1)) that together account for 0.2 per cent of the total in 2023.

Within livestock production, Manure Management (3B) at 55.78 kt in 2023 is the largest source of  $\text{NH}_3$ . In Ireland, approximately two-thirds of animal manure is excreted at pasture annually, reflecting the relatively short period that the main livestock categories (cattle and sheep) are housed.

Dairy Cattle (3B1a) and Non-Dairy Cattle (3B1b) account were the largest sources (39.7 per cent) of Agriculture sector ammonia emissions in 2023. Other livestock, which includes Sheep (3B2), Swine (3B3), Goats (3B4d), Horses (3B4e), Mules and asses (3B4f), Poultry (3B4g) and Other animals (3B4h), combined account for 8.6 per cent of total ammonia emissions from agriculture in 2023 (Figure 2.3).

Consequently, the trend in  $\text{NH}_3$  emissions is largely determined by cattle population and steadily increased up to 1998 to 139.98 kt, the highest emissions across the timeseries. Populations of cattle and sheep have declined after 1998, in combination with decreases in fertiliser use, which contributed to a decrease in  $\text{NH}_3$  emissions in the period 2000 to 2012. Subsequently, increases in cattle numbers and fertiliser use have led to increases in  $\text{NH}_3$  emissions, largely due to the removal of milk quotas in 2015. In 2023,  $\text{NH}_3$  emissions have dropped below 2015 levels for the first time since the abolition of milk quotas.  $\text{NH}_3$  emissions then increased up to 2018 to 130.43 kt in response to significant drought conditions in 2018 and increased nitrogen fertiliser and concentrate feeding to bovines.

In recent years, the increased use of abatement options, such as inhibited urea fertiliser and low emission slurry spreading techniques, are leading to emission reductions. Emissions in 2023 are 4.1 per cent below those in 2022. This is largely due to increased use of low emission slurry spreading techniques for cattle slurry, the proportion of which was applied by these techniques increased from 59.0 per cent in 2022 to 64.0 per cent in 2023 and a reduction in nitrogen fertiliser use of 18.2 per cent between 2022 and 2023.




**Figure 2.3 Emission Trend for Ammonia from agriculture 1990–2023**

## 2.2.4 Non-Methane Volatile Organic Compounds (NMVOCs)

Total non-methane volatile organic compound emissions have decreased by 34.7 per cent, from 161.80 kt in 1990 to 105.69 kt in 2023 (Figure 2.4). The NMVOC emissions are determined largely by the Agriculture sectors: 3B Manure Management and 3Da1 Inorganic N-fertilisers and emissions from solvents and other product use and the food and beverages industry. Agriculture categories combined accounted for 39.8 per cent of the national total and showed a decrease of 13.9 per cent between 1990 (48.89 kt) and 2023 (42.09 kt). The combined solvents use (2D and 2G) and fugitive emissions from the oil sector (1.B.2.a) produced 19.2 per cent of the 2023 total of NMVOC emissions in Ireland, having decreased by 26.7 per cent between 1990 (27.65 kt) and 2023 (20.27 kt). The Food and Beverage Industry (2H2) contributed to 27.7 per cent of total emissions in 2023, having increased by 204.6 per cent from 9.62 kt in 1990 to 30.95 kt in 2023.

Combustion sources in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors are also important sources, accounting for 6.4 per cent of national total NMVOC emissions in 2023, a reduction of 81.0 per cent between 1990 (35.53 kt) and 2023 (6.74 kt). Technological controls for volatile organic compounds (VOCs) emitted by motor vehicles have been more successful than in the case of NO<sub>x</sub> and have contributed to a substantial reduction in emissions from Road Transport (1A3b), with the total transport sector's contribution having decreased by 91.3 per cent between 1990 (35.83 kt) and 2023 (3.11 kt), equating to 22.1 per cent of the national total in 1990, falling to 2.9 per cent in 2023. A total of five NFR source categories make up the source Other NFR sectors (Figure 2.4) contributing 0.8 per cent of national total NMVOC emissions in 2023. The largest sources of emissions within this categorisation is Solid waste disposal on land sector (5A) and Fugitive emissions from natural gas exploration (1B2b).




**Figure 2.4 Emission Trend for Non-Methane Volatile Organic Compounds 1990–2023**

### 2.2.5 Carbon Monoxide (CO)

Carbon monoxide emissions continue to decline, driven by a large decrease in the use of solid fuels for space heating in the Residential (1A4b) sector, which is the principal source of CO, and major reductions due to three way exhaust catalysts in gasoline vehicles in Road Transport (1A3b) (Figure 2.5). National total CO emissions have reduced from 594.19 kt in 1990 to 88.97 kt in 2023, a reduction of 85.0 per cent.

Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined, account for 60.2 per cent of national total CO emissions in 2023, a reduction of 81.7 per cent between 1990 (292.81 kt) and 2023 (53.53 kt). The Transport sector is another important source and accounted for 20.7 per cent of national total emissions in 2023, a major reduction (93.1 per cent) from 267.94 kt in 1990 to 18.43 kt in 2023. Public Electricity and Heat Production (1A1a) sector reached its peak in 2001 (23.84 kt) and has decreased by 66.7 per cent to reach 7.94 kt in 2023, a reduction of 56.3 per cent on 1990 levels (18.15 kt). Emissions decreased in 2023 by 31.8 per cent compared to 2022 as a result of a decrease in coal and fuel oil use in electricity generation. Combustion sources from Manufacturing Industries and Construction (1A2) account for 8.1 per cent of the national total in 2023 and at 7.22 kt showed a 35.1 per cent decrease on their 1990 levels (11.12 kt). Agriculture/Forestry/Fishing (1A4c) combustion sources account for 1.5 per cent of the total CO emissions in 2023. Petroleum refining (1A1b), Manufacture of soil fuels and other energy industries (1A1c), Other Product Use (2G), Biological Treatment of waste (5B1) and Waste incineration (5C) emissions form the Other NFR sectors category and combined account for the remainder of CO emissions (0.5 per cent of the total) in 2023.



**Figure 2.5 Emission Trend for Carbon Monoxide 1990–2023**

## 2.3 Particulate Matter

Particulate matter (PM) emission estimates include PM with diameter less than 10 µm (PM<sub>10</sub>), PM with diameter less than 2.5 µm (PM<sub>2.5</sub>) and total suspended particulates (TSP).


### 2.3.1 Particulate Matter <10 µm Diameter (PM<sub>10</sub>)

Emissions of particulate matter <10 µm diameter amounted to 17.77 kt in 2023, a 53.3 per cent reduction from 38.09 kt in 1990 (Figure 2.6). The main determinant of the trend in PM<sub>10</sub> emissions are Other sectors, combustion in Residential (1A4b) and Commercial/Institutional (1A4a) combined and Agriculture with 35.6 per cent and 25.2 per cent and 17.2 per cent shares, respectively of the national total in 2023.

Petroleum refining (1A1b), Manufacture of solid fuels and other energy industries (1A1c), Fugitive emissions from soil fuels: Coal mining and handling (1B1a), Cement production (2A1), Lime production (2A2), Quarrying and mining of minerals other than coal (2A5a), Construction and demolition (2A5b), Storage, handling and transport of mineral products (2A5c), Storage, handling, transport of chemical products (2B10b), Road paving with asphalt (2D3b), Other solvent use (2D3i), Other product use (2G), Biological treatment of waste – solid waste (5A), Incineration (5C) and Other waste (5E) are included under the Other NFR sectors heading, which combined accounted for 35.6 per cent of national emission in 2023. The largest contributors are within the Other NFR grouping are Quarrying and mineral of minerals other than coal (2A5a) and Road paving with asphalt (2D3b). Part replacement of coal and peat in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors are the reason for reductions in the second largest contributor to the total PM<sub>10</sub> emissions. A 78.8 per cent reduction in emissions was seen from these sectors, from being the largest contributor to the PM<sub>10</sub> emissions (55.4 per cent of national total emissions) at 21.11 kt in 1990, emissions have fallen to 4.47 kt in 2023 (and 25.2 per cent of the national total emissions). Emissions from Agriculture arise from Farm-level agricultural operations including storage, handling and transport of agricultural products (3Dc), Off-

farm storage, handling and transport of bulk agricultural products (3Dd) and Manure Management (3B) categories that together in 2023 at 3.05 kt indicated a 7.7 per cent increase on their 1990 levels. Emissions from Transport (1A3), at 12.7 per cent share of the total in 2023 increased throughout the 1990s with increased total vehicle kilometre travel particularly of diesel vehicles. However, increases in vehicle numbers seen over the last decade has been partially offset by changes in both the age structure of the national fleet and developments in diesel fuel technology, resulting in Transport emissions decreasing by 23.3 per cent (from 2.95 kt in 1990 to 2.26 kt in 2023).


Manufacturing Industries and the Construction sector (1A2) used to account for an increasing percentage of the national total PM<sub>10</sub>, until reaching a peak of 2.48 kt in 2005. This trend parallels other pollutants and is due to the increase in cement production post-2000 following the entry into the market of two new plants. Emissions for this sector have decreased since 2005 and accounted for 7.2 per cent of the national total in 2023 (1.27 kt), representing a decrease across the 1990-2023 time series of 30.9 per cent. Public Electricity and Heat Production (1A1a) sector emissions accounted for 1.5 per cent of the national total in 2023 and at 0.27 kt reduced by 71.4 per cent from 1990 levels (0.96 kt). This reflects the considerable reduction of coal, oil and peat use in this sector due to the improvement in efficiency of modern gas fired power plants replacing older peat and oil-fired plants and the increased share of renewables, primarily wind power, and increased interconnection.



**Figure 2.6 Emission Trend for Particulate Matter <10 µm in Diameter 1990–2023**

### 2.3.2 Particulate Matter <2.5 µm Diameter (PM<sub>2.5</sub>)

National total emissions of particulate matter <2.5 µm diameter amounted to 9.31 kt in 2023, a 67.7 per cent reduction on 28.79 kt in 1990 (Figure 2.7). Emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined are the main determinant of the trend with their 47.7 per cent share of the national total PM<sub>2.5</sub> emissions in 2023. There has, however, been a reduction of 78.8 per cent in emissions from these sectors between 1990 (20.95 kt and 72.8 per cent share) and 2023 (4.44 kt). Reduced use of coal and peat, with increased use of gasoil, kerosene and natural gas in the two sectors has resulted in lower emissions and a reduction in the contribution to the national total.




**Figure 2.7 Emission Trend for Particulate Matter <2.5 µm in Diameter 1990–2023**

Emissions from Agriculture arise from Manure Management (3B), Farm level agricultural operations including storage, handling and transport of agricultural products (3Dc), Off-farm storage, handling and transport of bulk agricultural products (3Dd) sectors that together in 2023 at 0.84 kt accounted for 9.0 per cent of the national total and indicated a 12.0 per cent increase on their 1990 levels (0.75 kt). Transport (1A3) contributed 1.39 kt (14.9 per cent share) to the national total in 2023. Emissions from Transport sector (1A3), dominated by Road Transport (1A3b) increased from 1990 (2.58 kt) to a peak in 1996 (3.09 kt), but have been decreasing since 2005, with a 46.0 per cent reduction between 1990 and 2023 largely due to technological advances and the age structure of the national fleet, which in turn have been balanced by the increases in vehicle numbers over the time series (see comments in section 2.3.1 on the trends of PM<sub>10</sub> emissions). Emissions from Manufacturing Industries and Construction (1A2) at 1.18 kt in 2023 are 25.8 per cent below those in 1990.

Electricity and Heat Production (1A1a) sector accounted for 2.2 per cent of the national total emissions in 2023, a reduction of 67.7 per cent from 0.65 kt in 1990 and 0.21 kt in 2023. There was a reduction in coal and fuel oil use in 2023 which led to an 52.1 per cent reduction in emissions compared to 2022. Combustion from the Agriculture/Forestry/Fishing sector accounted for 1.1 per cent of national total PM<sub>2.5</sub> emissions in 2023 and have reduced by 88.6 per cent from 1990 when emissions were 0.92 kt, compared to 0.10 kt in 2023. The remainder of PM<sub>2.5</sub> emissions arise from Petroleum refining (1A1b), Manufacture of soil fuels and other energy industries (1A1c), Fugitive emissions from solid fuels: Coal mining and handling (1B1a), Cement production (2A1), Lime production (2A2), Quarrying and mining of minerals other than coal (2A5a), Construction and demolition (2A5b), Storage, handling and transport of mineral products (2A5c), Storage, handling, transport of chemical products (2B10b), Road paving with asphalt (2D3b), Other solvent use (2D3i), Other product use (2G), Biological treatment of waste – solid waste (5A), Incineration (5C) and Other waste (5E) are included under the other NFR sectors heading, which combined accounted for 12.3 per cent of national emission in 2023.

### 2.3.3 Total Suspended Particulates (TSP)

Total suspended particulate emissions have decreased by 33.1 per cent, from 74.13 kt in 1990 to 49.59 kt in 2023 (Figure 2.8). The main driver of the TSP trend is emissions in Other NFR sectors which includes a wide range of source categories namely Petroleum refining (1A1b), Manufacture of solid fuels and other energy industries (1A1c), Fugitive emissions from solid fuels: Coal mining and handling (1B1a), Cement production (2A1), Lime production (2A2), Quarrying and mining of minerals other than coal (2A5a), Construction and demolition (2A5b), Storage, handling and transport of mineral products (2A5c), Storage, handling, transport of chemical products (2B10b), Road paving with asphalt (2D3b), Other solvent use (2D3i), Other product use (2G), Biological treatment of waste – solid waste (5A), Incineration (5C) and Other waste (5E) which combined accounted for 70.24 per cent of the national total in 2023. Road paving with asphalt (2D3b) accounts for the majority of emissions within this subsector (86.1 per cent). Emissions from the Other NFR sectors have increased by 13.3 per cent from 40.20 kt in 1990 to 34.84 kt in 2023. Emissions from the agriculture (NFR 3) sector were the second largest contributor (9.83 per cent share of the total) with 4.87 kt emissions in 2023, a 27.5 per cent increase from 1990 levels (3.82 kt). Combined emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors were the third largest contributor to the total TSP emission in 2023. There has been, similar to emissions from both PM<sub>10</sub> and PM<sub>2.5</sub>, a reduction of 78.4 per cent in emissions from these sectors between 1990 (22.55 kt) and 2023 (4.87 kt). In the time series, the part replacement of coal and peat with natural gas, gasoil and kerosene has resulted in the contribution of these sectors falling from 30.4 per cent of the national total in 1990 to 9.82 per cent in 2023.




**Figure 2.8 Emission Trend for Total Suspended Particulates 1990–2023**

Emissions from Transport (1A3) decreased by 4.0 per cent to 2023 (3.19 kt and 6.43 per cent share of national total) compared to 1990 (3.32 kt and 4.5 per cent share). Manufacturing Industries and Construction (1A2) account for an increasing proportion of emissions post-2000 as a result of the entry into the market of two new cement production plants (Non-metallic minerals, 1A2f). Emissions from the 1A2 sector peaked in 2005 (2.80 kt) and have generally declined since. At 1.39 kt in 2023 (2.8 per cent of the total) emissions have decreased from 2005 by 50.6 per cent and by 33.3 per cent from

1990 (2.08 kt). Public Electricity and Heat Production sector (1A1a) emissions have decreased by 73.7 per cent over the time series, from 1.23 kt in 1990 to 0.32 kt and less than one per cent share of the total emissions in 2023. In 2023 the use of coal and oil decreased leading to a 55.9 per cent reduction in emissions compared to 2022. Emissions from Agriculture/Forestry/Fishing in 2023 (0.11 kt, 0.2 per cent of the national total) have decreased by 87.6 per cent since 1990 (0.93 kt).

### 2.3.4 Black Carbon (BC)

Black Carbon emissions have decreased by 70.0 per cent, from 4.37 kt in 1990 to 1.31 kt in 2023 (Figure 2.9). The main driver of the BC trend is Transport (1A3) emissions and combined emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors. In 2023, emissions from Transport sector (1A3) were the largest contributor to total BC emissions (34.7 per cent share). Emissions have reduced by 61.8 per cent between 1990 (1.19 kt) and 2023 (0.45 kt).



**Figure 2.9 Emission Trend for Black Carbon 1990–2023**

Combined emissions from the Residential (1A4b) and Commercial/Institutional (1A4a) sectors were 0.43 kt (32.9 per cent share) and have reduced by 80.4 per cent since 1990 (2.2 kt). Manufacturing Industry and Construction (1A2) emissions accounted for 23.6 per cent of the total in 2023 (0.31 kt) having decreased by 12.6 per cent since 1990. Emissions from Agriculture/Forestry/Fishing in 2023 were 0.05 kt, which equates to an 89.5 per cent reduction on 1990 (0.51 kt). Public Electricity and Heat Production (1A1a) emissions have decreased by 81.2 per cent over the time series. A reduction in the use of coal and oil in 2023 led to a 13.8 per cent decrease compared to 2022. The remaining 4.4 per cent in 2023 accounts for emissions from Petroleum refining (1A1b), Manufacture of solid fuels and other energy industries (1A1c), Cement production (2A1), Lime production (2A2), Road paving with asphalt (2D3b), Other product use (2G), and Industrial waste incineration (5C1bi).




## 2.4 Priority Metals

### 2.4.1 Lead (Pb)

Over the 1990–2023 time series, total national Pb emissions have decreased by 96.0 per cent, from 169.93 t in 1990 to 6.85 t in 2023 (Figure 2.10). The Pb emissions trend is largely determined by Road Transport (1A3b). Emissions of Pb have decreased considerably since 1990. There was a marked decrease between 1999 and 2000 when the lead content of petrol was reduced. In addition, there was an increase in the use of unleaded gasoline in road transport throughout the 1990s and the subsequent phasing out of leaded gasoline in 2000/2001. The contribution of Transport (1A3) to the much-reduced national total emissions has decreased by 96.7 per cent, from 157.61 t (92.7 per cent share) in 1990 to 5.13 t (74.9 per cent share) in 2023.

The second largest contributors to Pb emissions at 16.4 per cent share of national total in 2023 were the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined. The use of coal and peat in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors has been partly replaced with natural gas, gasoil or kerosene. Emissions from these sources have fallen from 6.26 t in 1990 to 1.12 t in 2023, a reduction of 82.1 per cent. Combustion in Manufacturing Industries and Construction (1A2) accounted for 4.0 per cent share of the total in 2023, and emissions from Public Electricity and Heat Production sector (1A1a) were responsible for 3.7 per cent share in 2023. Emissions from Metal Production (2C) have decreased in recent years due to the closure of a number of foundries and were negligible in 2023 (0.6 per cent). Similarly, emissions from Waste Incineration (5C) have also decreased to almost zero as a result of an outright ban on the incineration of clinical wastes in the mid-1990s. Incineration in 5C is now solely in relation to the destruction of vapours.




**Figure 2.10 Emission Trend for Lead 1990–2023**

### 2.4.2 Cadmium (Cd)

Total national emissions of Cd have decreased from 0.57 t in 1990 to 0.21 t in 2023 (Figure 2.11), a reduction of 62.6 per cent. Emissions of cadmium are largely determined by the Manufacturing Industries and Construction (1A2) sector, specifically combustion sources in Non-Ferrous Metals (1A2b). Manufacturing Industries and Construction (1A2) accounted for 31.1 per cent of total emissions in 2023 having increased by 22.2 per cent since 1990 (0.05 t and 9.5 per cent share). Public Electricity and Heat Production (1A1a) decreased from 1990 by 67.8 per cent and accounted for 17.4 per cent of the national total in 2023. Emissions from this source increased throughout the 1990s until

2001 as a result of the combustion of increasing quantities of coal and peat for electricity generation. However, 2023 saw a decrease of 39.6 per cent compared to 2022 due to the improvement in efficiency of modern gas fired power plants replacing older peat and oil-fired plants and the increased share of renewables, primarily wind power, and increased interconnection.

Residential (1A4b) and Commercial/Institutional (1A4a) combustion is also an important source of Cd emissions, with combined emissions from these sectors accounting for 14.7 per cent of the national total in 2023 due to the continued use of the fossil fuels (coal, peat and oil). However, emissions from the sector have decreased by 50.7 per cent, from 0.06 t in 1990 to 0.03 t in 2023. Transport (1A3) sector emissions have been increasing in the time series (by 128.5 per cent) and in 2023 it accounted for 10.6 per cent of the national total cadmium emissions. Combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 1.2 per cent of the total Cd emissions in 2023. As a result of the closure of the Irish Steel plant in late 2001, emissions from Metal Production (2C) accounted for a 24.5 per cent share in 2023 compared to a 54.3 per cent share (and main contributor to the total) in 1990.




**Figure 2.11 Emission Trend for Cadmium 1990–2023**

### 2.4.3 Mercury (Hg)

Total national emissions of Hg have decreased from 0.75 t in 1990 to 0.23 t in 2023, a reduction of 69.3 per cent (Figure 2.12). Emissions from Public Electricity and Heat Production (1A1a) decreased by 72.6 per cent in the trend and accounted for 21.4 per cent (the third largest source) of national total mercury emissions in 2023 (0.05 t). Combustion sources in Manufacturing Industries and Construction (1A2) account for the largest share of national total mercury emissions (29.2 per cent in 2023), having decreased by 15.2 per cent between 1990 and 2023. Emissions from this sector are largely dependent on the increased use of petroleum coke and coal as a fuel source in the cement industry after the entry of a number of additional cement producers into the Irish market post-2000. Emissions from the combined Residential (1A4b) and Commercial/Institutional (1A4a) are the second largest source and accounted for 21.9 per cent of total emissions in 2023, a 74.0 per cent decrease since 1990. Emissions

from this source are the result of an increase in the use of natural gas, kerosene and gasoil and a decrease in the use of coal and peat.

Biological treatment of waste – Solid waste disposal on land (5A) and Waste Incineration (5C) combined accounted for a 16.2 per cent share of total emissions in 2023. The Transport (1A3) sector accounted for 11.1 per cent and combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 0.3 per cent of the total Hg emissions in 2023. Glass Production (2A3) and Metal Production (2C) are no longer occurring due to plant closures (glass since 2010 and metal since 2002).



*Figure 2.12 Emission Trend for Mercury 1990–2023*


## 2.5 Other Metals

### 2.5.1 Arsenic (As)

Emissions of arsenic have decreased by 41.4 per cent from 1.81 t in 1990 to 1.06 t in 2023 (Figure 2.13). These emissions are largely dominated by incineration of hazardous and clinical wastes and crematoria in Waste Incineration sector (5C). Waste Incineration accounted for 66.3 per cent of national total arsenic emissions, having increased by 27.4 per cent from 0.55 t in 1990 to 0.70 t in 2023. The absolute and percentage contributions of this sector are increasing largely due to the increase in the number of cremations undertaken in Ireland which has increased since 1990. Continued (but decreasing) use of coal, peat and fuel oil as part of the fuel mix contributes largely (16.3 per cent share of the total in 2023) to the trend in emissions from Public Electricity and Heat Production (1A1a). However, arsenic emissions from the sector have decreased by 78.3 per cent over the time series due to the replacement of less-efficient peat plants with new plant, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.

The continued use of fossil fuels in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined results in the sectors accounting for 2.3 per cent of national total emissions in 2023, however, they have decreased by 78.5 per cent in the trend. Within the Manufacturing Industries and Construction (1A2) sector, the sub-sector Non-metallic minerals (1A2f) is responsible for the majority of total sector's emissions due to the increase in cement production and associated fuel use in the sector, in particular petroleum coke post-2000 with the entry into the market of new cement plants

and reflecting more recent post-recession decrease in production and consequential lower emissions from the sector. The sector's As emissions contributed 9.5 per cent the total in 2023, an increase of 9.1 per cent on the 1990 level. Metal Production (2C) is no longer an important source of As emissions, following the closure of the Irish Steel plant in 2001 and a reduction in emissions from Integrated Pollution Prevention and Control (IPPC)-licensed facilities. Emissions from this sector accounted for 12.0 per cent in 1990 (third largest contributor) and are reported as not occurring in 2023 due to the closure of the foundry which was a responsible for emissions from this sector.




**Figure 2.13 Emission Trend for Arsenic 1990–2023**

### 2.5.2 Chromium (Cr)

Emissions of Cr have decreased by 34.6 per cent from 5.08 t in 1990 to 3.33 t in 2023 (Figure 2.14). Transport (1A3) and Waste Incineration (5C) sectors are the main two drivers of the chromium trend. The Transport (1A3) sector accounted for 53.6 per cent of estimated national total emissions in 2023. Emissions from this source category have increased by 132.4 per cent over the time series between 1990 (0.77 t) and 2023 (1.78 t) due to the large increase in vehicle numbers in Ireland. The incineration of hazardous and clinical wastes and crematoria contribute to emissions from Waste Incineration (5C). The sector is the second largest source of chromium emissions and accounted for 20.4 per cent of 2023 national chromium emissions, an increase of 27.2 per cent in the trend. The absolute and percentage contributions of this sector are increasing largely due to an increase in industrial waste incineration (5C1bi) which includes the incineration of wood that has historically been treated with preservatives containing chromium.

Continued use of coal, peat and fuel oil as part of the fuel mix for Public Electricity and Heat Production (1A1a) means that the sector contributed 4.6 per cent to the emissions total in 2023. However, there has been a reduction in emission levels of 70.9 per cent between 1990 and 2023 as a result of the replacement of less-efficient peat plants with new plant, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy. Emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined accounted for 6.4 per cent of the national total in 2023, having decreased, by 77.0 per cent over the time series, reflecting the part


replacement of coal and peat with natural gas, gasoil and kerosene. Within the Manufacturing Industries and Construction (1A2) sector, the sub-sector Other (1A2f) is responsible for the majority of emissions largely due to the cement industry, as is evident with other heavy metal estimates. Emissions in 1A2 accounted for 9.7 per cent of national total chromium emissions in 2023 and have decreased by 38.8 per cent in the whole trend. Similar to other heavy metals estimates, the closure of the Irish Steel plant in 2001 has markedly reduced the effect of Iron and Steel Production (2C1) on emission trends, from 35.0 per cent share of the total (and the main contributor to chromium emissions) in 1990 to a 4.5 per cent share in 2023.



*Figure 2.14 Emission Trend for Chromium 1990–2023*

### 2.5.3 Copper (Cu)

Total copper emissions in Ireland increased over the 1990–2008 period but have been steadily decreasing since (Figure 2.15). Total emissions in 2023 (46.07 t) were 5.4 per cent lower than in their peak in 2008 (48.73 t) but they are 111.1 per cent higher than in 1990 (21.82 t). This trend is primarily driven by the Transport (1A3) sector that accounted for 83.5 per cent of estimated national total copper emissions in 2023 (38.47 t) and increased by 135.9 percent since 1990 (16.30 t).




**Figure 2.15 Emission Trend for Copper 1990–2023**

The trend can be attributed to increasing economic prosperity and population in Ireland, with a high reliance on private car travel as well as rapidly increasing road freight transport. Continued use of coal, peat and fuel oil as part of the fuel mix results in a contribution of 0.5 per cent to the total emissions from Public Electricity and Heat Production (1A1a) in 2023. However, a reduction in emission levels of 60.2 per cent is evident between 1990 and 2023 as a result of the replacement of older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy. Emissions from this source decreased by 42.4 per cent in 2023 due to a reduction in coal and fuel oil use compared to 2022. The Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined accounted for 0.5 per cent of emissions in 2023, showing a 81.2 per cent decrease in the trend since 1990. Waste Incineration (5C) emissions accounted for 0.8 per cent of the national total in 2023, having increased by 18.0 per cent since 1990 (0.32 t). Combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 0.8 per cent of the total Cu emissions in 2023 (3.2 per cent increase since 1990). The Manufacturing Industries and Construction (1A2) sector has decreased by 26.2 per cent between 1990 and 2023 (accounting for 0.6 per cent of the total in 2023). Emissions from the sector were increasing proportionately since 1990 reaching their peak in 2005 as a result of the entry into the market of new operators in the cement production sub-sector post-2000. The use of fossil fuels for combustion decreasing since 2005 as a result of economic circumstances and increasing used of waste as a fuel. Emissions from Industrial Processes, specifically Other solvent use (2D3i) and Other product use (2G) combined accounted for 13.2 per cent of the total in 2023. Emissions from 2D3i and 2G combined have increased by 137.3 per cent across the timeseries. The sources of emissions in these categories are lubricants in the case of category 2D3i and Tobacco and Fireworks in the case of category 2G.

#### 2.5.4 Nickel (Ni)

National total emission estimates of nickel were steadily increasing over the 1990–1999 period and have been generally decreasing from 1999 onwards (Figure 2.16). Total emissions in 2023 (4.82 t) were

86.0 per cent lower than at their peak in 1999 (34.49 t) and 78.3 per cent lower than in 1990 (22.18 t). The largest contributor to total Ni emissions in 2023 is the Manufacturing Industries and Construction (1A2) emission category with a 36.8 per cent share in 2023 (1.77 kt) having reduced by 83.1 per cent since 1990 (6.93 kt). The second largest contributor is emissions from the combined Residential (1A4b) and Commercial/Institutional (1A4a) sectors accounting for 30.4 per cent. A switch within these sectors from solid fuels (coal and peat) to less carbon intensive liquid fuels and natural gas has resulted in a reduction of 78.1 per cent in emissions since 1990 (6.68 kt). Public Electricity and Heat Production (1A1a) sector accounted for a 14.1 per cent share of the total nickel emissions in 2023. A reduction in emission levels of 95.6 per cent is evident between 1990 and 2023. This is because of the replacement of older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.



**Figure 2.16 Emission Trend for Nickel 1990–2023**


Combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 5.8 per cent in 2023. Similar to other heavy metals estimates, the closure of the Irish Steel plant in 2001 has significantly reduced the effect of Metal Production (2C) on emission trends, from 13.7 per cent share of the total (and fourth largest contributor to nickel emissions) in 1990. Emissions in 2023 were 0.26 t contributing to a 5.4 per cent share having reduced by 91.9 per cent since 1990.

### 2.5.5 Selenium (Se)

National total emission estimates of Se have decreased by 81.2 per cent, from 9.33 t in 1990 to 1.76 t in 2023 (Figure 2.17). The main contributor to the trend has been fuel combustion in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors which combined accounted for 67.7 per cent of selenium emissions in 2023, having decreased by 82.5 per cent since 1990. The second largest contributor to the trend is the Public Electricity and Heat Production (1A1.a) sector and in 2023 it accounted for 22.5 per cent share of total selenium emissions. Emissions from this sector have decreased by 83.2 per cent from their 1990 level of 2.35 t to 0.40 t in 2023 due to the replacement of

older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.

Transport (1A3) sector has been increasing in the time series and with its 2.4 per cent share of the national total in 2023, it has increased by 124.5 per cent since 1990 because of increasing the fuel use in this sector. Emissions from Manufacturing Industries and Construction accounted for 5.1 per cent of the total in 2023, having increased by 64.0 per cent from 1990 because of the increase in cement production and associated fuel use in this sector. The remainder of the selenium emissions arise from combustion in Agriculture/Forestry/Fishing (1A4c) sector with its 0.2 per cent share of the national total in 2023. Glass production (2A3) under Industrial Processes and Product Use (IPPU) sector used to be an important contributor to the selenium emissions trend throughout the 1990s and up to 2002, accounting for an average of 1.3 per cent of the national total in that period. In 2023 Industrial Processes accounted for 2.0 per cent of national total emissions.



*Figure 2.17 Emission Trend for Selenium 1990–2023*


## 2.5.6 Zinc (Zn)

National total emissions of Zn amounted to 55.19 t in 1990 and have fallen by 54.8 per cent to 24.96 t in 2023. In the 1990–2001 period, the main determinant of the trend in zinc emissions was Metal Production (2C), mostly iron and steel production, accounting on average for 51.7 per cent of national total emissions throughout that period (Figure 2.18). However, following the closure of Ireland’s only steel plant in 2001, emissions from this source sector are now almost negligible and limited to relatively small IPPC-licensed foundries and galvanising plants. The main determinant for the trend since 2003 has been the Transport (1A3) sector (53.3 per cent of national total zinc emissions in 2023) specifically, Road Transport (1A3b) sub-sector has increased significantly since 1990 as a result of the increase in the number of vehicles on Irish roads. As a result, emissions from Transport sector have increased substantially (144.2 per cent), from 5.45 t in 1990 to 13.31 t in 2023.



The second largest source in 2023 is emissions from combustion in Manufacturing Industries and Construction (1A2) which accounts for 15.4 per cent share of emissions in 2023 having increased from 3.26 t (5.9 per cent share) in 1990 to 3.85 t in 2023 (18.2 per cent increase). Emissions from combustion in Public Electricity and Heat Production (1A1a) emissions have decreased by 56.1 per cent since 1990 and accounted for a 4.5 per cent share of national total in 2023. 2023 saw a decrease of 49.8 per cent in emissions compared to 2022 due to a decrease in use of coal and oil for electricity generation.

Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined accounted for 11.7 per cent of the national total in 2023. However, reduced use of coal and peat through part replacement with natural gas and gasoil has resulted in the reduction of zinc emissions from these sectors by 77.0 per cent, from 12.74 t in 1990 to 2.93 t in 2023. The remainder of the zinc emissions arise from combustion in Agriculture/Forestry/Fishing (1A4c) sector with its 0.9 per cent share of the national total in 2023. Waste Incineration (5C) accounted for 0.01 per cent of the national total in 1990 however, following an outright ban on incineration of clinical waste in hospitals in 1997, emission from Clinical Waste Incineration (5Cbi) are no longer a source.



**Figure 2.18 Emission Trend for Zinc 1990–2023**


## 2.6 Persistent Organic Pollutants (POPs)

As part of Ireland's emission inventory improvement programme, the inventory agency tendered a project in 2007 to develop an inventory of persistent organic pollutants in Ireland. The project report (AEA/CTC, 2008) provides detailed information in relation to the methodological choice and activity data for the diverse range of sources that give rise to emissions of POPs within Ireland. For the purposes of identifying the major contributors to the trend in PCDD/F, PCBs, HCB and PAHs, some of this information is provided in the following sections. The approach has been updated subsequently as new versions of EMEP/EEA Inventory Guidebook have been published.

### 2.6.1 Dioxins and Furans (PCDD/F)

Dioxin and furan emission levels decreased from 44.32 g I-TEQ in 1990 to 11.47 g I-TEQ in 2023 (74.1 per cent reduction on 1990 levels). The main contributors to the trend are the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined with a 69.8 per cent share of total emissions in 2023; however, emissions from these sectors have reduced (by 74.3 per cent) from 31.17 g I-TEQ ( and 70.3 per cent share) in 1990 to 8.01 g I-TEQ in 2023 (Figure 2.19).

The second largest contributor to the trend is the Waste sector which is dominated by emissions from the Other Waste (5E) sector, where building and vehicle fires and residential burning of waste are the emission sources. Total emissions from the waste sector accounted for 19.9 per cent (2.28 g I-TEQ) of national total emissions in 2023, a reduction of 74.7 per cent on 1990 levels (9.01 g I-TEQ). The introduction of unleaded petrol and technological improvements in road vehicles have partially offset increased numbers of vehicles in the national fleet, with a 4.3 per cent share of national total emissions in Transport (1A3) in 2023, a reduction of 28.9 per cent on 1990 levels. Combustion in Agriculture/Forestry/Fishing (1A4c) sector accounted for 3.2 per cent of emissions in 2023 (0.36 g I-TEQ) a decrease of 68.9 per cent on the 1990 emissions (1.16 g I-TEQ). Public Electricity and Heat Production (1A1a) emissions have decreased by 63.5 per cent from their 1990 level of 0.83 g I-TEQ to 0.31 g I-TEQ and are responsible for a 2.6 per cent share of national total in 2023 due to the replacement of older less-efficient generation plants, reduction in the quantities of coal combusted overall and fuel switching from oil to natural gas and wind energy. Process emissions from the manufacture of cement (2A1) in Industrial Processes and Product Use sector, accounting for 0.1 per cent of national emissions in 2023 (0.01 g I-TEQ), a 99.0 per cent reduction on 1990 levels (1.45 g I-TEQ).




**Figure 2.19 Emission Trend for Dioxins and Furans 1990–2023**

## 2.6.2 Hexachlorobenzene (HCB)

Figure 2.20 outlines the trend in hexachlorobenzene emissions in Ireland across the 1990–2023 time series. The graph indicates that HCB emissions from Secondary Aluminium Processing (2C) which, for the period up to and including 1996, dominated the inventory with a contribution of 40 kg per year and is no longer a source of HCB emissions within Ireland due to the banning of hexachloroethane (HCE)-based cover gas use (HCB was present as a contaminant in such cover gases). Consequently, emissions since 1997 are more relevant to trend analysis up to 2023.

There is very limited information on the release of HCB to air for most source sectors in Ireland. The main source is the use of pesticides in agricultural practices significantly increasing up to 2004 (8.03 kg) and decreasing to 2.03 kg in 2005 following a ban on the use of certain pesticides. Subsequently, emissions have increased up to 2012 before decreasing thereafter and in 2023 were 2.12 kg representing an 86.1 per cent share. The Public Electricity and Heat Production (1A1a) sector was the second largest source in 2023 accounting for 0.28 kg and a 11.4 per cent share of the national total, having decreased by 42.3 per cent between 1990 and 2023.

Incineration of hazardous and clinical wastes and crematoria included in the waste incineration (5C) accounted for 0.3 per cent share of national emissions in 2023, a decrease of 87.9 per cent since 1990. The remainder of the HCB emissions arise mainly from combustion in the Residential (1A4b) and Commercial/Institutional (1A4a), Manufacturing Industries and Construction (1A2) and Agriculture/Forestry/Fishing (1A4c) sectors, which combined are presented as “Other” NFR sectors in Figure 2.20 and account for 1.8 per cent of the total in 2023. Transport (1A3) accounted for 0.3 per cent of national emissions in 2023.




**Figure 2.20 Emission Trend for Hexachlorobenzene 1990–2023**

## 2.6.3 Polychlorinated Biphenyls (PCBs)

Estimated national total emissions of polychlorinated biphenyls have decreased by 87.2 per cent from 39.35 kg in 1990 to 5.06 kg in 2023. Emissions peaked in 2003 (67.59 kg) (Figure 2.21). Other Waste (5E) is the largest contributor to the trend in PCB emissions in Ireland accounting for 2.70 kg in 2023. In 2023, the emissions from the Waste sector as a whole were 2.96 kg (58.5 per cent of national total emissions), a decrease of 81.4 per cent compared to 1990 (15.88 kg). Emissions from the Waste sector

peaked in 2003 (59.84 kg) due to an increase in the quantity of household waste that remain unaccounted for in national statistics and which is assumed to be burned. Combined emissions from the Residential (1A4b) and Commercial/Institutional (1A4a) sectors accounted for 33.3 per cent of the national total in 2023 (1.68 kg), a 82.9 per cent decrease from those estimated for 1990 (9.85 kg).



**Figure 2.21 Emission Trend for Polychlorinated Biphenyls 1990–2023**


Of particular note for PCB emissions is the contribution of the NFR Sector 2L (Other Production, consumption, storage, transportation or handling of bulk products), which in Ireland’s inventory covers PCB use as dielectric fluid in electrical equipment such as transformers and capacitors. However, through the introduction of Hazardous Waste Management Plans and the Waste Electrical and Electronic Equipment (WEEE) Regulations, emissions since 2006 have been decreasing in general.

Public Electricity and Heat Production (1A1a) emissions were 0.01 kg in 2023 and contributed 0.2 per cent of the total. Increases in cement production have led to increases in fuel combustion-based emissions in Manufacturing Industries and Construction (1A2) up until 2005. This source category’s emissions have decreased by 62.4 per cent between 1990 (1.06 kg) and 2023 (0.40 kg), with a contribution of 7.9 per cent to the national total in 2023.

## 2.6.4 Polycyclic Aromatic Hydrocarbons (PAHs)

For the purposes of this report, total PAHs in the form of the sum of emissions of benzo[a]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene, and indeno[1,2,3-cd] pyrene are presented in Figure 2.22. All together the emissions from these combined four pollutants decreased by 73.8 per cent between 1990 (29.63 t) and 2023 (7.75 t). For all four PAHs, the main source sectors are the same. The inventories are dominated by emissions from combustion in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors. In the Residential (1A4b) sector (main driver of the trend), the lack of combustion controls or abatement, together with relatively low-temperature combustion conditions, leads to high emissions of PAHs. Even though national total emissions of PAHs have generally declined across the time series, the Residential (1A4b) and Commercial/Institutional (1A4a)

sectors combined accounted for 92.2 per cent of the national total in 2023, having decreased by 74.6 per cent from 28.10 t in 1990 to 7.14 t in 2023. The decline in emission levels is due primarily to the decline in the use of coal and sod peat for residential space heating, as reported in the National Energy Balance.



**Figure 2.22 Emission Trend for Polycyclic Aromatic Hydrocarbons 1990–2023**

Combustion emissions in Manufacturing Industries and Construction (1A2) sector accounted for 3.1 per cent of the total in 2023 (0.24 t), a decrease of 81.6 per cent since 1990 (1.31 t). Emissions from Transport (1A3) have increased by 163.6 per cent to 0.29 t representing a 3.8 per cent share of the national total in 2023 for all four PAHs as compared to 0.4 per cent 1990 (0.11 t). Other Waste (5E) is the main driver in the Waste sector resulting in the Waste sector accounting for 0.02 per cent of the total PAHs emissions in 2023, a 96.9 per cent reduction since 1990 (0.05 t).

## Chapter - Three Energy

### 3.1 Overview of the Energy (NFR 1) Sector

The Energy sector covers combustion and fugitive sources of emissions associated with the production, transport, conversion, and use of fossil fuels. Emissions from combustion in this sector account for the bulk of total national emissions for most substances covered in this IIR. Estimates of the various pollutants are included for all emission sources that occur in the country and the required level of disaggregation is achieved for sufficiently detailed completion of the NFR tables.

Annual energy balance sheets published by Sustainable Energy Authority Ireland (SEAI) are the principal source of activity data for computing the emissions in the Energy sector. Ireland's energy statistics are compiled using a combination of top-down and bottom-up methods and the annual energy balances have undergone major improvement over recent years to take account of emission inventory requirements and more harmonised reporting to Eurostat and the International Energy Agency (IEA). The annual submission of up-to-date energy balances from SEAI to the inventory agency is one of the primary data inputs covered by a MoU in Ireland's national system (Chapter One). A fully consistent set of energy balance sheets for the years 1990–2023 underlies the time-series estimates of emissions for *Energy* in this submission. The 2023 energy balance is provided in Annex B.

Substantial plant-level fuel-use data are also available for many important categories in the Energy sector, especially for more recent years, which allows bottom-up estimates to be derived for some pollutants using Tier 3 methods. These data are obtained through direct arrangements with the operators of certain plants through their returns under Chapter III the Large Combustion Plant (LCP) of the Industrial Emissions Directive and the EU ETS and under environmental reporting related to their IPPC permits.

The emissions of SO<sub>2</sub> from fuel combustion are determined from the fuel properties, and fully representative emission factors are readily obtained for the fossil fuels used in most emission categories in Ireland. In general, other pollutants emitted from combustion sources are heavily reliant on emission factors from non-national sources and are taken from the Inventory guidebook (EMEP/EEA 2023). The Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP), which is aimed at supporting national experts in reporting PM emission inventories, serves as the reference for emission factors in the category 1A3a for the different forms of Particulate Matter (PM). Particulate Matter emission factors for other sectors are derived from the Inventory Guidebook (EMEP/EEA, 2023). In the past, emission inventories for heavy metals and POPs received little attention in Ireland and special studies were necessary to compile national emission estimates for reporting purposes. The separate detailed studies on emissions of heavy metals and POPs in Ireland were undertaken by consultants (AEA/CTC, 2008; Netcen/CTC, 2006) and they made use of the best available emission factors for many emission sources, with strong dependence on UK information sources. The results from these studies were the basis for developing time-series emissions of heavy metals and POPs in the Energy sector in previous submissions. In this submission, many of the EFs for heavy metals and POPs are from the Inventory Guidebook (EMEP/EEA, 2023).

## 3.2 Public Electricity and Heat Production (NFR 1A1a)


The production of electricity and heat from fossil fuels has traditionally been the most important source of key pollutants such as SO<sub>2</sub> and NO<sub>x</sub> in most countries. Approximately 64.1 per cent of electricity production in Ireland (SEAI, 2023) is dependent on fossil fuels and Category 1A1a, therefore, remains one of the major emission categories. Emissions of SO<sub>2</sub>, NO<sub>x</sub>, CO, three particulate matter pollutants, BC, all nine heavy metals and POPs have decreased significantly since 1990. NMVOCs were the only emissions to have increased in the time series due to the increased use of natural gas. The level of emissions in sector 1A1a depends heavily on the mix of fossil fuels and renewables used for electricity production. In 1990, heavy fuel oil (HFO), coal, peat and natural gas were the principal fuels used. The use of HFO, coal, and peat has declined as natural gas became the preferred fossil fuel during latter years. For example, 2019 represented an all-time low in the time series for coal used for electricity generation as consumption fell by 70 percent year on year following a 44 per cent decrease in 2018. However, 2020 saw consumption increase by 32 per cent compared to 2019, followed by a 246 per cent increase in 2021 compared to 2020 as gas fired plants were offline. While 2023 saw a 44 per cent fall in coal consumption for electricity generation from 2022 levels, it remains higher than the low recorded in 2019. Incineration emissions from Ireland's waste-to-energy plants, however small, have been increasing since the first such installation came into operation in 2011 and a second in 2017.

### 3.2.1 Emissions of Sulphur Dioxide and Nitrogen Oxides

Until 2000, the Electricity Supply Board (ESB) operated all public electricity power plants in Ireland. After 2000, several new gas-fired plants and one peat-fired plant were built by other operators, while the ESB replaced old peat-burning stations with new stations also burning peat and has been engaged in a major retrofit and improvement programme for plants in general. The shift to natural gas and the use of low-sulphur coal, combined with a decline in the sulphur content of fuel oil, have reduced SO<sub>2</sub> emissions by 99.3 per cent from 103.04 kt in 1990 to 0.71 kt in 2023. The sector's contribution to overall SO<sub>2</sub> emissions in 2023 was 9.6 per cent as opposed to 56.0 per cent share in 1990. In 2023, SO<sub>2</sub> emissions decreased by 61.7 per cent compared to 2022 levels, this is attributable to a decrease of coal and oil use in electricity generation and associated rise in natural gas consumption. Incineration of non-renewable waste accounted for 0.2 per cent of SO<sub>x</sub> emissions from power generation in 2023.

At the same time, the changed fuel mix, together with the application of extensive NO<sub>x</sub> emission control technology and more modern plants, has decreased NO<sub>x</sub> emissions by 90.7 per cent from 46.37 kt (and 27.6 per cent share of total NO<sub>x</sub> emissions) in 1990 to 4.33 kt (and 5.0 per cent share of the total) in 2023. In comparison to 2022, 1A1a NO<sub>x</sub> emissions decreased by 42.1 per cent and SO<sub>2</sub> emissions decreased by 61.7 per cent in 2023 due to marked decrease of coal and oil use in electricity generation. The ESB has supplied estimates of SO<sub>2</sub> and NO<sub>x</sub> on a plant-by-plant basis to the inventory agency for all years since 1990, mainly for the purpose of compiling SO<sub>2</sub> and NO<sub>x</sub> inventories under Chapter III of the IED Directive. The emissions for power plants operated by other companies are obtained either directly from their LCP, AER or PRTR submissions or estimated by the inventory agency from fuel data made available under the ETS. Ireland sought clarification from these power plants regarding the reporting of emission using validated average figures (after subtracting the value of the confidence interval). In this submission all relevant plants have reported correctly to the inventory


agency. Incineration of non-renewable waste in MSW accounted for 10.4 per cent of NO<sub>x</sub> emissions from power generation in 2023.



**Figure 3.1 Nitrogen Oxide Implied Emission Factors (IEF) for Category 1A1a**

The weighted average emission factors of SO<sub>2</sub> and NO<sub>x</sub> per fuel type (coal, peat, oil, natural gas, biomass and waste (MSW) in Category 1A1a are given in Figures 3.1 and 3.2 as implied emission factors (IEFs) to illustrate the level of decrease due to the factors mentioned above. The SO<sub>2</sub> emission factors reflect the sulphur content and net calorific value of the fuels used in the particular year and they account for sulphur retention levels in the fuel ash of 5 per cent and 10 per cent for coal and peat, respectively. The NO<sub>x</sub> emission factors (apart from landfill gas) are compiled from plant-level estimates that are determined from measurement, unit load factor and plant performance. Emission factors for landfill gas are default values as per the revised Inventory Guidebook (EMEP/EEA, 2023).





**Figure 3.2 Sulphur Dioxide Implied Emission Factors (IEF) for Category 1A1a**

### 3.2.2 Emissions of Other Substances

The emissions of all substances other than SO<sub>2</sub> and NO<sub>x</sub> in Category 1A1a are estimated by the inventory agency using the fuel-use energy data given by the national energy balance and appropriate emission factors taken from the Inventory Guidebook (EMEP/EEA, 2023) and plant-specific factors for non-renewable wastes from waste incineration. For the pollutants Cd, Hg, Pb, Dioxins, PCB, PAH and HCB Ireland utilises the emission factors presented in the 2023 EMEP/EEA Inventory Guidebook. Specifically, the emission factors for coal are taken from table 3-2, peat from table 3-3, fuel oil from table 3-6, natural gas and landfill gas from table 3-4. These emission factors are presented in Table C.1 Annex C. The activity data for 1A1a follows the format of that presented in Table B.1 Annex B for all years. Dioxins emissions for Peat power plants are estimated with plant specific emission factors. Furthermore, for Oil, there are no emission factors available for BaP, PCB and HCB in the 2023 EMEP/EEA Inventory guidebook. For natural gas and landfill gas there are also no emission factors available for PCB and HCB in the 2023 EMEP/EEA Inventory guidebook. Ireland uses the same activity data for all air pollutant and greenhouse gas emission calculations to maintain consistency in approaches across inventories (see section 3.1). The emission factors for 2023 (split by pollutants and fuel type) and their sources are listed in Table C.1 of Annex C.

### 3.3 Petroleum Refining (NFR 1A1b)

Emissions from fuel combustion at one small oil refinery in Ireland are estimated in this source category. Detailed information on the fuels used in different parts of the refinery is available through the company's AER, PRTR and ETS submissions. This allows for the selection of appropriate emission factors from national data and from international literature sources recognised as being fit for purpose by the emissions inventory community. In the case of heavy metals and POPs, the estimates are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2023). The emission factors used, and their sources are listed in Table C.2 of Annex C.

### 3.4 Manufacture of Solid Fuels and Other Energy Industries (NFR 1A1c)

Emissions from this source category refer to combustion emissions from the production of peat briquettes from milled peat at two plants (one of which closed in 2019 and the last of which closed in mid-year 2023), one natural gas production platform (which ceased production in July 2020) and one natural gas refinery in Ireland. As in the case of the oil refinery, the energy balance fuel-use data are supplemented by information reported by the plants under the ETS, which again allows for the selection of appropriate emission factors using national data and high-quality international sources. The estimates for heavy metals and POPs are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2023). The emission factors used, and their sources are listed in Table C.3 of Annex C.

### 3.5 Manufacturing Industries and Construction (NFR 1A2)

This category covers emissions from combustion in manufacturing industries and construction activities. Category 1A2 is split into the following seven subcategories:

- 1.A.2.a Iron and Steel
- 1.A.2.b Non-Ferrous Metals
- 1.A.2.c Chemicals
- 1.A.2.d Pulp, Paper and Print
- 1.A.2.e Food Processing, Beverages and Tobacco
- 1.A.2.f Non-metallic minerals
- 1.A.2.g Other Industry

Where it is possible to separate process emissions from those associated with fuel use, the process emissions associated with these industrial groups are reported in the Industrial Processes (NFR 2) sector (Chapter Four). The relevant process emissions in Ireland are those related to some metal industries.

Comparison of the Sustainable Energy Authority of Ireland (SEAI) Energy Balance data with ETS fuel-use data indicates that the combustion activities within the 1.A.2 category are dominated by a limited number of large industrial processing plants. It is assumed that all biomass reported as fuel use within the SEAI Energy Balance is clean untreated wood and wood-processing waste. The ETS data for major wood-processing facilities indicate the use of large-scale biomass boilers, fired using wood biomass, chippings, pulp and wood dust. Knowledge of these installations indicates that none of these wood-based fuels are pre-treated and hence emission factors applicable to clean wood use in large-scale boilers have been used to estimate POP emissions from these sources.

The Iron and Steel (1A2a) sector was dominated in the 1990s by fuel use and emissions from one electric arc furnace but, since its closure in 2001, the fuel use reported in 2002 in this sector relates to a small number of iron foundries. These iron foundries used only a very small amount of gasoil and LPG, and since 2003 combustion emissions are not occurring. For 1A2a, 1990 was the only year in the timeseries for which coal was combusted as a fuel in the category (29 ktoe). The emission factors used are those presented in Table 3.2 of the Inventory Guidebook (EMEP/EEA, 2023). The emission factor for coal is significantly higher than the emission factors associated with the other fuels combusted in the category, thus the combustion of coal in the category for the year 1990 has a marked influence on

emissions for that year. The process emissions from this industrial activity are described in chapter 4, section 4.4.1.

The Non-Ferrous Metal (1A2b) subcategory is dominated by the considerable fuel use (mainly natural gas replacing fuel oil) reported at a single large alumina plant, whilst the Chemicals (1A2c) subcategory includes natural gas, kerosene, fuel oil, gasoil, pet coke, and LPG use at large chemical plants. The Food Processing, Beverages and Tobacco (1A2e) sector covers a diverse range of industrial plants, many of which relate to agriculture, with under 30 of these installations reporting to the EU ETS and using predominantly natural gas and liquid fuels.

The bulk of fuel use reported in the energy balance under the 1A2f subcategory is accounted for by major cement works, lime producers, a small number of brickworks and fuel use at boiler plant within industries such as the pharmaceutical, glass and tile manufacturing sectors. There are at present four cement plants in Ireland, all of which use the dry kiln process currently fuelled by coal, petroleum coke and fuel oil, while meat and bone-meal, solid recovered fuels (SRF) and tyre derived fuels are co-incinerated at three of the four plants. The Inventory Guidebook (EMEP/EEA, 2023), provides some POP emission factors on an overall grams of pollutant per tonne of clinker produced basis. The guidebook states that these emissions should be allocated to combustion in cement manufacture. These emission factors have been used to determine the total emissions from cement plants.

Fuel-use data are available from plant operators as part of their reporting requirements under the EU ETS (Directive 2003/87/EC). Residual fuel not used in the cement manufacture sector but used in other manufacturing industries in 1A2f is calculated and emissions from fuel use are calculated using combustion emission factors in 1A2f, while the fuel use data from cement manufacture is removed to ensure there is no double counting. The same method is used for heavy metals in 1A2f. Emissions for individual cement plants were collected for all available years for Cadmium, Mercury and Lead and a plant specific g/t emission factor was determined using clinker production. This was then used to calculate emissions for years where plant emissions data was not available. For other metals As, Cr, Cu, Ni, Se, Zn the Tier 2 emission factors from Table 3-25 Chapter 1A2 of the Inventory guidebook (EMEP/EEA, 2023) were used with plant clinker production to estimate emissions.

Residual fuel not used in the cement manufacture sector but used in other manufacturing industries in 1A2f is calculated and emissions from this fuel use are calculated using combustion emission factors appropriate to 1A2f. All other industrial fuel use is reported under subcategory 1A2g. This sector covers a diverse range of manufacturing branches ranging from textile and leather, through machinery, transport equipment, wood products, mining (excluding fuel mining) and quarrying to other manufacturing businesses. Ireland's national Energy statistics do not provide an estimate of fuel used in mobile construction. All emissions associated with the category 1A2g are reported in 1A2gviii, and mobile emissions, 1A2gvii, are reported as "included elsewhere" (IE) in 1A2gviii. Currently all fuel is assumed to be stationary as the energy balance does not provide an estimate for fuels used in mobile machinery within category 1A2g. The inventory agency tendered and finalized research to be undertaken to quantify the extent and amounts of fuel used by mobile combustion on construction sites, but the COVID-19 pandemic had a negative impact on its outcome as it affected the collection of activity data. As a result, it was concluded that changing the methodology to estimate emissions

from off-road vehicles and other machinery to consider the results of the research project would not improve the accuracy of the inventory.

The revised and expanded energy balance sheets developed by SEAI incorporates a mapping of industrial fuel use in combustion into the NFR subcategories 1A2a through 1A2g under sector 1A2 Manufacturing Industries and Construction. This facilitates the complete disaggregation of emissions at subcategory level. In addition, information on fuel consumption was obtained in respect of a small number of energy-intensive industries (e.g. alumina production and cement manufacture) from their ETS returns, allowing their respective energy use amounts to be reconciled with the breakdown given in the national energy balance. Emissions in subcategories 1A2a through 1A2g are estimated on a top-down basis using disaggregated fuel use from the energy balance and the mix of country-specific and default emission factors as shown in Tables C.4 through C.10 of Annex C. The estimates for heavy metals and POPs are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2023). Emission factors for heavy metals and POPs (including references) for NFR 1A2 (a–g) are shown in Table C.11 of Annex C.

Ireland uses the emission factors from the Inventory Guidebook (EMEP/EEA, 2009) for liquid fuels. The Tier 1 EFs for liquid fuels in the latest version of the Inventory Guidebook are an average of Tier 2 EFs and are not considered representative considering one of the liquid fuel values for NO<sub>x</sub> is 942g/GJ based on reciprocating engines which are not present in Ireland. The Tier 1 value of 100 g/GJ used by Ireland is still valid, as all the guidebook editions reference Chapter B216 of older editions of the guidebook. A generic emission factor of 100 g/GJ was applied for all liquid fuels for sectors 1A2f, which was obtained from “Table 3-4: Tier 1 emission factors for 1.A.2 combustion in industry using liquid fuels”, for “‘Other’ Liquid Fuels” in page 16, 1.A.2 Manufacturing industries and construction (combustion), EMEP/EEA emission inventory guidebook 2009 (EMEP/CORINAIR B216). Table 1-1 of the Guidebook 2023 refers to the appropriate chapters for EFs, either 1A1, 1A2 or 1A4. Very few EFs were used from Chapter 1A2 as Ireland does not have these types of industry. Emission factors were again mainly sourced from the small combustion chapter (1A4) for boilers, as these are the most appropriate for the boiler types used in Ireland. The average of Tier 2 EFs are not applicable as Ireland has no reciprocating engines or turbines.

When estimating emissions for 1A2gviii, the fuel used for 3 large plants for all years where data is available (2005-2023, EU ETS), was subtracted from the total fuel use and a Tier 2 EF (EMEP/EEA 2023, Chapter 1.A.4 Small Combustion, Table 3-48, 'wood') was used for NMVOC and all other pollutants. The residual amount of biomass fuel, after the fuel used for 3 large plants is removed, uses the Tier 1 EFs. The emission factors used are presented in Annex C, Table C.10.

### 3.6 Transport (NFR 1A3)

As abatement measures continue to reduce emissions of key pollutants from major stationary combustion sources, transport in general, and road transport in particular, has become more important as a source of atmospheric emissions in many countries. The effects of technological emission controls for passenger cars and other vehicles in Ireland have been largely offset by the substantial increases in vehicle numbers, with the result that major reductions in the emissions of pollutants such as NO<sub>x</sub> did not occur until 2007 when the economic situation caused an overall

decrease in most emissions evident from 2008. In 2023 NO<sub>x</sub> emissions from transport were 32.18 kt (38.1 per cent share of total) and a decrease of 52.3 per cent is evident since 1990 (67.46 kt and 38.7 per cent share).


Road transport in Ireland is a larger source of NO<sub>x</sub> than electricity production, and road traffic also continues to be the major source of CO, BC, Pb, Cr, Cu and Zn, although BC, Pb, Cr, Cu and Zn emissions are very small in absolute terms.

### 3.6.1 Domestic and International LTOs (NFRs 1A3aii(i) and 1A3ai(i))

As a relatively small island state, aviation emissions are dominated by the international component. Under the LRTAP Convention, only the landing and take-off (LTO) component of emissions for both domestic and international flights is reported in the national total. The cruise component, domestic and international, is reported as a memo item. After the motorway network to Dublin from both Galway and Cork was completed around 2008/09, and in parallel upgrades of the Cork to Dublin rail line were completed, domestic air travel was no longer competitive, leading to the cessation of specific internal routes within Ireland.


The fuel consumption associated with domestic and international LTOs is estimated using a Tier 3b approach (Table 3.6.2, 2006 IPCC guidelines) based on origin and destination data for domestic air travel provided by EUROCONTROL (Eurocontrol, 2024), using an Advanced Emission Model (AEM) to estimate fuel burned and emissions for the full trajectory of each flight segment using aircraft and engine specific information. This approach was developed in 2019 and retrospectively applied from 2005 to 2023 where EUROCONTROL data is available. For the years 1990-2004, the number of flights for each airport was estimated based on domestic passenger and aircraft movement statistics as well as the relationship between all Irish airports and Dublin airport which is the principal destination of all domestic flights. Domestic LTO and Cruise fuel consumption rates for 1990-2004 are based on an average (2005 to 2011) for each departure airport using EUROCONTROL consumption rates.

Figures 3.3 and 3.4 and Table C.12 of Annex C present the number of LTOs, domestic and international, from Irish airports for all years from 1990 to 2023.



**Figure 3.3 Number of Domestic LTOs from Irish airports 1990-2023**

For data handling purposes, the inventory agency aggregated approximately 15 small regional airport/aerodrome pairs to “Other” which account for approximately 3.0 per cent of all domestic flights along with nine Irish airports which account for the remaining 97.0 per cent of all domestic flights in 2023. Emission factors associated with aviation emissions are presented in Annex C tables C.13 to C.18.



*Figure 3.4 Number of International LTOs from Irish airports 1990-2023*

### 3.6.2 Road Transport (NFR 1A3b)

The emissions of nine priority pollutants (SO<sub>2</sub>, NO<sub>x</sub>, NMVOCs, NH<sub>3</sub>, CO, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, BC) as well as seven heavy metals (Pb, Cd, Cu, Cr, Ni, Se, Zn) and POPs (Dioxins and Furans, PAHs) reported under sub-category 1.A.3.b Road Transport are estimated directly from the COPERT 5 model (Pastramas N. et al., 2014) developed within the CORINAIR programme for estimating a range of emissions from this important source.

The national total emissions in all cases are determined by the quantity of fuel sold in the country, as given by the energy balance. Approximately 65.82 petajoule (PJ) equivalent fuel energy from petrol, diesel, liquefied petroleum gas (LPG) and biofuels was consumed by road transport in Ireland in 1990. Consumption went up to 190.86 PJ in 2007 followed by a sharp decline to 142.33 PJ in 2012 and subsequently increasing to 160.16 PJ in 2018, with a significant reduction in 2020 to 132.93 PJ, mainly attributed to the COVID impact and travel restrictions. In 2021, consumption increased to 141.65 PJ due to easing of COVID-related travel restrictions and in 2023 a further increase in 2022 to 151.33 PJ and in 2023, 151.95 PJ. The energy share in biofuel has continued to increase since its introduction in 2005 and increased to 12.4 PJ in 2023.

Separate estimates of emissions are produced based on fuel types and amounts used for fuel purchased in Ireland and then subsequently used outside of Ireland. In 2023, 6.0 per cent of diesel and 0.5 per cent of petrol sold in Ireland was consumed outside the country. For some countries, including Ireland, the national totals determined by fuels used are the basis for assessment of their performance in relation to relevant Protocols under CLRTAP. Annex A.3 outlines the methodology

used to estimate the quantities of automotive fuels used in Ireland and includes the adjusted annual emissions for 1987 and the years 1990–2023.

The emissions of SO<sub>2</sub> for road transportation are computed from the amounts of petrol and diesel used by motor vehicles, as reported in the energy balance, and the sulphur content of the fuels. For the other substances, the COPERT 5 model estimates emissions based on distance travelled using a detailed bottom-up approach (Tier 3) that accounts for such factors as fuel type, fuel consumption, engine capacity, driving speed and a wide range of applicable technological emission controls that may be applied across the different vehicle categories according to the age of vehicles.

A total of 284 vehicle categories are determined by these variables in COPERT 5. Annex C shows the trend in emission factors for NO<sub>x</sub>, SO<sub>2</sub>, NH<sub>3</sub> and NMVOCs, CO and Particulate Matter per vehicle type for road transportation over the period 1990–2023 determined by the COPERT variables for the vehicle fleet in Ireland. For PM<sub>2.5</sub>, PM<sub>10</sub> and TSP, non-exhaust emission that includes brake and tyre wear emissions and road abrasion emissions for PM<sub>2.5</sub>, PM<sub>10</sub> and TSP are also presented as estimated using the COPERT model.

The primary model inputs for each year are the populations of vehicles in the relevant categories, their annual kilometres of travel in three selected speed classes (urban, rural and highway), total fuel amounts and the fuel specifications. The numbers of vehicles are taken from annual bulletins of vehicle and driver statistics (e.g. DTTAS, 2022) and these are allocated to the different control technologies (PRE ECE; ECE 15/00-01; ECE 15/02; ECE 15/03; ECE 15/04; Euro 1; Euro 2; Euro 3; Euro 4; Euro 5; Euro 6 a/b/c, Euro 6 d-temp & Euro 6 d) based on their age and the application dates of the controls in Ireland. Information to assign values of annual kilometres of travel for the three speed classes (corresponding to urban, rural and highway driving) used for the individual vehicle categories is taken from the National Roads Authority (NRA) and odometer records from the National Car Testing (NCT) and the Commercial Vehicle Roadworthiness Test (CVRT) Service. The following methodology was applied:

#### Step 1

- Balanced mileage linked across from last year's file for 1990-2022
- Mileage data is sourced from national testing centres and processed for all passenger cars, light goods vehicles & heavy goods vehicles so that average yearly mileages are allocated per engine size and euro class.
- Bus & coach annual mileage is sourced from national bus operator's annual publication or if unavailable, extrapolated subject to the fuel increase / decrease. The motorcycle and moped mileage is sourced from the national transport bulletin produced each year.

#### Step 2

- COPERT model ran unbalanced
- The differences in petrol are spread across all petrol vehicles i.e. the adjusted mileage was multiplied by the excess / deficit in fuel



- The diesel car mean mileages were left untouched whilst the excess or lower fuel is distributed amongst the commercial fleet to keep the passenger car fleet annual mileage as close as possible to the reported figures.

### Step 3

- Once the statistical fuel input is within approx. 0.5 per cent of the fuel output (energy consumption), the model can be run balanced.

COPERT 5.7.1. was then run in “energy balance” mode and in turn the small difference between the statistical quantities input and the energy consumption values was balanced.

The emissions are then estimated in COPERT 5, fuel balancing is undertaken in the model using inbuilt consumption rates for the different vehicle categories to ensure that the total fuel amounts calculated from annual kilometres and consumption rates for all vehicle categories match the input fuel quantities (the fuel sold as given in the energy balance or the fuel used as determined by Annex A.3) and, consequently, emissions relate to the relevant total fuel amount. This fuel balancing may involve some adjustment to kilometres travelled or to the shares of total kilometres in the different road classes.

#### 3.6.2.1 Trends in emissions in Road Transport

Liquid fuel consumption increased until it peaked in 2007 after which it declined until 2012 with a subsequent return to growth in emissions thereafter until 2016. The trends appear to follow Ireland’s economic growth patterns, whereas the from 2016 to 2019 emissions stabilised despite increases in vehicle population. This is a result of both improved vehicle technologies and increased biofuel penetration. In 2021 there was an increase in emissions of air pollutants due to the easing of restrictions on passenger car journeys in 2020 in response to the COVID-19 pandemic, proceeded by a small decline in emissions in 2022, followed by a return to 2021 levels in 2023.

The three most influential parameters for calculating air pollutant emissions include fleet, fuel inputs and allocation of mileage per technology type. Overall, the fleet population has increased by 181.2 per cent from 1990-2023 and diesel passenger cars in particular have exponentially increased by 1,604.8 per cent in the same period. The energy content (TJ) for all combustion fuels has more than doubled from 1990-2023. Biofuel incorporated into the mix of available fuel types in 2005 as a measure to reduce road transport emissions has increased from 45.98 TJ in 2005 to 12,385 TJ in 2023. In addition, overall mileage has more than doubled from 21.5 billion in 1990 to 54.9 billion in 2023.

NO<sub>x</sub> emissions rose from 55.9 kt in 1990 to a peak in 2000 of 66.1 kt because of increasing diesel cars and overall fleet population. These emissions remained relatively stable until 2007 but thereafter have consistently declined from 62.3 kt in 2007 to 25.0 kt in 2023. The introduction of improved technologies and the rollover of newer euro classes was a major contributing factor. Despite an increase in activity in HDVs, NO<sub>x</sub> emissions from HDVs have decreased in 2015-2023 mainly due to the move from older technologies into the newer Euro IV-VI classes for HDV which have improved emissions standards and lowered emissions.

PM<sub>2.5</sub> and PM<sub>10</sub> follow a similar trend throughout the timeseries. A best-fit line observation indicates a steady increase in emissions from 1990-2000, relatively stable until 2007, followed by a steady decline in emissions until 2023. The initial increase during the 1990s was due to an increase in the fleet and activity. 2000 to 2007 emissions remained relatively stable as the impact of an increasing fleet and activity within a growing economy was being offset by newer and cleaner technologies, ultimately restricting an upward trend. With the advancement of newer euro classes outweighing the impact of an increasing diesel fleet, emissions continued to decline thereafter until 2023, with 2021 the low point due to Covid related travel restrictions throughout the country.

SO<sub>2</sub> has seen significant reductions from a high in 1990 of 5.4 kt to 0.14 kt in 2008 because of reduced sulphur content within the fuels. From 2009-2022 there was a further decline, albeit relatively small in comparison to previous years i.e., 0.05 kt in 2009 to 0.04 kt in 2023. SO<sub>2</sub> emissions increased to 0.041 kt in 2021 due to an increase in diesel use as pandemic-related road travel restrictions lessened across the country. There was a further slight increase in 2022 to 0.043 kt as travel rebounds close to pre-covid levels. NMVOC emissions have fallen steadily from 32.8 kt in 1990 to 2.7 kt in 2023. CO emissions have decreased from 246.2 kt in 1990 to 16.8 kt in 2023.

In contrast, NH<sub>3</sub> emissions have increased from 0.04 kt in 1990 to 0.54 kt in 2023, which is over a 10-fold increase. NH<sub>3</sub> emissions largely arise from hot emissions and as a result are dependent on the mileage of each vehicle category. NH<sub>3</sub> emissions did rise steadily from 1990 up to 2008 coinciding with increased mileage. NH<sub>3</sub> emissions have been declining in recent years because of improved technologies and the removal of older vehicles from the national car fleet. Fluctuations in NH<sub>3</sub> emissions for Heavy Duty Vehicle (HDV) is due to an increase in mileage by stock from 1996-2000 of 20 per cent on average over these four years. This level of activity was maintained at this level from 2001-2007 after which there was a decrease from 2008-2012 of 12 per cent on average for these four years due to the economic recession in Ireland. As a result, emissions fluctuated with the increasing and decreasing HDV activity. A recovery can be seen in 2013, 2014, 2015 and 2016 where there was an increase in HDV mileage x stock (4 per cent, 3 per cent, 8 per cent and 11 per cent respectively).

### 3.6.3 Railways (NFR 1A3c)

The emissions under sub-category 1A3c Railways are calculated from the amounts of fuel used by these activities and the country specific SO<sub>2</sub> emission factors and default Inventory Guidebook (EMEP/EEA, 2023) emission factors for gasoil for all other pollutants. No solid fuels have been used in railways since 1970. All emission factors for railways are given in Table C.20 of Annex C.

### 3.6.4 National Navigation (NFR 1A3dii)

The emissions NO<sub>x</sub> and PM are estimated using a Tier 2 approach and emission factors from Table 3-5 & 3-7 of the Inventory Guidebook (EMEP/EEA, 2023). All other pollutants are estimated using default emission factors from Table 3-1 for bunker Fuel Oil and Table 3-2 for Marine Diesel of the Inventory Guidebook (EMEP/EEA, 2023). All emission factors for national navigation are given in Table C.20 of Annex C.

### 3.6.5 Other Transportation (NFR 1A3e)

The emissions reported in sub-category 1A3e Other Transportation refer to the use of natural gas in pipeline compressor stations and emission factors (country-specific and default values) for this sub-category are given in Table C.21 of Annex C. The fuel use is estimated as the difference between the value given for natural gas under own use/losses in the energy balance sheets (Annex B) and the amount of gas estimated to be lost from the distribution network.

## 3.7 Other Sectors (NFR 1A4)

The NFR Subcategory 1A4 Other Sectors covers combustion sources in the Commercial/Institutional (1A4a), the Residential (1A4b), and Agriculture/Forestry/Fishing (1A4c) sectors. The Residential sector remains the most important source of emissions for most substances in this subcategory in Ireland, while Agriculture/Forestry/Fishing is a major contributor to NO<sub>x</sub> emissions, largely because of the influence of agricultural (off-road) machinery. The activity data for all 1A4 subcategories are taken directly from the energy balance. As there are no fuels allocated to the sector 1A4aii Commercial/Institutional: Mobile or 1A4bii Residential: Household and Gardening (Mobile) in national energy statistics, it is accounted for in category 1A3b Road Transport where all gasoline fuel is fully accounted. Sector Residential (1A4b) uses Tier 2 emission factors from the Inventory guidebook (EMEP/EEA, 2023) for all fuels. The Tier 2 methodology uses an assumption for solid fuel fireplace and stove use. The split of stove/fireplace usage is based EPA and CSO data when available and on expert opinion. These splits and the updated Tier 2 emission factors for 1A4b are referenced in Tables C.24 and C.25 of Annex C. The emission factors for sub-category 1A4 Other Sectors are given in Tables C.22 through C.28 of Annex C.

### 3.7.1 Commercial / Institutional (NFR 1A4a) and Residential (1A4b)

Combustion in the residential sector is one of the principal sources of emissions, contributing 65.2 per cent of SO<sub>2</sub> emissions, 59.8 per cent of CO emissions, and 58.2 per cent of PM<sub>2.5</sub> of emissions in Ireland in 2023. This sector also contributed 14.4 per cent and 24.6 per cent of Irelands' Pb and Hg 2023 emissions, respectively. This sector is also one of the principal sources of emissions of Dioxins and PAHs contributing 85.7 per cent and 90.6 per cent, respectively, in 2023. Coal burning in the residential sector is also an important but declining source of NMVOC emissions as coal consumption decreases, with the sector accounting 5.8 per cent of national emissions in 2023. Significant fuel switching has been seen in the residential sector, with a continuous reduction in the use of coal (down 84.5 per cent) and peat (down 80.7 per cent) since 1990. Increased use of liquid fuels gasoil and kerosene (229.6 per cent), natural gas (296.4 per cent), and renewables (148.3 per cent) in this sector has resulted in significant decreases in emissions since 1990. Between 1990 and 2023, decreases have been seen in SO<sub>2</sub> emissions (87.1 per cent), NMVOC emissions (81.0 per cent), CO (81.7 per cent), PM<sub>2.5</sub> (78.8 per cent), Pb (82.1 per cent), Hg (74.0 per cent), Dioxins (74.3 per cent) and PAHs (66.8 per cent) in the Commercial/Institutional and Residential sectors.

### 3.7.2 Agriculture/Forestry/Fishing (NFR 1A4ci and ii)

This subcategory covers both stationary combustion (1A4ci) and mobile combustion (1A4cii) in Agriculture. The national energy balance does not currently provide information on the end use of gasoil in agriculture. However, based on information received from agricultural experts, 90 per cent of total gasoil in the sector is assigned to mobile machinery with the remainder assigned to stationary

combustion. For both stationary and mobile combustion, the emission factors used are presented in Table C.26 of Annex C.

Ireland utilises the Tier 2 approach outlined in the Inventory Guidebook (EMEP/EEA, 2023) to estimate emissions of NO<sub>x</sub>, NMVOC, CO, PM<sub>10</sub>, PM<sub>2.5</sub> and TSP from mobile combustion in this sector. The Tier 2 emission factors are based on data from the Danish Inventory (Winther and Nielsen, 2006). The emission factors are grouped according to the EU emission legislation stages, and three additional layers are added to cover the emissions from engines prior to the first EU legislation stages. The country specific SO<sub>x</sub> emission factors and default Inventory Guidebook (EMEP/EEA, 2023) emission factors for NH<sub>3</sub> and BC are used for this sub-category. The implied emission factors are presented in Table C.27 of Annex C. Emission Factors for Heavy Metals and POPs are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2009, 2023). For category 1A4ci Agriculture/Forestry/Fishing: Stationary, the NO<sub>x</sub> Tier 1 liquid fuel emission factors from the Inventory Guidebook (EMEP/EEA, 2023) are an average of Tier 2 EFs and are not considered representative for Ireland for the same reasons as described in section 3.5.

### 3.7.3 Agriculture/Forestry/Fishing: National Fishing (NFR 1A4ciii)

Emissions from this sub-category were reported for the first time in the 2013 submission. The national energy balance now includes marine diesel used in national fishing for all years from 1990 to 2023. The emission factors used for this sub-category are country specific (SO<sub>x</sub>) and Inventory Guidebook (EMEP/EEA, 2023) default values for all other pollutants and are presented in Table C.28 of Annex C.

## 3.8 Other Stationary and Mobile (including military) (NFR 1A5)

Ireland does not report any emissions under category 1A5. For NFR categories 1A5a and 1A5b the notation key IE is used to indicate that military emissions (mobile and stationary) are included elsewhere as emissions are allocated in 1A4a and 1A3 (aviation and road). All stationary emissions from military are reported in sector 1A4a and all mobile military emission are reported under 1A3 (aviation and road).

## 3.9 Fugitive emissions from Solid Fuels (NFR 1B1)

### 3.9.1 Coal mining and handling (1B1a)

Emissions from the NFR subcategory 1B1a Coal mining are reported in this submission. The national energy balance includes coal mined in the years 1990 to 1995 when the last commercial coal mine was closed in Ireland. The emission factors used and the resulting time series of NMVOC, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions are based on the Inventory Guidebook (EMEP/EEA, 2023) default values and are presented in Table 3.1.

Emissions of PM<sub>2.5</sub>, PM<sub>10</sub> and TSP from the handling and storage of coal other than domestic production has been estimated using an estimate of the area of coal storage and tonnes of coal handled in Ireland's only coal burning power plant and emission factors from the Inventory Guidebook (EMEP/EEA, 2023).

**Table 3.1. Fugitive Emissions from Coal mining and handling**

Activity Data		1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Coal mining	kt	25.00	1.00	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Area of coal storage	ha	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20
Coal handling	kt	1,867	2,248	2,145	2,130	1,305	1,373	1,745	1,459	1,414	1,693	1,658	1,304	736	222	293	293	848	473
<b>Emission Factors</b>																			
Coal mining	kg NMVOC/t	0.80	0.80	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal mining	kg TSP/t	0.09	0.09	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal mining	kg PM <sub>10</sub> /t	0.04	0.04	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal mining	kg PM <sub>2.5</sub> /t	0.01	0.01	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal storage	t PM <sub>2.5</sub> /ha	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.04	0.04	0.04	0.04	0.04	0.04	0.04
Coal storage	t PM <sub>10</sub> /ha	4.10	4.10	4.10	4.10	4.10	4.10	4.10	4.10	4.10	4.10	4.10	0.41	0.41	0.41	0.41	0.41	0.41	0.41
Coal storage	t TSP/ha	10.25	10.25	10.25	10.25	10.25	10.25	10.25	10.25	10.25	10.25	10.25	1.03	1.03	1.03	1.03	1.03	1.03	1.03
Coal handling	g PM <sub>2.5</sub> /t	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Coal handling	g PM <sub>10</sub> /t	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00
Coal handling	g TSP/t	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50
<b>Emissions</b>																			
Coal mining	kt NMVOC	0.020	0.001	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal mining	kt TSP	0.002	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal mining	kt PM <sub>10</sub>	0.001	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal mining	kt PM <sub>2.5</sub>	0.000	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal storage	kt PM <sub>2.5</sub>	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.001	0.001	0.001	0.001	0.001	0.001	0.001
Coal storage	kt PM <sub>10</sub>	0.128	0.128	0.128	0.128	0.128	0.128	0.128	0.128	0.128	0.128	0.128	0.013	0.013	0.013	0.013	0.013	0.013	0.013
Coal storage	kt TSP	0.320	0.320	0.320	0.320	0.320	0.320	0.320	0.320	0.320	0.320	0.320	0.032	0.032	0.032	0.032	0.032	0.032	0.032
Coal handling	kt PM <sub>2.5</sub>	0.001	0.001	0.001	0.001	0.000	0.000	0.001	0.000	0.000	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Coal handling	kt PM <sub>10</sub>	0.006	0.007	0.006	0.006	0.004	0.004	0.005	0.004	0.004	0.005	0.005	0.004	0.002	0.001	0.001	0.001	0.003	0.001
Coal handling	kt TSP	0.014	0.017	0.016	0.016	0.010	0.010	0.013	0.011	0.011	0.013	0.012	0.010	0.006	0.002	0.002	0.002	0.006	0.004

## 3.10 Fugitive emissions from Oil and Natural Gas (NFR 1B2)

### 3.10.1 Oil (NFR 1B2a)

The NFR subcategory 1B2a is an important source of fugitive NMVOC emissions. Emissions of NMVOCs are estimated from two sources, Refining/Storage (1B2aiv) and Distribution of Oil Products (1B2av). Emissions from Refining/Storage are estimated using a Tier 3 methodology using plant specific data for years 2007-2023 from Ireland's only refinery. An implied emission factor of 0.8812 kg/t is used for years before 2007 using an average of 2007-2016 implied emission factors and the crude oil throughput for the refinery.

The emissions from 1B2av, were estimated using a Tier 2 approach using emission factors and abatement efficiencies from the Inventory Guidebook (EMEP/EEA, 2023). The Stage I and Stage II abatement controls were applied as appropriate to the implementation of legislation in Ireland. The resulting time series of NMVOC emissions are presented in Table 3.2. Emissions of NMVOCs from these subcategories nearly doubled in the period from 1990 to 2007 in line with the increases in crude oil throughput for the refinery and petrol distribution for the transport sector but have since decreased due to the economic situation and the implementation of abatement controls in petrol distribution causing an overall decrease in most emissions. An increase in emissions in Refining/Storage source can be seen from 2014 to 2018 before decreasing again from 2019 to 2023 with such changes responding correlatively due to increased and decreased refinery crude oil throughput.

The refinery responded to a previous query from the inventory agency that the only fugitive emissions to air are for NMVOCs (emissions that occur through leakage, evaporation or other uncontrolled losses). The estimate is based on throughput, mass balance and industry (e.g. API) guideline emissions factors at the plant, tankage, jetty, road-loading and Wastewater Treatment Plant. Emissions from flaring are included in the totals for the other pollutants, as site emissions are calculated based on total metered fuel use. Therefore, reporting under 1A1b is complete for the plant and there should be no additional reporting under 1B2aiv except for NMVOC. The notation key "IE" included elsewhere is used for 1B2aiv for Cd, Hg and PCDD/F as all emissions from refineries including flaring are reported under category 1A1b. No heavy metal or dioxins to air are reported under LCP/PRTR/IPPC returns to the EPA in Ireland.

### 3.10.2 Natural Gas (NFR 1B2b)

Fugitive losses of NMVOC from pipelines and distribution systems were estimated for the first time in the 2021 submission. The methodology is continued for this the 2025 submission. These emissions are based on activity data of fugitive methane releases and gas analysis reports which in combination provide gas composition supplied by Gas Networks Ireland (GNI) in the greenhouse gas emission inventory. The description of this methodology is in section 3.3.2.2.3 of Ireland's NID (EPA, 2025). Ireland has used existing information about leakage from greenhouse gas fugitive emission reporting to the UNFCCC to estimate NMVOC emissions under this source category.

**Table 3.2. Fugitive Non-Methane Volatile Organic Compound Emissions**

Activity Data		1990	1995	2000	2005	2010	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Crude to refineries	kt	1,804	2,229	3,278	3,309	2,905	3,068	2,838	2,752	3,366	3,200	3,181	3,024	2,531	2,838	3,025	2,983	2,778
Petrol distribution	kt	885	1,037	1,493	1,711	1,387	1,195	1,124	1,064	1,009	942	849	774	733	543	575	655	698
<b>Emission Factors</b>																		
Crude to refineries	kg NMVOC/t	0.88	0.88	0.88	0.88	0.92	0.81	0.86	0.95	0.83	0.85	0.85	0.90	0.89	0.79	0.77	0.77	0.49
Petrol distribution	kg NMVOC/t	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.001	0.002	0.001
<b>Emissions</b>																		
Crude to refineries	kt NMVOC	1.59	1.96	2.89	2.92	2.66	2.48	2.45	2.62	2.80	2.73	2.72	2.73	2.25	2.25	2.32	2.30	1.36
Petrol distribution	kt NMVOC	1.79	2.10	3.30	3.81	3.15	2.60	2.22	2.12	1.95	1.75	1.77	1.37	1.30	1.00	0.83	1.11	1.02
Total emissions	kt NMVOC	3.37	4.06	6.18	6.72	5.80	5.08	4.67	4.74	4.75	4.49	4.49	4.10	3.56	3.25	3.15	3.41	2.38

### 3.10.3 Venting and flaring (oil, gas, combined oil and gas) (NFR1B2c)

Regarding the flares at refineries, all emissions have been included within the category 1A1b Petroleum Refining, as site emissions were calculated based on total metered fuel use.

## 3.11 Recalculations in the Energy Sector

The results of recalculations for the combustion categories 1A1, 1A2, 1A3 and 1A4 are given in Tables 3.3 through 3.7 below.

### *1A1a Public Electricity and Heat Production*

There were no recalculations to emission inventory from Public Electricity and Heat Production refining in this submission.

### *1A1b Petroleum refining*

There were no recalculations to emission inventory from petroleum refining in this submission.

### *1A1c Manufacture of Solid Fuels and Other Energy Industries*

There were recalculations in this sector for reporting year 2022 as a result of updated activity data which reduced milled peat for this year by 5.9 per cent. As a result, recalculations for NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, Pb, Cd and Hg ranged from -3.8 per cent to -5.6 per cent.

### *1A2 Manufacturing Industry and Construction sector*

In this submission, there are recalculations across all 1A2 'Manufacturing Industry and Construction' subsectors based on revisions to the historical energy balances going back as far as 1990. This is because the national energy balance provided by SEAI now incorporates a major data set on business energy use. This Business Energy Use Survey (BEUS), was first published by the Central Statistics Office (CSO) in December 2018. This valuable data source provides a new basis for the breakdown of energy use in the commercial services, public services and industrial sectors, at a level of detail not previously available. SEAI have revised the National Energy Balances from 1990 to 2022 incorporating this new improved data. In some cases, the revisions to estimates of business energy use have had knock on effects in other sectors, particularly the residential sector, leading to revisions there also.

There was a redistribution of fuel within this sector in the 2025 submission compared to the 2024 submission. Across the period 2001 to 2021 there was an average of 245 TJ of natural gas reallocated to 1A4 Other sectors except for 2020 when 37 TJ was added to 1A2 Manufacturing Industries and Construction.

There was a reallocation of oil approximately 196 TJ per annum across the time series to 1A2 from 1A4 other sectors with the largest of these additions occurring in 2022 when 885 TJ was added. On average, there is 229 TJ less fuel consumed per annum in the sector throughout the time series compared to the previous submission. The changes in 2022 in 1A2 were predominantly driven by 1A2b Non-Ferrous Metals and 1A2f non-metallic minerals across the major air pollutants.

### *1A2a Iron and steel*

There were no recalculations to emission inventory from Iron and steel in this submission.

### *1A2b Non-ferrous metals*



There were fuel reallocations across the time series 1991-2022 for oil averaging 0.11 ktoe with the largest reallocation recorded in 2022 with 0.55 ktoe. In addition, there was a total of 47.44 ktoe or annual average of 2.37 ktoe natural gas removed across the period 2001 – 2020, with 3.15 ktoe decrease on average from 2006 -2020.

There were also fuel reallocations in gasoil/diesel and DERV resulting in 3.32 ktoe being added with largest amounts added 0.1 and 0.29 ktoe in 2020 and 2022 respectively. Specifically, there were additions across the periods 1991-2009 and the period 2020- 2022 with the average added over these periods was 0.15 ktoe. The largest addition was recorded for the 2022 energy balance with a 0.29 ktoe addition. The fuel reallocations resulted in recalculations to NO<sub>x</sub> 7 per cent, SO<sub>2</sub> 4.7 per cent and 3.1 per cent increase in NMVOC and CO for 2022.

#### *1A2c Chemicals*

For the period 1991-2022 there were fuel reallocations in oil for the sub sector totalling 8.06 ktoe averaging 0.25 ktoe annually with the largest increase recorded at 1.29 ktoe in 2022. In addition, there was a reallocation of natural gas across the period 2001- 2022 totalling -14.43 ktoe with the largest reallocation occurring in 2022 with a decrease of 11.39 ktoe.

These updates in fuel allocations in the energy balance produced recalculations of NO<sub>x</sub> -6.6 per cent, SO<sub>2</sub> -2.7 per cent, NMVOC -4.7 per cent and CO -4.5 per cent for reporting year 2022.

#### *1A2d Pulp, Paper, and Print*

2022 saw recalculations due to reallocations involving removal of -1.37 ktoe of natural gas offset by a 0.14 ktoe addition of oil to this sub sector. There was on average 0.15 ktoe of natural gas removed from the sector from 2000 to 2022 each year with 2021 being an exception where 0.04 ktoe was added, causing recalculations across the time series.

Annual average recalculations for NO<sub>x</sub> -1.8 per cent, SO<sub>2</sub> -0.32 per cent, NMVOC 0.5 per cent, TSP 1.9 per cent, PM<sub>2.5</sub> -72.9 per cent, PM<sub>10</sub> 0.4per cent and BC -1.6 per cent on average across the time series. Heavy metals emissions were also recalculated, Pb -3.1 per cent and Ni -1.7per cent on average across the time period.

#### *1A2e Food Processing, Beverages, and Tobacco*

This sub category saw significant recalculations due to the inclusion of 5 - 6 ktoe of solid biomass in 2021 & 2022 when there was none in the 2024 submission.

There was on average 0.75 ktoe of oil per annum reallocated into the sector from 1991 to 2020 and on average 6.49 ktoe of oil added annually from 2021 to 2022 including 11.16 ktoe for reporting year 2022. Additionally, there was on average 1.7 ktoe of natural gas reallocated out of the sector from 2001 to 2022, generally increasing across the time series to a maximum of 7.58 ktoe taken out of the sector in 2022.

There were further reallocations for LPG over the period 1991-2022 totalling an increase of 5.91 ktoe, an average of 0.18 ktoe annually was added with the largest increase at 3.84 ktoe across the period occurring for reporting year 2022. In addition, there was an increase of 27.82 ktoe from 1991- 2022, an average of 0.87 ktoe with the largest addition at 6.64 ktoe across the period occurring for reporting year 2022.

In terms of emissions, there were recalculations for reporting year 2022 -0.5 per cent NO<sub>x</sub>, SO<sub>2</sub> -0.1 per cent, NMVOC -3.4 per cent, TSP -0.7 per cent, PM 2.5 -52.7 per cent and PM 10 -0.9 per cent.

#### *1A2f Non-Metallic Minerals*

There were fuel reallocations involving oil, natural gas, LPG and Gasoil/Diesel/DERV over the time period 1991-2022. In oil there was an addition of 23.58 ktoe into the sub sector, averaging 0.74 ktoe with the largest increase of 6.35 ktoe for the reporting year 2022. In natural gas there was a deduction totalling 6.93 ktoe from 2001-2022, averaging -0.32 ktoe over this period with the largest deduction -1.28 ktoe occurring in 2022.

Reallocations to gasoil/diesel/DERV over the period totalling an increase of 21.88 ktoe, an average of 0.68 ktoe with the largest increase 4.87 ktoe occurring in 2022.

This resulted in the following recalculations: <sup>SO<sub>2</sub></sup> 10.3 per cent, NMVOC -0.3 per cent, CO -0.3 per cent, TSP -21.4 per cent, PM<sub>2.5</sub> -1.4 per cent and PM<sub>10</sub> -0.7 per cent.

#### *1A2g Other*

There were fuel reallocations in oil, natural gas, LPG, gasoil/diesel/DERV, renewables and biomass. In oil there was an addition of 82.30 ktoe across the period 1990-2022, averaging 2.5 ktoe annually with largest of the increases occurring in the period 2009-2013 with an additional 32.21 ktoe. There was deductions for natural gas across the period 2001- 2022 totalling 28.78 ktoe, averaging 1.31 ktoe decrease annually with the largest of these deductions at 19.80 ktoe occurring in the period 2017-2020.

For gasoil/diesel/DERV there were additions totalling 88.69 ktoe across the period 1990-2022, averaging 2.69 ktoe with the largest of these additions at 32.12 ktoe occurring in the period 2009-2012. In LPG there were deductions totalling 6.96 ktoe across the period 1990-2022, averaging 0.21 ktoe annually with the largest decrease of 4.00 ktoe in 2022. In renewables and biomass there were reallocations relating to reporting year 2022 where 9.59 ktoe was removed.

This resulted in the following recalculations: NO<sub>x</sub> 2.7 per cent, SO<sub>2</sub> 0.4 per cent, NMVOC 3.6 per cent, TSP 0.3 per cent, PM 2.5 3.3 per cent and PM 10 0.2 per cent.

#### *1A3a (i) & (ii) International and Domestic Aviation*

There was a recalculation to emission inventory for reporting year 2022 on NMVOC 1.4 per cent versus reported inventory figures for 2018 to 2022 due to revised NMWVOC emission factors for both international and domestic flights. The average change to reported figures was 0.5 per cent and added 0.009 ktoe of NMVOC in total to the reporting period 2018-2022.

#### *1A3b Road Transport*

There was a recalculation to emission inventory road transport for reporting year 2018 on Pb -1.5 per cent versus reported inventory figure in 2022 due to revision in industry data received and incorporated for Pb content in fuel.

#### *1A3c Railways*

There were no recalculations to emissions from railways in this submission.

### *1A3dii National navigation*

There were no recalculations to emissions from national navigation in this submission.

### *1A4a Commercial/Institutional*

In this submission there are recalculations arising from a reallocation of oil, kerosene, LPG and gasoil/diesel/DERV. In oil across the time period 1990-2022 there was an overall decrease of 128.25 ktoe, averaging a decrease of 3.89 ktoe annually with the largest decrease occurring in 2019. In LPG there was a decrease of 2.79 ktoe across the same time period, averaging an annual decrease of 0.08 ktoe with the largest decrease occurring in reporting year 2022. For gasoil/diesel/DERV across the same time period there was an overall decrease of 58.75 ktoe, averaging a 1.78 ktoe decrease annually with the largest decrease 6.38 ktoe recorded in 2022.

There was a further reallocation in kerosene totalling a decrease of 77.52 ktoe over the reporting period 1991-2020, averaging a 2.58 ktoe decrease annually with the largest decrease 7.25 ktoe occurring in 2019. However, this decreasing trend was offset by 2021 and 2022 when 10.80 ktoe was added in these years. In natural gas there was a reallocation increasing natural gas by a total of 121.84 ktoe across the time period 2001-2020 averaging an annual increase of 5.54 ktoe with the largest increases occurring over the period 2018-2020 totalling 37.35 ktoe.

This resulted in the following recalculations across the 1990-2020 time period: NO<sub>x</sub> +0.4 per cent, SO<sub>2</sub> -1.2 per cent, NMVOC +1.1 per cent, NH<sub>3</sub> +1.8 per cent, PM<sub>2.5</sub> -1.2 per cent and Pb -1 per cent.

### *1A4b Residential*

In this submission there were minor recalculations arising from reallocations of oil, kerosene, gasoil/diesel/DERV, renewables and biomass. In oil there was an overall decrease of 17.71 ktoe, averaging a decrease of 0.53 ktoe annually over the time period 1990-2020 with the largest decrease of 11.517 ktoe in 2022. In kerosene there was an overall decrease of 64.58 ktoe, averaging a decrease of 1.96 ktoe annually with the largest decrease of 9.22 ktoe in 2022.

Across the time period 1991-2020 there was a decrease of gasoil/diesel/DERV totalling 82.08 ktoe, averaging a decrease of 2.57 ktoe with the largest decreases totalling 76.49 ktoe across the period of 1999-2012. In renewables there was a decrease totalling 47.97 ktoe across the time period 2012-2022, averaging 4.36 ktoe annually with the largest decrease of 17.99 ktoe in 2022. For biomass over the same period there was an increase of 6.88 ktoe, averaging 0.63 ktoe annually with the largest decrease of 1.79 ktoe in 2021

This has resulted in the following recalculations across the 1990-2020 time period: NH<sub>3</sub> 0.4 per cent and Cd 0.5 per cent.

### *1A4c Agriculture/Forestry/Fishing*

In this submission there were minor recalculations to the period 2014 to 2022 arising from reduction of gasoil/diesel/DERV resulting in the following changes over the same period: PM<sub>2.5</sub> -0.2 per cent and Cd -0.3 per cent and

Overall recalculations are presented in Tables 3.3-3.7 below

**Table 3.3. Recalculations in Energy NFR 1A1**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
1A1a Public Electricity and Heat Production	NO <sub>x</sub>	kt	46.37	41.39	39.72	32.38	11.92	8.37	10.53	9.09	7.81	9.82	8.31	8.12	6.74	5.99	5.59	8.53	7.48
	SO <sub>2</sub>	kt	103.04	91.63	79.87	42.50	9.78	9.73	9.85	8.95	6.20	5.51	3.96	3.98	2.84	2.24	2.00	3.06	1.86
	NMVOC	kt	0.19	0.25	0.36	0.37	0.37	0.31	0.30	0.27	0.26	0.26	0.30	0.30	0.29	0.29	0.29	0.30	0.32
	Pb	t	0.82	0.93	0.93	0.89	0.61	0.61	0.75	0.67	0.70	0.75	0.76	0.67	0.56	0.44	0.34	0.45	0.39
	Cd	t	0.11	0.13	0.14	0.13	0.08	0.07	0.09	0.08	0.08	0.09	0.09	0.08	0.06	0.05	0.04	0.06	0.05
	Hg	t	0.15	0.17	0.16	0.16	0.13	0.12	0.15	0.13	0.13	0.15	0.15	0.14	0.10	0.08	0.06	0.07	0.06
1A1b Petroleum Refining	NO <sub>x</sub>	kt	0.47	0.52	0.77	0.94	0.80	0.63	0.68	0.58	0.54	0.38	0.32	0.26	0.33	0.29	0.29	0.31	0.28
	SO <sub>2</sub>	kt	0.48	0.51	0.78	0.91	0.67	0.90	0.43	0.57	0.01	0.03	0.02	0.03	0.03	0.03	0.04	0.04	0.04
	NMVOC	kt	0.01	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Pb	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Cd	t	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1c Manufacture of Solid Fuels and Other Energy Industries	NO <sub>x</sub>	kt	0.16	0.12	0.15	0.29	0.32	0.21	0.18	0.22	0.22	0.24	0.32	0.11	0.53	0.21	0.18	0.17	0.16
	SO <sub>2</sub>	kt	0.13	0.10	0.12	0.16	0.21	0.25	0.24	0.19	0.03	0.09	0.25	0.11	0.37	0.13	0.14	0.14	0.12
	NMVOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Pb	t	0.02	0.02	0.02	0.03	0.03	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.02	0.02	0.02	0.03	0.03	0.02	0.03	0.03	0.03	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02
<b>Submission 2025</b>																			
1A1a Public Electricity and Heat Production	NO <sub>x</sub>	kt	46.37	41.39	39.72	32.38	11.92	8.37	10.53	9.09	7.81	9.82	8.31	8.12	6.74	5.99	5.59	8.53	7.48
	SO <sub>2</sub>	kt	103.04	91.63	79.87	42.50	9.78	9.73	9.85	8.95	6.20	5.51	3.96	3.98	2.84	2.24	2.00	3.06	1.86
	NMVOC	kt	0.19	0.25	0.36	0.37	0.37	0.31	0.30	0.27	0.26	0.26	0.30	0.30	0.29	0.29	0.29	0.30	0.32
	Pb	t	0.82	0.93	0.93	0.89	0.61	0.61	0.75	0.67	0.70	0.75	0.76	0.67	0.56	0.44	0.34	0.45	0.39
	Cd	t	0.11	0.13	0.14	0.13	0.08	0.07	0.09	0.08	0.08	0.09	0.09	0.08	0.06	0.05	0.04	0.06	0.05
	Hg	t	0.15	0.17	0.16	0.16	0.13	0.12	0.15	0.13	0.13	0.15	0.15	0.14	0.10	0.08	0.06	0.07	0.06
1A1b Petroleum Refining	NO <sub>x</sub>	kt	0.47	0.52	0.77	0.94	0.80	0.63	0.68	0.58	0.54	0.38	0.32	0.26	0.33	0.29	0.29	0.31	0.28
	SO <sub>2</sub>	kt	0.48	0.51	0.78	0.91	0.67	0.90	0.43	0.57	0.01	0.03	0.02	0.03	0.03	0.03	0.04	0.04	0.04
	NMVOC	kt	0.01	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Pb	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Cd	t	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1c Manufacture of Solid Fuels and Other Energy Industries	NO <sub>x</sub>	kt	0.16	0.12	0.15	0.29	0.32	0.21	0.18	0.22	0.22	0.24	0.32	0.11	0.53	0.21	0.18	0.17	0.15
	SO <sub>2</sub>	kt	0.13	0.10	0.12	0.16	0.21	0.25	0.24	0.19	0.03	0.09	0.25	0.11	0.37	0.13	0.14	0.14	0.11
	NMVOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Pb	t	0.02	0.02	0.02	0.03	0.03	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.02	0.02	0.02	0.03	0.03	0.02	0.03	0.03	0.03	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02

**Table 3.3. Recalculations in Energy NFR 1A1 (continued)**

<b>% Change in Emissions</b>	<b>Pollutant</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2005</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>	<b>2022</b>
1A1a Public Electricity and Heat Production	NO <sub>x</sub>	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO <sub>2</sub>	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NMVOC	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
1A1b Petroleum Refining	NO <sub>x</sub>	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO <sub>2</sub>	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NMVOC	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
1A1c Manufacture of Solid Fuels and Other Energy Industries	NO <sub>x</sub>	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-5.4%
	SO <sub>2</sub>	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-5.6%
	NMVOC	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-3.8%
	Pb	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-5.4%
	Cd	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-5.5%
	Hg	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-5.6%

**Table 3.4. Recalculations in Energy NFR 1A2**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
1A2 Manufacturing Industries and Construction	NO <sub>x</sub>	kt	9.08	7.98	10.25	16.42	9.02	7.53	9.37	9.55	10.39	10.31	10.61	9.82	9.21	8.22	8.17	8.34	8.16
	SO <sub>2</sub>	kt	33.95	40.42	42.34	14.13	6.52	5.73	4.95	5.38	4.00	3.04	3.08	3.20	3.27	1.88	1.49	1.80	1.61
	NMVOC	kt	1.69	1.26	1.72	2.70	2.48	2.23	2.18	2.24	2.71	2.78	2.84	3.06	3.12	2.95	3.24	2.96	3.01
	Pb	t	1.20	0.84	1.08	1.15	0.64	0.51	0.47	0.48	0.48	0.49	0.45	0.46	0.46	0.33	0.33	0.32	0.32
	Cd	t	0.05	0.05	0.08	0.11	0.09	0.08	0.07	0.07	0.08	0.08	0.08	0.09	0.09	0.09	0.09	0.07	0.08
	Hg	t	0.08	0.05	0.11	0.15	0.09	0.08	0.09	0.07	0.06	0.12	0.11	0.14	0.11	0.11	0.09	0.11	0.09
<b>Submission 2025</b>																			
1A2 Manufacturing Industries and Construction	NO <sub>x</sub>	kt	9.08	7.99	10.27	16.45	9.04	7.53	9.38	9.56	10.38	10.31	10.61	9.82	9.18	8.21	8.15	8.38	8.03
	SO <sub>2</sub>	kt	33.95	40.43	42.36	14.15	6.54	5.74	4.96	5.39	4.00	3.05	3.10	3.20	3.28	1.89	1.50	1.81	1.53
	NMVOC	kt	1.69	1.26	1.72	2.70	2.48	2.23	2.18	2.27	2.76	2.86	2.92	3.13	3.18	3.03	3.29	3.04	2.98
	Pb	t	1.20	0.84	1.08	1.15	0.65	0.52	0.47	0.48	0.49	0.50	0.46	0.46	0.47	0.34	0.34	0.32	0.33
	Cd	t	0.05	0.05	0.08	0.11	0.09	0.08	0.07	0.07	0.08	0.09	0.08	0.09	0.09	0.09	0.09	0.08	0.07
	Hg	t	0.08	0.05	0.11	0.15	0.09	0.08	0.09	0.07	0.06	0.12	0.11	0.14	0.11	0.11	0.09	0.11	0.09
<b>% Change in Emissions</b>																			
1A2 Manufacturing Industries and Construction	NO <sub>x</sub>	%	0.0%	0.2%	0.2%	0.2%	0.2%	0.1%	0.1%	0.1%	-0.1%	0.0%	0.0%	-0.1%	-0.2%	-0.1%	-0.3%	0.5%	0.1%
	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.4%	0.4%	0.3%	0.3%	0.6%	0.7%	0.6%	0.1%
	NMVOC	%	0.0%	0.1%	0.1%	0.0%	0.0%	-0.2%	-0.1%	1.0%	1.9%	3.1%	2.8%	2.2%	1.9%	2.6%	1.8%	2.6%	0.5%
	Pb	%	0.0%	0.2%	0.4%	0.5%	0.9%	0.9%	1.0%	1.1%	1.1%	1.7%	1.7%	1.5%	1.4%	2.3%	2.3%	2.3%	0.7%
	Cd	%	0.0%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	1.7%	3.2%	4.8%	4.7%	3.7%	3.4%	4.1%	3.6%	4.4%	0.8%
	Hg	%	0.0%	0.0%	0.0%	0.0%	-0.1%	-0.2%	-0.1%	-0.1%	-0.1%	0.0%	0.0%	0.0%	-0.1%	-0.1%	-0.2%	0.2%	0.0%

**Table 3.5. Recalculations in Energy NFR 1.A.3 (a & b)\***

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
1A3a(i) & (ii) International and Domestic Aviation	NO <sub>x</sub>	kt	4.80	5.15	8.10	11.08	10.00	8.80	7.44	8.74	9.81	10.50	10.79	12.79	13.62	13.80	5.05	5.74	12.79
	SO <sub>2</sub>	kt	0.30	0.32	0.50	0.68	0.63	0.56	0.47	0.54	0.60	0.68	0.69	0.81	0.88	0.89	0.32	0.36	0.81
	NMVOOC	kt	0.24	0.24	0.37	0.43	0.48	0.42	0.38	0.41	0.42	0.48	0.49	0.51	0.52	0.52	0.16	0.17	0.43
1A3b Road Transportation	NO <sub>x</sub>	kt	55.90	56.90	66.13	62.14	42.59	40.64	38.63	40.02	41.27	41.81	41.20	39.33	38.29	34.59	26.76	25.43	25.17
	SO <sub>2</sub>	kt	5.37	5.13	1.66	0.54	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04
	NMVOOC	kt	38.83	36.18	25.72	18.25	11.58	10.63	9.65	9.09	8.54	7.92	7.06	5.95	5.24	4.67	3.99	3.75	3.82
	NH <sub>3</sub>	kt	0.04	0.41	1.67	2.45	1.49	1.33	1.15	1.04	0.95	0.87	0.78	0.66	0.59	0.54	0.43	0.45	0.51
	PM <sub>2.5</sub>	kt	2.09	2.42	3.45	3.15	2.30	2.17	2.02	2.04	2.04	2.01	1.88	1.67	1.58	1.49	1.20	1.21	1.26
	Pb	t	145.22	89.03	8.51	4.95	4.49	4.41	4.26	4.38	4.50	4.72	4.82	4.84	4.97	4.94	4.18	4.44	4.79
	Cd	t	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
	Hg	t	0.01	0.01	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
<b>Submission 2025</b>																			
1A3a(i) & (ii) International and Domestic Aviation	NO <sub>x</sub>	kt	4.80	5.15	8.10	11.08	10.00	8.80	7.44	8.74	9.81	10.50	10.79	12.79	13.62	13.79	5.05	5.74	12.76
	SO <sub>2</sub>	kt	0.30	0.32	0.50	0.68	0.63	0.56	0.47	0.54	0.60	0.68	0.69	0.81	0.88	0.89	0.32	0.36	0.81
	NMVOOC	kt	0.24	0.24	0.37	0.43	0.48	0.42	0.38	0.41	0.42	0.48	0.49	0.51	0.52	0.53	0.16	0.17	0.43
1A3b Road Transportation	NO <sub>x</sub>	kt	55.90	56.90	66.13	62.14	42.59	40.64	38.63	40.02	41.27	41.81	41.20	39.33	38.29	34.59	26.76	25.42	25.20
	SO <sub>2</sub>	kt	5.37	5.13	1.66	0.54	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04
	NMVOOC	kt	38.83	36.18	25.72	18.25	11.58	10.63	9.65	9.09	8.54	7.92	7.06	5.95	5.24	4.67	3.98	3.74	3.83
	NH <sub>3</sub>	kt	0.04	0.41	1.67	2.45	1.49	1.33	1.15	1.04	0.95	0.87	0.78	0.66	0.59	0.54	0.43	0.46	0.51
	PM <sub>2.5</sub>	kt	2.09	2.42	3.45	3.15	2.30	2.17	2.02	2.04	2.04	2.01	1.88	1.67	1.58	1.49	1.20	1.21	1.26
	Pb	t	145.22	89.03	8.51	4.95	4.49	4.41	4.26	4.38	4.50	4.72	4.82	4.84	4.90	4.94	4.18	4.44	4.79
	Cd	t	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
	Hg	t	0.01	0.01	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
<b>% Change in Emissions</b>																			
1A3a(i) & (ii) International and Domestic Aviation	NO <sub>x</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.1%	0.0%	-0.1%	-0.2%
	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NMVOOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.2%	0.5%	0.4%	0.4%	1.4%
1A3b Road Transportation	NO <sub>x</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NMVOOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.1%	-0.2%	-0.1%	0.0%
	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.2%	0.3%	0.0%
	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-1.5%	0.0%	0.0%	0.0%	-0.1%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

\* Emissions from both LTO and cruise are included here to capture all possible recalculations. However, the cruise component is reported as a memo item under the LRTAP Convention.

**Table 3.6. Recalculations in Energy NFR 1A3 (c, d & e)**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
1A3c Railways	NO <sub>x</sub>	kt	2.20	1.84	2.03	2.02	2.01	2.03	1.95	1.94	1.78	1.81	1.85	1.91	1.93	2.02	1.61	1.74	1.94
	SO <sub>2</sub>	kt	0.25	0.14	0.12	0.11	0.06	0.06	0.06	0.05	0.02	0.02	0.01	0.01	0.00	0.00	0.00	0.00	0.00
	NM VOC	kt	0.20	0.16	0.18	0.18	0.18	0.18	0.17	0.17	0.16	0.16	0.16	0.17	0.17	0.18	0.14	0.15	0.17
	PM <sub>2.5</sub>	kt	0.06	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.05	0.05
	Pb	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1A3d(ii) National Navigation	NO <sub>x</sub>	kt	2.24	2.42	3.76	4.42	3.48	3.01	3.20	3.07	3.85	3.79	4.56	4.00	4.42	4.77	5.94	6.33	5.28
	SO <sub>2</sub>	kt	1.16	1.26	1.22	0.67	0.10	0.08	0.09	0.08	0.05	0.03	0.03	0.09	0.11	0.11	0.13	0.14	0.12
	NM VOC	kt	0.05	0.05	0.08	0.11	0.11	0.09	0.10	0.10	0.12	0.12	0.15	0.13	0.14	0.15	0.18	0.20	0.17
	PM <sub>2.5</sub>	kt	0.09	0.10	0.13	0.12	0.05	0.05	0.05	0.05	0.06	0.06	0.07	0.06	0.07	0.07	0.09	0.10	0.08
	Pb	t	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3e(i) Pipeline Compressors	NO <sub>x</sub>	kt	0.06	0.11	0.05	0.13	0.14	0.13	0.12	0.12	0.12	0.12	0.12	0.11	0.12	0.12	0.13	0.13	0.13
<b>Submission 2025</b>																			
1A3c Railways	NO <sub>x</sub>	kt	2.20	1.84	2.03	2.02	2.01	2.03	1.95	1.94	1.78	1.81	1.85	1.91	1.93	2.02	1.61	1.74	1.94
	SO <sub>2</sub>	kt	0.25	0.14	0.12	0.11	0.06	0.06	0.06	0.05	0.02	0.02	0.01	0.01	0.00	0.00	0.00	0.00	0.00
	NM VOC	kt	0.20	0.16	0.18	0.18	0.18	0.18	0.17	0.17	0.16	0.16	0.16	0.17	0.17	0.18	0.14	0.15	0.17
	PM <sub>2.5</sub>	kt	0.06	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.05	0.05
	Pb	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1A3d(ii) National Navigation	NO <sub>x</sub>	kt	2.24	2.42	3.76	4.42	3.48	3.01	3.20	3.07	3.85	3.79	4.56	4.00	4.42	4.77	5.94	6.33	5.28
	SO <sub>2</sub>	kt	1.16	1.26	1.22	0.67	0.10	0.08	0.09	0.08	0.05	0.03	0.03	0.09	0.11	0.11	0.13	0.14	0.12
	NM VOC	kt	0.05	0.05	0.08	0.11	0.11	0.09	0.10	0.10	0.12	0.12	0.15	0.13	0.14	0.15	0.18	0.20	0.17
	PM <sub>2.5</sub>	kt	0.09	0.10	0.13	0.12	0.05	0.05	0.05	0.05	0.06	0.06	0.07	0.06	0.07	0.07	0.09	0.10	0.08
	Pb	t	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3e(i) Pipeline Compressors	NO <sub>x</sub>	kt	0.06	0.11	0.05	0.13	0.14	0.13	0.12	0.12	0.12	0.12	0.12	0.11	0.12	0.12	0.13	0.13	0.13
<b>% Change in Emissions</b>																			
1A3c Railways	NO <sub>x</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1A3d(ii) National Navigation	NO <sub>x</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%



**Table 3.7. Recalculations in Energy NFR 1A4**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
1A4a Commercial/Institutional	NO <sub>x</sub>	kt	2.91	2.69	2.50	2.38	2.05	1.88	1.98	2.13	1.99	2.13	2.08	1.99	2.11	2.09	1.82	1.96	1.94
	SO <sub>2</sub>	kt	11.40	5.38	1.81	1.25	0.31	0.28	0.23	0.25	0.14	0.13	0.13	0.21	0.19	0.15	0.14	0.17	0.13
	NM VOC	kt	0.46	0.42	0.49	0.55	0.49	0.45	0.48	0.50	0.46	0.50	0.48	0.47	0.50	0.50	0.43	0.48	0.48
	NH <sub>3</sub>	kt	0.00	0.00	0.00	0.00	0.02	0.02	0.03	0.04	0.04	0.03	0.04	0.03	0.03	0.02	0.02	0.03	0.03
	PM <sub>2.5</sub>	kt	0.54	0.36	0.26	0.31	0.15	0.14	0.14	0.17	0.16	0.16	0.17	0.15	0.16	0.14	0.13	0.13	0.13
	Pb	t	0.56	0.35	0.25	0.32	0.13	0.12	0.12	0.15	0.14	0.14	0.14	0.13	0.14	0.12	0.11	0.11	0.11
	Cd	t	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01
	Hg	t	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4b Residential	NO <sub>x</sub>	kt	4.85	4.79	5.72	6.95	7.62	6.56	6.13	5.94	5.26	5.69	5.98	5.57	6.01	5.78	6.37	5.93	4.97
	SO <sub>2</sub>	kt	26.76	17.52	15.72	13.09	9.29	8.27	8.07	8.46	7.50	7.57	8.26	7.36	7.31	6.61	6.99	7.48	5.42
	NM VOC	kt	35.07	22.18	15.17	13.49	13.17	11.93	11.91	12.70	11.40	11.70	11.89	10.08	10.84	9.70	10.10	9.62	7.51
	NH <sub>3</sub>	kt	0.40	0.24	0.15	0.12	0.11	0.10	0.10	0.10	0.09	0.09	0.09	0.08	0.08	0.07	0.07	0.07	0.06
	PM <sub>2.5</sub>	kt	20.41	13.31	9.32	8.54	8.57	7.80	7.85	8.40	7.58	7.84	8.02	6.84	7.39	6.65	6.95	6.54	5.12
	Pb	t	5.71	3.60	2.47	2.19	2.15	1.95	1.94	2.07	1.86	1.91	1.94	1.64	1.76	1.58	1.64	1.56	1.22
	Cd	t	0.05	0.04	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.02
	Hg	t	0.18	0.12	0.09	0.09	0.09	0.08	0.08	0.09	0.08	0.08	0.08	0.07	0.08	0.07	0.08	0.07	0.06
1A4c Agriculture/Forestry/Fishing	NO <sub>x</sub>	kt	8.59	14.00	12.69	12.47	7.28	6.44	6.06	5.40	4.70	4.15	3.94	4.07	4.33	3.96	3.52	3.29	3.69
	SO <sub>2</sub>	kt	1.41	1.34	0.94	0.86	0.37	0.06	0.09	0.08	0.05	0.04	0.04	0.05	0.03	0.02	0.02	0.02	0.03
	NM VOC	kt	1.35	1.66	1.31	1.04	0.58	0.51	0.46	0.38	0.32	0.28	0.27	0.27	0.27	0.25	0.23	0.21	0.24
	NH <sub>3</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	PM <sub>2.5</sub>	kt	0.92	1.03	0.75	0.58	0.31	0.28	0.24	0.20	0.17	0.15	0.14	0.13	0.13	0.12	0.11	0.10	0.12
	Pb	t	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.02
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Submission 2025</b>																			
1A4a Commercial/Institutional	NO <sub>x</sub>	kt	2.91	2.68	2.49	2.38	2.04	1.90	1.99	2.13	2.01	2.14	2.10	2.01	2.15	2.13	1.88	2.02	1.99
	SO <sub>2</sub>	kt	11.40	5.37	1.80	1.24	0.31	0.27	0.22	0.24	0.14	0.13	0.13	0.20	0.18	0.15	0.13	0.18	0.13
	NM VOC	kt	0.46	0.42	0.49	0.56	0.49	0.46	0.49	0.50	0.47	0.51	0.50	0.48	0.51	0.52	0.46	0.49	0.49
	NH <sub>3</sub>	kt	0.00	0.00	0.00	0.00	0.02	0.02	0.03	0.04	0.04	0.03	0.04	0.03	0.03	0.02	0.03	0.03	0.03
	PM <sub>2.5</sub>	kt	0.54	0.36	0.26	0.31	0.14	0.13	0.13	0.16	0.16	0.15	0.16	0.15	0.15	0.14	0.13	0.14	0.13
	Pb	t	0.56	0.35	0.25	0.32	0.12	0.12	0.11	0.14	0.14	0.13	0.14	0.13	0.13	0.12	0.11	0.12	0.11
	Cd	t	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01
	Hg	t	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4b Residential	NO <sub>x</sub>	kt	4.85	4.78	5.71	6.94	7.62	6.56	6.13	5.95	5.28	5.71	6.00	5.60	6.03	5.81	6.39	5.92	4.94
	SO <sub>2</sub>	kt	26.76	17.51	15.71	13.07	9.29	8.27	8.06	8.46	7.51	7.57	8.27	7.37	7.31	6.61	6.99	7.47	5.41
	NM VOC	kt	35.07	22.18	15.17	13.49	13.17	11.94	11.91	12.70	11.40	11.70	11.89	10.08	10.85	9.72	10.12	9.64	7.53
	NH <sub>3</sub>	kt	0.40	0.24	0.15	0.12	0.11	0.10	0.10	0.10	0.09	0.09	0.09	0.08	0.08	0.07	0.08	0.08	0.06
	PM <sub>2.5</sub>	kt	20.41	13.31	9.32	8.54	8.57	7.81	7.85	8.41	7.58	7.84	8.02	6.84	7.40	6.67	6.98	6.57	5.15
	Pb	t	5.71	3.60	2.47	2.19	2.15	1.95	1.94	2.07	1.86	1.91	1.94	1.64	1.77	1.58	1.64	1.56	1.22
	Cd	t	0.05	0.04	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.02
	Hg	t	0.18	0.12	0.09	0.09	0.09	0.08	0.08	0.09	0.08	0.08	0.08	0.07	0.08	0.07	0.08	0.07	0.06
1A4c Agriculture/Forestry/Fishing	NO <sub>x</sub>	kt	8.59	14.00	12.69	12.47	7.28	6.44	6.06	5.40	4.70	4.15	3.94	4.07	4.33	3.96	3.52	3.29	3.69
	SO <sub>2</sub>	kt	1.41	1.34	0.94	0.86	0.37	0.06	0.09	0.08	0.05	0.04	0.04	0.05	0.03	0.02	0.02	0.02	0.03
	NM VOC	kt	1.35	1.66	1.31	1.04	0.58	0.51	0.46	0.38	0.32	0.28	0.27	0.27	0.27	0.25	0.23	0.21	0.24
	NH <sub>3</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	PM <sub>2.5</sub>	kt	0.92	1.03	0.75	0.58	0.31	0.28	0.24	0.20	0.17	0.15	0.14	0.13	0.13	0.12	0.11	0.10	0.12
	Pb	t	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.02
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>% Change in Emissions</b>	<b>Pollutant</b>	<b>Unit</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2005</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>	<b>2022</b>
1A4a Commercial/Institutional	NO <sub>x</sub>	%	0.0%	-0.2%	-0.4%	-0.4%	-0.6%	0.8%	0.6%	0.3%	1.0%	0.6%	1.0%	0.9%	1.8%	1.9%	3.4%	2.9%	0.0%
	SO <sub>2</sub>	%	0.0%	-0.1%	-0.3%	-0.7%	-2.2%	-3.5%	-2.6%	-2.4%	-2.1%	-2.0%	-3.9%	-4.3%	-2.2%	-2.7%	-2.5%	2.9%	-1.1%
	NM VOC	%	0.0%	-0.1%	-0.2%	0.2%	0.4%	2.2%	1.9%	1.7%	2.6%	2.0%	2.6%	2.9%	3.7%	3.8%	5.6%	2.7%	0.7%
	NH <sub>3</sub>	%				0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	4.1%	10.6%	9.9%	7.1%	0.3%
	PM <sub>2.5</sub>	%	0.0%	-0.2%	-0.7%	-0.9%	-3.1%	-3.7%	-3.3%	-3.1%	-2.4%	-2.8%	-2.5%	-3.7%	-2.2%	-2.4%	-1.3%	3.7%	-1.3%
	Pb	%	0%	0%	-1%	-1%	-4%	-5%	-4%	-4%	-3%	-3%	-3%	-5%	-3%	-4%	-3%	4%	-1.6%
	Cd	%	0%	0%	-1%	-1%	-1%	-1%	-1%	-1%	0%	-1%	-1%	-1%	3%	7%	7%	7%	-0.5%
	Hg	%	0%	0%	0%	0%	0%	-3%	-1%	-1%	0%	0%	0%	0%	1%	1%	3%	1%	-0.1%
1A4b Residential	NO <sub>x</sub>	%	0.0%	-0.1%	-0.2%	-0.2%	-0.1%	0.0%	0.0%	0.1%	0.4%	0.4%	0.3%	0.5%	0.4%	0.5%	0.3%	-0.1%	0.0%
	SO <sub>2</sub>	%	0.0%	0.0%	-0.1%	-0.1%	-0.1%	0.0%	-0.1%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.1%	0.1%	0.0%	0.0%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.2%	0.2%	0.3%	0.0%
	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%	2.2%	3.1%	3.3%	0.1%
	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.3%	0.4%	0.5%	0.0%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.1%	2.4%	3.2%	3.8%	0.1%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.1%	0.0%	0.0%
1A4c Agriculture/Forestry/Fishing	NO <sub>x</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.2%	-0.6%	0.0%
	Pb	%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	-1%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.2%	-0.6%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.1%	0.0%

### 3.12 Quality Assurance/Quality Control

The time series spreadsheet system developed for individual categories, together with direct linking to the energy balance, allows for simple and efficient checking of activity data, emission factors, annual emissions and aggregated totals. Year-on-year changes immediately highlight any omissions, anomalies or internal errors. Initial checks are conducted by the inventory compiler as part of the calculation process, which is followed by a second check and completion of the QA/QC sheets which are integral to the calculation workbooks by another member of the inventories team. Cross-checks are performed for fuel data against the available supplementary sources for categories, such as 1A1a Public Electricity and Heat Production and some industrial processes, while maintaining consistency with fuel-use application in the estimation of GHG emissions. When new versions of the COPERT model are introduced for calculations in 1A3b, the previous year's activity data is run in the new model to compare with the current year, and the current year's activity data is run in the old version of the model to compare with the previous year. This allows firstly for identification of changes due to the model, and secondly for identification of changes due to the activity data.

### 3.13 Planned Improvements

This submission included checks on emission factors based on the latest Inventory Guidebook and this work will continue in future submissions with changes made where appropriate. All emissions associated with the category 1A2g are reported in 1A2gviii, and mobile emissions, 1A2gvii, are reported as "included elsewhere" (IE) in 1A2gviii. Currently all fuel is assumed to be stationary as the energy balance does not provide an estimate for fuels used in mobile machinery. The inventory agency had previously tendered and finalized research to be undertaken to quantify the extent and amounts of fuel used in off-road mobile combustion, but the COVID-19 pandemic had a negative impact on its outcome as it affected the collection of activity data. As a result, it was concluded that changing the methodology to estimate emissions from off-road vehicles and other machinery to take into account the results of the research project would not improve the accuracy of the inventory. The use of BEUS data has focused on data from 2009 to 2018 with the 2018 to 2023 splits based on the latest BEUS data splits. Review recommendations and responses are included in this report in Annex H.

### 3.14 Memo Items

The memo items of the NFR reporting format refer to activities for which the emissions are excluded from national totals. The use of fuels in domestic and international aviation (cruise phase) and marine bunkers are the most important of these activities. Some of the associated emissions, particularly from international aviation, are increasing very rapidly and it is therefore important that they are closely monitored for comparison with other sources and for the benefit of the international organisations that will have to develop control strategies for them in the future. The estimation of emissions for memo items is described here because they are calculated as part of the general estimation procedures for the Energy sector. The national energy balance sheets include marine bunkers as a specific item and the emissions may be calculated directly. The approach used to for the cruise element of aviation is explained in Section 3.6.1 and this data is provided to SEAI for inclusion in the national energy balance. Emission factors for international cruise aviation and navigation are documented in Tables C.19 and C.29 of Annex C.

## Chapter Four - Industrial Processes and Product Use

### 4.1 Overview of the Industrial Processes and Product Use (NFR 2) Sector

The Industrial Processes sector has historically not been a large source of emissions in Ireland. Indeed, major industrial processes within the chemical sector and metal production that are common to many other developed countries have never been part of the economy in Ireland. Hence, many of the production processes within this sector are not relevant to the inventories of air pollutants in Ireland. Also, of note is the fact that for a number of pollutants, it has not been possible to separate emissions from the combustion of fuel within industry and those associated with production processes. For all industries, fuel-based estimates of emissions have been collated and are reported under NFR Sector 1A2 (Manufacturing Industries and Construction). Where specific information is available in relation to process emissions as distinct from those associated with fuel combustion, they are reported under NFR Sector 2 (Industrial Processes and Product Use) and are discussed in this chapter. In most of these cases, process-specific information is sourced from Annual Environmental Reports, which form part of the reporting obligations under IPPC/IED permits in Ireland. In some cases, production data (estimated and/or calculated) are also used where available.

Relevant subcategories under Mineral Products (2A), Chemical Industry (2B), Metal Production (2C), Solvent and other product use (2D-2H) and Other Production Processes (2L) for which process emissions of various pollutants are reported under Industrial Processes and Product Use in Ireland are described in the following sections.

### 4.2 Mineral Products (NFR 2A)

The industrial processes for which estimates are included in Ireland's air pollution inventory under NFR 2.A are as follows:

- 2.A.1 Cement Production
- 2.A.2 Lime Production
- 2.A.3 Glass Production
- 2.A.5.a Quarrying and mining of minerals other than coal
- 2.A.5.b Construction and demolition
- 2.A.5.c Storage, handling and transport of mineral products
- 2.A.6 Other mineral products

#### 4.2.1 Cement Production (NFR 2A1)

Cement manufacture is a major mineral industry. During cement manufacture, raw materials, such as limestone, are finely ground and then transformed in a kiln at high temperatures (calcination) to produce clinker. Gypsum is then blended with clinker to produce cement. The combustion process in the cement kiln is an integral part of the production process, where the fuel ash becomes part of the cement clinker. It is therefore not possible in most cases to distinguish the process and combustion emissions from one another. As a result, because most of the pollutants originate from the fuels used, emissions are generally reported under NFR Category 1A2f (Non-metallic minerals) and notation keys IE and NA are reported under 2A1 for these pollutants.

There are at present four cement plants in Ireland, all of which use the dry kiln process, and they are currently fuelled by coal, petroleum coke and fuel oil, with meat and bone-meal, solid recovered fuels (SRF) and tyre derived fuels co-incinerated at three of the four plants. Literature sources, in particular, the EMEP/EEA Guidebook (2023), provide some POP emission factors on an overall grams of pollutant per tonne of clinker produced basis. The guidebook states that these emissions are allocated to combustion in cement manufacture and so emissions of POPs are reported as NA for category 2A1. These emission factors have been used to determine the total emissions from cement plants. Fuel-use data are available from plant operators as part of their reporting requirements under the EU ETS (Directive 2003/87/EC) since 2005. Residual fuel not used in the cement manufacture sector but in other manufacturing industry in 1A2f is calculated and emissions from fuel use are calculated using combustion emission factors in 1A2f, while the fuel use data from cement manufacture is removed to ensure there is no double counting, as is discussed in Chapter 3, section 3.5 of this report. In this submission emissions of particulates are reported (see table 4.0 in Annex D) using EFs from Table 3.1 EMEP/EEA Guidebook (2023).

#### 4.2.2 Lime Production (NFR 2A2)

The lime production process involves the grinding and “burning” of limestone ( $\text{CaCO}_3$ ) to produce what is commonly termed “quicklime” ( $\text{CaO}$ ). It can then be further treated by the addition of water, a process called slaking, to produce slaked lime ( $\text{CaOH}$ ), which generates large amounts of heat and steam. The finished product can then be packaged and distributed for use. Currently, there are two lime plants in Ireland and a third that operated until 1999. It is understood that all three utilised limestone quarries and kilns to burn the limestone raw material. The nature of the fuel used and the abatement in place varies from plant to plant.

Process emissions from lime production are obtained as the difference between total emissions on a “per unit production” basis and those estimated from fuel combustion. In this case, it was found that fuel combustion estimates of emissions were generally larger than those estimated on a “per unit production” basis and therefore all emissions from lime production, with the exception of particulates, are assumed to be included in those reported within Sector 1A2f and the notation key IE (included elsewhere) is used for 2A2. In this submission emissions of particulates are reported, see Table 4.0, using EFs from Chapter 2.A.2 Table 3.3. Lime production (controlled) (EMEP/EEA Guidebook (2023)). Use of controlled emission factors are appropriate because both remaining lime facilities have BAT abatement techniques in use to minimise/prevent diffuse dust emission from dusty operations, bulk storage areas and other operations on site.

#### 4.2.3 Glass Production (NFR 2A3)

The manufacture of glass was not a predominant industry in Ireland, being limited to three sub-sectors: lead crystal, container glass, and glass wool. The only container glass plant closed in 2002, one of the lead crystal plants closed in early 2006, the glass wool plant closed in 2008 and the last one, (second of the two) lead crystal plant closed in 2009. Therefore, after this period, emissions of all pollutants are reported as ‘NO’ (not occurring). The pollutants for which process emission estimates have been made are particulate matter (TSP,  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$ , BC) priority metals (Pb, Cd and Hg), other metals (As, Cr, Cu, Ni, Se and Zn), and PCDD/F. In addition, fuel-derived emissions from the glass industry for 1990-2009 are already accounted for within the Manufacturing Industries and Construction (1A2) sector.

Heavy Metal emissions can occur from glass processes from the metals contained in fossil fuels burnt to melt the glass and from metal additives to the glass. Metal emissions from industry fossil fuel use are accounted for within NFR Sector 1A2g (Other Manufacturing Industries). The metals emitted from glass production processes depend on the type of glass produced. Lead oxide and sometimes arsenic trioxide are used in the production of lead crystal glass and both metals can be emitted to air. Selenium and chromium compounds are used as colouring agents for container glass. Metal compounds are not believed to have been used to any great extent in the production of glass wool. Of the glass processes outlined, all the plants were regulated under Integrated Pollution Prevention Control IPPC licences. Therefore, there is some information available from their AERs until they closed. Other licence information includes some details of plant design and operation, including capacities, fuel types and operating hours. In addition, confidential information in relation to production statistics has also been supplied to the inventory team.

Emission data for individual metals reported in AERs are limited to Pb emissions from the two lead crystal installations. These data have been used to estimate emissions of Pb from lead crystal production, while emission factors are used for the other glass processes. Literature emission factors are used from the Inventory Guidebook (EMEP/EEA, 2023) and, where deemed more appropriate, emission factors from the UK National Atmospheric Emissions Inventory (NAEI) database. The emission factors used are presented in Table D.1, Annex D. Total emissions for each metal from glass production are presented in Table 4.2.

Ireland includes process emissions for PM<sub>2.5</sub> from 2A3 Glass production using confidential production data and the Tier 2 emission factors presented in Tables 3.3, 3.5 and 3.6 from Chapter 2A3 of the 2023 EMEP/EEA Inventory guidebook (EMEP/EEA 2023). However, the Guidebook (chapter 1.A.2) notes that, Tier 1 emission factors from combustion in the industrial sector (NFR 1A2) would overlap at least partly with the estimate from process emissions in NFR 2A3. Therefore, Ireland estimates the emissions from category NFR 2A3 and subtracts those estimates from the total emissions calculated for category NFR 1A2f to avoid double counting of these emissions. The emission factors used are presented in Table D.1, Annex D.

The potential for PCDD/F emissions from glass production is generally low because of the long residence times in high-temperature conditions, although chlorine can be introduced via fuels and raw materials, and therefore there is some potential for PCDD/F emissions. However, in the plants in Ireland, the main energy sources used were gas and electricity, and therefore PCDD/F emissions from fuel combustion were likely to be low. The information on abatement technology is uncertain for the glass manufacturing plants in Ireland. The URS Dames & Moore (2000) PCDD/F inventory report implies that one furnace would be fitted with abatement by 2005, but it is assumed that this is the large facility that closed in 2002. IPPC licence information implies that the environmental performance at the glass wool plant was improved in 1999, but the nature of the improvements is not clear.

Emission factors for PCDD/F are provided in the UNEP Toolkit (2013) for two different classes of facility: 0.2 µg I-TEQ/t of glass produced for a facility with no dust control, and 0.015 µg I-TEQ/t of glass produced for a facility with abatement. However, there have been improvements in environmental performance; therefore, the emission factor of 0.2 µg I-TEQ/t is used to estimate emissions in 1990, with a linear decrease to 0.11 µg I-TEQ/t in 2000 (URS Dames & Moore, 2000) and with a subsequent decrease to 0.015 µg I-TEQ/t by 2003 (when the container plant had closed) and

remaining at this level up to 2009 when the last plant ceased its operation. Dioxin and furan emission estimates for glass production are presented in Table 4.1 of electronic Annex D.

#### 4.2.4 Quarrying and Mining of Minerals Other than Coal (NFR 2A5a)

Emissions of PM<sub>2.5</sub>, TSP and PM<sub>10</sub> have been estimated for all years of the time series 1990- 2023. A tier 2 method was implemented in 2025 based on the proxy method provided from the NECD Emissions inventory review 2023. Activity data was taken from national statistics from the CSO and emission factors from "Consideration of best practice in emission inventory reviews: Appendix 1, p 21 A Proxy solution for mining and quarrying". Activity data and emissions are presented in Table 4.2 of electronic Annex D.

#### 4.2.5 Construction and Demolition (NFR 2A5b)

Emissions of PM<sub>2.5</sub>, TSP and PM<sub>10</sub> have been estimated for all years of the time series 1990- 2023. Activity data was taken from national statistics from the CSO and the US EPA Tier 1 methodology and emission factors from Chapter 2.A.5.b section 3.2 of the guidebook (EMEP/EEA, 2023). Ireland checked for sources of required data to implement a Tier 2 method and determined it is not possible to source any of the required data. The methodologies provided by US EPA with AP-42 require very detailed local data e.g. material silt content, road surface silt content, material moisture content, medium wind speed, mean vehicle weight, mean vehicle speed, vehicle kilometre travelled (VKT)etc. The 2023 EEA/EMEP guidebook states that collection of such data is likely to be possible only for individual large point sources. Ireland does not have this data available for any construction and demolition sites and will continue to use the Tier 1 method to estimate emissions from this source. The emission estimates are presented in Table 4.3 of electronic Annex D.

#### 4.2.6 Storage, Handling and Transport of Mineral Products (NFR 2A5c)

Emissions of PM<sub>2.5</sub> have been estimated for all years of the time series 1990- 2023. Activity data was taken from national statistics from the CSO and emission factors from Table 3.4 Chapter 2.A.5.a of the guidebook (EMEP/EEA, 2023). Activity data and emission estimates are presented in Table 4.4 of electronic Annex D.

#### 4.2.7 Other (NFR 2A6)

The industrial processes included within NFR Sector 2A6 are Bricks and Ceramics Production and Asphalt Production. Each of these subcategories is described in the following sections in terms of the pollutants for which emission estimates are made.

##### 4.2.7.1 Bricks and Ceramics Production

The production of bricks and ceramics is a small sector in Ireland with a total of four IPPC-licensed facilities in operation. Emission estimates are only made for PCDD/F as there are no data available in relation to process emissions of other pollutants and, furthermore, they are expected to be negligible (AEA/CTC, 2008). Direct production information in relation to the bricks and ceramics sector is not available; however, raw material input data are provided by the companies under the EU ETS. For the purposes of inventory estimates, as a worst-case scenario it is assumed that raw material input equals product output. Emission factors are sourced from the UNEP Toolkit (2013) in which two classes of facility are suggested: 0.2 µg I-TEQ/t of brick produced for a facility with no dust control and 0.02 µg I-TEQ/t of brick produced for a facility with abatement. The URS Dames & Moore (2000) report suggests an emission factor of 0.11 µg I-TEQ/t, which is the average of the two emission factors, and



this value, has been used across the time series. The UNEP Toolkit (2013) does not include emission factors for ceramics production and therefore the emission factor for bricks is also applied to ceramics production. Dioxin and furan emission estimates for bricks and ceramics production are presented in Table 4.5 of electronic Annex D.

#### 4.2.7.2 Asphalt Production

In the context of this inventory, the term “asphalt” is used to describe a bituminous product that may contain varying amounts of aggregate, used to build and maintain roads, whilst “bitumen” is assumed to be a heavy oil tar product which is used at elevated temperatures particularly in roofing materials for some buildings. Currently, only PCDD/F emission estimates from asphalt are included in Ireland’s air pollution emission inventory.

Information in relation to the production of asphalt in Ireland is sourced from the European Asphalt Pavement Association (EAPA, 2001, 2007, 2012, 2016, 2018), and the Irish Asphalt Pavement Producers Association (IAPA 2019, 2020, 2021, 2022, 2023, 2024) which generates an annual report outlining the quantity and end use of asphalt produced in European countries and Ireland. Production data are available from 1994 onwards, with pre-1994 production estimates assumed to be equal to those in 1994. The production levels until 2006 show an upward year-on-year trend due to increased road building in Ireland, from 2007 to 2014 the trend was decreasing but a slight increase is seen from 2015 to 2019 and a decrease was noted in 2020 which was partly related to COVID restrictions. An increase has been seen in 2021 and 2022 with an increase in road building activity.

In Ireland, bag filters were fitted to most asphalt production facilities prior to 2000 and it was suggested that all facilities would have bag filters by 2001 (URS Dames & Moore, 2000). The UNEP Toolkit (2013) gives a range in emission factors of 0.007 to 0.07 µg I-TEQ/t asphalt produced. Given the above information, the emission factor for PCDD/F from asphalt production of 0.07 µg I-TEQ/t is adopted for 1990. A linear decrease in the emission factor is then assumed to 0.039 µg I-TEQ/t by 2000, and a further linear decrease is assumed to 0.007 µg I-TEQ/t by the end of 2002. The emission factor is assumed to be 0.007 µg I-TEQ/t from 2003 and onwards. Dioxin and furan emission estimates for asphalt production are presented in Table 4.6 of electronic Annex D.

### 4.3 Chemical Industry (NFR 2B)

The chemical industry is not a dominant industry in Ireland in relation to industrial processes and is not an important source of emissions. The only source of emissions for which estimates are collated are NO<sub>x</sub> emissions from Nitric Acid Production for the years 1990-2002.

#### 4.3.1 Nitric Acid Production (NFR 2B2)

Nitric acid is used as a raw material mainly in the manufacture of nitrogen-based fertiliser. It may also be used in the production of adipic acid and explosives, for metal etching, and in the processing of ferrous metals. In the manufacture of nitrogenous fertilisers, the Haber Bosch process is utilised in which NH<sub>3</sub> is made by combining nitrogen from the air with hydrogen from natural gas and water, using the energy from the gas and a catalyst. Nitric acid is produced by burning (oxidising) the NH<sub>3</sub> over a catalyst. The nitric acid is combined with more NH<sub>3</sub> to produce ammonium nitrate, which is solidified into granules or bead-like prills for application to land using a fertiliser spreader. Up to its



closure in 2002, there was one such plant in Ireland, which utilised the above process to produce calcium ammonium nitrate and other nitrogenous fertiliser blends. The inventory agency received direct correspondence from the plant in relation to the quantities of nitric acid produced and the measured emissions of NO<sub>x</sub>. Emission estimates and associated activity data for NO<sub>x</sub> emissions from nitric acid production are presented in Table 4.7 in electronic Annex D. Abatement measures were installed at the plant in the mid-1990s and they are reflected in emission estimates from 1995.

#### 4.3.2 Storage, handling and transport of chemical products (NFR 2B10b)

Emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are reported in category 2.B.10.b following a 2022 NECD review recommendation. Emissions reported from storage, handling and transport of fertilisers were reallocated from category 3Dc. Emissions are estimated using emission factors from CEPMEIP database for SNAP 040415 storage and handling of NPK fertilisers. Emission factors of 100 g/t, 32 g/t and 4 g/t from CEPMEIP (2001) are utilised in the calculation of emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub>, respectively. Activity data and emissions are presented in Table 4.7 of Annex D.

### 4.4 Metal Production (NFR 2C)

This category includes a wide range of processes such as primary and secondary iron and steel production, aluminium production and other non-ferrous production. In this category, emissions are estimated for the following subcategories and pollutants:

- 2C1 Iron and Steel Production – As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, PCDD/F;
- 2C2 Ferroalloys Production – As, Cd, Cr, Ni, Pb, Zn, HCB;
- 2C3 Aluminium Production – Zn;
- 2C5 Lead production – Pb;
- 2C7 Other metal production – Cd, Cr, Cu, Ni, Pb, Zn;

Ireland is a major European producer of Zn and Pb ores. The preparation of Pb and Zn concentrates does not produce emissions and the concentrates are exported for further processing.

#### 4.4.1 Iron and Steel Production (NFR 2C1)

This sector covers the manufacture of iron and steel, an energy-intensive process likely to generate high emissions to air from the use of furnaces and sintering processes, as well as the manual handling of the raw material to finished goods, which can include hot and cold rolling, and turning, temping and cutting of metal to reach a desired end product. Steel production in Ireland has been limited to a single large electric arc furnace installation, which closed in 2001 but was operational throughout the period 1990–2001. One small foundry remained in operation contributing a negligible amount of emissions after the large plant's closure. The main plant produced up to 360 kt of steel per annum mainly from recycled scrap steel. It received an IPPC licence to operate just months before its closure, therefore no AERs were filed by the plant. However, some emission testing was carried out with respect to heavy metal emissions as part of its licence application.

Heavy metal emission estimates have been calculated using the aforementioned emission testing results for Cd, Cr, Pb, Ni and Zn, whilst for the remaining pollutants (i.e. As, Cu, Hg, and Se) Inventory Guidebook (EMEP/EEA, 2023) emission factors have been used, assuming no abatement at the plant. The emission factors used are presented in Table D.3, Annex D. Emission estimates for the 1990–2001 time series are shown in Table 4.9 of electronic Annex D. Metal production data are available from the

site for the period 1994–2001, with pre-1994 production assumed to be equal to that in 1994. Emission estimates are calculated by multiplying the production data by the relevant pollutant emission factor and assume that no abatement was in place at the plant.

Electric arc furnaces are significant sources of POPs. The overall approach to report emissions of POPs from the iron and steel category has been to account for emissions from fuel combustion within Sector 1A2a, with the process emissions reported (where possible) under Sector 2C1, which have been estimated using the approach described with respect to Cement Production (2A1). Emission factors based on per unit production were used to calculate initial estimates of total emissions. These factors are sourced from the Inventory Guidebook (EMEP/EEA, 2023). The difference between estimates determined on this basis and those reported for fuel combustion sector 1A2a is then reported in sector 2C. Emission estimates for the sector are presented in Table 4.9 of Annex D for PCDD/F. The notation key 'IE' (included elsewhere) is used for PCB and HCB emissions for years 1990-2002 and the notation key NO is used for the years 2003-2022. Emissions of HCB and PCB are reported and accounted for under NFR 1A2a as it has not been possible to separate process and fuel combustion related to these emissions for the period 1990-2002 for which there was an iron and steel industry in Ireland.

#### 4.4.2 Ferroalloys Production (NFR 2C2)

This sector covers several secondary sites engaged in iron and steel manufacture. Two types of installation are distinguished. The first type covers installations involved in the manufacture of ductile iron for use in street furniture, public benches, waste bins and manhole covers, and the second is the manufacture of cast iron for appliances. The process of creating ductile iron utilises electric arc furnaces to smelt the raw materials, iron and magnesium. In the manufacture of cast iron, ferrous and non-ferrous metals, including scrap metal, are used within the process. Since 1990, there have been three relevant facilities in Ireland. Due to a change in operations, one of these plants reported negligible emission estimates from 2003 onwards; one facility closed in 2014 leaving a single operating facility from 2014 onwards.

A number of the larger metal processing sites are regulated under IPPC. Some metal emission estimates and particulate emission estimates have been reported in AERs; however, not all installations report emissions in all years. In some cases, only production data and emissions of TSP are available. Where production data only were provided by the plant operator, they were used to calculate emissions of TSP using USEPA factors for dust emissions from abated/unabated iron foundry cupola processes. Estimates for metal emissions were then obtained from the TSP estimates based on Inventory Guidebook (EMEP/EEA, 2023) dust composition data for foundry dust. Abatement techniques are also taken into account in emission calculations at a plant-specific level, where this applies. Emission estimates of HCB for the time series 1990-1996 are presented in Table 4.10. The only source of HCB is the secondary manufacture of aluminium, for which the Inventory Guidebook (EMEP/EEA, 2023) indicates a factor of 5 g/t of aluminium. This factor has been used to estimate HCB emissions across the time series until use of the HCE-based cover gas was banned in 1996 and emissions are reported using the notation key NO for years after 1996.

Estimates for TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC have also been made for this category. Total Particulate Matter data reported in AERs were used to estimate emissions from TSP. Fractionation profiles based on the

emission factors within the Inventory Guidebook (EMEP/EEA, 2023) were used to estimate PM<sub>10</sub>, PM<sub>2.5</sub> and BC emissions.

Where production data are not available, TSP estimates have been used to estimate metal production across the time series using the BiPRO waste report (2005), other POPs like PCDD/F, PCB and PAHs are reported under fuel combustion sector 1A2b. The HCB emission factor used is presented in Table D.3, Annex D, and process emission estimates for the 1990-1996 time series are presented in Table 4.10 of Annex D.

#### 4.4.3 Aluminium Production (NFR 2C3)

Ireland is an important producer of alumina at one large plant using the Bayer process (extraction of AlO<sub>3</sub> using NaOH). The production of alumina using the Bayer process does not give rise to significant metal emissions and therefore process emissions are not estimated for this source. Ireland has some secondary aluminium processing for which estimates of Zn have been made following reports from the plant involved. The plant closed in late 2006 and therefore estimates are only provided for the 1990–2006 time series as presented in Table 4.11 of Annex D. Production data for the plant are not available and therefore estimates were made using PM as an indicator. The UK NAEI emission factor of 2.725 g/t is then applied (Table D.3, Annex D).

#### 4.4.4 Lead Production (NFR 2C5)

A significant quantity of Lead is mined in Ireland, but such mining is assumed not to be a significant source of emissions to air. Estimates at facility level of Lead emissions have been obtained from AERs and PRTR data, and this facility reporting is a metal processing facility which carries out secondary lead production and processes lead and lead alloys from scrap lead. Abatement using bag filters is in place in this facility and no fugitive releases of lead have been reported since 2009. No other pollutants are reported by the facility for the years 1990-2009. Emission estimates for the time series are presented in Table 4.12 and in electronic Annex D and are reported as not occurring since 2009.

**Table 4.12 Emission Time Series for Lead Production**

Pollutant (unit)	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Pb (kg)	7.845	7.845	7.845	7.845	0.018	0.008	0.013	0.006	0.025	0.034	0.030	NO

#### 4.4.5 Other Metal Production (NFR 2C7)

This category covers all other metal manufacture and manipulation, including any emissions from the mining of raw materials. A significant quantity of Zn is mined in Ireland, but such mining is assumed not to be a significant source of emissions to air. Ireland has a number of small aluminium casting companies in addition to facilities for wire manufacture and the manufacture of refined or secondary Pb and Cu products, as well as a number of Zn galvanising plants. The emissions reported in 2C7c Other metal production (Cd, Cr, Cu, Pb, Ni and Zn) have been obtained from AERs and PRTR reporting with respect to these secondary metal operations. The notation key “NA” is used for other metals in category 2C7c as the activity occurs, but the individual metals are reported as not emitted by the facilities. Installation of abatement equipment resulted in some emissions to air falling below detection limits, hence reporting of various metals ceases for some facilities in certain years. In the case of 2C7c, Cu is reported from metal plating facilities until 2008, table 4.13 gives total Cu emissions reported under 2C7 which includes those reported under 2C7a and 2C7c which is why the table does

not match 2C7c emission values. Emission estimates for the time series are presented in Table 4.13 of electronic Annex D. Emissions arise from 3 plants in the latter parts of the time series and are therefore very sensitive to changes in activity of individual plants. The emissions reported (Cu only) in 2C7a Copper Production were taken from the annual reporting of monitoring results from a licenced facility that granulates copper cables. Installation of abatement equipment resulted in all emissions to air falling below detection limits, hence reporting of Cu ceased in 2007. As no roasting, smelting, converting, refining or electro-refining of copper is carried out at this facility, the emissions associated with Primary and Secondary copper production are not occurring. The notation key NA is used for other metals in category 2C7a for years 1990-2006 and notation key “NO” is used for years 2007-2023 as emissions do not occur.

Process emissions of POPs (where applicable) are included in combustion emissions and reported as included elsewhere (IE) for category 2C7.

## 4.5 Overview of NMVOC emissions from Solvent and Other Product Use (NFR 2D-2L) Sector

The emission estimates presented in Solvent and Other Product Use (NFR 2D-2H) include Domestic solvent use including fungicides (2D3a), Road Paving with asphalt (2D3b), Coating Applications (2D3d), Degreasing and surface cleaning (2D3e), Dry Cleaning (2D3f), Chemical Products, Manufacture and Processing (2D3g), Printing (2D3h), Other Solvent Use (Fat, Edible and non-edible oil extraction) (2D3i), Other Product use (Use of Tobacco products and Fireworks) (2G) and Food and Beverages industry (2H2). Emissions are the result of continuing improvement of NMVOC emission estimates for Ireland through the outsourcing of tendered projects by the EPA. For 2D3c Asphalt roofing it has been determined that the production of asphalt roofing materials does not occur in Ireland. The Irish Asphalt Pavement Producers Association (IAPA) and the Bitumen suppliers were contacted with regards to the production of asphalt roofing in Ireland. They confirmed that asphalt roofing materials are imported from the UK and not manufactured in Ireland. The CSO was contacted to determine if any production statistics exist and only import and export figures are reported for asphalt roofing materials. This category is reported as NO.

In 2012, the inventory agency commissioned a research project to update the NMVOC emission inventory for 2006-2013. This was a follow-on project to CTC/AEA (2005) and Finn et al. (2001) and resulted in a revised dataset where new data and methodologies had become available. This approach was taken in accordance with the Inventory Guidebook (EMEP/EEA 2016) methodology for NMVOC emissions for Solvent and Other Product Use (NFR 2D-2L). The results of this project were provided in the 2016 submission. This project continued into 2016 and resulted in further improvements to the NMVOC emission inventory from Solvent Usage.

Emissions data were gathered using a similar methodology to previous approaches. Bottom-up data was mainly obtained from submissions of Annual Environmental reports (AERs) which detail emissions in a variety of reporting formats ranging from the Solvent Mass Balance Summary, Solvent Management Plan (SMP), Pollution Release and Transfer Register (PRTR), or the Annual Environmental Report returns Workbook. In addition, new data sources were used from legislation designed to limit and report solvent usage (Solvent Directive 1999/13/EC). In conjunction with these data, the number of operators within each category was estimated using NACE codes provided by the Central Statistics Office (CSO) or from expert opinion.

Top-down methods were used for activities not covered by the IPPC licensing system nor under the Solvent Directive (1999/13/EC). The most significant included the use of non-industrial paints, metal degreasing and the use of domestic solvents. Input in the form of activity data, solvent usage or VOC emissions data for each individual activity were collated into spreadsheets. Emissions were estimated by applying the Inventory Guidebook (EMEP/EEA 2016 and since updated to EMEP/EEA 2023) methods, default emission factors and general guidance as appropriate. Scaling up to national level was applied where necessary. The emission factors used are presented in Table D.4, Annex D and the estimated emissions are also presented in tables 4.14- 4-46 of Annex D.

Emissions reported in NFR 2019 format are aggregated from the Selected Nomenclature for Air Pollutants (SNAP) categories. SNAP codes are used in the Inventory Guidebook (EMEP/EEA 2023) where sectoral emission sources and emission factors are provided in this system. Therefore, SNAP codes are adopted in these categories as it ensures that reporting of emissions is consistent with the guidebook and therefore other Parties submissions. Additionally, the use of SNAP codes facilitates a sub-sectoral analysis of drivers and trends.


For a number of sources, it was not possible to obtain reliable country-specific data (SNAP 060107: Paint Application: Wood, SNAP 060109: Non-Industrial Paint Application, SNAP 060408: Domestic Solvent Use). As a consequence of this, UK and other Parties' emission factors, and in some cases activity data (scaled by surrogate data), were used in the estimation methodology.

Obtaining country-specific data has been identified as an important issue in the past (Barry, S. and Regan, B., 2014). While new activity data were obtained for sectors that previously relied upon proxy sources, a number of sectors are still estimated using proxy information sources. Further reducing the dependence of these would require substantial investment, and the improvement that this would bring, over the use of proxy-based data is thought to be relatively small due to the similarity in lifestyle behaviour between the countries operating within the EU (and in particular the United Kingdom) and therefore a common market place.

The main drivers associated with trends in implied emission factors relate to reduced solvent content of products, and paints. The trends in activity data reflect the fact that Ireland experienced rapid economic growth from the late 1990s to 2007. As a result, there was a substantial increase in the number of vehicles, growth in the number of individual households, and generally a higher per capita consumption of paints, cosmetics, toiletries, and other solvent containing products. Since 2007, there has been a rapid economic downturn, which has had a marked impact on consumption, and therefore emissions of NMVOC. As economic conditions began to improve emissions have also increased over the period 2012 to 2023.

Figure 4.1 illustrates the overall trend and shows a 39.33 per cent increase in total emissions between 1990 (33.9 kt of NMVOC) and 2023 (47.2 kt of NMVOC). The main contributor to the trend is sector 2H2 Food and Beverages industry, with 204.6 per cent increase between 1990 and 2023 and was responsible for 62.1 per cent share of emissions from solvent and other product use in 2023. In 2023 a decrease of 5.3 percent was seen in this sector compared to 2022 emissions mainly due to a decrease in emissions from spirit production. The second largest contributor is sector 2D3a (Domestic Solvent Use including fungicides) with a contribution of 25.3 per cent of emissions from solvent and other

product use (having increased by 50.7 per cent since 1990). Sector 2D3d (Coating Applications) accounted for 4.5 per cent in 2023 showing a decrease of 70.8 per cent between 1990 and 2023.



**Figure 4.1 NMVOC Emission Trend for 2D-2H Other Solvent and Product 1990–2023**

Sector 2D3a, (Domestic Solvent Use including fungicides) is the second largest contributor to NMVOC emissions (Table 4.14, Annex D). A Tier 2 method was implemented which uses population data obtained from the CSO and per capita emission factors for product use from the Inventory guidebook (EMEP/EEA, 2023). This approach was used as the statistics required for the use of Tier 2b approach were not complete in terms of the product types covered by domestic solvent use. Further work is currently underway by the Inventory team to collect suitable activity data to improve the estimation of emissions from Domestic Solvent use including fungicides. ESIG data was requested in 2024 and the team is working to assess the appropriateness of using this dataset for the 2026 submission. For Hg no Tier 2 emission estimate is available for fluorescent tubes. The 2023 Inventory Guidebook does not contain an emission factor for this source and states “Due to the uncertainty around these releases, this source is currently not considered in the Guidebook.” Therefore, this emission is reported as not applicable (NA).

Sector 2D3b (Road paving with asphalt) emissions contribute less than 0.1 per cent of 2023 emissions. Emissions from the sector have decreased by 9.1 per cent since 1990.

Emissions from Coating applications (2D3d) decreased between 2007 and 2012 before increasing in 2013 and 2014 and decreasing again from 2015 to 2019 and reaching their lowest level in 2020 (17.24 kt) they have since increased to 21.14 kt in 2023. The main driver of NMVOC emissions from this emission category is the application of decorative paint (SNAP codes: 060103/060104). A number of factors contributed to the ongoing decrease in emissions including the substantial reduction in the solvent content of paint in recent years to comply with the Deco-Paints Directive (EP and CEU, 2004b), a greater awareness of environmental issues from the general public in addition to the economic downturn in Ireland. From discussions with industry, pressure from some of the larger retailers is noted to be one of the key drivers for the decrease in solvent use in architectural paint. The sales of water-based paints have decreased by 7.1 per cent between 1990 and 2023 whereas solvent-based paint sales have decreased by 44.5 per cent over the same period.



Emissions from 2D3e (Degreasing) decreased by 53.2 per cent between 1990 and 2023. Emissions peaked in 1996 at 2.15 kt. The methodology is based on net consumption of solvents (imports minus exports) provided by the CSO. The analysis showed that the main solvent used in this sector is Dichloromethane. The reductions are assumed to be driven by improved management practices and abatement technologies (open-top tanks have been phased out in the European Union as a result of the Solvents Emissions Directive 1999/13/EC). Emissions from this emission source accounted for 1.5 per cent of the total emissions from solvent and other product use in 2023.

Data obtained under the reporting requirements of the Solvent Directive (1999/13/EC) was used to estimate emissions from 2D3f (Dry Cleaning). Solvent usage, emissions data and national statistics were used to estimate emissions from this emission source. Emissions decreased by 82.9 per cent over the 1990-2023 period. Emissions from 2D3f accounted for 0.1 per cent of the total emissions from solvent and other product use in 2023.

2D3g (Chemical products) accounts for 1.2 per cent of emissions in solvent and other product use in 2023. In 1990, this sector accounted for 8.9 per cent. This emission category consists of fourteen emission sources; however, the majority of the emissions sources contribute very little to the overall emissions from the Chemical products sector. The diversity within these sectors is very large in terms of the type of process, the products made and the scale involved. The main driver of emissions from this emission source is Pharmaceutical Production (SNAP code 060306). Emissions from pharmaceutical production accounted for 82.0 per cent of emissions in the 2D3g category in 2023. Emissions from 2D3g decreased by 81.4 per cent between 1990 and 2023. Emissions decreased as a result of the introduction of new management practices or through the use of abatement technology (CTC/AEA, 2005 and Barry S. and O'Regan B., 2014). This indicates that current policy strategies are having an impact on solvent use and emissions. In addition, large reductions in emissions were found in several emission sources between 1990 and 2023. For instance, emissions reduced significantly from SNAP code 060303 (Polyurethane Processing) and SNAP code 060305 (Rubber Processing). This was mainly a result of plant closures with the last rubber processing facility closing in 2016. 2D3h (Printing) emissions decreased 92.1 per cent over the 1990 to 2023. In 2023, the sector accounted for 0.5 per cent of total emissions from solvent and other product use. Emissions from this sector continue to decrease since 2017 and decreased by 47.8 per cent in 2023 (0.23 kt) compared to 2022 (0.44 kt). This was due to a change to water-based solvents and outsourcing of solvent based printing to the UK by some of the larger companies in recent years. It is important to note that the print industry is included under the Solvent Directive (1999/13/EC) and is subject to IPPC licencing where applicable which is most likely responsible for the decrease in emissions.

2D3i (Other Solvent use) emissions increased 70.8 per cent over the 1990 to 2023 timeseries. In 2023, the sector accounted for 4.8 per cent of total emissions from solvent and other product use. This emission category consists of 8 sources which include SNAP codes: 060401 (Glass Wool Enduction), 060402 (Mineral Wool Induction), 060404 (Fat, edible and non-edible oil extraction), 060405 (Application of Adhesives and Glues), 060406 (Preservation of Wood), 060407 (Underseal Treatment and Conservation of Vehicles), 060409 (Vehicle Dewaxing) and 060412 (Other). Application of Glues and Adhesives contribute 2.8 per cent of total emissions from solvent and other product use in 2023. Emissions from this sub-category have increased 142.9 per cent between 1990 and 2023 due to

increased consumption of glue products. The methodology for reporting 060404 (Fat, edible and non-edible oil extraction) includes emissions from 1990 to 2023, the methodology is a Tier 2 method based on Oilseed rape crop yield data provided by the CSO and emission factor from the Inventory guidebook (EMEP/EEA 2023).

Emission sources from 2G4 include 060602 (Use of tobacco) and 060303 (Use of shoes). Emissions from SNAP code 060602 (Use of tobacco) has been included in this submission using Tier 2 emission factor from Inventory guidebook (EMEP/EEA 2023) and excise volumes data obtained from The Revenue Commissioners. Emissions from category 2G4 accounted for 0.03 per cent of emissions from solvent and other product use in 2023 and have decreased by 64.5 per cent between 1990 and 2023.

Sector 2H2 (Food and Beverage industry) is the largest contributor (62.1 per cent in 2023) to NMVOC emissions (Table 4.41-4.46 in Annex D). Tier 2 methodologies were applied to SNAP codes 040605 Bread, 040607 Production of Beer and 040608 Production of spirits, 040627 Meat frying and meat rendering, coffee roasting and feedstock, using activity data from CSO and Eurostat. There has been an annual average 6.9 per cent increase in spirit production since 2014 with a notable increase in production (+32 per cent) seen in 2017 followed by a decrease of 6.2 per cent in 2018 and a 16 per cent increase in 2019 and decrease of 9 per cent in 2020 followed by a 17.4 per cent increase in 2021 and a 3.6% increase in 2022. Spirit sales dropped by approximately 4.8 per cent in Ireland in 2020 while exports dropped an estimated 16 per cent due to the impact of COVID-19 hospitality closures and travel restrictions (Drinks Ireland, 2021). A marked recovery was seen in 2021 with exports up by approximately 25 per cent and a further increase of 17.3 per cent in 2022. Emissions from spirit production decreased 6.2 per cent in 2023 compared to 2022.

#### 4.5.1 Domestic Solvent Use including fungicides (NFR 2D3a)

This subcategory covers SNAP sector 060408. This category addresses NMVOC emissions from the general use of products containing solvents by members of the public in their homes but does not include the use of decorative paints. Many domestic products are also used in industry and commerce and in many cases, it is difficult or impossible to separate total sales into domestic and industrial components. Products that contain VOCs can be divided into a number of categories such as Cosmetic and Personal Care Products, Household Products, DIY products, Car Care Products, Varnish remover, Sealant and filling agents, Pharmaceutical Products Use and Pesticides. In this submission, a Tier 2 methodology was used with per-capita emission factors. This is the recommended approach to use where product statistics for the use of the Tier 2b approach are not complete in terms of the product types covered by domestic solvent use. Further study is planned to source appropriate product statistics. Estimates of NMVOC emissions from domestic solvent use are provided in Table 4.14 of electronic Annex D.

#### 4.5.2 Road Paving with asphalt (NFR 2D3b)

This sector covers the use of asphalt for road paving and covers SNAP sector 040611. This source is estimated using a Tier 2 methodology using annual weight of warm and hot mix asphalt used in Ireland for years 1993-2023 and the Tier 2 emission factor from the Inventory guidebook (EMEP/EEA 2023). PM<sub>2.5</sub>, PM<sub>10</sub>, TSP and BC are estimated for all years 1990-2023. Estimates of NMVOC, PM<sub>10</sub>, TSP, PM<sub>2.5</sub> and BC emissions from Road Paving with asphalt as well as activity data are provided in Table 4.15 of electronic Annex D.



### 4.5.3 Coating Application (NFR 2D3d)

This sector covers the use of paints within the industrial, trade and domestic sectors. The term paint includes pigmented coatings and clear coatings such as lacquers and varnishes, with the exception of glues, adhesives and inks. Unless captured on release and either recovered or destroyed, the solvent content of paint can be considered to be emitted to the atmosphere. Due to the range and variation of activity data for coating applications, it cannot be aggregated and included in the NFR tables. Activity data sources are described for the subcategories in sections 4.5.3.1 to 4.5.3.6. The subcategories covered in this source category are presented below, with the relevant SNAP code in parentheses. SNAP codes not included below are deemed not to occur in Ireland.

Paint Application – Car Repairing (060102)

Paint Application – Construction and Buildings (060103)

Paint Application – Domestic Use (060104)

Paint Application – Boat Building (060106)

Paint Application – Wood (060107)

Paint Application – Other Industrial Paint Application (060108)

Paint Application – Other Non-Industrial Paint Application (060109)

Dependent on the SNAP code of interest, both bottom-up and top-down approaches have been used in emission estimates. Where there is an absence of country-specific data, per-capita emission factors derived from a number of EU member states national inventories were used to estimate emissions in Ireland using population statistics as a proxy. Further details of the methodological choices for this source category are provided in Barry and O'Regan. (2014), CTC/AEA (2005) and in Finn et al. (2001).

#### 4.5.3.1 Paint Application: Car Repairing (SNAP 060102)

Activity data was obtained from a number of sources. From 2006-2023, sales data was obtained from a large supplier and data was scaled up based on market share and expert opinion. Data used in 1998 was calculated by Finn et al. (2001), data for 2000 and 2001 was provided by the British Coating Federation. Data was extrapolated and interpolated for the intervening years using passenger car numbers reported by the Department of Transport, each year (DoT, 2022). Emission factors were obtained using survey data from AEA/CTC (2005) and Barry and O'Regan (2014), default emission factors provided by the Inventory Guidebook (EMEP/EEA 2023) and where necessary, emission factors were calculated based on the average decrease in VOC content in known coating applications. The emission estimate includes thinners (EF 1000-835g/L), body fillers (EF 249-175g/L), top coat (720-420g/L) and primers (720-540g/L). This is considered a Tier 2 method. Emission estimates and activity data are provided in Table 4.16 of electronic Annex D.

#### 4.5.3.2 Paint Application: Construction and Building (SNAP 060103) and Domestic Use (SNAP 060104)

Activity data was obtained from the Irish Decorative Surface Coating Association (IDSCA) for the period 2006-2023. The Irish Business and Employers' Confederation (IBEC) collated the total product sales for both water-based and solvent-based paints and provided the information to the inventory agency for the period 2000-2004. Following the experience in the UK (CTC/AEA, 2005), total product sales are

proportioned between trade (Construction and Buildings) and retail (Domestic Use) use, assuming a 44:56 split in 1998, reaching 40:60 in 2003, and 30:70 in 2013. The split for the period 2014-2023 is assumed to be the same as 2013. Estimates of paint sales prior to 1998 were extrapolated using GDP ( $R=0.70$ ).

A number of emission factors were used to calculate NMVOC emissions from decorative coating applications. A survey of products found in popular retail chain stores was completed to establish a realistic emission factor for decorative surface coating products for recent years:

- Interior matt walls and ceiling paint was found to be 30g of VOC/l for solvent based paints and 22.5g of VOC/l for water based paints.
- Interior glossy walls and ceiling paint were found to have 76g of VOC/l for solvent based paint and 50g of VOC/l for water based paints.
- Exterior wall paint of mineral substrate were found to have 126g of VOC/l of solvent based paints and 9g of VOC/l for water based paints.
- Interior/exterior trim and cladding paints for wood and metal have an average solvent content of 324g of VOC/l of solvent based paints and 43g of VOC/l of water based paints.
- Primers were found to have on average for solvent based primers 201g of VOC/l and 45g of VOC/l for water based primers.

These emission factors were used for 2010-2023 while emission factors prior to this were assumed to be similar to 2007 limits outlined in the decorative paints directive. This is considered a Tier 2 method. Emission estimates and activity data for this category are provided in Table 4.17 of electronic Annex D.

#### 4.5.3.3 Paint Application: Boat Building (SNAP 060106)

Paint application in the Marine Sector includes a diverse range of products designed to prevent corrosion and protect ships hulls against damage from fouling. The formulation varies depending on the area being coated and application techniques also vary ranging from spraying to brushing and application by roller.

Activity data were obtained from a major marine coating supplier from 2010-2023 and was upscaled based upon the company's market share. Previous annual emissions were assumed to be the same as 2010. The supplier also provided an estimated industry product breakdown. Emission factors between products are relatively similar with Top coats, primers and anti-corrosion products having an estimated VOC content of 400g of VOC per kg of product while anti-fouling products are estimated to contain 440g of VOC per kg of product.

Paint Application in the Marine Sector in Ireland can be divided into domestic sector, cargo or freight sector and the fishery sector. Larger vessels which require more product application are unable to dry dock in Ireland due to a lack of facilities to handle larger vessels. Therefore, sales data are adjusted based upon expert opinion to account for this (50 per cent of paint sales are applied elsewhere). This sector is a minor emission source for this reason. The methodology is considered a Tier 1 method. Estimates of NMVOC emissions and activity data from this source category are provided in Table 4.18 of electronic Annex D.

#### 4.5.3.4 Paint Application: Wood (SNAP 060107)

This subcategory refers to all paints used for the wood and wooden products sector but excludes the use of wood preservatives and creosote. Some activity data were available; however, no indication of the number of operators or market size was obtainable. Therefore, the emissions estimate was downscaled from UK data where consumption patterns and product range were considered to be comparable to Ireland. This involved using UK emissions data from the UK National Atmospheric Emissions Inventory (NAEI) and calculating per capita emissions (kg/per person) and applying this to Ireland using national population statistics. The methodology is considered a Tier 1 method. Estimates of NMVOC emissions from this source sector and proxy activity data are provided in Table 4.19 of Annex D.

#### 4.5.3.5 Paint Application: Other Industrial Sources (SNAP 060108)

The methodology for this source category involves the use of IPPC emissions data and scaling up to account for emissions in the non-IPPC sector based on information obtained from reporting under the Solvent Directive 1999/13/EC. This category covers paints applied in industrial activities other than those already described in previous sections. Products painted include agricultural, construction and earth-moving equipment, aircraft, cans and drums, domestic appliances, electrical components, freight containers, machine tools, military vehicles, motor-vehicle components, office equipment, paper and plastics, and toys.

The scale of operation varies considerably from large operations employing automated roller coating to small-scale spraying painting. Processes may be enclosed or open air, and both air-dried and stove coatings are used. The emission estimate was up-scaled based on information obtained as a result of Solvents Directive 1999/13/EC. Estimates of NMVOC emissions from this source category are provided in Table 4.20 of Annex D. This methodology is considered a Tier 3 method.

#### 4.5.3.6 Paint Application: Other Non-Industrial Sources (SNAP 060109)

This category refers to the use of high performance protective and/or anti corrosive paints applied to structural steel, concrete and other substrates and any other non-industrial coatings. The sector includes coatings for offshore drilling rigs, production platforms and similar structures as well as road marking paints and non-decorative floor paints. Finn et al. (2001) obtained the activity data for this category as the difference between total paint sales in Ireland according to CSO data on paint sales and that used in other SNAP sectors under SNAP 0601. However, as no other data are available, emissions have been calculated following the advice of CTC/AEA (2005) using extrapolation from UK per capita estimates.

In order to establish whether the use of UK data is appropriate, per capita emissions were compared to other reporting parties. It was found that per capita emissions from this category range from 0.08 to 0.45 kg/person. The UK's per capita estimate is calculated at 0.14 kg per person. This is considered to be a realistic estimate for Irish emissions and was used to estimate emissions. This involved calculating UK per capita emissions (kg/per person) and applying this to Ireland using national population statistics. Estimates of NMVOC emissions from this source category are provided in Table 4.21 of Annex D. The methodology is considered a Tier 1 method.

#### 4.5.4 Degreasing and Dry Cleaning (NFR 2D3e and 2D3f)

Degreasing and Dry Cleaning (2D3e and 2D3f) covers the four subcategories that constitute SNAP Sector 0602. The subcategories for which emission estimates have been made are as follows with the relevant SNAP code in parentheses:

Metal Degreasing (060201)

Dry Cleaning (060202)

Electronic Components (060203)

Other Industrial Cleaning (060204)

Activity data were obtained in the form of net consumption statistics (import minus exports) supplied by the CSO. Solvents included in the emissions estimate include perchloroethylene, dichloromethane, trichloroethylene and hydrocarbons from 1992-2023 before which time data are not available. The methodologies outlined in the Inventory Guidebook (EMEP/EEA 2023) and emissions data collected under the Solvents Directive 1999/13/EC are used to derive emission estimates. Further details of the methodological choices for this source category are provided in Barry, S. and O'Regan B. (2014).

##### 4.5.4.1 Metal Degreasing (SNAP 060201), Electronics Manufacture (SNAP 060203) and Other Industrial Cleaning (SNAP 060204)

Degreasing is a process for cleaning water-insoluble substances, such as grease, fats, oils, waxes, carbon deposits, fluxes and tars, primarily from various metal products, but plastic, fibreglass, printed circuit boards and other products may also be treated by the same process. Therefore, a wide range of activities is covered.

The metalworking industries are the major users of solvent degreasing. Many manufacturers of electronic components also employ degreasing, but it is difficult to differentiate between the emissions emanating from degreasing and those from other sources. As a result, for the purposes of inventory estimates, emissions from Other Industrial Cleaning (060204) and Electronic Manufacture (060203) are included with Metal Degreasing (060201) as national statistics do not facilitate disaggregation of individual sectors.

The Inventory Guidebook (EMEP/EEA 2023) Tier 1 methodology is used for inventory estimates, and solvent consumption statistics (import minus exports) are used as the activity data. The default emission factor of 460g of VOC per kg of cleaning product is used. As data are not available for 1990-1991, the annual emission estimates for these years are assumed to be the same as 1992. Estimates of consumption activity data and NMVOC emissions from this source category are provided in Table 4.22 of Annex D.

##### 4.5.4.2 Dry Cleaning (SNAP 060202)

Dry cleaning refers to any process to remove contamination from furs, leather, down leathers, textiles or other objects made of fibres, using organic solvents. Dry cleaning can be defined as the use of chlorinated organic solvents, principally perchloroethylene, to clean clothes and other textiles.

Emissions and usage data were obtained from the Solvents Directive 1999/13/EC for the years 2008-2010. In addition, the CSO provides information directly to the inventory agency in relation to

perchloroethylene imports and exports. It is assumed that the net consumption (imports minus exports) in any year are used in that year for inventory estimates, even if there is some carryover of stock between years. Data are available for the years 1992 to 2023. Based on the percentage of perchloroethylene used in Dry Cleaning compared to national consumption in 2008-2010 and 2012, emissions were calculated for 1990-2007, 2011 and for 2013-2023. Consumption activity data and estimates of NMVOC emissions from this source sector are provided in Table 4.23 of Annex D. The methodology is considered a Tier 3 methodology.

#### 4.5.5 Chemical Products, Manufacture and Processing (NFR 2D3g)

The mapping of Chemical Products, Manufacture and Processing (2D3g) to SNAP covers 14 subcategories in SNAP Code 0603. These subcategories are all industrial applications and, similar to Coating Application (2D3d), emission sources not included including Asphalt blowing and Leather tanning activities do not occur in Ireland and the notation key “NO” is used for emissions from these categories. The 9 subcategories for which emission estimates are made are as follows with the relevant SNAP code in parentheses:

PVC Processing (060302)  
Polyurethane Processing (060303)  
Rubber Processing (060305)  
Pharmaceutical Products Manufacturing (060306)  
Paints Manufacturing (060307)  
Inks Manufacturing (060308)  
Adhesives Manufacturing (060309)  
Adhesive and Magnetic Tapes, Films and Photographs Manufacturing (060311)  
Textile Finishing (060312)

Activity data sources are described for the subcategories in sections 4.5.5.1 to 4.5.5.8. Information on activity data pertaining to these subcategories has been obtained from IPPC licenced companies with the exclusion of PVC processing (060302) which is based upon expert opinion from Finn et al. (2001). Estimates were up-scaled to reflect national emissions using the number of companies for each sector classified under European industrial activity classifications (NACE Rev.2) provided by the CSO. Emissions from Adhesive and Magnetic Tapes, Films and Photographs Manufacturing (060311) are included under SNAP code (060405) Industrial adhesive usage in section 4.5.7.3 Other Use of Solvents and Related Activities (2D3i-2G). Further details as to the exact methodological choices and the use of Inventory Guidebook (EMEP/EEA 2023) methodologies applied in estimating emissions can be found in Finn et al. (2001), CTC/AEA (2005), and Barry and O'Regan (2014). As emissions data is sourced from a range of IPPC licenced companies the aggregation and reporting of activity data in the NFR for category 2D3g is not possible.

##### 4.5.5.1 Polyvinyl Chloride (PVC) Processing (SNAP 060302)

The manufacture of polyvinyl chloride plastic involves an enclosed reaction or polymerisation step using the basic monomer to produce the resin, a drying step, and a final treating and forming step. Plastics are polymerised in completely enclosed vessels. Treatment of the resin after polymerisation varies with the proposed use. The major sources of air emission in plastics manufacture are the raw

materials or monomers, solvents, or other volatile liquids emitted during the reaction, sublimed solids such as phthalic anhydride emitted in alkyd production, and solvents lost during storage and handling of thinned resins. Processing of PVC is not significant in Ireland. Emission data have been sourced from the installations involved which suggest an emission of 5 t/annum (Finn et al., 2001). The methodology is considered a Tier 1 method.

#### 4.5.5.2 Polyurethane Processing (SNAP 060303)

This category deals with the application and subsequent discharge of organic compounds as blowing agents for creating polyurethane foams. Emissions are from the release of these blowing agents during foaming, or subsequently by the long-term release over several years. Polyurethane is used in building construction, for heat insulation, and for packaging material. For soft polyurethane foams, water may be used. Hard polyurethane foams utilise organic liquids as blowing agents.

Emission data have been sourced from IPPC-licensed companies involved in the manufacture of polyurethane and other foams. Estimates of NMVOC emissions from this category are provided in Table 4.24 of Annex D. The methodology is considered a Tier 1 method.

#### 4.5.5.3 Rubber Processing (SNAP 060305)

No detailed information for rubber processing is available within the Inventory Guidebook (EMEP/EEA 2023). Therefore, it is assumed in inventory estimates that this category includes processes such as moulding and mixing of natural and synthetic rubbers. Operations involving trimming and cutting are ignored since NMVOC emissions would not be associated with such operations.

Emission data have been sourced from IPPC-licensed companies involved in rubber processing that utilise organic solvents. Estimates of NMVOC emissions from this category sector are provided in Table 4.25 of Annex D. Emissions from this sector were dominated by the manufacture of tennis balls from two companies both of which have ceased operation and account for the steep decline in emissions. The last rubber processing facility closed in 2016 and emissions thereafter are reported using the notation key NO. The methodology is considered a Tier 2 method.

#### 4.5.5.4 Pharmaceutical Products Manufacturing (SNAP 060306)

Depending on the nature of the pharmaceutical manufacturing facility, organic chemicals are used in the synthesis, extraction, fermentation and purification of Active Pharmaceutical Ingredients. Solvents are also used in the dilution of liquids, granulation, packaging and film coating. Thousands of individual products are categorised as pharmaceuticals. These products are usually produced in modest quantities in relatively small plants using batch processes. A typical pharmaceutical plant will use the same equipment to make several different products at different times.

The pharmaceutical industry is well established in Ireland and subject to IPPC licence requirements. Emission estimates have been made for 1998 and 2004 using an emission factor of 2 per cent of usage data (Finn et al. 2001, CTC/AEA, 2005) and for 2006-2023 using reported fugitive emissions data supplied by IPPC licenced facilities to the EPA. Other years (1990-1997 and 2005 emissions estimates) are interpolated or extrapolated from these estimates. The methodology is considered a Tier 3 method. Estimates of NMVOC emissions from this source category are provided in Table 4.26 of electronic Annex D.

#### 4.5.5.5 Coating Manufacture: Paint (SNAP 060307)

The manufacture of paint involves the dispersion of coloured oil or pigments in a vehicle, usually an oil or resin, followed by the addition of an organic solvent for viscosity adjustment. Only the physical processes of weighing, mixing, grinding, tinting, thinning and packaging take place. No chemical reactions are involved.

Input and usage data have been sourced from a number of installations for 1998 and 2004 and from 2007 to 2023 emissions data was obtained from AERs. Emissions data were upscaled based on national statistics to reflect national emissions. Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data was available. Other years in the time series are estimated by interpolation and extrapolation. This methodology is considered a Tier 3 method. Emission estimates for NMVOC are provided in Table 4.27 of electronic Annex D.

#### 4.5.5.6 Inks Manufacturing (SNAP 060308)

There are four major classes of printing ink: letterpress and lithographic inks, commonly called oil or paste inks, and flexographic and rotogravure inks, which are referred to as solvent inks. These inks vary considerably in physical appearance, composition, method of application, and drying mechanism. Flexographic and rotogravure inks have many elements in common with the paste inks but differ in that they are of very low viscosity, and they almost always dry by evaporation of highly volatile solvents.

Emissions data were obtained from IPPC licensed facilities for 2008-2023. Where emissions estimates are based on usage data they are calculated based on an assumed emission factor of 2.5 per cent, which is the UK NAEI emission factor for this category (CTC/AEA, 2005). Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data were available. Gaps in the time series were then filled by interpolation and extrapolation. The methodology is considered a Tier 2 method. Estimates of NMVOC emissions from this source category are provided in Table 4.28 of electronic Annex D.

#### 4.5.5.7 Adhesives Manufacturing (SNAP 060309)

This category includes the manufacture of glues and adhesives as it was difficult to derive separate activity data for glues and adhesives from those obtained from IPPC-licensed installations.

Emissions and usage data were supplied for a number of years in the time series (1998, 2006-2023). Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data were available for that period. The methodology is considered a Tier 3 method. Estimates of NMVOC emissions from this source category are provided in Table 4.29 of Annex D.

#### 4.5.5.8 Textile Finishing (SNAP 060312)

Textile fabric finishing is part of the textile finishing industry. In fabric printing, a decorative pattern or design is applied to constructed fabric by roller, flat-screen or rotary-screen methods. Pollutants of interest in fabric printing are VOCs from mineral spirit solvents in print pastes or inks. Solvent use in this sector is usually associated with dry processing rather than wet processing of textiles.



Very little information is available for this activity. Two IPPC-regulated companies provided information to allow estimates to be made for a limited number of years (1998, 2004 and 2006-2015). Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data were available. The remaining years were extrapolated. This is considered a Tier 1 method. The last company closed in 2016 and emissions are reported using the notation key NO for 2016 -2023. Estimates of NMVOC emissions from textile finishing are provided in Table 4.30 of Annex D.

#### 4.5.6 Printing (2D3h)

Printing involves the use of various types of inks, which may contain a proportion of organic solvents which may be diluted before use. Different inks have different proportions of organic solvents and require dilution to different extents. Printing can also require the use of cleaning solvents and organic dampeners. The main printing techniques identified include offset, cold-set web offset, heat-set web offset, sheet-fed offset, rotogravure, flexography, letterpress, and screen-printing.

Activity data information includes usage and emission data, sourced from IPPC-regulated companies and scaled for those not regulated by IPPC based on national statistics and average emissions. Estimates of NMVOC emissions from printing are provided in Table 4.31 of Annex D. The large decrease in emissions is due to abatement measures introduced by the companies operating in Ireland due to the Solvent Directive (1999/13/EC) and a general greater awareness of environmental issues by the print industry. While the printing industry was affected by the economic recession from 2008 in Ireland and resulted in a decreasing emission trend, emissions increased in 2009 and 2010 due to emissions from two IPPC licenced facilities. An overall decrease in emissions has been seen since then. The methodology is considered a Tier 3 method.

#### 4.5.7 Other Use of Solvents and Related Activities (2D3i-2G)

This sector consists of 11 subcategories, which are a mixture of industrial and non-industrial activities, only 8 of which are applicable to Ireland. In NFR 2D3i (Other solvent use) the categories; Mineral wool enduction (060402) and Other (060412) are not considered to occur in Ireland. In NFR 2G4 Other product use the category Use of shoes (060602) is not estimated for Ireland. The 8 subcategories for which emission estimates are made are as follows with the relevant SNAP code in parentheses:

NFR 2D3i;

- Glass Wool Blowing/Enduction (060401)
- Fat, edible and non-edible oil extraction (060404)
- Application of Glues and Adhesives (060405)
- Preservation of Wood (060406)
- Underseal Treatment and Conservation of Vehicles (060407)
- Vehicle Dewaxing (060409)
- 4-stroke engine oil burning

NFR 2G4;

- Use of Fireworks (060601)
- Use of Tobacco (060602)



Both bottom-up and top-down approaches are used in the estimation of emissions from the subcategories outlined, depending on the availability of data for each subcategory. Similar to the other categories, further information in relation to subcategory estimation methodologies can be found in Finn et al. (2001), CTC/AEA (2005) and Barry and O'Regan (2014).

#### 4.5.7.1 Glass Wool Blowing/Enduction (SNAP 060401)

Glass fibre manufacturing is the high-temperature conversion of various raw materials into a homogeneous melt, followed by the fabrication of this melt into glass fibres. The two basic types of glass fibre products, textile and wool, are manufactured by similar processes. Within the category in Ireland, formaldehyde and phenol are used. Usage and emission data have been sourced from one IPPC-regulated company which ceased operation in 2009. Estimates of NMVOC emissions from this source category are provided in Table 4.32 of electronic Annex D.

#### 4.5.7.2 Fat, edible and non-edible oil extraction (SNAP 060404)

This sector covers solvent extraction of edible oils from oilseeds and drying of leftover seeds before resale as animal feed. The extraction of oil from oil seeds is performed either mechanically or through the use of solvents, or both. Where solvent is used, it is generally recovered and cleaned for reuse. The seed may be subjected to solvent treatment many times before all the oil is extracted. The remaining seed residue is then dried and may be used as an animal feed. Emissions of NMVOC, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> for this sector were estimated using statistics obtained from the CSO on the national yield of oilseed. The Inventory Guidebook (EMEP/EEA, 2023) Tier 2 emission factors from Table 3-4 were applied. Activity data and estimates of NMVOC, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are given in Table 4.33 of Annex D.

#### 4.5.7.3 Application of Glues and Adhesives (SNAP 060405)

This sector covers the use of all adhesives excluding domestic adhesive usage and includes adhesive and magnetic tape production (SNAP 060311). These data include adhesives used for publications and packaging, footwear, construction, transport equipment, rubber and plastic products, abrasives, engineering, laminating and other sectors.

This estimate is based upon net consumption statistics (import minus export data) and the Inventory Guidebook (EMEP/EEA, 2023) default emission factor of 522 g/kg of adhesive. The methodology is considered a Tier 2 method. Activity data and estimates of NMVOC emissions from the application of glues and adhesives are provided in Table 4.34 of Annex D.

#### 4.5.7.4 Preservation of Wood (SNAP 060406)

This section refers to emissions from the industrial use of wood preservatives. It does not include emissions from the surface coating of timber with paints, varnishes or lacquer (which are covered under SNAP 060107), and it does not cover the use of wood preservatives by the public at large (which is covered under SNAP 060408). Wood preservation is carried out using solvent-based preservatives, water-based preservatives or creosote. Creosote is an oil product, prepared from coal tar distillation, and contains a high proportion of aromatic compounds such as PAHs. Regulations banning the sale of

creosote took effect from June 2003. However, creosote may still be used for industrial applications, e.g. railway sleepers, telegraph poles and fencing, but with tougher restrictions on its composition and how it is applied. Creosote is gradually being replaced by water-borne preservatives. Preservatives based on organic solvents have a wide-ranging content of organic solvent, usually white spirit or other petroleum-based hydrocarbons. Water-borne preservatives consist of solutions of inorganic salts in water, with Cu, Cr and As (CCA)-based preservatives being the most widely used. Water-borne preservatives are not of concern to this inventory, as they do not contain VOCs.

In addition to bottom-up IPPC-licensed data, usage data was provided by the sole Creosote using company in Ireland. The Inventory Guidebook (EMEP/EEA, 2023) emission factor of 105 g/litre creosote applied is used. The methodology is considered a Tier 3 method. Estimates of NMVOC emissions from wood preservation are provided in Table 4.35 of Annex D.

#### 4.5.7.5 Underseal Treatment and Conservation of Vehicles (SNAP 060407)

The application of coatings to the underside of car bodies is conducted for protection from stone chips and for sound deadening. In the aftermarket sector, coatings are applied to the underside of cars only during repair of damaged bodywork. Finn et al. (2001) stated that sources within the trade suggested that application of underseal in Ireland was zero or minimal. However, further contact with suppliers revealed that a market of 650 l/annum existed at the time (1998). It is assumed that this market existed for all years prior to 1998. However, CTC/AEA (2005) suggested that this market no longer exists in Ireland and that emissions decreased in a linear fashion up to 2003, after which emissions from the activity no longer occur. The approach uses an average solvent content of 20 per cent, a density of 1,000 kg/m<sup>3</sup>, and assumes that 100 per cent of the solvent is emitted. Estimates of NMVOC emissions from this source category are provided in Table 4.36 of Annex D.

#### 4.5.7.6 Vehicle Dewaxing (SNAP 060409)

In the past, some manufacturers of new cars applied a protective covering to parts of the car body after painting to provide protection during transport. Removal of this coating was carried out at the import centres using solvents. However, car manufacturers now invariably use either water-soluble wax that can be removed using hot water or self-adhesive film instead of wax. Consequently, it is assumed that emissions from this activity are now zero. Discussion with car distributors suggested that, historically, 20 per cent of new cars in Ireland were dewaxed and that the practice was discontinued after 2003. An emission factor of 1 kg/car is applied to estimate emissions using vehicle statistics provided by the CSO (Finn et al., 2001). Estimates of NMVOC emissions from vehicle dewaxing are provided in Table 4.37 of Annex D.

#### 4.5.7.7 Use of Tobacco (SNAP 060602)

This category comprises NMVOC, NO<sub>x</sub>, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC emissions from the combustion (smoking) of tobacco products. Activity data was obtained from The Office of the Revenue Commissioners regarding the excise volumes of tobacco and includes an estimation of Illegal tobacco imported to Ireland from an illegal products research report produced by the Office of the Revenue Commissioners (Office of the Revenue Commissioners, 2021). A decrease of 16.3 per cent consumption of tobacco products was seen in 2023. The Inventory guidebook (EMEP/EEA, 2023) Tier

2 emission factors in table 3-15 are applied. Activity data and estimates of emissions from Use of Tobacco are provided in Table 4.38 of electronic Annex D.

#### 4.5.7.8 Use of Fireworks (SNAP 060601)

This category comprises NO<sub>x</sub>, CO, SO<sub>2</sub>, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions from the use of fireworks. Activity data was obtained from The Department of Justice and Equality and relates to the professional use of fireworks in displays. The Inventory guidebook (EMEP/EEA, 2023) Tier 2 emission factors in Table 3-14 are applied. Estimates of emissions from Use of Fireworks are provided in Table 4.39 of Annex D.

#### 4.5.7.9 4-stroke engine lubricant consumption

The COPERT model used for estimating emissions of air pollutants from road transport includes heavy metal emissions from lubricant oil. These metals are assumed to be emitted to the atmosphere when oil burning occurs in the engine combustion chamber. Lubricant consumption in 2-stroke engines is considered intentional, therefore emissions must be reported under 1A3b. On the other hand 4-stroke engine lubricant consumption is undesirable and should not take place. However small amounts are consumed in the combustion chamber and their emissions are reported in 2D3i. Emissions of heavy metals are shown in Table 4.40 of Annex D.

### 4.5.8 Food and Beverage Industry (NFR 2H2)

According to the EMEP/EEA Guidebook (EMEP/EEA, 2023) this sector includes emissions from all processes in the food production chain which occur after the slaughtering of animals and the harvesting of crops as well as drink manufacturing including production of alcoholic beverages. For Ireland, Wine production (040606) is not occurring, Sugar production (040625) has not occurred since 2005, when the last sugar factory was closed (no activity data available), and Flour production (040626) does not occur on a large scale (no activity data available), margarine and fats production is not thought to occur (no activity data available). Emissions include Spirit production, Animal feed production and Bread production which are the most significant source of emissions in the Food and Beverage industry in Ireland. The 6 subcategories for which emission estimates are made for Ireland are as follows with the relevant SNAP code in parentheses, where applicable:

Bread (SNAP 040605)

Beer (SNAP 040607)

Spirits (SNAP 040608)

Meat fish etc. frying/curing (SNAP 040627)

Coffee Roasting

Feedstock

#### 4.5.8.1 Bread (SNAP 040605)

This sector includes bread, cakes and baking products. Activity data on white bread, bread products and baking products production was obtained from EUROSTAT for years 1995 to 2023. The data for years 1990-1994 was taken to be the same as 1995 as no data was available for these years. Tier 2 emission factors from the Inventory guidebook (EMEP/EEA, 2023) were used for bread and cakes, biscuits and breakfast cereals i.e. baking goods. The NMVOC emissions from Bread is given in table 4.41 of Annex D.

#### 4.5.8.2 Beer (SNAP 040607)

This includes mainstream beer production and craft beer production which has seen a steady increase since 2005 and a significant increase in Ireland since 2010. Activity data was obtained from a variety of sources including the Irish Brewers Association reports (ABFI, 2013, 2014, 2015, 2017, 2018, 2020, 2021, 2022, 2023, 2024) and Independent Craft Brewers of Ireland and Bord Bia Report (Feeney, 2015). The Inventory Guidebook (EMEP/EEA, 2023) Tier 2 emission factor of 0.035 kg/hL was used. The NMVOC emissions from Beer production is given in table 4.42 of Annex D.

#### 4.5.8.3 Spirits (SNAP 040608)

Spirit production is a significant source of NMVOC emissions within the Food and Beverage industry due to the growth of the Whiskey production industry in Ireland. In the current submission activity data is based on national statistics for the years 1999-2023. Other years were extrapolated using this data. A Tier 2 emission factor of 15kg/hl alcohol was used from the Inventory Guidebook (EMEP/EEA, 2023). NMVOC emissions from Spirit production in Ireland is given in table 4.43 of Annex D.

#### 4.5.8.4 Meat, fish etc., frying/curing (SNAP 040627)

Emissions mainly occur from the cooking of meat, fish and poultry, releasing fats and oils and their degradation products. Emissions from fish frying and curing were not estimated due to absence of accurate activity data. Activity data for fish frying is under investigation and included in the planned improvement section 4.10. Activity data was obtained from the CSO on tonnes of animal slaughterings in Ireland which is taken to be the equivalent of meat rendered in Ireland. The Inventory guidebook (EMEP/EEA, 2023) emission factor of 0.33 kg/Mg of meat rendered was used. Activity data on human consumption of meat from the CSO was taken to equate to meat frying and using the Inventory Guidebook (EMEP/EEA, 2023) emission factor of 0.3 kg/Mg product. This is considered a Tier 2 method. The NMVOC emissions from meat frying/curing is given in table 4.44 of electronic Annex D.

#### 4.5.8.5 Coffee Roasting

The roasting of coffee beans is a source of NMVOC emissions. This activity does not have a relevant SNAP code. Activity data for unroasted coffee imports was obtained from the UN Comtrade Database and the Tier 2 emission factor from the Inventory Guidebook (EMEP/EEA, 2023) was used to estimate emissions as can be seen in table 4.45 of Annex D.

#### 4.5.8.6 Feedstock

The processing of by-products to produce animal feeds is a source of NMVOC emissions in Ireland. The tonnage of animal feed produced was sourced from the CSO and the Inventory Guidebook (EMEP/EEA, 2023) emission factor of 1 kg/Mg feed was used to estimate emissions from this source as can be seen in table 4.46 of Annex D.

### 4.6 Consumption of POPs and heavy metals 2K (e.g. electrical and scientific equipment)

Ireland does not estimate emissions for 2K for the entire timeseries, even though there is a Tier 1 method in the 2023 EMEP/EEA Guidebook. Ireland has reviewed the limited activity data available by contacting the European Recycling Platform (ERP) Ireland. ERP manage compliance of WEEE and waste batteries and have provided an estimate for 2018 of 0.95kg of batteries in the mercuric oxide category as placed on the market in Ireland by their Battery Producer members in Ireland. WEEE Ireland also provided information from the main waste collection facility of hazardous waste including batteries.

They estimate approximately 4,000kg per year of button cells are collected and sent outside of Ireland for recycling, however it is not known what proportion of button cells contain mercury. Ireland has a functional waste collection system, and all batteries are collected and sent to hazardous waste collection facilities which export this waste to hazardous recycling or disposal facilities outside of Ireland. We do not believe that this waste stream results in mercury emissions to air and any mercury containing products which are placed on the market are collected in the battery mix.

#### 4.7 Other production, consumption, storage, transportation or handling of bulk product (NFR 2L)

The Other production, consumption, storage, transportation or handling of bulk product category in Ireland's air pollutant inventory includes emissions of PCDD/F and PCBs from leakage from electrical equipment and emissions of PCBs from fragmentisers and shredders. The main use of PCBs since the 1970s, when open uses were banned, has been as dielectric fluids in electrical equipment such as transformers and capacitors. However, the production and use of dielectric fluid containing PCBs has been highly regulated since 1986. Releases to the environment have decreased since 1990 as older PCB-containing equipment is taken out of service and is replaced by PCB-free equipment, which reduces the stocks that may lead to PCB emissions. It is also taken into consideration that, in some cases, trace PCDD/F may be present in PCB dielectric fluid. These arise from the original PCB synthesis process and from oxidation during dielectric breakdown events.

Electrical equipment, including white goods and electronic equipment, is partly recycled by breaking down the products in fragmentisers and shredders. Fragments are separated into ferrous scrap, a fraction containing non-ferrous scrap (which would then be processed separately), and a waste fraction that is typically disposed to landfill. Polychlorinated biphenyls are present in the capacitors of old electrical equipment. Hence, there is potential for PCBs to be released to air during fragmentiser operations.

##### 4.7.1 Leakage from Electrical Equipment (NFR 2L)

The release of PCBs to the environment from electrical equipment is very difficult to estimate with any accuracy due to the large number of components potentially containing PCBs, the range of lifetime and replacement rates for PCB components, and the difficulties for users in identifying such components. Polychlorinated biphenyls have never been manufactured in Ireland. Production ceased in the UK in 1977 and in the rest of Europe and North America in 1986. Manufacturers of electrical equipment were then supplied with alternative dielectric media and replacement products entered the market. However, some countries outside the EU and North America continued to produce these substances until recently. Hence, products from those countries may have continued to contain PCBs until the mid-1990s. Current releases to the environment arise principally from the closed electrical appliances that still exist, as their useful life could be up to 40 years.

Activity data are very difficult to obtain on quantities of PCBs in existing transformers and associated leakage rates. A National Inventory of PCB Holdings for Ireland was originally prepared in 2001. This inventory has been updated a number of times, the most recent data corresponding to 2023. The report for this inventory provides an estimate of the total volume of PCB oil (confirmed and suspected) for 2023 of 8.29 m<sup>3</sup>. This estimate includes both inventoried (confirmed) large and small holdings and

estimated non-inventoried (suspected) holdings. Indications are that this is an overestimate and that many of the suspected holdings do not contain any PCBs. The estimate of holdings for 2023 represents a substantial decrease on the peak value in 2009 (522.06 m<sup>3</sup>) following a large decrease in 2008 (114.29 m<sup>3</sup>). This is partly due to methodological changes in the inventory compilation, which has given rise to a step change in the emission estimates. This issue requires further investigation to determine the level of inconsistency that may have been introduced across the current time series. The European Union's Chemical Legislation European Enforcement Network (CLEEN) initiated a project to compare inventories of PCBs in Member States of the EU. The CLEEN project documents summarise a large amount of information held within the EU offices (on PCB stocks) that have been reported by Member States to the EC but have not to date been published or synthesised by the Commission itself. Analysis of the CLEEN data indicates that Ireland has a lower than average PCB per-capita stock when compared with other Member States. All of this qualitative information points towards a lower than average prevalence of PCB-containing materials within electrical equipment in Ireland and this has been taken into consideration in the estimation method used for category 2L.

The derivation of activity data outlined above provides a time series of estimates of PCB-containing oil stocks in Ireland, based on a worst-case assumption that all of the as yet unreported transformer stocks do contain PCBs. The estimates range from 417,620 dm<sup>3</sup> of oil in 1990 to 8,296 dm<sup>3</sup> of oil (as reported by the EPA) in 2023. Data from the UK NAEI indicate that annual emissions of PCBs derived from dielectric fluid stocks can be estimated as 0.5 g PCBs/kg fluid, of which emissions to air comprise 0.06 g PCBs/kg emitted, with the remainder emitted to land. In the absence of source activity and monitoring data, these factors have been used to estimate Ireland's PCB emission estimates. The time series of PCB emissions from leakage of electrical equipment is presented in Table 4.47 of electronic annex D. As noted above, the current data gives rise to a sharp increase in emissions for 2009 to decrease again in 2010 and further in each consecutive year, caused by a change in the methodology used for estimating the volume of dielectric fluid containing PCBs.

The data on PCDD/F concentrations in dielectric PCB fluid from Dyke (1997) give a concentration in PCB dielectric fluid of 83.5 µg I-TEQ/kg of PCBs. It is assumed that the evaporation rate is the same for PCBs and PCDD/F so that for every kilogram of PCBs that is emitted to air, 83.5 µg I-TEQ of PCDD/F are emitted. Using this factor, estimates for PCDD/F emissions to air from dielectric fluid stocks in Ireland have been made. The time series of PCDD/F emissions from leakage from electrical equipment is presented in Table 4.47 of Annex D.

#### 4.7.2 Fragmentisers and Shredders (NFR 2L)

The practice of fragmenting or shredding electrical equipment currently occurs in a small number of IPPC-licensed facilities, where any suspected POP-containing components (e.g. capacitors) are removed and the residual material is then exported. White goods are also exported for recovery or treatment. The recycling of electrical and electronic goods has also been improved since the introduction of the WEEE Regulations in 2005. However, prior to the commencement of the WEEE Regulations and the All Island Fridge & Freezer Collection and Export Scheme in 2004, it is possible that white goods may have been shredded within Ireland, although there is little evidence that such practice was widespread. To provide a worst-case estimate for this potential emission, the UK NAEI activity data have been scaled on a per-capita basis to prepare estimates for Ireland for 1990–2005.



The shredding of End-of-Life Vehicles (ELVs) is another operation that may result in the possible emissions of POPs, and it has been found to be a relatively significant source in other European countries. Currently, two companies operate ELV shredders at three locations. The larger company operates two shredders but undertakes no monitoring of POPs on incoming vehicles or auto residue post-shredding. However, due to the ELV regulations all vehicles are “de-polluted” either on-site or prior to receipt from dismantlers, with all suspected contaminated materials being removed. In addition, the de-polluting process is expected to further improve in future years. Approximately 30,000 tonnes of auto residue (de-polluted vehicles) are shredded annually. Prior to the implementation of the ELV regulations the entire intact vehicle was shredded. Therefore, it can be assumed that for earlier years the shredding of ELVs would have resulted in larger quantities of shredded auto residue, with a higher potential for release of POPs.

Very limited data are available on emissions of POPs to air from fragmentisers, especially for the early part of the time series. Emission estimates for POPs are based on the Inventory Guidebook (2000) factor of 0.004 g/capita/year for PCB emissions from fragmentisers, which is considered to apply in the early part of the time series, around the time of the banning of PCBs (1985). The starting point for the time series of estimates of emissions from fragmentisers in Ireland is the estimated emissions in 1986 using population data and the factor of 0.004 g/capita/year, and this leads to an initial estimate of 14 kg PCBs emitted to air in Ireland in 1986. Assuming a 20-year lifespan of electronic equipment, it is reasonable to assume that 5 per cent of the 1986 emissions are removed each year, as old PCB-containing equipment is disposed of and new PCB-free equipment is used as replacements. This assumption leads to an estimated time series of PCB emissions to air of 11.9 kg in 1990, falling to zero emissions by 2006 and are reported using the notation key “NO” for the period 2006 to 2023. Although this is a very broad “top-down” approach and is subject to significant uncertainty, there are very little additional data available to inform more accurate estimates. Emission estimates for the time series are presented in Table 4.48 of Annex D.

## 4.8 Recalculations in the Industrial Processes Sector

### 4.8.1 Heavy Metals recalculations

There were no recalculations to emission estimates for 1990-2022. Recalculations are as shown in Table 4.50.

### 4.8.2 Particulate matter recalculations

Recalculations occurred in particulate matter emissions, TSP, PM<sub>2.5</sub> and PM<sub>10</sub> in 2A5a Quarrying and mining of minerals other than coal from 1990-2022 due to correction of transcription errors for the years 2021 and 2022.

Recalculations also occurred in emissions of PM<sub>2.5</sub> from 2A5c Storage and Handling of mineral products due to updated activity data in some years over the last decade.

Recalculations to emission estimates for 1990-2022 can be seen in table 4.51.

### 4.8.3 Nitrogen Oxides (NO<sub>x</sub>) recalculations

There were no recalculations to emission estimates for 1990-2022 as can be seen in table 4.51.

#### 4.8.4 Non- Methane Volatile Organic Compounds (NMVOC) recalculations

Recalculations occurred in 2D3d Coating applications NMVOC emissions for 1990-2022 due to the correction of transcription errors in the activity data file. An increase occurred of 0.03 per cent on average across the time series. Emissions in 2D3g Chemical products manufacturing or processing for 1990-2022 decreased 0.14 per cent across the time series due to the correction of an error in calculation methodology in this submission. Recalculations also occurred in subcategory 2D3h Printing due to the updated industry information causing a increase of 3.1 per cent on average across the time series. Recalculations also occurred in subcategory 2G Other product use and 2H2 Food and Beverages Industry causing a minor increase of 0.09 per cent and 0.03 percent on average across the time series due to the updated industry information. Further QAQC checks have been included in solvents files to avoid further transcription errors in emissions calculations. Recalculations can be seen in Table 4.52.

### 4.9 Quality Assurance/Quality Control

Section 4.7 outlines the recalculations which were undertaken in the Industrial Processes sector in this reporting round. The inventory agency will continue to implement QA/QC procedures with respect to the estimates from the Industrial Processes sector in future submissions. The time series spreadsheet system developed for individual categories as described in Section 1.3.5 now allows for simple and efficient checking of activity data, emission factors, annual emissions and aggregated totals. Year-on-year changes immediately highlight any omissions, anomalies or internal errors. Initial checks are conducted by the inventory compiler as part of the calculation process, which is followed by a second check by another member of the inventories team and completion of the QA/QC sheets in calculation workbooks. Further checks have been introduced to activity data files to ensure that the update of IPCC facility emissions activity data from internal databases is accurate.

### 4.10 Overview of Notation Keys

Table 4.49 describes the notation keys used in the 2022 NMVOC emission inventory for Other Solvent and Product Use (2D-2L).

*Table 4.49 Notation keys used in 2023 NMVOC inventory*

NFR	Snap	Description	Notation Key	Reason
2D3a	060411	Domestic use of pharmaceutical products	IE	Included in 060408
2D3e	060203	Electronic components manufacturing	IE	Included under metal degreasing
2D3e	060204	Other industrial cleaning	IE	Included under metal degreasing
2D3g	060311	Adhesive, magnetic tapes, films and photographs manufacturing	IE	Included under 060405

### 4.11 Planned Improvements

The inventory team will continue to review emission estimates for this sector in light of any new information that may become available for future submissions. In addition, the inventory team also plans to continue to outsource contracts on a periodic basis to re-examine and extend the inventory time series with respect to emissions of heavy metals and POPs.



For category 2H2 and PM<sub>10</sub> emissions, Ireland utilises the default Tier 2 emission factors from Table 3-28 on the basis that activity data statistics are available. Table 3-28 does not include emission factors for PM<sub>10</sub>, PM<sub>2.5</sub>, TSP, BC. In fact, the text at the top of page 21 of Chapter 2H2 states that the default Tier 2 emission factors are expected to be more useful than the background emission factors because of the availability of activity data statistics. Ireland estimates handling of agricultural products in NFR category 3.D.d Off-farm storage, handling and transport of bulk agricultural products and so considers additional estimates included in 2H2 may be double counting these emissions.

Ireland is aware of the EFs in the 2023 Guidebook for NFR sector 2D3c Asphalt roofing. Data was requested from the Irish Asphalt Producers association (IAPA), the PRODCOM data was requested from the national statistics agency (CSO) however, neither of these organizations could provide a suitable data source. It has been determined that the production of asphalt roofing materials does not occur in Ireland. This category is thus reported using the notation key “NO”. In relation to emissions of Hg from NFR 2K, Ireland notes that a Tier 1 approach is available in the 2023 EMEP/EEA Inventory Guidebook. Ireland has a functional waste collection system, and all batteries are collected and sent to hazardous waste collection facilities which export this waste to hazardous recycling or disposal facilities outside of Ireland. We do not believe that this waste stream results in mercury emissions to air and any mercury containing products which are placed on the market are collected in the battery mix.

The levels of solvent use and the emissions from solvents are changing substantially in response to product replacement and reformulation and emission controls being implemented under IPPC and the Solvents Directive (1999/13/EC). The reduction of solvent content has been captured in the methodologies, but this has relied on a number of assumptions, and the collection of real data is required to determine emissions with improved confidence.

In addition, liaison with industry will allow refinement of the estimates for activities subject to licensed controls and to reflect abatement measures in the time series. However, the per capita approach to estimating NMVOC emissions will remain the only option in several important categories, such as SNAP code 060107 (Paint Application: Wood). For category 2D3a Domestic Solvent Use including fungicides) Hg from the use of fluorescent tubes, the 2023 Guidebook (EMEP/EEA, 2023) does not have an emission factor for this source and states “Due to the uncertainty around these releases, this source is currently not considered in the Guidebook.” and so it is reported as NA.

Review recommendations and responses are included in this report as Annex H.

**Table 4.50 Recalculations for Industrial Processes 1990–2022 (Heavy Metals)**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
2A3 Glass Production	Pb	t	0.311	0.311	0.311	0.129	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Cd	t	0.007	0.007	0.007	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Hg	t	0.001	0.001	0.001	0.001	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2C Metal Production	Pb	t	3.473	3.387	2.127	0.041	0.035	0.016	0.007	0.016	0.001	0.000	0.000	0.000	0.000	0.000	0.001	0.000	NO
2C Metal Production	Cd	t	0.249	0.238	0.241	0.001	0.008	0.001	0.000	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	NO
2C Metal Production	Hg	t	0.033	0.031	0.036	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	NO
<b>Submission 2025</b>																			
2A3 Glass Production	Pb	t	0.311	0.311	0.311	0.129	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Cd	t	0.007	0.007	0.007	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Hg	t	0.001	0.001	0.001	0.001	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2C Metal Production	Pb	t	3.473	3.387	2.127	0.041	0.035	0.016	0.007	0.016	0.001	0.000	0.000	0.000	0.000	0.000	0.001	0.000	NO
2C Metal Production	Cd	t	0.249	0.238	0.241	0.001	0.008	0.001	0.000	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	NO
2C Metal Production	Hg	t	0.033	0.031	0.036	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	NO
<b>% Change in Emissions</b>																			
2A3 Glass Production	Pb	%	0.0%	0.0%	0.0%	0.0%	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Cd	%	0.0%	0.0%	0.0%	0.0%	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Hg	%	0.0%	0.0%	0.0%	0.0%	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2C Metal Production	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2C Metal Production	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2C Metal Production	Hg	%	0.0%	0.0%	0.0%	-	-	-	-	-	-	-	-	-	-	-	-	-	-

**Table 4.51 Recalculations for Industrial Processes 1990–2022 (Particulates and NOx)**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
2A3 Glass Production	TSP	kt	0.023	0.023	0.022	0.006	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM <sub>10</sub>	kt	0.020	0.020	0.019	0.005	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM <sub>2.5</sub>	kt	0.018	0.018	0.017	0.005	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A5a Quarrying and mining of minerals other than coal	TSP	kt	3.603	3.461	9.834	14.461	5.508	3.773	4.066	3.788	4.090	4.566	5.518	6.043	6.632	7.504	6.884	5.995	5.994
2A5a Quarrying and mining of minerals other than coal	PM <sub>10</sub>	kt	1.154	1.108	3.071	4.508	1.765	1.232	1.323	1.231	1.320	1.461	1.737	1.892	2.070	2.342	2.149	1.876	1.876
2A5a Quarrying and mining of minerals other than coal	PM <sub>2.5</sub>	kt	0.143	0.137	0.373	0.549	0.213	0.148	0.161	0.151	0.160	0.178	0.214	0.233	0.255	0.288	0.263	0.236	0.236
2A5b Construction and Demolition	TSP	kt	1.739	1.928	1.552	6.247	12.492	0.527	0.747	1.845	1.171	1.374	1.138	3.263	0.951	2.819	2.989	3.066	3.012
2A5b Construction and Demolition	PM <sub>10</sub>	kt	0.522	0.578	0.465	1.871	3.733	0.159	0.225	0.553	0.351	0.413	0.343	0.977	0.287	0.845	0.896	0.919	0.902
2A5b Construction and Demolition	PM <sub>2.5</sub>	kt	0.052	0.058	0.047	0.187	0.373	0.016	0.022	0.055	0.035	0.041	0.034	0.098	0.029	0.085	0.090	0.092	0.090
2A5c Storage and Handling of mineral products	PM <sub>2.5</sub>	kt	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2B10b Storage, handling and transport of chemical products	TSP	kt	0.179	0.192	0.173	0.148	0.142	0.120	0.123	0.149	0.140	0.140	0.141	0.155	0.171	0.155	0.159	0.169	0.138
2B10b Storage, handling and transport of chemical products	PM <sub>10</sub>	kt	0.057	0.061	0.055	0.047	0.046	0.039	0.039	0.048	0.045	0.045	0.045	0.050	0.055	0.050	0.051	0.054	0.044

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
2B10b Storage, handling and transport of chemical products	PM <sub>2.5</sub>	kt	0.007	0.008	0.007	0.006	0.006	0.005	0.005	0.006	0.006	0.006	0.006	0.006	0.007	0.006	0.006	0.007	0.006
2C Metal Production	TSP	kt	0.071	0.071	0.006	0.001	0.001	0.001	0.000	0.001	NO	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2C Metal Production	PM <sub>10</sub>	kt	0.060	0.060	0.005	0.001	0.001	0.001	0.000	0.001	NO	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2C Metal Production	PM <sub>2.5</sub>	kt	0.042	0.042	0.003	0.001	0.001	0.000	0.000	0.000	NO	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2D3b Road Paving	TSP	kt	33.000	25.500	43.500	51.000	34.500	27.000	28.500	27.000	27.000	28.500	28.500	31.500	30.750	30.000	28.500	30.000	30.750
2D3b Road Paving	PM <sub>10</sub>	kt	4.400	3.400	5.800	6.800	4.600	3.600	3.800	3.600	3.600	3.800	3.800	4.200	4.100	4.000	3.800	4.000	4.100
2D3b Road Paving	PM <sub>2.5</sub>	kt	0.220	0.170	0.290	0.340	0.230	0.180	0.190	0.180	0.180	0.190	0.190	0.210	0.205	0.200	0.190	0.200	0.205
2D3i Other solvent use	TSP	kt	0.022	0.014	0.009	0.016	0.031	0.061	0.065	0.054	0.038	0.044	0.038	0.046	0.045	0.042	0.050	0.057	0.086
2D3i Other solvent use	PM <sub>10</sub>	kt	0.018	0.012	0.008	0.013	0.025	0.050	0.053	0.044	0.031	0.036	0.031	0.038	0.037	0.035	0.041	0.047	0.070
2D3i Other solvent use	PM <sub>2.5</sub>	kt	0.012	0.008	0.005	0.009	0.017	0.034	0.035	0.030	0.021	0.024	0.020	0.025	0.025	0.023	0.027	0.031	0.047
<b>Submission 2025</b>																			
2A3 Glass Production	TSP	kt	0.023	0.023	0.022	0.006	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM <sub>10</sub>	kt	0.020	0.020	0.019	0.005	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM <sub>2.5</sub>	kt	0.018	0.018	0.017	0.005	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2A5a Quarrying and mining of minerals other than coal	TSP	kt	3.603	3.461	9.834	14.461	5.508	3.773	4.066	3.788	4.090	4.566	5.518	6.043	6.632	7.504	6.884	0.304	0.296
2A5a Quarrying and mining of minerals other than coal	PM <sub>10</sub>	kt	1.154	1.108	3.071	4.508	1.765	1.232	1.323	1.231	1.320	1.461	1.737	1.892	2.070	2.342	2.149	0.133	0.130
2A5a Quarrying and mining of minerals other than coal	PM <sub>2.5</sub>	kt	0.143	0.137	0.373	0.549	0.213	0.148	0.161	0.151	0.160	0.178	0.214	0.233	0.255	0.288	0.263	0.014	0.014

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
2A5b Construction and Demolition	TSP	kt	1.739	1.928	1.552	6.247	12.492	0.527	0.747	1.845	1.171	1.374	1.138	3.263	0.951	2.819	2.989	3.066	3.012
2A5b Construction and Demolition	PM <sub>10</sub>	kt	0.522	0.578	0.465	1.871	3.733	0.159	0.225	0.553	0.351	0.413	0.343	0.977	0.287	0.845	0.896	0.919	0.902
2A5b Construction and Demolition	PM <sub>2.5</sub>	kt	0.052	0.058	0.047	0.187	0.373	0.016	0.022	0.055	0.035	0.041	0.034	0.098	0.029	0.085	0.090	0.092	0.090
2A5c Storage and Handling of mineral products	PM <sub>2.5</sub>	kt	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2B10b Storage, handling and transport of chemical products	TSP	kt	0.179	0.192	0.173	0.148	0.142	0.120	0.123	0.149	0.140	0.140	0.141	0.155	0.171	0.155	0.159	0.169	0.138
2B10b Storage, handling and transport of chemical products	PM <sub>10</sub>	kt	0.057	0.061	0.055	0.047	0.046	0.039	0.039	0.048	0.045	0.045	0.045	0.050	0.055	0.050	0.051	0.054	0.044
2B10b Storage, handling and transport of chemical products	PM <sub>2.5</sub>	kt	0.007	0.008	0.007	0.006	0.006	0.005	0.005	0.006	0.006	0.006	0.006	0.006	0.007	0.006	0.006	0.007	0.006
2C Metal Production	TSP	kt	0.071	0.071	0.006	0.001	0.001	0.001	0.000	0.001	NO	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2C Metal Production	PM <sub>10</sub>	kt	0.060	0.060	0.005	0.001	0.001	0.001	0.000	0.001	NO	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2C Metal Production	PM <sub>2.5</sub>	kt	0.042	0.042	0.003	0.001	0.001	0.000	0.000	0.000	NO	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2D3b Road Paving	TSP	kt	33.000	25.500	43.500	51.000	34.500	27.000	28.500	27.000	27.000	28.500	28.500	31.500	30.750	30.000	28.500	30.000	30.750
2D3b Road Paving	PM <sub>10</sub>	kt	4.400	3.400	5.800	6.800	4.600	3.600	3.800	3.600	3.600	3.800	3.800	4.200	4.100	4.000	3.800	4.000	4.100
2D3b Road Paving	PM <sub>2.5</sub>	kt	0.220	0.170	0.290	0.340	0.230	0.180	0.190	0.180	0.180	0.190	0.190	0.210	0.205	0.200	0.190	0.200	0.205
2D3i Other solvent use	TSP	kt	0.022	0.014	0.009	0.016	0.031	0.061	0.065	0.054	0.038	0.044	0.038	0.046	0.045	0.042	0.050	0.057	0.086
2D3i Other solvent use	PM <sub>10</sub>	kt	0.018	0.012	0.008	0.013	0.025	0.050	0.053	0.044	0.031	0.036	0.031	0.038	0.037	0.035	0.041	0.047	0.070

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
2D3i Other solvent use	PM <sub>2.5</sub>	kt	0.012	0.008	0.005	0.009	0.017	0.034	0.035	0.030	0.021	0.024	0.020	0.025	0.025	0.023	0.027	0.031	0.047
<b>% Change in Emissions</b>																			
2A3 Glass Production	TSP	%	0.0%	0.0%	0.0%	0.0%	-	-	-	-	-	-	-	-	-	-	-	-	-
2A3 Glass Production	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	-	-	-	-	-	-	-	-	-	-	-	-	-
2A3 Glass Production	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	-	-	-	-	-	-	-	-	-	-	-	-	-
2A5a Quarrying and mining of minerals other than coal	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-94.9%	-95.1%
2A5a Quarrying and mining of minerals other than coal	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-92.9%	-93.1%
2A5a Quarrying and mining of minerals other than coal	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-93.9%	-94.2%
2A5b Construction and Demolition	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2A5b Construction and Demolition	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2A5b Construction and Demolition	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2A5c Storage and Handling of mineral products	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	28.9%	1.1%	10.6%	6.7%	0.0%	0.8%	-0.9%	-0.6%	0.1%	0.0%	-5.0%	-2.6%
2B10b Storage, handling and transport of chemical products	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2B10b Storage, handling and transport of	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
chemical products																			
2B10b Storage, handling and transport of chemical products	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2C Metal Production	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2C Metal Production	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2C Metal Production	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3b Road Paving	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3b Road Paving	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3b Road Paving	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3i Other solvent use	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3i Other solvent use	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3i Other solvent use	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
2B2 Nitric Acid Production	NOx	kt	0.960	0.280	0.302	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
<b>Submission 2025</b>																			
2B2 Nitric Acid Production	NOx	kt	0.960	0.280	0.302	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
<b>% Change in Emissions</b>																			
2B2 Nitric Acid Production	NOx	%	0.0%	0.0%	0.0%	-	-	-	-	-	-	-	-	-	-	-	-	-	-

**Table 4.52 Recalculations for Industrial Processes 1990–2022 (NMVOC)**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
2D3a Domestic solvent use																			
including fungicides	NMVOC	kt	7.927	8.143	8.568	9.347	10.298	10.344	10.386	10.434	10.503	10.599	10.716	10.836	10.982	11.128	11.254	11.331	11.532
2D3b Road Paving	NMVOC	kt	0.035	0.027	0.046	0.054	0.037	0.029	0.030	0.029	0.029	0.030	0.030	0.034	0.033	0.032	0.030	0.032	0.033
2D3c Asphalt Roofing	NMVOC	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2D3d Coating Applications	NMVOC	kt	7.233	6.961	6.625	6.300	3.375	3.243	3.026	3.106	3.405	2.891	2.348	2.107	2.047	1.986	1.719	1.938	1.809
2D3e Degreasing	NMVOC	kt	1.485	2.034	0.722	0.812	0.754	0.788	0.856	0.677	0.754	0.585	0.754	1.013	1.455	1.427	1.070	1.027	0.900
2D3f Dry Cleaning	NMVOC	kt	0.282	0.270	0.090	0.085	0.053	0.058	0.075	0.057	0.069	0.044	0.062	0.064	0.072	0.058	0.070	0.056	0.063
2D3g Chemical products manufacturing or processing	NMVOC	kt	3.023	3.024	2.786	0.989	0.750	0.809	0.795	0.775	0.660	0.606	0.749	0.887	0.946	0.556	0.426	0.295	0.425
2D3h Printing	NMVOC	kt	2.912	2.912	1.946	0.812	2.182	1.496	0.973	0.993	1.116	1.295	1.268	1.054	0.944	0.448	0.456	0.403	0.336
2D3i Other solvent use	NMVOC	kt	1.323	1.075	1.444	3.642	1.907	2.052	1.712	1.430	1.749	1.930	1.890	1.983	2.242	2.260	2.310	1.981	2.009
2G Other product use	NMVOC	kt	0.034	0.037	0.039	0.032	0.024	0.025	0.023	0.020	0.019	0.020	0.017	0.022	0.011	0.017	0.016	0.017	0.014
2H2 Food and Beverages Industry	NMVOC	kt	9.617	10.285	9.751	13.531	18.198	18.281	20.788	22.457	19.133	20.444	21.618	26.959	27.034	29.597	26.695	30.086	30.949
<b>Submission 2025</b>																			
2D3a Domestic solvent use																			
including fungicides	NMVOC	kt	7.927	8.143	8.568	9.347	10.298	10.344	10.386	10.434	10.503	10.599	10.716	10.877	11.045	11.211	11.373	11.474	11.721
2D3b Road Paving	NMVOC	kt	0.035	0.027	0.046	0.054	0.037	0.029	0.030	0.029	0.029	0.030	0.030	0.034	0.033	0.032	0.030	0.032	0.033
2D3c Asphalt Roofing	NMVOC	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2D3d Coating Applications	NMVOC	kt	7.233	6.961	6.625	6.300	3.375	3.243	3.026	3.106	3.405	2.891	2.348	2.110	2.050	1.991	1.724	1.938	1.816
2D3e Degreasing	NMVOC	kt	1.485	2.034	0.722	0.812	0.754	0.788	0.856	0.677	0.754	0.585	0.754	1.013	1.455	1.427	1.070	1.027	0.900
2D3f Dry Cleaning	NMVOC	kt	0.282	0.270	0.090	0.085	0.053	0.058	0.075	0.057	0.069	0.044	0.062	0.064	0.072	0.058	0.070	0.056	0.063
2D3g Chemical products manufacturing or processing	NMVOC	kt	3.023	3.024	2.786	0.991	0.751	0.810	0.795	0.777	0.662	0.608	0.744	0.880	0.942	0.556	0.425	0.285	0.425
2D3h Printing	NMVOC	kt	2.912	2.912	1.946	0.943	1.865	1.609	0.852	0.898	0.980	1.023	1.028	0.844	0.730	0.414	0.411	0.290	0.442
2D3i Other solvent use	NMVOC	kt	1.323	1.075	1.444	3.642	1.907	2.052	1.712	1.430	1.749	1.930	1.890	1.983	2.242	2.260	2.310	1.981	2.009
2G Other product use	NMVOC	kt	0.034	0.037	0.039	0.032	0.024	0.025	0.023	0.020	0.019	0.020	0.017	0.022	0.011	0.017	0.016	0.017	0.014



NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
2H2 Food and Beverages Industry	NMVOC	kt	9.617	10.285	9.751	13.531	18.198	18.281	20.788	22.457	19.133	20.444	21.618	26.960	27.035	29.598	26.697	30.088	30.951
<b>% Change in Emissions</b>																			
2D3a Domestic solvent use including fungicides	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.8%	1.1%	1.3%	1.6%
2D3b Road Paving	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3c Asphalt Roofing	NMVOC	%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2D3d Coating Applications	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.2%	0.2%	0.3%	0.0%	0.4%
2D3e Degreasing	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3f Dry Cleaning	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3g Chemical products manufacturing or processing	NMVOC	%	0.0%	0.0%	0.0%	0.2%	0.2%	0.1%	0.0%	0.3%	0.3%	0.3%	-0.7%	-0.7%	-0.4%	-0.2%	-0.3%	-3.6%	0.1%
2D3h Printing	NMVOC	%	0.0%	0.0%	0.0%	16.2%	-14.5%	7.6%	-12.4%	-9.5%	-12.1%	-21.0%	-18.9%	-19.9%	-22.6%	-7.6%	-9.9%	-27.9%	31.5%
2D3i Other solvent use	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2G Other product use	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	3.1%
2H2 Food and Beverages Industry	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

## Chapter Five - Agriculture

### 5.1 Overview of the Agriculture (NFR 3) Sector

The Agriculture sector is the largest source of NH<sub>3</sub> in Ireland and at 115.55 kt accounted for 99.3 per cent of national total NH<sub>3</sub> emissions (116.36 kt) in 2023. The majority of the remainder of national total emissions is attributable to Road Transport (1A3b) and smaller fractions to combustion in the Manufacturing Industry and Construction (1A2), Commercial (1A4a) and Residential (1A4b), Agriculture/Forestry/Fishing: Off-road vehicles and other machinery (1A4cii) sectors, Other product Use (2G) and Biological treatment of waste – Composting (5B1).

Emissions of NH<sub>3</sub> from agriculture in Ireland's inventory are calculated using a Tier 2 approach developed by a member of the inventory team. The methodologies employed follow those utilised by Misselbrook et al. (2004, 2010, 2016), Misselbrook and Gilhespy (2023), and the Inventory Guidebook (EMEP/EEA, 2023) and are described in the following sections. The methodology is based largely on the UK National Ammonia Reduction Strategy Evaluation System (NARSES) model for emissions from livestock and where required the Inventory Guidebook (EMEP/EEA, 2023) is also used (poultry and minor livestock species). For the calculation of NH<sub>3</sub> emissions from nitrogen fertilizer application the Tier 2 approach provided in the Inventory Guidebook (EMEP/EEA, 2023) is adopted.

The trend in emissions of NH<sub>3</sub> from agricultural sources is shown in Figure 5.1. Management of animal manures (3B) produced 48.3 per cent of NH<sub>3</sub> emissions from agriculture in 2023, with the application of inorganic fertiliser, sewage sludge and animal manures deposited and applied to soils (51.7%) accounting for the remainder. The NH<sub>3</sub> emission trend is largely determined by the cattle population and shows a steady increase up to 138.64 kt in 1998 (an increase by 8.0 per cent from 1990). There has been some decline in the populations of cattle and sheep since 1999, as well as a decrease in fertiliser use, which contributed to a downturn in NH<sub>3</sub> emissions in the 1999–2014 period. Emissions for the period 2015 onwards are on an upward trajectory in response to growth plans for the agricultural sector in Ireland and the removal of milk quotas in 2015. Emissions peaked again in 2018 at 129.50 kt. The NH<sub>3</sub> emissions from the agriculture sector in 2023 were 10.0 per cent lower than the emission levels in 1990 and 10.8 per cent lower than the 2018 level.


NMVOCs from agriculture are estimated to be 42.09 kt in 2023, accounting for 39.8 per cent of the NMVOC inventory total (110.25 kt). Emissions from manure management (3B) make up 89.5 per cent of agriculture related NMVOC in 2023, with the remaining 10.5 per cent from cultivated crops (3De). For the calculation of NMVOC emissions, a Tier 2 methodology is adopted for cultivated crops (3De) included in Chapter 3D Table 3.4 of the 2023 Inventory Guidebook (EMEP/EEA, 2023) which exist in Ireland (namely Wheat, Rape and Grass). The Inventory Guidebook (EMEP/EEA, 2023) Tier 1 approach is adopted for cultivated crops (3De) not included in Table 3.4. A Tier 2 approach is utilised for NMVOC emissions from manure management (3B). The trend in NMVOC emissions from agriculture is shown in Figure 5.2.

NO<sub>x</sub> emissions from agriculture are estimated to be 29.44 kt in 2023, accounting for 34.9 per cent of the national total (84.42 kt). Emissions from manure management (3B) make up 2.8 per cent of the agriculture related NO<sub>x</sub> with Inorganic N-fertilizers (3Da1), Animal manure applied to soils (3Da2a), Sewage sludge applied to soils (3Da2b), and Urine and dung deposited by grazing animals (3Da3)


accounting for 37.4 per cent, 22.9 per cent, 0.3 per cent and 36.7 per cent of total agriculture emissions, respectively. Emissions in 2023 were 11.8 per cent below those in 1990. Estimates from manure management (3B) are derived from the Tier 2 mass flow approach to estimate NH<sub>3</sub> emissions, whilst emissions from agricultural soils (3D) utilise the Tier 1 emission factor presented in the Inventory Guidebook. The trend in NO<sub>x</sub> emissions from agriculture is shown in Figure 5.3.

Estimates for agriculture emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> contribute significantly to national totals for these pollutants, accounting for 9.8 per cent, 17.2 per cent and 9.0 per cent of national totals respectively. For these calculations, Tier 1 and 2 approaches provided in the Inventory Guidebook (EMEP/EEA, 2023) are adopted. For PM<sub>2.5</sub> Dairy cattle (3B1a) and Non-dairy cattle (3B1b) are the main drivers of the trend accounting for 81.7 per cent of emissions from agriculture in 2023. For PM<sub>10</sub> the main driver of the trend in emissions from agriculture is Dairy cattle (3B1a) and Non-dairy cattle (3B1b) which accounted for 34.3 per cent of emissions from agriculture in 2023. A similar trend is evident for TSP with Swine (3B3) Non-dairy cattle (3B1b) and Dairy cattle (3B1a) accounting for 26.0 per cent, 20.1 per cent, and 16.7 per cent respectively of the agriculture sector total in 2023. The trend in PM<sub>2.5</sub> is presented in Figure 5.4.


The use of some pesticides in agriculture can be a source of POP emissions, notably due to the trace content of HCB within some pesticides as a contaminant from the manufacturing process. Emissions of HCB from NFR Sector 3Df (Use of pesticides) are a key category in 2023, accounting for 86.1 per cent of national total HCB emissions.




**Figure 5.1. Emission Trend for Ammonia from Agriculture 1990–2023**



**Figure 5.2. Emission Trend for NMVOC from Agriculture 1990–2023**



**Figure 5.3. Emission Trend for NO<sub>x</sub> from Agriculture 1990–2023**



**Figure 5.4. Emission Trend for PM<sub>2.5</sub> from Agriculture 1990–2023**

## 5.2 Manure Management (NFR 3B)

The following sections outline the activity data, assumptions and calculations utilised in estimating NH<sub>3</sub>, NMVOC, NO<sub>x</sub>, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions from agriculture in Ireland. For NH<sub>3</sub> and NO<sub>x</sub>, the Tier 2 methodology uses a mass flow approach based on the concept of the flow of Total Ammoniacal Nitrogen (TAN) through the manure management system. Emissions are calculated for the same animal sub-categories as those utilised in Ireland’s national greenhouse gas inventory (Table E.1 Annex E). The first step in the mass flow approach is the estimation of total annual nitrogen excretion by the animals. For dairy cows and other cattle, Ireland utilises the method described in IPCC (2006), chapter 10, further enhanced by country-specific data on feeding practices and milk production (O’Mara, 2007, Duffy et al., 2025) to estimate N excretion. For all other categories of livestock, national values are utilised. Total nitrogen excretion is then apportioned to that which is deposited in buildings, collection yards (only applicable to dairy cows during lactation) and grazing. Nitrogen excretion values are presented in Table E.3 Annex E.

The method used to estimate NMVOC emissions is the tier 2 approach based on emission factors from Table 3.11 and Table 3.12 of the Inventory Guidebook (EMEP/EEA, 2023) for all livestock categories. The tier 2 approach considers NMVOC emissions from the following: silage stores, silage for feeding, housing, outdoor manure stores, manure application and from grazing animals as outlined in section 3.4 Chapter 3.B of the Inventory Guidebook (EMEP/EEA, 2023).

For TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions are estimated based upon animal numbers using the adapted Tier 1 method and emission factors from the Inventory Guidebook (EMEP/EEA, 2023). Additional information with respect to the EFs used for NH<sub>3</sub>, NMVOC, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emission estimation is provided in Table E.7 of Annex E.

### 5.2.1 Cattle (NFR 3B1)

A Farm Facilities Survey conducted in 2003 (Hyde et al., 2008) and recent analysis by Teagasc (Buckley et al., 2024) provides the basis for the calculation of the number of days housed and the number of days spent grazing by cattle on farms in Ireland across the timeseries. National averages are used for the purpose of inventory calculations. Data for the number of days housed and proportion of manure and therefore nitrogen deposited in the manure in housing systems are presented in Table E.2.1. The number of days spent grazing and the proportion of nitrogen excreted at pasture is then calculated by subtracting these values from the total number of days in the year and the total nitrogen excreted.

Two housing types are distinguished for cattle production systems in Ireland – liquid (slurry-based) and solid manure (deep bedding)-based housing. As a result of differing management practices on farms, a proportion of each of the cattle subdivisions is not housed (out-wintered) and therefore graze pasture for the full year, however this proportion has reduced across the timeseries. The proportion of each sub-category of cattle that is managed in this manner is accounted for in Table E.2.1. For liquid manure-based housing an emission factor of 27.7 per cent of the TAN available in liquid-based housing is applied (Misselbrook and Gilhespy, 2023). The TAN in cattle manure is assumed to be 60 per cent. For solid manure-based housing, emission factors of 16.8 per cent and 4.2 per cent of the TAN available in solid manure-based housing are applied to cattle housed on straw and calves housed on straw, respectively (Misselbrook and Gilhespy, 2023).

In addition to animal housing, emissions are estimated for cow collecting yards used during milking, utilising an emission factor of 22.5 per cent of the TAN available (Misselbrook and Gilhespy, 2023).

The storage of both liquid and solid manure is considered. Liquid manure is stored either below the animals in slatted floor housing or removed from the house to outdoor storage. Emissions are calculated separately for indoor and outdoor storage. It is assumed that a crust will form in the indoor under slat storage. In addition, a fraction of the organic nitrogen in liquid manure is mineralised to TAN before emissions are calculated. A value of 0.1 (Dammgen et al., 2007) is applied as suggested in the Inventory Guidebook (EMEP/EEA, 2023). To fully account for all losses of N from liquid manure during storage, estimates are made of  $N_2O$ , NO and  $N_2$  losses during storage utilising the emission factors provided in the Inventory Guidebook (EMEP/EEA, 2023). An emission factor of 5 per cent of the TAN available in liquid manure stores is applied to estimate  $NH_3$  emissions from liquid manure storage in covered stores and 10 per cent of the TAN available in liquid manure stored in uncovered stores (Misselbrook and Gilhespy, 2023).

Solid manure is generally stored in the shed or outside in heaps. The contribution of the nitrogen content of straw used for bedding is accounted for based on national data on straw used for bedding and the length of the housing period (Hyde et al., 2008, Buckley et al., 2023). Where manures are managed as a solid, a fraction of the TAN is immobilised in organic matter. Immobilisation of nitrogen reduces the potential for  $NH_3$  emissions from solid manures during storage and after landspreading. The value proposed in the Inventory Guidebook (EMEP/EEA, 2023) of  $0.0067 \text{ kg kg}^{-1}$  (Kirchmann and Witter, 1989) is applied. To fully account for all losses of N from solid manure during storage, estimates are made of  $N_2O$ , NO and  $N_2$  losses during storage utilising the emission factors provided in the Inventory Guidebook (EMEP/EEA, 2023). An emission factor of 26.3 per cent of the TAN available in solid manure stores is utilised (Misselbrook and Gilhespy, 2023).

Landspreading emissions are calculated by estimating the quantity of TAN available post storage of the manure (both liquid and solid) and accounting for the period of the year in which it is spread (i.e. spring, summer, autumn and winter) as outlined in Table E.5 Annex E. Additionally, information on the use of low emission spreading techniques (Buckley et al., 2024) is included in emission estimates. Data on broadcast spreading (splashplate), trailing hose application, trailing shoe application and injection are included. For liquid manure, a dry matter value of 6 per cent is assumed and two emission factors are applied for the proportion that is broadcast spread, 48.4 per cent of the TAN available for the proportion applied in summer and 26.1 per cent for proportion applied in spring, autumn and winter (Misselbrook and Gilhespy, 2023). With respect to low emission spreading techniques the values presented in the UNECE Guidance document on preventing and abating ammonia emissions from agricultural sources (UNECE, 2014) are adopted which equate to a 30 per cent, 45 per cent and 70 per cent reduction in emission levels with the use of trailing hose, trailing shoe and injection, respectively. The trailing shoe abatement factor in the TFRN Ammonia guidance document suggests 30-60% reduction in emissions, the half point i.e. 45% reduction is similar to values suggested by Irish work to date (Owusu-Twum et al., 2024). For solid manure, an emission factor of 68.3 per cent of the TAN available is applied regardless of the period of the year in which it is spread (Misselbrook and Gilhespy, 2023). The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2023) of 0.04 kg NO per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from manure (solid and liquid) application to soil.

For cattle (both dairy and non-dairy cattle) grazing an emission factor of 6 per cent of the TAN available at grazing is applied for all cattle categories (Misselbrook and Gilhespy, 2023). The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2023) of 0.04 kg NO per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from excreta deposited during grazing.

NMVOC emissions are estimated using Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2023). Estimates are made for housed cattle, silage feeding, silage store, manure storage, manure application and grazing. The emission factors for silage feeding and silage storage, housing and grazing are combined with the feed intake values (MJ feed intake) from Ireland's Tier 2 approach to the estimation of CH<sub>4</sub> emissions from enteric fermentation and manure management. The feed intake values for silage, grass and concentrates are presented in Table E.6, Annex E with the emission factors used presented in Table E.7. Emissions of NMVOC from manure storage and manure application use the Inventory Guidebook (EMEP/EEA, 2023) approach whereby the NMVOC emissions from livestock buildings are coupled with the ratio of NH<sub>3</sub> emissions from storage and application with those in livestock buildings, respectively.

Emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are estimated using adapted Tier 1 emission factors in table A1.7 of the Annex to the Inventory Guidebook (EMEP/EEA, 2023) split by cattle type and housing category (slurry/solid housing). The emission factors used are presented in Table E.7, Annex E.

### 5.2.2 Sheep (3B2)

Sheep in Ireland are categorised into those on upland and those on lowland areas. Four subcategories exist within both upland and lowland areas, namely ewes, rams, lambs and other sheep more than 1 year old. The CSO publishes sheep population statistics on an annual basis and, to derive the number of heads on both lowland and upland areas, several assumptions are made as follows based on expert opinion. Total ewe and ram numbers are taken as the mean of the June and December CSO censuses. On this basis, the number of ewes and rams are subdivided using hill and lowland flock data from

National Farm Surveys (lowland/upland) for years 2010-2022. The lowland/upland split for years 1990-2009 is extrapolated from the trendline of the 2010-2022 data. The total number of lambs slaughtered in any one-year period is used as the activity data for lambs. Monthly lamb slaughtering figures are available from the CSO. From 2001 onwards, these numbers are adjusted for the number of lambs that originate in Northern Ireland and that are slaughtered in the Republic of Ireland and for the number of lambs that are reared in the Republic but slaughtered in Northern Ireland. National totals are then subdivided similarly to the other categories of sheep. The numbers of other sheep over 1 year old are calculated from unpublished CSO data. Population statistics for each subcategory of sheep are presented in Table E.1, Annex E. Input data with respect to manure management practices are presented in Table E.2 Annex E. Nitrogen excretion coefficients for all subcategories of sheep are provided in Table E.3 Annex E.

The proportion of TAN in the nitrogen excreted by sheep is assumed to be 50 per cent based on the default value in the EMEP/EEA Guidebook 2023. An emission factor of 21.6 per cent of the TAN available in sheep housing (Misselbrook and Gilhespy, 2023) is applied. Information on the number of days that sheep are housed during the winter period is derived from the Farm Facilities Survey (Hyde et al., 2008), which suggests sub-category specific housing period lengths (Table E.2.2 Annex E). More specifically, lowland and upland ewes are assumed to be housed for 84 and 85 days/year. No differentiation is made for upland and lowland rams, lambs and other sheep >1 year old - being housed for 56, 28 and 67 days/year respectively. In Ireland, sheep are generally housed in solid-manure-based housing systems.

The NH<sub>3</sub> emission factors used for the storage and landspreading of solid manure used for Cattle (3B1) are also considered appropriate for sheep manure (Misselbrook and Gilhespy, 2023). For storage of solid manure and subsequent landspreading the emission factors for solid manure from cattle are used (26.3 per cent of TAN available in the manure store and 68.3 per cent of TAN available at landspreading). Account is also taken of the nitrogen added from straw used for bedding and the immobilisation of TAN in organic matter when solid manures are managed following the approach adopted for solid manure from cattle. The emission factors presented in the Inventory Guidebook (EMEP/EEA, 2023) are used to estimate NO, N<sub>2</sub>O and N<sub>2</sub> emissions from manure management. The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2023) of 0.04 kg NO per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from excreta deposited during grazing and the application of solid manure to soil. For sheep grazing, the emission factor applied for cattle of 6 per cent of the TAN available at grazing is utilised to estimate NH<sub>3</sub> emissions from grazing sheep (Misselbrook and Gilhespy, 2023).

NM VOC emissions are estimated using Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2023) based on volatile solid (VS) excretion data as estimated in the national greenhouse gas inventory for agriculture. Estimates are made for housed sheep, manure storage, manure application and grazing sheep. The emission factors used are presented in Table E.7, Annex E.

Emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are estimated using Tier 1 emission factors in the Inventory Guidebook (EMEP/EEA, 2023) The emission factors used are presented in Table E.7, Annex E.

### 5.2.3 Swine (NFR 3B3)

Detailed population statistics are available for seven subcategories of pigs in Ireland using national statistics published by the CSO as follows: sows in pig, gilts in pig, other breeding sows, boars, gilts not



yet served and two categories of fattening pigs (<20 kg and >20 kg live weight). The CSO undertakes and publishes two censuses per year, one in June and one in December. The average of the two census values is used in deriving the average annual pig populations in the seven subcategories, thus providing an appropriate measure of the number of pigs on farms for the purposes of the annual NH<sub>3</sub> emission inventory.

For the NH<sub>3</sub> inventory, it is assumed that all pigs are housed and that the housing systems are liquid/slurry-based. Furthermore, it is assumed that the proportion of TAN in nitrogen excreted is 70 per cent. An emission factor of 29.2 per cent of the TAN in slurry produced by gilts in pig, gilts not yet served and pigs greater than 20 kgs is applied. For sows in pig, other sows for breeding and boars, an emission factor of 28.6 per cent of the TAN produced in the slurry of these pig sub-categories is used and an emission factor of 12.9 per cent of the TAN in slurry produced by pigs under 20 kg (Misselbrook and Gilhespy, 2023).

For slurry storage, emissions of NH<sub>3</sub> are calculated separately for covered and uncovered stores based on the proportion of slurry stored in covered and uncovered storage (Hyde et al., 2008). Emission factors of 13.0 per cent of the TAN available in covered storage and 52.0 per cent of the TAN available in uncovered slurry stores are applied (Misselbrook and Gilhespy, 2023). As is the case with cattle slurry, account is taken of the losses of N<sub>2</sub>O, NO and N<sub>2</sub> using the emission factors presented in the Inventory Guidebook (EMEP/EEA, 2023) and the mineralisation of organic nitrogen in the liquid manure is also accounted for (Dammgen et al., 2007). It is assumed that all pig slurry is applied using the splashplate method up to and including 2019 as no information is currently available on the use of low emission spreading techniques for pig slurry up to that point. Survey data for 2020 suggests that 47 per cent of pig slurry was applied with low emission techniques in that year and assumed to equal that proportion for each subsequent year. An emission factor of 19.2 per cent (Misselbrook and Gilhespy, 2023) of the TAN available post storage is applied on the basis that pig slurry contains less than 4 per cent DM. With respect to low emission techniques an emission reduction coefficient of 30 per cent (equivalent to that from bandspreading) is applied to the proportion of slurry deemed to spread by low emission techniques.

The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2019) of 0.04 kg NO per kg of nitrogen is applied to calculate NO emissions from the application of liquid manure to soil.

NMVOC emissions are estimated using Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2023) based on volatile solid (VS) excretion data as estimated in the national greenhouse gas inventory for agriculture. Estimates are made for housed pigs, manure storage and manure application. The emission factors used are presented in Table E.7, Annex E.

Emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are estimated using Tier 1 emission factors from the Inventory Guidebook (EMEP/EEA, 2023). The emission factors used are presented in Table E.7, Annex E.

#### 5.2.4 Poultry (NFR 3B4g)

Detailed population statistics are available for eight subcategories of poultry in Ireland using national statistics collated by the Department of Agriculture and the CSO as follows: layers, broilers, layer breeders, broiler breeders, turkeys, turkey breeders, geese and ducks. The population statistics are provided in Table E.1, Annex E. The estimation of NH<sub>3</sub> emissions from poultry production utilises bird places as opposed to bird numbers so that production cycles are considered. The number of bird

places is estimated from the annual bird population assuming that all bird places are full throughout the year after rest periods have been taken into account (rest periods are those periods after a production cycle in which the housing systems are emptied of all manure and bedding, thoroughly washed and prepared for the next batch of birds). In the case of broilers, there are 5.5 production cycles per year, with a 3- to 4-week rest period between production cycles. In the case of turkeys, there are 2.5 production cycles per year of approximately 120 days in length, with a 3- to 4-week rest period also applied. All other poultry subcategories have production cycles of over 1 year, and therefore no adjustments to population statistics are made.

It is assumed that all poultry are housed in some form of solid manure housing system and that the proportion of TAN in nitrogen excreted by poultry is 70 per cent. For a proportion of laying birds, free-range systems are in use, which consist of an area of grassland beside the bird house which the birds are allowed onto for a period of hours during the day. The percentage of laying birds that are housed in this type of system is based on statistics supplied by the Department of Agriculture, Food and Marine. An emission factor of 35 per cent of TAN is applied to the quantity of TAN which is deposited outdoors. Emission factors for housing of 20.0 per cent, 21.0 per cent, and 35.0 per cent, 57.0 per cent and 24.0 per cent of TAN are applied to layer, broiler, turkey, geese and duck housing, respectively (EMEP/EEA, 2023). The storage of poultry manure is separated into two classes, litter and layer manure. An emission factor of 30.0 per cent of TAN is applied to broiler manure, 8.0 per cent for layer manure, 24.0 per cent for turkey manure, 16.0 per cent for geese manure and 24.0 per cent for duck manure (EMEP/EEA, 2023). To account for the loss of other nitrogen compounds, account is taken of the losses of  $N_2O$ , NO and  $N_2$  using the emission factors presented in the Inventory Guidebook (EMEP/EEA, 2023). It is assumed that all poultry manure is broadcast spread and emission factors of 45.0 per cent, 38.0 per cent, 54.0 per cent, 45.0 per cent and 54.0 per cent of the TAN available post storage, are adopted for layers, broilers, turkeys, geese and ducks respectively (EMEP/EEA, 2023). The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2023) of 0.04 kg NO per kg of nitrogen is applied to calculate NO emissions from excreta deposited during grazing and the application of manure to soil.

NMVOC, TSP,  $PM_{10}$  and  $PM_{2.5}$  emissions are estimated using Tier 1 emission factors from the Inventory Guidebook (EMEP/EEA, 2023). The emission factors used are presented in Table E.7, Annex E.

#### 5.2.5 Other livestock – Goats (NFR 3B4d), Horses (3B4e), Mules and Asses (NFR 3B4f) and Other Animals (3B4h)

The remaining livestock categories include goats, horses, mules and asses, deer, mink and foxes and estimates are made for emissions of  $NH_3$ , NO, NMVOC, TSP,  $PM_{10}$  and  $PM_{2.5}$ .

For  $NH_3$  emissions the emission factors used are the default Tier 2 factors presented in the Inventory guidebook (EMEP/EEA, 2023) for each step in the manure management chain. As part of the mass flow approach, inventory guidebook emission factors for NO,  $N_2O$  and  $N_2$  emissions from manure management are also estimated. The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2023) of 0.04 kg NO per kg of nitrogen is applied to calculate  $NO_x$  emissions from excreta deposited during grazing and the application of manure to soil.

NMVOC emissions are estimated for goats (3B4d), Horses (3B4e) and Mules and asses (3B4f) using the Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2023). In the case of other animals

(3B4h) the Tier 1 emission factors presented in the inventory guidebook (EMEP/EEA, 2023) are used. The emission factors applied are presented in Table E.7, Annex E.

Tier 1 emission factors from the Inventory Guidebook (EMEP/EEA, 2023) are applied to estimate emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> from the livestock categories described except for deer for which no emission factor is supplied in the Inventory Guidebook (EMEP/EEA, 2023).

In the period 2010 to 2011 there was a change in the Agricultural Waste Management System (AWMS) for mink (3B4h) production from 100% solid manure-based in the period 1990-2010 to 52% liquid/slurry-based and 48% solid manure-based AWMS resulting in a decrease in associated NH<sub>3</sub> and NO<sub>x</sub> emissions for category 3B4h in the period.

### 5.2.6 Uncertainties

There is extensive and up-to-date statistical data on all aspects of the agriculture sector in Ireland. Most of this data is compiled and published by the Central Statistics Office and is the official source of the basic data for inventory purposes. The exception is for statistics on synthetic fertiliser use and the poultry population which are obtained from the Department of Agriculture Food and the Marine (DAFM). The CSO and DAFM are key data providers whose annual statistical inputs to the inventory agency are covered by Memorandum of Understanding (MOU) in Ireland's national inventory system. As a result, the uncertainty associated with animal population statistics is low, at 1 per cent. The emission factor uncertainty associated with NH<sub>3</sub> emission factors for dairy and other cattle is 50 per cent, and for all other livestock categories 100 per cent. Much of NH<sub>3</sub> emission research is aimed at dairy cattle and other cattle therefore emissions from these categories are relatively well quantified in comparison to the other livestock categories. In comparison, the uncertainties associated with NMVOC and PM<sub>10</sub> emissions in agriculture are largely due to the uncertainty (300 per cent) associated with the emission factors for both pollutants as discussed in the Inventory Guidebook (EMEP/EEA, 2023).

## 5.3 Agricultural Soils (NFR 3D)

### 5.3.1 Direct Soil Emissions - Inorganic N-fertilizers (NFR 3Da1)

The calculation of NH<sub>3</sub> emissions from nitrogen fertilizer application to agricultural soils utilises the Tier 2 approach outlined in the Inventory Guidebook (EMEP/EEA, 2023). Total fertilizer sales and emission estimates for each year of the time series 1990-2022 (Table E.4, Annex E) are apportioned into the categories, Ammonium sulphate, CAN, NK mixtures, NPK mixtures, NP mixtures, Other straight N compounds, Urea, and protected urea products, according to the known sales of these compounds in each year as supplied to the inventory agency by the DAFM. For the fertiliser types NPK mixtures, NP mixtures and NK mixtures an emission factor of 24 g NH<sub>3</sub> per kg N applied is used as provided in Annex E.8 of the IIR, this differs from Table 3.2 of the 2023 EMEP/EEA Guidebook, Chapter 3D where the emission factor for NPK and NP mixtures is 84 g NH<sub>3</sub> per kg N applied. This is because as stated in footnote (d) to the emission factor table in the 2023 EMEP/EEA Guidebook "NK mixtures are equivalent to AN and that NPK and NP mixtures are 50% MAP and 50% DAP". In Ireland the NK, NP and NPK mixtures used are AN based and thus the use of an emission factor for NPK and NP which is based on MAP and DAP is not appropriate, hence the use of 24 g NH<sub>3</sub> per kg N applied. Table E.8 of Annex E includes additional information on the methodological approach, quantities of individual fertilizer types, the emission factors used and resultant emissions. The emission factor based on the

original values from Stehfest and Bouwman (2006) as referenced in the Inventory Guidebook (EMEP/EEA, 2023) of 0.038 kg NO<sub>2</sub> per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from the application of inorganic N-fertilizers to soil. Ireland, in previous submissions, reported emissions of PM<sub>10</sub>, PM<sub>2.5</sub> and TSP from farm-level agricultural operations under 3Da1, to separate these emissions from those previously reported under 2B10b. These are now reported under 3Dc following a recommendation in the NECD expert review of Ireland's 2019 submission. The default emission factors presented in the Inventory Guidebook (EMEP/EEA, 2023) are applied (Table E.8, Annex E) coupled with the total utilisable agricultural area (Table E.9, Annex E).

### 5.3.2 Direct Soil Emissions – Livestock manure applied to soils (NFR 3Da2a)

The calculation of NH<sub>3</sub> emissions from livestock manure applied to soil is discussed in sections 5.2.1 to 5.2.5 inclusive. Emissions for each of the livestock species are summed and reported under 3Da2a. The emission factor based on the original values from Stehfest and Bouwman (2006) as referenced in the Inventory Guidebook (EMEP/EEA, 2023) of 0.038 kg NO<sub>2</sub> per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from the application of livestock manure to soils. Emissions of NMVOC from 3Da2a are estimated based on the Tier 2 approach and have been reported along with emissions from manure management in NFR 3B. The notation key included elsewhere (IE) is used for NMVOC for 3Da2a.

From 2018-2022 a significant decrease in NH<sub>3</sub> emissions from application of animal manure on soils is seen, and information regarding the use of low emission spreading techniques (LESS) is sourced from Teagasc's national Farm Survey (Buckley et al., 2023).

### 5.3.3 Direct Soil Emissions – Sewage sludge applied to soils (NFR 3Da2b)

Estimates of NH<sub>3</sub> are estimated from the application of sewage sludge (3Da2b) in this submission. The quantity of sewage sludge applied to land is estimated as part of the calculations for emissions of CH<sub>4</sub> and N<sub>2</sub>O from wastewater in Ireland's Greenhouse Gas Inventory. The fraction of nitrogen volatilised is 0.13 kg NH<sub>3</sub> per kg N applied (EMEP/EEA, 2023). With respect to NO emissions, the emission factor based on the original values from Stehfest and Bouwman (2006) as referenced in the Inventory Guidebook (EMEP/EEA, 2023) of 0.038 kg NO<sub>2</sub> per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from the application of sewage sludge to soils.

### 5.3.4 Direct Soil Emissions – Other organic fertilizers applied to soils (NFR 3Da2c)

Emissions of NH<sub>3</sub> and NO from this source category are currently reported as NE. Information on the quantities of other organic fertilizers applied to agricultural soils does not exist in Ireland and is considered negligible compared to the total quantity of nitrogen applied in manures or deposited during grazing.

### 5.3.5 Direct Soil Emissions – Urine and dung deposited during grazing (NFR 3Da3)

The calculation of NH<sub>3</sub> emissions from urine and dung deposited on soil during grazing is discussed in sections 5.2.1 to 5.2.5 inclusive. Emissions for each of the livestock species are summed and reported under 3Da3. The emission factor based on the original values from Stehfest and Bouwman (2006) as referenced in the Inventory Guidebook (EMEP/EEA, 2023) of 0.038 kg NO<sub>2</sub> per kg of nitrogen is applied to calculate NO<sub>x</sub> emissions from urine and dung deposited on soil during grazing. Emissions of NMVOC from 3Da3 are estimated based on the Tier 2 approach and have been reported along with emissions from manure management in NFR 3B. The notation key included elsewhere (IE) is used for NMVOC for 3Da3.

### 5.3.6 Farm-level agricultural operations (NFR 3Dc)

Ireland reports emissions of PM<sub>10</sub> and PM<sub>2.5</sub> from farm level agricultural operations under 3Dc in this submission following a recommendation from the NECD review in 2018. These emissions are estimated using Tier 2 emission factors for crop operations in wet climate conditions Table 3.6 and Table 3.8, Chapter 3.D of the inventory guidebook (EMEP/EEA, 2023)

### 5.3.7 Off-farm storage, handling and transport (NFR 3Dd)

In this category, fugitive PM emissions in the form of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are estimated from the bulk handling of cereal grains. The general method for estimating fugitive PM emissions involves multiplying the amount of material, which in this case is cereal grain (barley, wheat and oats), by an emission factor. Given the importance of agriculture to Ireland's economy, production statistics are freely available (Table E.9, Annex E). Data in relation to the production of cereal grains are collated and provided by the Central Statistics Office (CSO). Emission factors of 100 g/t, 25 g/t and 4 g/t from CEPMEIP (2001) are utilised in the calculation of emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub>, respectively.

### 5.3.8 Cultivated crops (NFR 3De)

Emissions of NMVOCs are estimated using the default emission factor of 0.86 kg/ha for those crops not listed in Table 3.4 Chapter 3.D of the inventory guidebook (EMEP/EEA, 2023) and the crop areas presented in Table E.9, Annex E. For wheat, rape and grass (15°C), the Tier 2 emission factors presented in Table 3.5 Chapter 3.D of the inventory guidebook (EMEP/EEA, 2023) are used. The areas of these crops and grassland area are presented in Table E.9, Annex E.

### 5.3.9 Use of Pesticides (NFR 3Df)

The main source of POPs from pesticides is HCB contamination of currently used pesticides. Where available, annual pesticides usage data have been used. Pesticide usage surveys are available for 2003, 2004, 2011, 2012, 2013 and 2014 - 2018 from the DAFM. Interpolation and extrapolation are used for the remaining years up to the date at which usage stopped for particular pesticides as described in Table 4 annex to chapter 3.D.f-3.I of the inventory guidebook (EMEP/EEA, 2023). Both the activity data used and impurity factors for each active ingredient are presented in Table E.10 of Annex E.

**Table 5.1 Emission Estimates for Hexachlorobenzene Emissions from Pesticide Use**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
HCB (kg)	7.163	7.247	7.299	2.030	2.151	2.172	2.152	2.124	2.090	2.089	2.041	2.039	2.056	2.072	2.177	2.123

## 5.4 Field burning of agricultural residues (NFR 3F)

Field burning of agricultural residues is the practice whereby crop residues are burnt as a means of clearing land to allow tillage operations to proceed. As stated in chapter 3F of the EMEP/EEA Guidebook 2023 the practice is largely outlawed. Prescribed burning of agricultural residues in Ireland may only be allowed in the case of disease or maintenance of soil organic matter as detailed in GAEC 6 page 52 of the cross-compliance handbook (DAFM, 2015). In addition, the inventory agency has undertaken some analysis on this issue (Zimmerman, 2014) which suggests that fires on agricultural land are in general fires that have occurred due to fires in adjoining forestry or peatland, for example, and are not the anthropogenic burning of agricultural residues. The inventory agency continues to discuss with relevant agencies and research projects with respect to the identification of fires due to

its importance in the estimation of non-CO<sub>2</sub> gases within the Land Use, Land Use Change and Forestry sector.

### 5.5 Uncertainties associated with Synthetic Fertilizer (3Da1), Organic fertilizers (3Da2c) and urine and dung deposited by grazing animals (3Da3).

Although losses of NH<sub>3</sub> from N-fertilisers applied to grass grazed by livestock are difficult to distinguish from subsequent NH<sub>3</sub> emissions from urine patches produced by grazing animals, those two emissions are calculated separately with emissions from grazing reported in 3Da3. The sources making the largest contributions to the overall uncertainty are ammonia losses from synthetic fertiliser use and animal manures deposited to pasture, range and paddock. The emission factors for these sources are currently assigned an uncertainty of  $\pm 200\%$ , and they contribute to 95.6 per cent of the overall uncertainty.

### 5.6 Recalculations in the Agriculture Sector

Recalculations in the agriculture sector are highlighted in tables 5.2 to 5.6. For NH<sub>3</sub> emissions from agriculture, there is a 0.7 per cent decrease annually on average across the timeseries 1990-2022, largely due to the update of the non dairy cattle model and the use of a Tier 2 methodology for sheep resulting in changes to N excretion rates which resulted in a 17.0 per cent decrease on average across the timeseries for category Manure Management Sheep (3B2) and a 5.2 per cent on average increase annually in emissions from Manure Management-Non dairy cattle (3B1b)..

A 6.7 per cent increase for NMVOC emissions across the time series 1990-2022 is evident, due to effect of revised feed intake values for other cattle in the revision to the Tier 2 CH<sub>4</sub> model used to estimate greenhouse gas emissions and the adoption of a Tier 2 model for sheep, this resulted in an increase in emissions for other cattle of 12.8 per cent and a decrease in emissions of 45.7 per cent on average across the time series in Manure Management Sheep (3B2).

For TSP, PM<sub>10</sub> and PM<sub>2.5</sub>, recalculations are negligible and confined to most recent years. Updated harvest statistics for 3Dc, correction of a transcription error for 3B4d are the main recalculations.

### 5.7 Quality Assurance/Quality Control

The general QA/QC procedures set down in Ireland's QA/QC plan have been undertaken for the Agriculture sector. The spreadsheets incorporate transparent linking between input data statistics and calculations, as well as internal checks on the calculations and the outputs. The inventory experts are actively involved in assessing the outcomes of NH<sub>3</sub> emission research in Ireland and continually re-examine the underlying assumptions in inventory estimates with sector-specific experts in the Department of Agriculture and other related bodies.

### 5.8 Planned Improvements

A large number of input variables determine emissions in the Agriculture sector and the final results are very sensitive to changes in many of these variables. Assumptions relating to some parameters have an important bearing on the outcome. Whilst methodologies for the agricultural emission sources that are relevant in Ireland are now very comprehensive, they remain generalised and

necessarily simplified considering the complex systems and processes involved. The key to developing better estimates and reducing uncertainty is to take full account of national circumstances of climate, soil types, livestock- and crop-production practices, manure management systems and other influencing factors in a robust and justifiable manner when applying these methodologies. This requires detailed data from research programmes, and large amounts of statistical data. Nevertheless, the inventory agency is continually developing emission estimates so that they fully reflect national circumstances within the availability of reliable statistics and research studies.

The inventory agency will consider revising the ammonia emission factors, if necessary, based on any updates in the Inventory Guidebook and the NARSES model and following the publication of any relevant country-specific research for the next annual submission. The inventory agency also continues to engage with the DAFM and various other bodies including the research community to identify new mechanisms for the capture of the use of abatement technologies.

Review recommendations and responses are included in this report as Annex H.



**Table 5.2. NH<sub>3</sub> Recalculations for Agriculture 1990–2023**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
Submission 2024																				
3B1a Manure management - Dairy cattle	NH3	kt	14.16	13.05	12.24	10.72	10.45	10.75	10.84	10.97	11.31	12.21	12.89	13.40	14.21	14.98	15.49	15.52	15.67	
3B1b Manure management - Non-dairy cattle	NH3	kt	25.68	29.19	31.10	34.41	32.06	31.16	35.54	35.56	32.81	34.28	35.25	36.82	39.00	34.86	34.32	35.53	35.60	
3B2 Manure management - Sheep	NH3	kt	1.47	1.55	1.45	1.22	0.90	0.93	0.95	0.96	0.96	0.94	0.96	1.03	1.01	0.98	1.08	1.10	1.13	
3B3 Manure management - Swine	NH3	kt	3.79	4.69	5.09	4.92	4.54	4.57	4.51	4.41	4.48	4.42	4.59	4.69	4.68	4.71	4.81	4.97	4.73	
3B4d Manure management - Goats	NH3	kt	0.03	0.03	0.02	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	
3B4e Manure management - Horses	NH3	kt	0.53	0.58	0.60	0.68	0.91	0.91	0.95	0.87	0.81	0.80	0.79	0.73	0.72	0.70	0.75	0.71	0.71	
3B4f Manure management - Mules and asses	NH3	kt	0.05	0.04	0.03	0.03	0.04	0.05	0.06	0.05	0.05	0.05	0.05	0.06	0.05	0.06	0.06	0.06	0.06	
3B4gi Manure mangement - Laying hens	NH3	kt	0.34	0.24	0.23	0.26	0.29	0.29	0.35	0.38	0.40	0.44	0.45	0.47	0.49	0.50	0.51	0.53	0.52	
3B4gii Manure mangement - Broilers	NH3	kt	1.08	1.49	1.67	1.73	1.60	1.55	1.55	1.45	1.63	1.65	1.66	1.64	1.62	1.73	1.85	1.79	1.76	
3B4giii Manure mangement - Turkeys	NH3	kt	1.00	1.07	0.87	0.84	0.58	0.71	0.81	0.74	0.79	0.81	0.84	0.84	0.91	0.86	0.91	1.01	0.99	
3B4giv Manure management - Other poultry	NH3	kt	0.11	0.11	0.11	0.16	0.08	0.08	0.08	0.08	0.09	0.09	0.10	0.11	0.12	0.13	0.14	0.14	0.14	
3B4h Manure management - Other animals (please specify in IIR)	NH3	kt	0.30	0.18	0.19	0.19	0.21	0.21	0.23	0.23	0.23	0.20	0.18	0.16	0.14	0.12	0.09	0.07	0.07	
3Da1 Inorganic N-fertilizers (includes also urea application)	NH3	kt	19.91	19.73	19.86	15.31	19.53	15.59	12.32	13.77	13.99	14.82	16.64	17.30	18.92	16.89	17.18	17.11	18.02	
3Da2a Animal manure applied to soils	NH3	kt	33.96	35.95	36.82	37.88	35.13	35.01	38.81	39.06	37.49	39.72	41.35	43.02	45.37	40.05	36.93	36.26	33.76	
3Da2b Sewage sludge applied to soils	NH3	kt	0.02	0.02	0.10	0.39	0.54	0.38	0.44	0.34	0.28	0.30	0.29	0.30	0.29	0.34	0.25	0.25	0.25	
3Da3 Urine and dung deposited by grazing animals	NH3	kt	12.95	13.74	14.01	13.46	12.37	12.13	12.49	12.63	12.55	12.65	13.06	13.40	13.77	13.52	13.75	14.25	14.40	
Total	NH3	kt	115.36	121.67	124.39	122.21	119.25	114.35	119.94	121.52	117.87	123.42	129.11	133.98	141.31	130.43	128.12	129.32	127.81	
Submission 2025																				
3B1a Manure management - Dairy cattle	NH3	kt	14.16	13.05	12.24	10.72	10.45	10.75	10.84	10.97	11.31	12.21	12.89	13.40	14.21	14.98	15.49	15.52	15.67	
3B1b Manure management - Non-dairy cattle	NH3	kt	34.02	35.20	34.34	34.77	32.13	31.60	32.66	32.45	33.30	33.26	33.98	34.89	33.61	33.81	32.28	31.37	31.19	
3B2 Manure management - Sheep	NH3	kt	1.44	1.50	1.41	1.16	0.83	0.84	0.89	0.90	0.90	0.88	0.89	0.95	0.93	0.91	0.98	1.00	1.02	
3B3 Manure management - Swine	NH3	kt	3.79	4.69	5.09	4.92	4.54	4.57	4.51	4.41	4.48	4.42	4.59	4.69	4.68	4.71	4.81	4.97	4.73	
3B4d Manure management - Goats	NH3	kt	0.03	0.03	0.01	0.01	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.01	0.02	0.01	
3B4e Manure management - Horses	NH3	kt	0.53	0.58	0.60	0.68	0.91	0.91	0.95	0.87	0.81	0.80	0.79	0.73	0.72	0.70	0.75	0.71	0.71	
3B4f Manure management - Mules and asses	NH3	kt	0.05	0.04	0.03	0.03	0.04	0.05	0.06	0.05	0.05	0.05	0.05	0.06	0.05	0.06	0.06	0.06	0.06	
3B4gi Manure mangement - Laying hens	NH3	kt	0.34	0.24	0.23	0.26	0.29	0.29	0.35	0.38	0.40	0.44	0.45	0.47	0.49	0.50	0.51	0.53	0.52	
3B4gii Manure mangement - Broilers	NH3	kt	1.08	1.49	1.67	1.73	1.60	1.55	1.55	1.45	1.63	1.65	1.66	1.64	1.62	1.73	1.85	1.79	1.76	
3B4giii Manure mangement - Turkeys	NH3	kt	1.00	1.07	0.87	0.84	0.58	0.71	0.81	0.74	0.79	0.81	0.84	0.84	0.91	0.86	0.91	1.01	0.99	
3B4giv Manure management - Other poultry	NH3	kt	0.11	0.11	0.11	0.16	0.08	0.08	0.08	0.08	0.09	0.09	0.10	0.11	0.12	0.13	0.14	0.14	0.14	
3B4h Manure management - Other animals (please specify in IIR)	NH3	kt	0.30	0.18	0.19	0.19	0.21	0.21	0.23	0.23	0.23	0.20	0.18	0.16	0.14	0.12	0.09	0.07	0.07	
3Da1 Inorganic N-fertilizers (includes also urea application)	NH3	kt	19.91	19.73	19.86	15.31	19.53	15.59	12.32	13.77	13.99	14.82	16.64	17.30	18.92	16.89	17.18	17.11	18.02	
3Da2a Animal manure applied to soils	NH3	kt	39.53	39.77	38.66	37.70	34.73	34.94	36.25	36.29	37.42	38.46	39.95	41.02	40.64	39.23	36.02	34.33	32.36	
3Da2b Sewage sludge applied to soils	NH3	kt	0.02	0.02	0.10	0.39	0.54	0.38	0.44	0.34	0.28	0.30	0.29	0.30	0.29	0.34	0.34	0.36	0.36	
3Da3 Urine and dung deposited by grazing animals	NH3	kt	12.13	12.66	12.82	12.25	11.14	10.98	11.33	11.17	11.14	11.34	11.76	11.92	12.16	12.22	12.55	12.89	12.92	
Total	NH3	kt	128.43	130.37	128.23	121.12	117.61	113.49	113.29	114.11	116.82	119.76	125.07	128.50	129.50	127.18	123.95	121.87	120.52	
% Change in Emissions																				
3B1a Manure management - Dairy cattle	NH3	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	
3B1b Manure management - Non-dairy cattle	NH3	%	32.5%	20.6%	10.4%	1.0%	0.2%	1.4%	-8.1%	-8.7%	1.5%	-3.0%	-3.6%	-5.3%	-13.8%	-3.0%	-6.0%	-11.7%	-12.4%	



NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
3B2 Manure management - Sheep	NH <sub>3</sub>	%	-1.6%	-2.6%	-2.9%	-4.8%	-8.2%	-9.6%	-7.1%	-6.5%	-6.6%	-6.5%	-7.6%	-7.3%	-7.9%	-7.2%	-8.8%	-9.4%	-9.2%
3B3 Manure management - Swine	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4d Manure management - Goats	NH <sub>3</sub>	%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-16.7%	-27.0%
3B4e Manure management - Horses	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gii Manure management - Broilers	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giii Manure management - Turkeys	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giv Manure management - Other poultry	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da2a Animal manure applied to soils	NH <sub>3</sub>	%	16.4%	10.6%	5.0%	-0.5%	-1.1%	-0.2%	-6.6%	-7.1%	-0.2%	-3.2%	-3.4%	-4.6%	-10.4%	-2.1%	-2.5%	-5.3%	-4.1%
3Da2b Sewage sludge applied to soils	NH <sub>3</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	35.2%	45.4%	43.8%
3Da3 Urine and dung deposited by grazing animals	NH <sub>3</sub>	%	-6.3%	-7.9%	-8.5%	-9.0%	-9.9%	-9.4%	-9.2%	-11.6%	-11.2%	-10.3%	-10.0%	-11.1%	-11.7%	-9.6%	-8.7%	-9.6%	-10.3%
<b>Total</b>	<b>NH3</b>	<b>%</b>	<b>11.3%</b>	<b>7.1%</b>	<b>3.1%</b>	<b>-0.9%</b>	<b>-1.4%</b>	<b>-0.8%</b>	<b>-5.5%</b>	<b>-6.1%</b>	<b>-0.9%</b>	<b>-3.0%</b>	<b>-3.1%</b>	<b>-4.1%</b>	<b>-8.4%</b>	<b>-2.5%</b>	<b>-3.3%</b>	<b>-5.8%</b>	<b>-5.7%</b>

**Table 5.3. NMVOC Recalculations for Agriculture 1990–2023**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
3B1a Manure management - Dairy cattle	NMVOC	kt	10.32	9.63	9.16	8.16	7.92	8.38	8.16	8.13	8.85	9.75	10.20	10.46	10.17	10.66	11.78	11.64	11.59
3B1b Manure management - Non-dairy cattle	NMVOC	kt	21.96	23.48	23.76	25.08	23.67	22.95	23.79	24.59	24.38	23.98	24.91	24.91	24.31	23.76	23.30	22.28	22.10
3B2 Manure management - Sheep	NMVOC	kt	0.57	0.60	0.57	0.46	0.31	0.32	0.35	0.35	0.36	0.35	0.35	0.37	0.37	0.36	0.38	0.38	0.40
3B3 Manure management - Swine	NMVOC	kt	0.95	1.15	1.30	1.27	1.10	1.12	1.09	1.11	1.13	1.09	1.13	1.13	1.14	1.14	1.15	1.19	1.08
3B4d Manure management - Goats	NMVOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	NMVOC	kt	0.11	0.13	0.13	0.15	0.20	0.20	0.20	0.19	0.18	0.17	0.17	0.16	0.16	0.15	0.16	0.15	0.15
3B4f Manure management - Mules and asses	NMVOC	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gi Manure management - Laying hens	NMVOC	kt	0.31	0.23	0.26	0.32	0.35	0.35	0.43	0.47	0.48	0.54	0.55	0.57	0.59	0.60	0.62	0.64	0.63
3B4gii Manure management - Broilers	NMVOC	kt	0.87	1.20	1.34	1.38	1.29	1.24	1.24	1.16	1.31	1.32	1.33	1.31	1.30	1.39	1.48	1.44	1.41
3B4giii Manure management - Turkeys	NMVOC	kt	0.74	0.79	0.65	0.62	0.43	0.53	0.60	0.55	0.58	0.60	0.62	0.62	0.67	0.64	0.67	0.75	0.73
3B4giv Manure management - Other poultry	NMVOC	kt	0.18	0.18	0.18	0.26	0.14	0.14	0.14	0.13	0.14	0.15	0.16	0.18	0.19	0.21	0.23	0.23	0.23
3B4h Manure management - Other animals	NMVOC	kt	0.41	0.25	0.29	0.29	0.36	0.36	0.38	0.38	0.38	0.35	0.31	0.27	0.23	0.20	0.14	0.11	0.11
3Da1 Inorganic N-fertilizers (includes urea application)	NMVOC	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3De Cultivated crops	NMVOC	kt	4.58	4.56	4.59	4.43	4.75	4.74	4.68	4.65	4.63	4.60	4.62	4.66	4.68	4.70	4.69	4.52	4.50
<b>Total</b>	<b>NMVOC</b>	<b>kt</b>	<b>41.01</b>	<b>42.20</b>	<b>42.22</b>	<b>42.43</b>	<b>40.52</b>	<b>40.34</b>	<b>41.07</b>	<b>41.73</b>	<b>42.43</b>	<b>42.91</b>	<b>44.36</b>	<b>44.66</b>	<b>43.83</b>	<b>43.81</b>	<b>44.62</b>	<b>43.34</b>	<b>42.93</b>
<b>Submission 2025</b>																			
3B1a Manure management - Dairy cattle	NMVOC	kt	10.32	9.63	9.16	8.16	7.93	8.38	8.16	8.13	8.85	9.75	10.20	10.47	10.18	10.69	11.83	11.70	11.60
3B1b Manure management - Non-dairy cattle	NMVOC	kt	30.10	29.54	27.66	25.94	25.01	24.43	24.73	25.42	25.93	25.84	26.83	26.94	24.97	25.59	25.48	23.19	22.45
3B2 Manure management - Sheep	NMVOC	kt	0.32	0.33	0.30	0.25	0.18	0.18	0.19	0.19	0.19	0.19	0.19	0.20	0.20	0.19	0.21	0.21	0.22
3B3 Manure management - Swine	NMVOC	kt	0.95	1.15	1.30	1.27	1.10	1.12	1.09	1.11	1.13	1.09	1.13	1.13	1.14	1.14	1.11	1.14	1.04
3B4d Manure management - Goats	NMVOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	NMVOC	kt	0.11	0.13	0.13	0.15	0.20	0.20	0.20	0.19	0.18	0.17	0.17	0.16	0.16	0.15	0.16	0.15	0.15
3B4f Manure management - Mules and asses	NMVOC	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gi Manure management - Laying hens	NMVOC	kt	0.31	0.23	0.26	0.32	0.35	0.35	0.43	0.47	0.48	0.54	0.55	0.57	0.59	0.60	0.62	0.64	0.63
3B4gii Manure management - Broilers	NMVOC	kt	0.87	1.20	1.34	1.38	1.29	1.24	1.24	1.16	1.31	1.32	1.33	1.31	1.30	1.39	1.48	1.44	1.41
3B4giii Manure management - Turkeys	NMVOC	kt	0.74	0.79	0.65	0.62	0.43	0.53	0.60	0.55	0.58	0.60	0.62	0.62	0.67	0.64	0.67	0.75	0.73
3B4giv Manure management - Other poultry	NMVOC	kt	0.18	0.18	0.18	0.26	0.14	0.14	0.14	0.13	0.14	0.15	0.16	0.18	0.19	0.21	0.23	0.23	0.23
3B4h Manure management - Other animals	NMVOC	kt	0.41	0.25	0.29	0.29	0.36	0.36	0.38	0.38	0.38	0.35	0.31	0.27	0.23	0.20	0.14	0.11	0.11
3Da1 Inorganic N-fertilizers (includes urea application)	NMVOC	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3De Cultivated crops	NMVOC	kt	4.58	4.56	4.59	4.43	4.75	4.74	4.68	4.65	4.63	4.60	4.62	4.66	4.68	4.70	4.68	4.54	4.55
<b>Total</b>	<b>NMVOC</b>	<b>kt</b>	<b>48.89</b>	<b>47.99</b>	<b>45.86</b>	<b>43.08</b>	<b>41.73</b>	<b>41.68</b>	<b>41.86</b>	<b>42.39</b>	<b>43.81</b>	<b>44.61</b>	<b>46.12</b>	<b>46.52</b>	<b>44.32</b>	<b>45.50</b>	<b>46.62</b>	<b>44.11</b>	<b>43.13</b>
<b>% Change in Emissions</b>																			
3B1a Manure management - Dairy cattle	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.4%	0.5%	0.1%
3B1b Manure management - Non-dairy cattle	NMVOC	%	37.0%	25.8%	16.4%	3.4%	5.7%	6.5%	4.0%	3.4%	6.4%	7.8%	7.7%	8.1%	2.7%	7.7%	9.4%	4.1%	1.6%

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
3B2 Manure management - Sheep	NMVOC	%	44.5%	45.5%	46.6%	46.3%	43.7%	43.1%	45.3%	46.0%	47.0%	46.6%	45.9%	46.5%	46.6%	46.4%	45.9%	45.2%	45.6%
3B3 Manure management - Swine	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-3.7%	-3.7%	-3.7%
3B4d Manure management - Goats	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-12.2%
3B4e Manure management - Horses	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gii Manure management - Broilers	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giii Manure management - Turkeys	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giv Manure management - Other poultry	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	NMVOC	%	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3De Cultivated crops	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.2%	0.5%	1.3%
<b>Total</b>	<b>NMVOC</b>	<b>%</b>	<b>19.2%</b>	<b>13.7%</b>	<b>8.6%</b>	<b>1.5%</b>	<b>3.0%</b>	<b>3.3%</b>	<b>1.9%</b>	<b>1.6%</b>	<b>3.3%</b>	<b>4.0%</b>	<b>4.0%</b>	<b>4.2%</b>	<b>1.1%</b>	<b>3.8%</b>	<b>4.5%</b>	<b>1.8%</b>	<b>0.5%</b>

**Table 5.4. TSP Recalculations for Agriculture 1990–2023**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
3B1a Manure management - Dairy cattle	TSP	kt	0.64	0.64	0.60	0.54	0.54	0.56	0.58	0.59	0.62	0.66	0.69	0.71	0.74	0.78	0.81	0.79	0.80
3B1b Manure management - Non-dairy cattle	TSP	kt	0.99	0.98	1.00	1.02	0.99	0.96	1.00	1.05	1.04	1.04	1.08	1.12	1.11	1.06	1.06	1.00	1.01
3B2 Manure management - Sheep	TSP	kt	0.10	0.11	0.10	0.08	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.07	0.06	0.06	0.07	0.07	0.07
3B3 Manure management - Swine	TSP	kt	0.98	1.24	1.36	1.32	1.22	1.24	1.23	1.20	1.22	1.20	1.25	1.28	1.28	1.30	1.32	1.37	1.32
3B4d Manure management - Goats	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	TSP	kt	0.01	0.02	0.02	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4f Manure management - Mules and asses	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	TSP	kt	0.35	0.26	0.30	0.37	0.41	0.41	0.49	0.54	0.55	0.62	0.63	0.66	0.68	0.69	0.71	0.74	0.72
3B4gii Manure management - Broilers	TSP	kt	0.32	0.44	0.50	0.51	0.48	0.46	0.46	0.43	0.49	0.49	0.49	0.49	0.48	0.51	0.55	0.53	0.52
3B4giii Manure management - Turkeys	TSP	kt	0.17	0.18	0.15	0.14	0.10	0.12	0.13	0.12	0.13	0.14	0.14	0.14	0.15	0.14	0.15	0.17	0.16
3B4giv Manure management - Other poultry	TSP	kt	0.05	0.05	0.05	0.08	0.04	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.06	0.06	0.07	0.07	0.07
3B4h Manure management - Other animals	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	TSP	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	TSP	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dd Off-farm storage, handling and transport of bulk agricultural products	TSP	kt	0.20	0.18	0.22	0.19	0.20	0.25	0.21	0.24	0.26	0.26	0.23	0.24	0.19	0.24	0.20	0.25	0.25
<b>Total</b>	<b>TSP</b>	<b>kt</b>	<b>3.82</b>	<b>4.11</b>	<b>4.29</b>	<b>4.28</b>	<b>4.07</b>	<b>4.13</b>	<b>4.24</b>	<b>4.29</b>	<b>4.44</b>	<b>4.54</b>	<b>4.64</b>	<b>4.78</b>	<b>4.77</b>	<b>4.87</b>	<b>4.95</b>	<b>5.01</b>	<b>4.95</b>
<b>Submission 2025</b>																			
3B1a Manure management - Dairy cattle	TSP	kt	0.64	0.64	0.60	0.54	0.54	0.56	0.58	0.59	0.62	0.66	0.69	0.71	0.74	0.78	0.81	0.79	0.81

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
3B1b Manure management - Non-dairy cattle	TSP	kt	0.99	0.98	1.00	1.02	0.99	0.96	1.00	1.05	1.04	1.04	1.08	1.12	1.11	1.06	1.06	1.00	1.00
3B2 Manure management - Sheep	TSP	kt	0.10	0.11	0.10	0.08	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.07	0.06	0.06	0.07	0.07	0.07
3B3 Manure management - Swine	TSP	kt	0.98	1.24	1.36	1.32	1.22	1.24	1.23	1.20	1.22	1.20	1.25	1.28	1.28	1.30	1.32	1.37	1.32
3B4d Manure management - Goats	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	TSP	kt	0.01	0.02	0.02	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4f Manure management - Mules and asses	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	TSP	kt	0.35	0.26	0.30	0.37	0.41	0.41	0.49	0.54	0.55	0.62	0.63	0.66	0.68	0.69	0.71	0.74	0.72
3B4gii Manure management - Broilers	TSP	kt	0.32	0.44	0.50	0.51	0.48	0.46	0.46	0.43	0.49	0.49	0.49	0.49	0.48	0.51	0.55	0.53	0.52
3B4giii Manure management - Turkeys	TSP	kt	0.17	0.18	0.15	0.14	0.10	0.12	0.13	0.12	0.13	0.14	0.14	0.14	0.15	0.14	0.15	0.17	0.16
3B4giv Manure management - Other poultry	TSP	kt	0.05	0.05	0.05	0.08	0.04	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.06	0.06	0.07	0.07	0.07
3B4h Manure management - Other animals	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	TSP	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	TSP	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dd Off-farm storage, handling and transport of bulk agricultural products	TSP	kt	0.20	0.18	0.22	0.19	0.20	0.25	0.21	0.24	0.26	0.26	0.23	0.24	0.19	0.24	0.20	0.25	0.25
<b>Total</b>	<b>TSP</b>		<b>3.82</b>	<b>4.11</b>	<b>4.29</b>	<b>4.28</b>	<b>4.07</b>	<b>4.13</b>	<b>4.24</b>	<b>4.29</b>	<b>4.44</b>	<b>4.54</b>	<b>4.64</b>	<b>4.78</b>	<b>4.77</b>	<b>4.87</b>	<b>4.95</b>	<b>5.01</b>	<b>4.95</b>
<b>% Change in Emissions</b>																			
3B1a Manure management - Dairy cattle	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%
3B1b Manure management - Non-dairy cattle	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.9%
3B2 Manure management - Sheep	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B3 Manure management - Swine	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4d Manure management - Goats	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-12.4%
3B4e Manure management - Horses	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gii Manure management - Broilers	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giii Manure management - Turkeys	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giv Manure management - Other poultry	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	TSP	%																	
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	TSP	%																	
3Dd Off-farm storage, handling and transport of bulk agricultural products	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
<b>Total</b>	<b>TSP</b>	<b>%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>

**Table 5.5. PM<sub>10</sub> Recalculations for Agriculture 1990–2023**

NFR Category	Pollutan t	Uni t	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
3B1a Manure management - Dairy cattle	PM <sub>10</sub>	kt	0.33	0.34	0.32	0.28	0.28	0.30	0.30	0.31	0.33	0.35	0.36	0.37	0.39	0.41	0.42	0.42	0.42
3B1b Manure management - Non-dairy cattle	PM <sub>10</sub>	kt	0.60	0.61	0.63	0.63	0.64	0.62	0.64	0.67	0.67	0.66	0.68	0.71	0.71	0.68	0.67	0.63	0.63
3B2 Manure management - Sheep	PM <sub>10</sub>	kt	0.04	0.04	0.04	0.03	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03
3B3 Manure management - Swine	PM <sub>10</sub>	kt	0.15	0.18	0.20	0.19	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.19	0.19	0.19	0.19	0.20	0.19
3B4d Manure management - Goats	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM <sub>10</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4f Manure management - Mules and asses	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM <sub>10</sub>	kt	0.07	0.05	0.06	0.08	0.09	0.09	0.10	0.11	0.12	0.13	0.13	0.14	0.14	0.15	0.15	0.16	0.15
3B4gii Manure management - Broilers	PM <sub>10</sub>	kt	0.16	0.22	0.25	0.26	0.24	0.23	0.23	0.22	0.24	0.24	0.25	0.24	0.24	0.26	0.27	0.27	0.26
3B4giii Manure management - Turkeys	PM <sub>10</sub>	kt	0.17	0.18	0.15	0.14	0.10	0.12	0.13	0.12	0.13	0.14	0.14	0.14	0.15	0.14	0.15	0.17	0.16
3B4giv Manure management - Other poultry	PM <sub>10</sub>	kt	0.05	0.05	0.05	0.08	0.04	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.06	0.06	0.07	0.07	0.07
3B4h Manure management - Other animals	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM <sub>10</sub>	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM <sub>10</sub>	kt	1.21	1.12	1.11	1.11	1.05	1.14	1.20	1.17	1.15	1.11	1.08	1.06	1.02	1.04	1.03	1.05	1.10
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM <sub>10</sub>	kt	0.05	0.04	0.05	0.05	0.05	0.06	0.05	0.06	0.06	0.07	0.06	0.06	0.05	0.06	0.05	0.06	0.06
<b>Total</b>	<b>PM<sub>10</sub></b>	<b>kt</b>	<b>2.83</b>	<b>2.85</b>	<b>2.87</b>	<b>2.86</b>	<b>2.70</b>	<b>2.81</b>	<b>2.92</b>	<b>2.91</b>	<b>2.96</b>	<b>2.95</b>	<b>2.97</b>	<b>3.01</b>	<b>2.97</b>	<b>3.01</b>	<b>3.04</b>	<b>3.06</b>	<b>3.09</b>
<b>Submission 2025</b>																			
3B1a Manure management - Dairy cattle	PM <sub>10</sub>	kt	0.33	0.34	0.32	0.28	0.28	0.30	0.30	0.31	0.33	0.35	0.36	0.37	0.39	0.41	0.42	0.42	0.42
3B1b Manure management - Non-dairy cattle	PM <sub>10</sub>	kt	0.60	0.61	0.63	0.63	0.64	0.62	0.64	0.67	0.67	0.66	0.68	0.71	0.71	0.68	0.67	0.63	0.63
3B2 Manure management - Sheep	PM <sub>10</sub>	kt	0.04	0.04	0.04	0.03	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03
3B3 Manure management - Swine	PM <sub>10</sub>	kt	0.15	0.18	0.20	0.19	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.19	0.19	0.19	0.19	0.20	0.19
3B4d Manure management - Goats	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM <sub>10</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4f Manure management - Mules and asses	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM <sub>10</sub>	kt	0.07	0.05	0.06	0.08	0.09	0.09	0.10	0.11	0.12	0.13	0.13	0.14	0.14	0.15	0.15	0.16	0.15
3B4gii Manure management - Broilers	PM <sub>10</sub>	kt	0.16	0.22	0.25	0.26	0.24	0.23	0.23	0.22	0.24	0.24	0.25	0.24	0.24	0.26	0.27	0.27	0.26
3B4giii Manure management - Turkeys	PM <sub>10</sub>	kt	0.17	0.18	0.15	0.14	0.10	0.12	0.13	0.12	0.13	0.14	0.14	0.14	0.15	0.14	0.15	0.17	0.16
3B4giv Manure management - Other poultry	PM <sub>10</sub>	kt	0.05	0.05	0.05	0.08	0.04	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.06	0.06	0.07	0.07	0.07

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
3B4h Manure management - Other animals	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM <sub>10</sub>	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM <sub>10</sub>	kt	1.21	1.12	1.11	1.11	1.05	1.14	1.20	1.17	1.15	1.11	1.08	1.06	1.02	1.04	1.03	1.07	1.11
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM <sub>10</sub>	kt	0.05	0.04	0.05	0.05	0.05	0.06	0.05	0.06	0.06	0.07	0.06	0.06	0.05	0.06	0.05	0.06	0.06
<b>Total</b>	<b>PM<sub>10</sub></b>	<b>kt</b>	<b>2.83</b>	<b>2.85</b>	<b>2.87</b>	<b>2.86</b>	<b>2.70</b>	<b>2.81</b>	<b>2.92</b>	<b>2.91</b>	<b>2.96</b>	<b>2.95</b>	<b>2.97</b>	<b>3.01</b>	<b>2.97</b>	<b>3.01</b>	<b>3.05</b>	<b>3.07</b>	<b>3.10</b>
<b>% Change in Emissions</b>																			
3B1a Manure management - Dairy cattle	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%
3B1b Manure management - Non-dairy cattle	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%
3B2 Manure management - Sheep	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B3 Manure management - Swine	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4d Manure management - Goats	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-12%
3B4e Manure management - Horses	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gii Manure management - Broilers	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giii Manure management - Turkeys	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giv Manure management - Other poultry	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	PM <sub>10</sub>	%																	
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM <sub>10</sub>	%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	1%	2%	1%
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
<b>Total</b>	<b>PM<sub>10</sub></b>	<b>%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.2%</b>	<b>0.5%</b>	<b>0.4%</b>

**Table 5.6. PM<sub>2.5</sub> Recalculations for Agriculture 1990–20223**

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																			
3B1a Manure management - Dairy cattle	PM <sub>2.5</sub>	kt	0.22	0.22	0.21	0.18	0.19	0.19	0.20	0.20	0.21	0.23	0.24	0.24	0.25	0.27	0.28	0.27	0.27
3B1b Manure management - Non-dairy cattle	PM <sub>2.5</sub>	kt	0.39	0.40	0.41	0.42	0.42	0.40	0.42	0.44	0.44	0.44	0.45	0.47	0.46	0.44	0.44	0.41	0.42
3B2 Manure management - Sheep	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B3 Manure management - Swine	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4d Manure management - Goats	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4f Manure management - Mules and asses	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM <sub>2.5</sub>	kt	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gii Manure management - Broilers	PM <sub>2.5</sub>	kt	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03
3B4giii Manure management - Turkeys	PM <sub>2.5</sub>	kt	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
3B4giv Manure management - Other poultry	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4h Manure management - Other animals	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM <sub>2.5</sub>	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM <sub>2.5</sub>	kt	0.05	0.05	0.05	0.05	0.04	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
<b>Total</b>	<b>PM<sub>2.5</sub></b>	<b>kt</b>	<b>0.75</b>	<b>0.76</b>	<b>0.76</b>	<b>0.74</b>	<b>0.73</b>	<b>0.73</b>	<b>0.76</b>	<b>0.78</b>	<b>0.80</b>	<b>0.80</b>	<b>0.83</b>	<b>0.85</b>	<b>0.86</b>	<b>0.85</b>	<b>0.86</b>	<b>0.84</b>	<b>0.84</b>
<b>Submission 2025</b>																			
3B1a Manure management - Dairy cattle	PM <sub>2.5</sub>	kt	0.22	0.22	0.21	0.18	0.19	0.19	0.20	0.20	0.21	0.23	0.24	0.24	0.25	0.27	0.28	0.27	0.28
3B1b Manure management - Non-dairy cattle	PM <sub>2.5</sub>	kt	0.39	0.40	0.41	0.42	0.42	0.40	0.42	0.44	0.44	0.44	0.45	0.47	0.46	0.44	0.44	0.41	0.41
3B2 Manure management - Sheep	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B3 Manure management - Swine	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4d Manure management - Goats	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4f Manure management - Mules and asses	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM <sub>2.5</sub>	kt	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gii Manure management - Broilers	PM <sub>2.5</sub>	kt	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03
3B4giii Manure management - Turkeys	PM <sub>2.5</sub>	kt	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
3B4giv Manure management - Other poultry	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4h Manure management - Other animals	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM <sub>2.5</sub>	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM <sub>2.5</sub>	kt	0.05	0.05	0.05	0.05	0.04	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM <sub>2.5</sub>	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
<b>Total</b>	<b>PM<sub>2.5</sub></b>	<b>kt</b>	<b>0.75</b>	<b>0.76</b>	<b>0.76</b>	<b>0.74</b>	<b>0.73</b>	<b>0.73</b>	<b>0.76</b>	<b>0.78</b>	<b>0.80</b>	<b>0.80</b>	<b>0.83</b>	<b>0.85</b>	<b>0.86</b>	<b>0.85</b>	<b>0.86</b>	<b>0.84</b>	<b>0.84</b>
<i>% Change in Emissions</i>																			
3B1a Manure management - Dairy cattle	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%
3B1b Manure management - Non-dairy cattle	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-1.0%
3B2 Manure management - Sheep	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B3 Manure management - Swine	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4d Manure management - Goats	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-12.4%
3B4e Manure management - Horses	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gii Manure management - Broilers	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giii Manure management - Turkeys	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4giv Manure management - Other poultry	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	PM <sub>2.5</sub>	%																	
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%	1.6%	1.3%
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
<b>Total</b>	<b>PM<sub>2.5</sub></b>	<b>%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.1%</b>	<b>-0.1%</b>



## Chapter Six - Waste

### 6.1 Overview of the Waste (NFR 5) Sector

Emissions from the Waste sector cover several different source categories and pollutants. These are detailed below in Table 6.1. All sources are considered in detail in this chapter.

**Table 6.1 Pollutant Emissions by Waste Source Category**

NFR Source Category	Pollutants
5A Biological treatment of waste - Solid waste disposal on land	NMVOC, Hg, PCDD/F, PCB, TSP, PM <sub>10</sub> , PM <sub>2.5</sub>
5B1 Biological treatment of waste - Composting	NH <sub>3</sub> , CO
5B2 Biological treatment of waste - Anaerobic digestion at biogas facilities	NO, NA, NE
5C1a Municipal waste incineration	NO
5C1bi Industrial waste incineration	NO <sub>x</sub> , SO <sub>2</sub> , NMVOC, CO, TSP, PM <sub>2.5</sub> , PM <sub>10</sub> , BC, Pb, Cd, Hg, As, Cr, Cu, Ni, PCDD/F, B[a]P, B[b]F, B[k]F, HCB, PCB
5C1bii Hazardous waste incineration	IE (5C1bi)
5C1biii Clinical waste incineration	NO <sub>x</sub> , SO <sub>2</sub> , NMVOC, CO, TSP, PM <sub>2.5</sub> , PM <sub>10</sub> , BC, Pb, Cd, Hg, As, Cr, Cu, Ni, PCDD/F, B[a]P, B[b]F, B[k]F, HCB, PCB
5C1biv Sewage sludge incineration	NO, NA, IE
5C1bv Cremation	NO <sub>x</sub> , SO <sub>2</sub> , NMVOC, CO, TSP, PM <sub>2.5</sub> , PM <sub>10</sub> , BC, Pb, Cd, Hg, As, Cr, Cu, Ni, PCDD/F, B[a]P, B[b]F, B[k]F, HCB, PCB
5C1bvi Other waste incineration	NO, NA, NE
5C2 Open burning of waste	PCDD/F, B[a]P, B[b]F, B[k]F, PCB
5D Waste-water handling	NA, NE
5E Other waste	PM <sub>2.5</sub> , PM <sub>10</sub> , TSP, PCDD/F, B[a]P, B[b]F, B[k]F, I[123-cd]P, PCB

The Waste sector contains four key categories for five pollutants. Industrial waste incineration (5C1bi) is a key category for two pollutants: As and Cr, accounting for 66.3 per cent and 20.4 per cent of national total emissions, respectively. This category is the largest source of As in Ireland's 2023 inventory. The second key category in the Waste sector is Other waste (5E) for two pollutants: PCDD/F and PCBs. This category accounts for 53.5 per cent of national total PCB emissions and for 19.9 per cent of national total PCDD/F emissions in Ireland's 2023 inventory. Solid waste disposal on land (5A) is a key category for Mercury (Hg) accounting for 9.3 per cent of the national total in 2023. Cremation (5C1bv) is a key category for mercury (Hg) and accounts for 6.6 percent of the national total in 2023.

### 6.2 Biological treatment of waste - Solid waste disposal on land (NFR 5A)

#### 6.2.1 Main Pollutants

Landfill gas generated at solid waste disposal sites is a source of NMVOC emissions. In Ireland sector 5A has been responsible on average for 0.3 per cent of national total emissions across the time series 1990-2023, showing a decrease of 59.0 per cent from 0.83 kt in 1990 to 0.34 kt in 2023. Emission factors for NMVOC were sourced from the Inventory Guidebook (EMEP/EEA, 2019) and are listed in Table F1 of Annex F. The activity data used is the net fugitive methane emissions from SWDS (kt) as calculated in the national greenhouse gas inventory, which are converted to volume of landfill gas (Gm<sup>3</sup>) using standard temperature and pressure (STP) molar conversion factors.

Emission factors from the Inventory Guidebook (EMEP/EEA, 2019) are also applied to annual MSW data to estimate emissions of TSP, PM<sub>10</sub> and PM<sub>2.5</sub>. Resulting emission estimates are included in Table 6.2 below.

**Table 6.2 Emission Time Series for NMVOC TSP, PM<sub>10</sub> and PM<sub>2.5</sub> from Solid Waste Disposal on Land**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
NMVOC (kt)	0.83	1.01	0.80	0.64	0.19	0.20	0.30	0.41	0.45	0.45	0.43	0.40	0.38	0.36	0.36	0.34
TSP (t)	0.89	0.90	1.01	0.88	0.69	0.51	0.36	0.29	0.30	0.37	0.31	0.21	0.19	0.17	0.23	0.24
PM <sub>10</sub> (t)	0.42	0.43	0.48	0.42	0.33	0.24	0.17	0.13	0.14	0.18	0.15	0.10	0.09	0.08	0.11	0.11
PM <sub>2.5</sub> (t)	0.06	0.06	0.07	0.06	0.05	0.04	0.03	0.02	0.02	0.03	0.02	0.02	0.01	0.01	0.02	0.02

### 6.2.2 Heavy Metals

The relevant emissions in the Category 5A Solid Waste Disposal on Land sector in Ireland's air pollutant inventory include emissions of Hg from the disposal of batteries, electrical equipment, fluorescent lighting tubes, and measurement and control equipment in solid waste disposal sites (landfills). There are no direct estimates of the scales of disposal of these items at landfills in Ireland. However, Netcen/CTC (2006) provides a methodology to estimate emissions in Ireland using UK emission estimates, scaling by population for batteries and by household numbers for electrical equipment, fluorescent lighting and measurement and control equipment. Emission estimates for the above sources of Hg are presented in Table 6.3.

**Table 6.3 Emission Time Series for Mercury from Solid Waste Disposal on Land**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Hg (kg)	28.96	18.27	18.68	20.15	22.00	22.35	22.28	22.08	21.98	22.02	21.88	21.79	21.78	21.60	21.57	21.33

No data are available on emissions of POPs to air from municipal solid waste (MSW) disposal, although the emissions are likely to be negligible. Currently, there is no information in Ireland on the release of POPs from landfill gas and no emission factors are given in the Inventory Guidebook (EMEP/EEA, 2023).

## 6.3 Biological treatment of waste - Composting (NFR 5B1) and Anaerobic Digestion Biogas (NFR 5B2)

Composting of organic waste, such as food waste, garden and park waste has taken place in Ireland since 2001. It consists of organic waste collected at kerbside and brought to civic amenity/temporary collections sites, as well as organic material composted at households. Activity data is sourced from National Waste Database Reports published by the EPA on a regular basis. Composting is a source of emissions of NH<sub>3</sub> and CO and the Tier 2 emission factors in the Inventory Guidebook (EMEP/EEA, 2023) of 0.66 kg/Mg waste and 0.56 kg/Mg waste are used, respectively. Emission estimates and activity data are presented in Table 6.4.

**Table 6.4 Time Series of Activity Data and Emissions of NH<sub>3</sub> and CO from Composting**

Year	2001	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Waste composted (kt)	22	271	285	283	257	259	239	231	228	255	244	259	247	216	199	259
NH <sub>3</sub> (kt)	0.01	0.07	0.07	0.07	0.06	0.06	0.06	0.06	0.05	0.06	0.06	0.06	0.06	0.05	0.05	0.06
CO (kt)	0.01	0.15	0.16	0.16	0.14	0.15	0.13	0.13	0.13	0.14	0.14	0.15	0.14	0.12	0.11	0.15

Anaerobic digestion is a natural process by which biomass is broken down by naturally occurring microorganisms in the absence of oxygen. These microorganisms digest the biomass and release a methane-rich gas (biogas) that, if collected in a biogas plant, can be used to generate renewable heat and power. The remaining material (digestate) is rich in nutrients, so it can be used as a fertiliser.

There are small number of anaerobic digestion facilities in Ireland. The tonnage of material (on a dry matter basis) is obtained each year. Wastes imported and treated at facilities in Ireland are also included, and wastes generated in Ireland but treated in another country are excluded. Activity data has been obtained for 2010 to 2022 and 2023 data is estimated based on per capita rates in 2022 and population data for 2023. Emissions from 1990 to 2009 are reported as Not Occurring (NO). The Tier 1 emission factor from Table 3-1 Chapter 5B2 of the Inventory Guidebook (EMEP/EEA, 2023) is applied to estimate NH<sub>3</sub> emissions from this source.

## 6.4 Waste Incineration (NFR 5C)

Ireland carries out less waste incineration compared with some European countries and there is currently no incineration of MSW without energy recovery occurring in Ireland. The Waste Incineration (5C) category includes estimates of air pollutant emissions from the incineration of clinical waste (5C1biii), industrial waste (5C1bi), including hazardous waste (5C1bii), emissions from crematoria (5C1bv), and the open burning/combustion of waste materials such as farm plastics (5C2). There is incineration of municipal wastes in Ireland following the commissioning of one incinerator in 2011 and a further one in 2017. However, these are waste-to-energy facilities and as such emissions are reported under Public electricity and heat production (1A1a). Approximately 50 per cent of health-care waste was incinerated during the 1990s, with a total of 150 incinerators in operation. By 1999, only two of these remained in operation and both closed the following year. Most of the industrial installations that incinerate hazardous industrial wastes are in the pharmaceutical sector. The practice of cremation is also less common in Ireland than in other countries but has increased in recent years due to the decrease in available burial plots, particularly in larger cities and towns in Ireland.

All the above are sources of heavy metals, POPs and combustion pollutants (NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, CO, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC) in Ireland and are discussed in detail in the following sections.

### 6.4.1 Clinical Waste (5C1biii)

The incineration of Clinical Waste is no longer carried out in Ireland. The bulk of hazardous clinical waste in Ireland is now treated using non-incineration technologies (namely sterilisation and shredding), with the remaining waste disposed of through landfilling, exported for incineration or used as a fuel in cement kilns. In the early 1990s, most hospitals operated on-site incinerator units where hazardous clinical waste was incinerated. Several hospitals operated the practice of incinerating both hazardous and non-hazardous waste. Due to the implementation of stricter standards on incineration

and the requirement for facilities to be licensed by the EPA, all incinerators were closed by the mid-to late-1990s. Prior to the closure of these facilities, several applications were made to the EPA in respect of IPPC licences. National reports and Government records contain some information on the quantity of health-care waste incinerated during the period of operation of the incinerators. From these sources, it was determined that an estimated 4,000 t of health-care waste was incinerated per annum. This value was used across the time series for the period 1990–1997, after which negligible quantities of health-care waste were incinerated up until the closure of the two remaining incinerators in 2000.

Emission estimates were derived for heavy metals using the quantity of health-care waste determined to be incinerated and Inventory Guidebook (EMEP/EEA, 2023) emission factors for As, Cd, Cr, Cu, Pb, Hg and Ni, assuming controlled air flow with no abatement. The emission factor for Zn was sourced from the UK NAEI. Emission factors are provided in Table F.1 of Annex F. Emission estimates for heavy metals are presented in Table 6.5.

Emission factors for NO<sub>x</sub>, CO, NMVOC, SO<sub>x</sub>, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> were sourced from the Inventory guidebook (EMEP/EEA, 2023), using a Tier 1 approach, and are listed in Table F1 of Annex F. Resulting emission estimates 1990–1997 are included in Table 6.5.

**Table 6.5 Time Series of Emissions from the Incineration of Clinical Waste**

Year	1990	1991	1992	1993	1994	1995	1996	1997
NO <sub>x</sub> (kt)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO (t)	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76
NMVOC (t)	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80
SO <sub>x</sub> (t)	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16
TSP (kt)	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07
PM <sub>10</sub> (kt)	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
PM <sub>2.5</sub> (kt)	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
As (kg)	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40
Cd (t)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Cr (kg)	1.60	1.60	1.60	1.60	1.60	1.60	1.60	1.60
Cu (kg)	24.00	24.00	24.00	24.00	24.00	24.00	24.00	24.00
Pb (t)	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14
Hg (t)	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22
Ni (kg)	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20
Zn (t)	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07
PCDD/F (g-I-TEQ)	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49
PCBs (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HCB (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
B[a]p (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
B[b]F (kg)	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.04
B[k]F (kg)	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.04

Emissions of POPs from clinical wastes have been estimated using emission factors sourced from the UK NAEI. Dioxin and furan emission estimates are made utilising an emission factor of 372.1 µg I-TEQ/t health-care waste incinerated. This emission factor is used in the inventory for the period 1990–1997 until closure of all major plants. The PCB emission factor of 3.15 kg/Mt in 1990 reduces to 2.87 kg/Mt by 1995 and 2.36 kg/Mt by 1997, the last year of clinical waste incineration on the basis that environmental performance at the plants would have improved as in the UK. The emission factors for

HCB from health-care waste incinerators have been estimated by taking the UK NAEI factor of 0.5 kg/Mt for 2006 and estimating the historical emission factors for 1990 and 1995 in proportion to those for PCBs to take account of the improvements in environmental performance that would have been introduced at some incinerators. Emission factors for intervening years are interpolated. Emission factors for 2006 are also available from the UK NAEI for benzo[a]pyrene, benzo[b]fluoranthene and benzo[k]fluoranthene, but there are no data for indeno[1,2,3-cd]pyrene. The emission factors for 2006 have been used to estimate emission factors for 1990 and 1995, scaling back in proportion to the emission factors for PCBs like that undertaken for HCB emission factors. Emission factors are given in Table F.1 of Annex F and the estimates for POPs are presented in Table 6.5.

#### 6.4.2 Industrial Waste (5C1bi), Hazardous Waste (5C1bii) and Sewage Sludge (5C1biv)

The category Hazardous Waste Incineration (5C1bii) is reported in Industrial Waste Incineration (5C1bi) and reported as IE under the latter category. EU Directives on waste management have set the basis for strict regulatory control on the environmental performance of hazardous industrial waste incinerators. The incineration of Industrial Waste (5C1bi) (including hazardous waste) is now highly regulated in Ireland. There are currently only a small number of facilities based in the pharmaceutical and chemical sectors that operate incinerators for the treatment of hazardous waste.

**Table 6.6 Time Series of Emissions from the Incineration of Industrial (incl. Hazardous & Sludge) Waste**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
NO <sub>x</sub> (kt)	0.02	0.02	0.02	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO (t)	1.90	1.90	1.40	2.54	1.28	1.01	0.92	0.93	0.52	0.58	0.48	0.65	0.67	0.76	0.84	0.84
NMVOG (kt)	0.20	0.20	0.15	0.27	0.13	0.11	0.10	0.10	0.06	0.06	0.05	0.07	0.07	0.08	0.09	0.09
SO <sub>x</sub> (t)	1.27	1.27	0.94	1.70	0.86	0.68	0.62	0.62	0.35	0.39	0.32	0.44	0.45	0.51	0.56	0.56
TSP (t)	0.27	0.27	0.20	0.36	0.18	0.14	0.13	0.13	0.07	0.08	0.07	0.09	0.10	0.11	0.12	0.12
PM <sub>10</sub> (t)	0.19	0.19	0.14	0.25	0.13	0.10	0.09	0.09	0.05	0.06	0.05	0.07	0.07	0.08	0.08	0.08
PM <sub>2.5</sub> (t)	0.11	0.11	0.08	0.14	0.07	0.06	0.05	0.05	0.03	0.03	0.03	0.04	0.04	0.04	0.05	0.05
As (t)	0.55	0.56	0.58	0.62	0.65	0.65	0.65	0.65	0.65	0.66	0.66	0.67	0.67	0.68	0.69	0.70
Cd (kg)	2.71	2.71	2.00	3.62	1.82	1.44	1.31	1.33	0.75	0.82	0.68	0.93	0.96	1.08	1.19	1.19
Cr (t)	0.53	0.54	0.56	0.60	0.63	0.63	0.63	0.63	0.63	0.63	0.64	0.65	0.65	0.66	0.67	0.68
Cu (t)	0.29	0.30	0.31	0.33	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.36	0.36	0.36	0.37	0.38
Pb (t)	0.04	0.04	0.03	0.05	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02
Hg (kg)	1.52	1.52	1.12	2.03	1.02	0.81	0.73	0.74	0.42	0.46	0.38	0.52	0.53	0.60	0.67	0.67
Ni (kg)	3.79	3.79	2.80	5.07	2.55	2.02	1.84	1.86	1.05	1.15	0.96	1.30	1.34	1.51	1.67	1.67
Zn (t)	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
PCDD/F (g-I-TEQ)	0.03	0.03	0.08	0.02	0.01	0.01	0.00	0.01	0.01	0.61	1.31	0.35	0.01	0.00	0.00	0.00
PCBs (kg)	0.09	0.08	0.03	0.03	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HCB (kg)	0.05	0.04	0.02	0.02	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01
B[a]p (kg)	0.02	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B[b]F (kg)	0.07	0.05	0.02	0.03	0.01	0.01	0.01	0.01	0.00	0.01	0.00	0.01	0.01	0.01	0.01	0.01
B[k]F (kg)	0.07	0.05	0.02	0.03	0.01	0.01	0.01	0.01	0.00	0.01	0.00	0.01	0.01	0.01	0.01	0.01

The facilities that operate these units report emissions to the atmosphere to the EPA as part of IPPC licensing requirements. The disposal of CCA treated wood (CCA being a preservative containing copper, chromium and arsenic) by burning is also included as a source in this sector.

Estimates of the quantity of hazardous waste incinerated at the relevant facilities, determined from returns to the National Waste Database (Carey et al, 1996; Crowe et al, 2000; Meaney et al, 2003; Collins et al, 2004a; Collins et al, 2004b; Collins et al, 2005; Le Bolloch et al, 2006; Le Bolloch et al, 2007; Le Bolloch et al, 2009; McCoole et al, 2009; McCoole et al, 2011; McCoole et al, 2012; McCoole et al, 2013, [Circular Economy and Waste Statistics | Environmental Protection Agency](#)), and information supplied by the facilities involved allows for the calculation of heavy metal emission estimates. Emission factors sourced from the Inventory Guidebook (EMEP/EEA 2023) for As, Cd, Pb, Hg, Ni and the UK NAEI for Cr and Cu are used to estimate emissions. Emission estimates are presented in Table 6.6, while the emission factors used are presented in Table F.1 of Annex F. The hazardous waste incinerators currently in use in Ireland are relatively modern units designed to optimise the burning process, with wet scrubber abatement systems in place to reduce the emissions of POPs to air. Further to the use of incinerators, there are also several facilities that use thermal oxidisers, which are subject to emission limit values. Annual Environmental Reports and IPPC Licence Applications provide adequate information in relation to the monitoring of PCDD/F emissions to air with limited information on the other relevant POPs. With respect to emissions of PAHs, emission factors sourced from the UK NAEI were applied to the tonnage of waste incinerated for each year. Pollutant-specific emission factors are presented in Table F.1 of Annex F. Emission estimates for POPs are presented in Table 6.6.

Emission factors for: NO<sub>x</sub>, CO, NMVOC, SO<sub>x</sub>, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> were sourced from the Inventory Guidebook (EMEP/EEA, 2023), using a Tier 1 approach, and are listed in Table F.1 of Annex F. Resulting emission estimates are included in Table 6.6.

### 6.4.3 Crematoria (5C1bv)

The practice of cremation is less popular in Ireland than in other countries. However, due to the decrease in the number of burial plots available, particularly in larger cities and towns, the number of cremations in Ireland has been steadily increasing. There are currently eight crematoria operating in Ireland. Cremation has been in operation in Ireland for a significant period, with one of the crematoria open since the early 1980s. A pet crematorium is also currently operating in Ireland; however, emissions from this source are regarded as negligible. Data on the number of cremations in Ireland have been obtained via correspondence with crematoria operators.

Heavy metals emissions are estimated using Inventory Guidebook (EMEP/EEA, 2023) emission factors for As, Cd, Pb, Cr, Hg, Ni, Cu Se and Zn and are presented in Table F.1 of Annex F. Emissions of POPs from crematoria include PCDD/F, HCB and benzo[a]pyrene. Inventory Guidebook (EMEP/EEA, 2019) emission factors are used to derive emission estimates for the years 1990–2023 (Table F.1 of Annex F). Emission factors for: NO<sub>x</sub>, CO, NMVOC, SO<sub>x</sub>, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> were sourced from the Inventory guidebook (EMEP/EEA, 2019), using a Tier 1 approach, and are listed in Table F.1 of Annex F. Resulting emission estimates are included in Table 6.7.

**Table 6.7 Time Series of Emissions from Crematoria**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
NO <sub>x</sub> (t)	1.24	1.24	1.82	2.03	2.54	3.37	3.73	4.11	4.52	4.92	5.34	5.81	6.81	7.51	7.90	8.31
CO (t)	0.21	0.21	0.31	0.34	0.43	0.57	0.63	0.70	0.77	0.84	0.91	0.99	1.16	1.27	1.34	1.41
NM VOC (t)	0.02	0.02	0.03	0.03	0.04	0.05	0.06	0.06	0.07	0.08	0.08	0.09	0.11	0.12	0.12	0.13
SO <sub>x</sub> (t)	0.17	0.17	0.25	0.28	0.35	0.46	0.51	0.56	0.62	0.67	0.73	0.80	0.93	1.03	1.08	1.14
TSP (t)	0.06	0.06	0.08	0.09	0.12	0.16	0.17	0.19	0.21	0.23	0.25	0.27	0.32	0.35	0.37	0.39
PM <sub>10</sub> (t)	0.05	0.05	0.08	0.09	0.11	0.14	0.16	0.17	0.19	0.21	0.22	0.24	0.29	0.32	0.33	0.35
PM <sub>2.5</sub> (t)	0.05	0.05	0.08	0.09	0.11	0.14	0.16	0.17	0.19	0.21	0.22	0.24	0.29	0.32	0.33	0.35
As (g)	20.4	20.4	29.9	33.4	42.0	55.5	61.5	67.8	74.5	81.2	88.2	95.9	112.3	123.9	130.2	137.1
Cd (g)	7.5	7.5	11.1	12.4	15.5	20.5	22.7	25.1	27.5	30.0	32.6	35.4	41.5	45.8	48.1	50.7
Cr (g)	20.3	20.3	29.8	33.3	41.8	55.3	61.3	67.6	74.2	80.9	87.8	95.5	111.9	123.4	129.8	136.6
Cu (g)	18.6	18.6	27.3	30.5	38.3	50.7	56.1	62.0	68.0	74.1	80.5	87.6	102.6	113.2	119.0	125.2
Pb (g)	45.0	45.0	66.1	73.8	92.6	122.6	135.6	149.7	164.4	179.1	194.5	211.6	247.8	273.4	287.4	302.4
Hg (kg)	2.2	2.2	3.3	3.7	4.6	6.1	6.7	7.4	8.2	8.9	9.7	10.5	12.3	13.6	14.3	15.0
Ni (kg)	0.03	0.03	0.04	0.04	0.05	0.07	0.08	0.09	0.09	0.10	0.11	0.12	0.14	0.16	0.17	0.17
PCDD/F (g-I-TEQ)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
HCBs (g)	0.23	0.23	0.33	0.37	0.46	0.61	0.68	0.75	0.82	0.89	0.97	1.06	1.24	1.37	1.44	1.51
B[a]p (g)	0.02	0.03	0.03	0.03	0.04	0.05	0.06	0.07	0.07	0.08	0.09	0.09	0.11	0.12	0.13	0.13

#### 6.4.4 Open Burning of Agricultural Wastes (5C2)

Open Burning of Agricultural Wastes, which includes the burning of crop residues, animal carcasses and poultry litter is a practice generally not undertaken in Ireland. This is because of requirements imposed on farmers who are in receipt of payments under the Common Agricultural Policy and national agri-environmental schemes<sup>4</sup>. All biomass burning on agricultural land is classified as wildfire and are thus non-anthropogenic in origin. Specific pieces of legislation which refer to burning of agricultural waste are contained in the following legislation: Wildlife Act, 1976; Wildlife (Amendment) Act, 2000; Birds and Habitats Directive; Air Pollution Act, 1987; The Safety, Health and Welfare at Work Act, 2005 and Safety, Health and Welfare at Work (General Application) Regulations, 2007; Forestry Act, 1946; Waste Management Act, 1996; Waste Management Act, 2008; SI No. 286 of 2009 – Waste Management Regulations; Fire Services Act, 1981; Fire Services Act, 2003; Criminal Damages Act, 1991. The various pieces of legislation do not refer to the three types of burning identified in the EMEP guidebook 2023 definitions. In the various pieces of legislation, reference is made to "a wood" or "forestry plantation" or "vegetation" or "straw" or "substance" or "waste". Therefore, emission estimates from these sources are reported as "NO" (not occurring).

Emissions from the open burning of farm plastics are the only source of emissions from agricultural wastes for which estimates are made. Information on the quantity of waste farm plastics that are burned in open fires is difficult to obtain. One of the largest sources of waste farm plastic is waste plastic silage wrap and to a lesser extent synthetic fertiliser bags. The increased replacement of conventional silage with plastic wrapped silage bales, which use substantially more plastic, has seen an increase in the quantity of this waste stream. Several different sources of information were utilised in the derivation of emission estimates. Information on the quantities of silage plastic on the market was obtained from the Irish Farm Film Producers Group (IFFPG) and national agricultural statistics

<sup>4</sup> <http://www.agriculture.gov.ie/farmerschemespayments/crosscompliance/>



were provided by the CSO and the Teagasc National Farm Survey. Using the area of land utilised for silage for each year of the time series, an estimate of the plastic used for conventional (pit) silage and baled silage is made. Account is taken of plastic recovery under the silage plastics collection service operated by both the IFFPG and the Farm Relief Services. The plastic collected is recycled and used to make products such as park benches, plastic bags, garden furniture and plastic piping. The remaining plastic is assumed to be burned. In the period 2006 to 2007, there were separate silage plastic collections organised by the then Department of Environment. These were subsequently discontinued. As a result, the IFFPG increased the quantity collected significantly which is the reason emissions have decreased significantly from 2011.

Dioxin and furan emissions from the open burning of farm plastics are determined using estimates of the quantities of material burned and the UNEP Toolkit (2013) emission factor of 300 µg I-TEQ/t burned for the open burning of municipal wastes. The UK NAEI provides an emission factor of 510 kg/Mt burned for the estimation of PCB emissions. There is minimal data available on emission factors for PAHs; however, the emission factors from the UK NAEI for small-scale waste burning are used as a best estimate. Emission factors of 89.5 kg/Mt for benzo[a]pyrene, 405 kg/Mt for benzo[b]fluoranthene and 405 kg/Mt for benzo[k]fluoranthene are applied. No data is available for indeno[1,2,3-cd] pyrene. Emission factors are compiled in Table F.2 of Annex F and the emission estimates are presented in Table 6.8.

Emissions of NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, CO and particles from this small source are not estimated and are therefore reported as “NE”.

**Table 6.8 Time Series of Emissions from the Open Burning of Farm Plastics**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
PCDD/F (g-I-TEQ)	0.94	1.10	0.65	0.23	1.15	0.28	0.45	0.49	0.46	0.50	0.59	0.83	0.48	0.45	0.14	0.14
PCBs (kg)	1.59	1.86	1.11	0.39	1.96	0.48	0.77	0.84	0.78	0.84	1.01	1.41	0.82	0.76	0.24	0.24
B[a]p (kg)	0.28	0.33	0.20	0.07	0.34	0.08	0.13	0.15	0.14	0.15	0.18	0.25	0.14	0.13	0.04	0.04
B[b]F (kg)	1.26	1.48	0.88	0.31	1.55	0.38	0.61	0.67	0.62	0.67	0.80	1.12	0.65	0.60	0.19	0.19
B[k]F (kg)	1.26	1.48	0.88	0.31	1.55	0.38	0.61	0.67	0.62	0.67	0.80	1.12	0.65	0.60	0.19	0.19

## 6.5 Wastewater handling (NFR 5D)

Emissions of NMVOC from wastewater treatment plants may in some cases be significant in urban areas. In Ireland’s case, based on the use of the emission factor of 15mg/m<sup>3</sup> of wastewater handled from EMEP/EEA 2023 guidebook and total wastewater quantities handled in Ireland, the emissions were estimated and reported under 5D1. These emissions include industrial wastewater handling and the notation key IE included elsewhere is used for category 5D2 Industrial wastewater handling. Ireland does not have dry toilets, including latrines, as a wastewater treatment system over the complete time series, hence there are no NH<sub>3</sub> emissions from this source. Domestic wastewater is defined as wastewater from household water use. Domestic wastewater is either treated in centralized treatment plants or in septic tanks. Centralised wastewater treatment plants also treat commercial and industrial wastewater and for that reason emissions from *Industrial Wastewater*



(5.D.2) are included in *Domestic Wastewater* (5.D.1). A description of wastewater flows can be found in Annex 3.5.G of Ireland's National Inventory Document.

## 6.6 Other Waste (NFR 5E)

This NFR category includes emissions from accidental vehicle and building fires and other burning, which constitutes bonfires, domestic burning of MSW and burning of construction wastes. These are all sources of POPs, TSP, PM<sub>10</sub> and PM<sub>2.5</sub>. Each of these combustion sources is described in the following sections. The activity data associated with each type of fire is shown in Table 6.9.

### 6.6.1 Accidental Fires

Accidental fires are poorly controlled combustion events that can release large quantities of POPs and particulates into the environment. These include accidental fires of houses, other buildings and cars. A variety of materials can be burned in accidental fires, which can lead to some difficulty in obtaining detailed activity data and applying emission factors correctly. However, there are some data available in Ireland in relation to accidental building and vehicle fires from the Fire Services Department (Table 6.9).

**Table 6.9 Time Series of activity data from Accidental Fires from Vehicles and Buildings**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Vehicle fires	4130	4130	7749	4907	4534	1609	1637	2681	1607	2137	2667	2316	1885	1827	1812	1987
Building fires-detached	1344	1344	1344	1263	1671	800	770	1392	621	860	1099	946	989	878	934	909
Building fires-undetached	2015	2015	2015	1894	2506	1200	1155	2088	931	1290	1649	1420	1484	1317	1400	1364
Apartment fires	426	426	426	470	245	242	214	240	192	211	229	209	217	204	191	194
Industrial fires	508	508	508	433	453	270	288	458	234	317	400	386	284	318	321	297

Vehicle fire statistics are only available since 2000, with the number of fires ranging from 1,600 to 7,700 per annum. With respect to earlier years, the URS Dames & Moore (2000) report suggests that the number of vehicle fires in 1998 was 4,130. It is assumed that, in the absence of any other information, the number of vehicle fires per year in the period 1990–1997 is equal to that in 1998. Dioxin and furan emissions from vehicle fires are estimated using the UNEP Toolkit 2013 emission factor of 100 µg I-TEQ/vehicle fire. An emission factor of 25.5 mg/vehicle fire based on the NAEI emission factor of 510 kg/Mt burned for small scale domestic waste burning (UK NAEI) for PCB emissions from the open burning of MSW is used for accidental vehicle fires, assuming that on average 50 kg of material are burnt per fire (Dyke, 1997), while those in relation to PAHs, also based on the mass of material burnt, sourced from the UK NAEI, suggest values for benzo[a]pyrene of 0.06 mg/vehicle fire, for benzo[b]fluoranthene of 0.095 mg/vehicle fire, for benzo[k]fluoranthene of 0.03 mg/vehicle fire, and for indeno[1,2,3-cd]pyrene of 0.065 mg/vehicle fire. With respect to TSP, PM<sub>10</sub> and PM<sub>2.5</sub>, the Tier 2 emission factors presented in the EMEP/EEA 2023 Guidebook are adopted.

The Fire Services Department also provides information in relation to building fires, which is disaggregated into the type of building and the number of fires that are chimney fires. Information is only available for the years 2000–2023 at this level of disaggregation. For data prior to 2000, no differentiation was made between chimney fires and other types of building fires. The proportion of chimney fires to the total number of building fires post-2000 is therefore used to estimate the number

of chimney fires annually prior to 2000. Limited information is available on the quantity of material burnt in accidental fires both in Ireland and internationally. The assumed quantity of material burnt in each building fire is 2.28 t per fire (Lorenz et al., 1996) and approximately 10 kg in each chimney fire. Dioxin and furan emissions are estimated using an emission factor of 400 µg I-TEQ/t of material burned (UNEP Toolkit, 2013). For PCB emissions, the emission factor of 510 kg/Mt burned (UK NAEI) for the open burning of MSW is applied, while, for PAH emissions (UK NAEI), the emission factors equate to 1.2 kg/Mt for benzo[a]pyrene, 1.9 kg/Mt for benzo[b]fluoranthene, 0.67 kg/Mt for benzo[k]fluoranthene and 1.3 kg/Mt for indeno[1,2,3-cd]pyrene. Accidental vehicle fires and building fires emission estimates are summed to provide an estimate of the total emissions from accidental fires. Emission estimates for the 1990–2023 time series are presented in Table 6.10. Emission factors are compiled in Table F.2 of Annex F. Ireland has included estimates of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions in this using the emission factors from the Inventory Guidebook (EMEP/EEA, 2023).

**Table 6.10 Time Series of Emissions from Accidental Fires from Vehicles and Buildings**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
PCDD/F (g-I-TEQ)	5.40	5.40	3.87	3.39	4.28	2.00	1.93	3.46	1.59	2.19	2.78	2.40	2.45	2.08	2.32	2.28
PCBs (kg)	6.47	6.47	4.14	3.82	5.00	2.39	2.30	4.13	1.86	2.57	3.28	2.82	2.94	2.47	2.77	2.70
B[a]p (kg)	1.5E-05	1.5E-05	9.7E-06	9.0E-06	1.2E-05	5.6E-06	5.4E-06	9.7E-06	4.4E-06	6.0E-06	7.7E-06	6.6E-06	6.9E-06	5.8E-06	6.5E-06	6.4E-06
B[b]F (kg)	2.4E-05	2.4E-05	1.5E-05	1.4E-05	1.9E-05	8.9E-06	8.6E-06	1.5E-05	6.9E-06	9.6E-06	1.2E-05	1.1E-05	1.1E-05	9.2E-06	1.0E-05	1.0E-05
B[k]F (kg)	8.5E-06	8.5E-06	5.4E-06	5.0E-06	6.6E-06	3.1E-06	3.0E-06	5.4E-06	2.4E-06	3.4E-06	4.3E-06	3.7E-06	3.9E-06	3.2E-06	3.6E-06	3.6E-06
I(123-cd)P (kg)	1.6E-05	1.6E-05	1.1E-05	9.7E-06	1.3E-05	6.1E-06	5.9E-06	1.1E-05	4.7E-06	6.6E-06	8.4E-06	7.2E-06	7.5E-06	6.3E-06	7.1E-06	6.9E-06
TSP (kt)	0.36	0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.23	0.24	0.24
PM10 (kt)	0.36	0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.23	0.24	0.24
PM2.5 (kt)	0.36	0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.23	0.24	0.24

## 6.6.2 Other Burning

This section includes the emission of POPs from domestic bonfires, the burning of domestic waste both indoors and outdoors and the open burning of construction waste. Domestic bonfires normally include a variety of garden wastes (e.g. wood, leaves, etc.), and their importance with respect to POP emissions is greatly increased in cases where other wastes are added to the bonfires (e.g. plastics). Some households are not covered, or opt not to be covered, by waste collection systems and may burn household waste. Combustion of treated wood that has been used for construction, fencing and furniture can be a particularly significant source of POP emissions. For example, where wood is pre-treated with chlorinated fungicides, such as lindane or pentachlorophenol, its combustion can be a potentially significant source of PCDD/F emissions to air, whilst wood pre-treatment with creosote is a potential source for PAH emissions. However, the use of these chemicals has been significantly reduced in Ireland since the early 1990s.

For domestic bonfires, activity data are determined on a per-capita basis using the UK inventory as the reference, as no information is available in Ireland. For the burning of household waste, estimates for uncollected household waste were obtained for each of the years 2001 through to 2013, as well as for 1998 and 1995 from National Waste Reports (Carey et al., 1996; Crowe et al., 2000; Collins et al., 2004a, 2004b; Le Bolloch et al., 2006, 2007, 2009, McCoolle et al, 2009; McCoolle et al, 2011; McCoolle et al, 2012; McCoolle et al, 2013), with data calculated for other years using waste statistical

data ([Circular Economy and Waste Statistics | Environmental Protection Agency](#)). “Uncollected waste” refers to the waste produced by the portion of the population not provided with, or not availing of, a waste collection service, corrected to take account of local conditions. This is calculated according to a standard methodology at the local authority level, based on total numbers of households, numbers of households served with waste collection, and quantities of waste collected per household in each local authority area. In addition, a proportion of households share waste collections services. Only the fraction of household waste that is combustible is burned. Compositional statistics at a national level are applied to estimate the quantities of combustible materials burnt.

Information on construction and demolition waste is available from National Waste Reports. The proportion of wood within this waste stream is estimated using data collected but not published in the National Waste Reports for the years 2004 and 2006, based on estimates of both authorised and unauthorised construction and demolition waste disposal. These values have been used for all other years in the absence of any other information. The URS Dames & Moore study (2000) suggests that 5 per cent of construction and demolition waste wood arising is burned on construction sites, whereas the UK NAEI suggests a value of 0.1 per cent. The value of 5 per cent is applied for the years 1990–1998, linearly decreasing for the years 1999–2003, with the value of 0.1 per cent applied for the period 2004–2021, based on correspondence with representatives from the National Construction and Demolition Waste Council who indicate that they would expect virtually no uncontrolled burning in urban areas. The UNEP toolkit for open burning of construction and demolition waste wood (60 µg-I-TEQ/t burned) is applied to estimate PCDD/F emissions from bonfires on the basis that bonfires contain mainly wood and garden waste. Domestic burning of MSW contains material that varies and that often includes plastics and sometimes specific chemicals that potentially affect PCDD/F emissions. The UNEP 2013 toolkit emission factor of 40 µg-I-TEQ/t burned is used for burning of household waste. This emission factor not only considers the wide range of materials in household waste but also other materials such as treated and untreated wood. In relation to PCDD/F emissions from wood burning, an emission factor of 60 µg I-TEQ/t is applied.

The estimated emission factor of 1.14 kg/Mt burned for PCBs from bonfires has been taken as the average of the UK NAEI emission factors for domestic wood combustion (e.g. fireplaces, 1.99 kg/Mt burned) and open burning of crop residues (0.29 kg/Mt). For the open burning of domestic wastes, the UK NAEI emission factor of 510 kg/Mt burned has been adopted for PCBs. There are no specific data on PCB emissions from the open burning of construction wood, but emission factors from the NAEI for industrial combustion of wood indicate no difference for treated and untreated wood. Emission factors for domestic wood combustion from the NAEI and the Inventory Guidebook range from 1.99 to 6 kg/Mt burned. The emission factor of 1.99 kg/Mt has been adopted for open burning of construction waste wood in Ireland.

Emission factors with respect to PAH emission estimates are also sourced from the UK NAEI. For bonfires, the emission factors are 1,300 kg/Mt for benzo[a]pyrene, 1,500 kg/Mt for benzo[b]fluoranthene, 500 kg/Mt for benzo[k]fluoranthene and 90 kg/Mt for indeno[1,2,3-cd] pyrene. For the open burning of domestic wastes, emission factors for small-scale waste burning are applied as follows: 89.5 kg/Mt for benzo[a]pyrene, 405 kg/Mt for both benzo[b]fluoranthene and benzo[k]fluoranthene. No data are available for indeno[1,2,3-cd] pyrene. These emission factors are also used to estimate emissions from the open burning of wood at construction sites. The three

sources of emissions described are summed to provide total emission estimates for Category 5.E Other Waste. Emission factors are compiled in Table F.2 of Annex F. Emission estimates for the 1990–2023 time series are presented in Table 6.11.

**Table 6.11 Time Series of Emissions from Other Waste Burning**

Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
PCDD/F (g-I-TEQ)	1.15	1.29	1.72	2.24	0.72	0.66	0.63	0.63	0.63	0.63	0.64	0.65	0.65	0.65	0.67	0.67
PCBs (kg)	7.73	9.46	14.75	21.21	1.43	0.70	0.24	0.24	0.25	0.25	0.26	0.27	0.18	0.15	0.22	0.22
B[a]p (t)	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
B[b]F (t)	0.02	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
B[k]F (t)	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
l(123-cd)P (kg)	0.82	0.83	0.85	0.87	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

## 6.7 Recalculations in the Waste Sector

In category Solid Waste Disposal (5A), there were minor recalculations in Hg for the years 2017-2022 due to a revision of activity data resulting in average increase in emissions of 0.7 per cent with a range of 0.1 per cent to 1.3 per cent over this period.

In the Biological Treatment of Solid Waste (5B) category, there was a recalculation for the year 2022 with new activity data provided by the EPA's Waste Statistics team. Emissions of NH<sub>3</sub> and CO decreased by 23.5 per cent.

In the Other waste (5E) category, there minor recalculations in PAHs due to revised activity data for the years 2017-2022 resulting in an average increase of 0.7 per cent with a range of 0.1 per cent to 1.6 per cent over this period.

Recalculations in the waste sector are provided in Table 6.12.

## 6.8 Quality Assurance/Quality Control

As identified in section 6.7 the implementation of QA/QC activities ensures that the estimates in the Waste sector are now fully consistent with estimates of greenhouse gas emissions from the sector.

## 6.9 Planned Improvements

The inventory team will continue to review emission estimates for this sector considering any new information that may become available for future submissions.

National inventory experts are currently investigating the relevant sources of information with respect to 5B2 Anaerobic digestion which is very much in its infancy in Ireland. This will include estimates of emissions for the relevant pollutants as and when sufficient activity data and knowledge of feedstocks is gathered. It is envisaged that the likely level of emissions from this category will be of the order of tonnes of NH<sub>3</sub> and thus negligible. There is no further update on this category in this submission.

In practice there is a small number of test/pilot digesters in Ireland. However, there is no regulation regarding the collection of data in respect of the feedstocks of these digesters. The inventory agency is working with the Department of Agriculture, Food and Marine to get this information. The activity data issue relates primarily to feedstocks that are not classified as wastes and do not have waste codes, mainly animal manures/slurries and crop residues or grass inputs. We have managed to collect the food wastes input data, but to estimate emissions from this category requires the animal manure amounts, otherwise a double count of emissions would occur in agriculture. It seems that the manure/slurry and cellulose material input is much greater than the food waste.

Review recommendations and responses are included in this report as Annex H.

**Table 6.12 Recalculations for Waste 1990–2022**

NFR Category	Pollutant	Unit	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																
5A Solid waste disposal on land	NMVOC	kt	1.01	0.80	0.64	0.19	0.30	0.41	0.45	0.45	0.43	0.40	0.38	0.36	0.33	0.36
5A Solid waste disposal on land	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	Hg	t	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5A Solid waste disposal on land	PCDD/F	g-I-TEQ	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
5A Solid waste disposal on land	PCBs	kg	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
<b>Submission 2025</b>																
5A Solid waste disposal on land	NMVOC	kt	1.01	0.80	0.64	0.19	0.30	0.41	0.45	0.45	0.43	0.40	0.38	0.36	0.33	0.36
5A Solid waste disposal on land	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM <sub>10</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM <sub>2.5</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	Hg	t	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5A Solid waste disposal on land	PCDD/F	g-I-TEQ	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
5A Solid waste disposal on land	PCBs	kg	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
<b>% Changes in emissions</b>																
5A Solid waste disposal on land	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.8%	1.1%	1.3%	0.1%
5A Solid waste disposal on land	PCDD/F	%	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
5A Solid waste disposal on land	PCBs	%	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

NFR Category	Pollutant	Unit	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																
5B Biological treatment of waste - Composting	NH3	kt	NO	NO	0.07	0.07	0.06	0.06	0.06	0.05	0.06	0.06	0.06	0.06	0.05	0.06
5B Biological treatment of waste - Composting	CO	kt	NO	NO	0.15	0.16	0.15	0.13	0.13	0.13	0.14	0.14	0.15	0.14	0.12	0.15
<b>Submission 2025</b>																
5B Biological treatment of waste - Composting	NH3	kt	NO	NO	0.07	0.07	0.06	0.06	0.06	0.05	0.06	0.06	0.06	0.06	0.05	0.05
5B Biological treatment of waste - Composting	CO	kt	NO	NO	0.15	0.16	0.15	0.13	0.13	0.13	0.14	0.14	0.15	0.14	0.12	0.11
<b>% Change in Emissions</b>																
5B Biological treatment of waste - Composting	NH3	%	NO	NO	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-23.5%
5B Biological treatment of waste - Composting	CO	%	NO	NO	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-23.5%

NFR Category	Pollutant	Unit	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																
5C Waste Incineration	NO <sub>x</sub>	kt	0.03	0.02	0.03	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.02	0.02	0.02
5C Waste Incineration	NMVOC	kt	0.20	0.15	0.27	0.13	0.11	0.10	0.10	0.06	0.06	0.05	0.07	0.07	0.08	0.09
5C Waste Incineration	SO <sub>2</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	TSP	kt	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM <sub>10</sub>	kt	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM <sub>2.5</sub>	kt	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Cd	t	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Pb	t	0.18	0.03	0.05	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02
5C Waste Incineration	Hg	t	0.22	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste Incineration	PCDD/F	g-I-TEQ	2.62	0.73	0.25	1.16	0.29	0.46	0.51	0.48	1.10	1.90	1.18	0.50	0.45	0.15
5C Waste Incineration	PCBs	kg	1.95	1.15	0.42	1.97	0.49	0.78	0.85	0.79	0.85	1.02	1.42	0.83	0.77	0.25
5C Waste Incineration	HCB	t	0.05	0.02	0.02	0.01	0.01	0.01	0.01	0.00	0.01	0.00	0.01	0.01	0.01	0.01
5C Waste Incineration	B[a]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[b]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[k]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Submission 2025</b>																
5C Waste Incineration	NO <sub>x</sub>	kt	0.03	0.02	0.03	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.02	0.02	0.02
5C Waste Incineration	NMVOC	kt	0.20	0.15	0.27	0.13	0.11	0.10	0.10	0.06	0.06	0.05	0.07	0.07	0.08	0.09

NFR Category	Pollutant	Unit	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
5C Waste Incineration	SO <sub>2</sub>	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	TSP	kt	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM <sub>10</sub>	kt	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM <sub>2.5</sub>	kt	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Cd	t	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Pb	t	0.18	0.03	0.05	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02
5C Waste Incineration	Hg	t	0.22	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste Incineration	PCDD/F	g-I-TEQ	2.62	0.73	0.25	1.16	0.29	0.46	0.51	0.48	1.10	1.90	1.18	0.50	0.45	0.15
5C Waste Incineration	PCBs	kg	1.95	1.15	0.42	1.97	0.49	0.78	0.85	0.79	0.85	1.02	1.42	0.83	0.77	0.25
5C Waste Incineration	HCB	t	0.05	0.02	0.02	0.01	0.01	0.01	0.01	0.00	0.01	0.00	0.01	0.01	0.01	0.01
5C Waste Incineration	B[a]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[b]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[k]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>% Change in Emissions</b>																
5C Waste Incineration	NO <sub>x</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	SO <sub>2</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	PCDD/F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%
5C Waste Incineration	PCBs	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	HCB	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	B[a]P	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	B[b]F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	B[k]F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	I[123-cd]P	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%



NFR Category	Pollutant	Unit	1995	2000	2005	2010	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>Submission 2024</b>																
5E Other Waste	TSP		0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.22	0.24
5E Other Waste	PM <sub>10</sub>		0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.22	0.24
5E Other Waste	PM <sub>2.5</sub>		0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.22	0.24
5E Other Waste	PCDD/F	g-I-TEQ	6.69	5.59	5.63	5.01	2.66	2.56	4.08	2.22	2.82	3.42	3.04	3.10	2.72	2.97
5E Other Waste	PCBs	kg	15.93	18.89	25.04	6.43	3.09	2.54	4.38	2.11	2.82	3.54	3.09	3.12	2.62	2.99
5E Other Waste	B[a]P	t	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	B[b]F	t	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5E Other Waste	B[k]F	t	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Submission 2025</b>																
5E Other Waste	TSP		0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.23	0.24
5E Other Waste	PM <sub>10</sub>		0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.23	0.24
5E Other Waste	PM <sub>2.5</sub>		0.36	0.37	0.34	0.43	0.21	0.20	0.36	0.17	0.23	0.29	0.25	0.26	0.23	0.24
5E Other Waste	PCDD/F	g-I-TEQ	6.69	5.59	5.63	5.01	2.66	2.56	4.08	2.22	2.82	3.43	3.05	3.10	2.73	2.98
5E Other Waste	PCBs	kg	15.93	18.89	25.04	6.43	3.09	2.54	4.38	2.11	2.82	3.54	3.09	3.12	2.62	2.99
5E Other Waste	B[a]P	t	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	B[b]F	t	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5E Other Waste	B[k]F	t	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>% Change in Emission</b>																
5E Other Waste	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	5.4%	0.0%
5E Other Waste	PM <sub>10</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	5.4%	0.0%
5E Other Waste	PM <sub>2.5</sub>	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	5.4%	0.0%
5E Other Waste	PCDD/F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.2%	0.2%	0.3%	0.4%
5E Other Waste	PCBs	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5E Other Waste	B[a]P	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.7%	1.0%	1.3%	1.6%
5E Other Waste	B[b]F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.7%	1.0%	1.2%	1.6%
5E Other Waste	B[k]F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.7%	1.0%	1.2%	1.6%
5E Other Waste	I[123-cd]P	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.7%	1.0%	1.3%	1.6%

## Chapter Seven - Gridded and LPS data

### 7.1 Overview of Gridded and LPS data reporting

It is mandatory to report gridded emissions and emissions from large point sources every four years both under the Convention on Long-Range Transboundary Air Pollution and under Directive (EU) 2016/2284 on the reduction of national emissions of certain atmospheric pollutants (the revised National Emission Ceilings Directive).

The Guidelines for Reporting Emissions and Projections Data under the Convention on Long-Range Transboundary Air Pollution (ECE/EB.AIR/125), adopted in 2014, specify the scope, methodologies, formats and deadlines for annual inventory submissions by Parties to the Convention. The guidelines specify that at four yearly intervals, starting in 2017, Parties shall report updated aggregated sectoral (GNFR) gridded emissions and LPS emissions. The EMEP grid is defined in paragraph 14 of the reporting guidelines and refers to a 0.1°x0.1° latitude-longitude projection in the geographic coordinate World Geodetic System (WGS) latest revision, WGS 84. The EMEP domain covers the geographic domain between 30°N-82°N latitude and 30°W-90°E longitude.

The reporting guidelines under the Convention refers to the EMEP/EEA Guidebook for technical guidance on the spatial distribution of emissions. Directive (EU) 2016/2284 refers to the reporting guidelines under the Convention and as such the requirements under the two reporting requirements are identical.

The development of a high-resolution model for distribution of emissions was part of a research project funded by the EPA ("National mapping of GHG and non-GHG emissions sources". Ref: 2015-CCRP-MS.26). The project has developed a model for distributing emissions at a resolution of 1 km x 1 km covering all sectors and pollutants included in the official Irish emission inventory. The generated spatial emissions data (GNFR) is fully consistent with the reported emission inventories (NFR) under the LRTAP Convention. A list of GNFR categories is presented in Table 7.1 and the relationship between NFR and GNFR is presented in Table 7.2.

**Table 7.1. List of GNFR categories**

<b>GNFR</b>
A_PublicPower
B_Industry
C_OtherStationaryComb
D_Fugitive
E_Solvents
F_RoadTransport
G_Shipping
H_Aviation
I_Offroad
J_Waste
K_AgriLivestock
L_AgriOther
M_Other
<b>'MEMO' ITEMS - NOT TO BE INCLUDED IN NATIONAL TOTALS</b>
O_AviCruise
P_IntShipping
z_Memo
N_Natural

**Table 7.2. Correspondence list for GNFR and NFR categories**

<b>NFR</b>	<b>NFR name</b>	<b>GNFR</b>
1A1a	Public electricity and heat production	A_PublicPower
1A1b	Petroleum refining	B_Industry
1A1c	Manufacture of solid fuels and other energy industries	B_Industry
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	B_Industry
1A2b	Stationary combustion in manufacturing industries and construction: Non-ferrous metals	B_Industry
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	B_Industry
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	B_Industry
1A2e	Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	B_Industry
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	B_Industry
1A2gvii	Mobile Combustion in manufacturing industries and construction	I_Offroad
1A2gviii	Stationary combustion in manufacturing industries and construction: Other	B_Industry
1A3ai(i)	International aviation LTO (civil)	H_Aviation
1A3aii(i)	Domestic aviation LTO (civil)	H_Aviation
1A3bi	Road transport: Passenger cars	F_RoadTransport
1A3bii	Road transport: Light duty vehicles	F_RoadTransport
1A3biii	Road transport: Heavy duty vehicles and buses	F_RoadTransport
1A3biv	Road transport: Mopeds & motorcycles	F_RoadTransport
1A3bv	Road transport: Gasoline evaporation	F_RoadTransport
1A3bvi	Road transport: Automobile tyre and brake wear	F_RoadTransport
1A3bvii	Road transport: Automobile road abrasion	F_RoadTransport
1A3c	Railways	I_Offroad
1A3di(ii)	International inland waterways	G_Shipping
1A3dii	National navigation (shipping)	G_Shipping

<b>NFR</b>	<b>NFR name</b>	<b>GNFR</b>
1A3ei	Pipeline transport	I_Offroad
1A3eii	Other (please specify in the IIR)	I_Offroad
1A4ai	Commercial/institutional: Stationary	C_OtherStationaryComb
1A4aii	Commercial/institutional: Mobile	I_Offroad
1A4bi	Residential: Stationary	C_OtherStationaryComb
1A4bii	Residential: Household and gardening (mobile)	I_Offroad
1A4ci	Agriculture/Forestry/Fishing: Stationary	C_OtherStationaryComb
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	I_Offroad
1A4ciii	Agriculture/Forestry/Fishing: National fishing	I_Offroad
1A5a	Other stationary (including military)	C_OtherStationaryComb
1A5b	Other, Mobile (including military, land based and recreational boats)	I_Offroad
1B1a	Fugitive emission from solid fuels: Coal mining and handling	D_Fugitive
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	D_Fugitive
1B1c	Other fugitive emissions from solid fuels	D_Fugitive
1B2ai	Fugitive emissions oil: Exploration, production, transport	D_Fugitive
1B2aiv	Fugitive emissions oil: Refining / storage	D_Fugitive
1B2av	Distribution of oil products	D_Fugitive
1B2b	Fugitive emissions from natural gas (exploration, production, processing, transmission, storage, distribution and other)	D_Fugitive
1B2c	Venting and flaring (oil, gas, combined oil and gas)	D_Fugitive
1B2d	Other fugitive emissions from energy production	D_Fugitive
2A1	Cement production	B_Industry
2A2	Lime production	B_Industry
2A3	Glass production	B_Industry
2A5a	Quarrying and mining of minerals other than coal	B_Industry
2A5b	Construction and demolition	B_Industry
2A5c	Storage, handling and transport of mineral products	B_Industry
2A6	Other mineral products	B_Industry
2B1	Ammonia production	B_Industry
2B10a	Chemical industry: Other (please specify in the IIR)	B_Industry
2B10b	Storage, handling and transport of chemical products (please specify in the IIR)	B_Industry
2B2	Nitric acid production	B_Industry
2B3	Adipic acid production	B_Industry
2B5	Carbide production	B_Industry
2B6	Titanium dioxide production	B_Industry
2B7	Soda ash production	B_Industry
2C1	Iron and steel production	B_Industry
2C2	Ferroalloys production	B_Industry
2C3	Aluminium production	B_Industry
2C4	Magnesium production	B_Industry
2C5	Lead production	B_Industry
2C6	Zinc production	B_Industry
2C7a	Copper production	B_Industry
2C7b	Nickel production	B_Industry
2C7c	Other metal production	B_Industry
2C7d	Storage, handling and transport of metal products	B_Industry
2D3a	Domestic solvent use including fungicides	E_Solvents
2D3b	Road paving with asphalt	E_Solvents
2D3c	Asphalt roofing	B_Industry
2D3d	Coating applications	B_Industry
2D3e	Degreasing	E_Solvents
2D3f	Dry cleaning	E_Solvents
2D3g	Chemical products	E_Solvents
2D3h	Printing	E_Solvents
2D3i	Other solvent use	E_Solvents
2G	Other product use	E_Solvents
2H1	Pulp and paper industry	B_Industry
2H2	Food and beverages industry	B_Industry
2H3	Other industrial processes	B_Industry

NFR	NFR name	GNFR
2I	Wood processing	B_Industry
2J	Production of POPs	B_Industry
2K	Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)	B_Industry
2L	Other production, consumption, storage, transportation or handling of bulk products	B_Industry
3B1a	Manure management - Dairy cattle	K_AgriLivestock
3B1b	Manure management - Non-dairy cattle	K_AgriLivestock
3B2	Manure management - Sheep	K_AgriLivestock
3B3	Manure management - Swine	K_AgriLivestock
3B4a	Manure management - Buffalo	K_AgriLivestock
3B4d	Manure management - Goats	K_AgriLivestock
3B4e	Manure management - Horses	K_AgriLivestock
3B4f	Manure management - Mules and asses	K_AgriLivestock
3B4gi	Manure management - Laying hens	K_AgriLivestock
3B4gii	Manure management - Broilers	K_AgriLivestock
3B4giii	Manure management - Turkeys	K_AgriLivestock
3B4giv	Manure management - Other poultry	K_AgriLivestock
3B4h	Manure management - Other animals	K_AgriLivestock
3Da1	Inorganic N-fertilizers (includes also urea application)	L_AgriOther
3Da2a	Animal manure applied to soils	L_AgriOther
3Da2b	Sewage sludge applied to soils	L_AgriOther
3Da2c	Other organic fertilisers applied to soils (including compost)	L_AgriOther
3Da3	Urine and dung deposited by grazing animals	L_AgriOther
3Da4	Crop residues applied to soils	L_AgriOther
3Db	Indirect emissions from managed soils	L_AgriOther
3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products	L_AgriOther
3Dd	Off-farm storage, handling and transport of bulk agricultural products	L_AgriOther
3De	Cultivated crops	L_AgriOther
3Df	Use of pesticides	L_AgriOther
3F	Field burning of agricultural residues	L_AgriOther
3I	Agriculture other	L_AgriOther
5A	Biological treatment of waste - Solid waste disposal on land	J_Waste
5B1	Biological treatment of waste - Composting	J_Waste
5B2	Biological treatment of waste - Anaerobic digestion at biogas facilities	J_Waste
5C1a	Municipal waste incineration	J_Waste
5C1bi	Industrial waste incineration	J_Waste
5C1bii	Hazardous waste incineration	J_Waste
5C1biii	Clinical waste incineration	J_Waste
5C1biv	Sewage sludge incineration	J_Waste
5C1bv	Cremation	J_Waste
5C1bvi	Other waste incineration	J_Waste
5C2	Open burning of waste	J_Waste
5D1	Domestic wastewater handling	J_Waste
5D2	Industrial wastewater handling	J_Waste
5D3	Other wastewater handling	J_Waste
5E	Other waste	J_Waste
6A	Other (included in national total for entire territory)	M_Other
<b>'MEMO' ITEMS - NOT TO BE INCLUDED IN NATIONAL TOTALS</b>		
1A3ai(ii)	International aviation cruise (civil)	O_AviCruise
1A3aii(ii)	Domestic aviation cruise (civil)	O_AviCruise
1A3di(i)	International maritime navigation	P_IntShipping
1A5c	Multilateral operations	z_Memo
1A3	Transport (fuel used)	z_Memo
6B	Other not included in national total of the entire territory	z_Memo
11A	Volcanoes	N_Natural
11B	Forest fires	N_Natural
11C	Other natural emissions	N_Natural

## 7.2 Mapping methodology

The methodology used in the emissions mapping follow the guidelines in the 2023 EMEP/EEA emission inventory guidebook. The overall approach aims to allocate the national total emissions to the geographical location where they occur as accurately as possible. A distinction is made between point sources and area sources. Point sources are sources that can be treated individually and have an exact location, e.g. industrial plants. Area sources cover a group of minor emission sources with similar characteristics that cannot be treated individually because of the number of sources, e.g. residential plants. Some sectors are covered only by point sources or only by area sources, but many sectors cover both point and area sources. In the latter case point and area sources are treated separately in the data processing and following they are combined on sectoral level in the spatial emission mapping.

Emissions from point sources can be allocated to an exact location, e.g. the location of a power plant or an industrial plant. Activity data and/or emissions are available for a number of large plants e.g. from PRTR/E-PRTR reporting. These data are used either directly (emissions) or indirectly (activity data) to allocate point source emissions. Both locations and emissions are generally very accurate for point sources.


The individual source contribution cannot be determined for area sources, and emissions allocations are based on a number of spatial data sets. For each area source related available spatial data are evaluated and the closest related are used for emission mapping, taking into account completeness (must cover the entire national area), spatial resolution, accuracy, update frequency etc.

The 2023 EMEP/EEA Guidebook describes a tiered approach for spatial distribution of emissions, depending on the data availability and level of detail for the individual emission sources/sectors. Furthermore, different methodological tiers can be used for different pollutants from a source, e.g. point source emission data are most often available only for some pollutants, while emission mapping for remaining pollutants follow a lower tier method.

The concept of tiered mapping is summarised as follows:

- **Tier 3 methods** are based on closely related spatial emission or activity data, e.g. data for regulated processes and industries, and road traffic flows by vehicle type derived from surveys.
- **Tier 2 methods** are based on the use of proxy statistics relate to the sector, e.g. heat demand for the residential sector, agricultural animal statistics, and land parcel identification system data
- **Tier 1 methods** are based on loosely related proxy statistics, e.g. building use, population density, and land use.

The tiered methodology is outlined in the decision tree in Figure 7.1



**Figure 7.1 General decision tree for emissions mapping (EMEP/EEA 2023)**

Mapping of Ireland's emissions is done on a highly disaggregated level both regarding sectoral and spatial resolution. Spatial distribution keys (GeoKeys) are set up for each NFR category with a spatial resolution of 1 km x 1 km. GeoKeys are normalised tables including the share of an emission source that should be allocated to each grid cell. Some GeoKeys are used for all pollutants from a sector, while others are pollutant specific. Further, some sectors have been disaggregated into different sources and GeoKeys have been set up on source level and afterwards combined to create one overall GeoKey for the NFR sector. In this way, a high level of accuracy is ensured in the emission mapping as the highest tier level methodology is applied for all sources.

GeoKeys for the individual sectors/sources are built from a number of different spatial data. Some of the spatial data sets describe the emission allocation very accurate, e.g. E-PRTR reporting, while others are proxies for activity level or other related parameters; some being good proxies, e.g. mileage data

for emissions from road transport, others being less good proxies, e.g. population density for domestic solvent use.

The common methodological approach is to make an overlay analysis of one or more spatial layers and the 1 km x 1 km grid in a geographical information system (GIS).

Preparation of the GeoKey for railway transport is described here as an example, and related maps are shown in Figure 7.2. Railway transport is an area source, and the emissions are spatially allocated to the railway network. The railway network is available as a digital map including the network as lines (Figure 7.2 a), which allow the emissions to be distributed evenly to the railway network. Activity data based on railway statistics are added to the map's attribute data, and are used to improve the spatial distribution to allocate emissions according to the activity levels. An overlay of the railway network including activity data and the 1 km x 1 km grid is made in GIS (Figure 7.2 b), and the layers are intersected to cut the railway lines by the grid (Figure 7.2 c). The length of each line segment is calculated using standard GIS tools, and the share of each railway line in each grid cell is calculated. As more line segments can occur in the same grid cell, e.g. when two railway lines meet, the shares are summarised by grid cell to generate the GeoKey, which holds the share of the national emission by grid cell (Figure 7.2 d).





*Figure 7.2 Example of GeoKey preparation*

The emission mapping is made using an orthogonal grid with a resolution of 1 km x 1 km in the Irish projection TM65 (EPSG 29902). The spatial emissions are redistributed into the reporting grid, the EMEP/EEA grid with a spatial resolution of 0.1 degree x 0.1 degree, using the share of each 1 km x 1 km grid cell that intersects the individual 0.1 degree x 0.1 degree grid cells.

### 7.3 Mapping methods for each GNFR

A summary of the data and mapping approaches used in compilation of the spatial inventory for Ireland are outlined in the tables below by GNFR sector. An indication of the tier 1-3 categorisation has also been provided as a simple measure of uncertainty in the approach applied. A more detailed description of the methodologies applied is available in Plejdrup et al. (2017)

### 7.3.1 Public Power (GNFR A\_PublicPower)

Detailed location information for this sector were available on the individual large point sources for the NFR sector 1A1a Public Power. The emissions for some pollutants were available for the individual point sources, and in these cases, the data were used directly. For the pollutants where plant specific data were not available, the distribution is based on the activity data. The use of plant specific data and exact location of the emissions corresponds to a tier 3 method.

### 7.3.2 Industry (GNFR B\_Industry)

As shown in Table 7.1, this GNFR covers many different source categories and hence the available spatial data vary across sectors. The categories include both combustion related categories and categories where the emissions are related to the process.

Where detailed emissions and location information were available, e.g. from the E-PRTR or the EU ETS on the individual point source emissions for the NFR sectors in GNFR B Industrial Combustion sector, these were used to map emissions to the known location. This is the case for e.g. emissions from refining and other energy industries (NFR categories 1A1b and 1A1c respectively).

For other source categories, some data are available at point source level, but the coverage does not match the national total. In these cases the emissions covered by point sources are allocated to the relevant point sources and the residual emission is distributed according to a more general spatial distribution key, e.g. industrial heat demand. These two distributions are then combined to one GeoKey covering the total sectoral emission.

This approach is considered a tier 2 or tier 3 method.

### 7.3.3 Other stationary combustion (GNFR C\_OtherStationaryComb)

This GNFR category covers combustion in three subsectors, i.e. commercial/institutional, residential and agriculture. The most important sector in terms of emission contribution is residential combustion.

For commercial/institutional plants, the distribution is based on heat demand for commercial and public buildings as calculated by the Irish Heat Map. The Heat Map is based on a study from 2015 commissioned by SEAI to fulfil Ireland's requirements under article 14 of the Energy Efficiency Directive (2012/27/EU). As part of this study a spatial representation of Ireland's heat demand was developed.

For residential plants, the distribution is based on information from the 2011 census on primary fuel types in households combined with an estimated unit consumption calibrated with the estimated national residential fuel consumption and the emission factors used in the emission inventory.

For the agricultural sector, the spatial data on farmyards and buildings from the Land Parcel Information System (LPIS) were used.

This is considered tier 2/3 methodologies.

### 7.3.4 Fugitive Emissions (GNFR D\_Fugitive)

This sector covers both categories estimated as point sources (e.g. coal mining/handling, service stations and flaring) and area sources (e.g. natural gas distribution).

The point source data have been used to allocate emissions and, where available, activity data have been incorporated to further improve the distribution of emissions. Information on coal mining areas

as well as coal consumers were provided by the EPA, while a list of service stations was provided by the CSO. For natural gas distribution, the spatial information included for gas use in the Heat Map was utilised to distribute emissions.

This is considered tier 2/3 methodologies.

#### 7.3.5 Solvents (GNFR E\_Solvents)

The national emissions from domestic solvent use were mapped across the country using population density as spatial proxy. This approach is a tier 2 method.

For the remaining solvent use categories, there was some spatial information available, e.g. location of dry cleaners. However, both population density and industrial heat demand were used as spatial proxies to map emissions from coating applications, chemical products and printing.

This is considered tier 1/2/3 methodologies.

#### 7.3.6 Road transport (GNFR F\_RoadTransport)

Spatial mileage data for national roads (NR) provided by TII for total mileage and % heavy vehicles are used to allocate emissions from road transport on NR. Road transport on other roads is estimated as the residual of the national total mileage used in the inventory, and emissions are allocated to roads other than NR. As mileage data is not available for other roads than NR a polygon map of the road network is applied for mapping, thereby using road area as a proxy for the activity level. Separate GeoKeys are prepared for passenger cars including vans and 2-wheelers (PC), heavy vehicles including busses (HV), and all vehicles (PC+HV).

The approaches used are tier 3 for NR and tier 2 for remaining roads.

#### 7.3.7 National navigation (GNFR G\_Shipping)

The estimates of the emissions from national navigation were mapped using a buffer zone of six nautical miles around the coast of Ireland. The buffer zone was adjusted to take into account the shortest path between headlands.

This approach is a tier 2 method.

#### 7.3.8 Aviation (LTO) (GNFR H\_Aviation)

National total emissions from aircraft operating on the ground and in the air over Ireland, up to an altitude of 1000 m (equating to the take offs and landing – LTO) were mapped at the locations of the airports including a five-kilometre buffer zone. The number of LTOs at each airport was used to further improve the distribution of emissions.

This approach is a tier 3 method.

#### 7.3.9 Off road mobile sources (GNFR I\_Offroad)

This GNFR category comprises several different activities such as railways, fishing and agricultural machinery.

For railways, the railway network and data for annual passages were provided by Irish Rail and this information has been used to develop a GeoKey for this sector.

For fishing, the emissions have been distributed based on data for fishing areas within the Irish exclusive economic zone and fishing statistics.

For agricultural machinery, data on the number of different types of machinery at county level were obtained from the CSO and this information was combined with the land information from LPIS on cropland and improved grassland.

This is considered tier 2/3 methodologies.

#### 7.3.10 Waste handling and treatment (GNFR J\_Waste)

The estimates of the emissions from solid waste disposal on land were mapped at the locations of landfill sites. For composting 75 % of the emission were allocated to the licensed facilities while the remaining 25 % were allocated to non-urban residential buildings.

Emissions from clinical waste incineration, industrial waste incineration and cremation were mapped at the locations of the known facilities. Activity and location data for industrial waste incinerators and crematoria were available and used to weight emissions to areas of known activity proportionally.

Estimates of the national emissions from other waste handling (e.g. accidental fires) were mapped according to population density.

This is considered tier 1/2/3 methodologies.

#### 7.3.11 Agricultural livestock (GNFR K\_AgriLivestock)

National emissions from pigs and poultry were distributed based on detailed data on farms and animal numbers from the 2010 agricultural census provided by UCD.

National emissions from mink were distributed based on farm locations and animal numbers provided by EPA.

National emissions from cattle, sheep and horses were distributed based on data from the 2010 agricultural census provided by CSO on the number of animals per electoral district combined with the Land Parcel Identification System (LPIS) data on location of farmyards and buildings.

National emissions from goats, mules and asses, and deer were distributed based on data from the 2010 agricultural census provided by CSO on the number of animals per county combined with the Land Parcel Identification System (LPIS) data on location of farmyards and buildings.

This approach is a tier 2/3 method.

#### 7.3.12 Agricultural soils (Other emissions) (GNFR L\_AgriOther)

National emissions from sources related to agricultural soils, e.g. application of fertiliser and manure as well as grazing animals were distributed on cropland and/or grassland from LPIS, taking into account the animal density when distributing emissions from animal manure.

This approach is a tier 1/2 method.

#### 7.3.13 Aviation (Cruise) (GNFR O\_AviCruise)

This category includes cruise emissions from both national and international aviation. For national cruise emissions, the distribution is based on information on the number of flights between the major Irish airports and emissions are allocated to great circle lines between these airports.

For international cruise emissions, the majority of emissions will occur outside the Irish territory, but for the purposes of the submission, emissions are allocated evenly across the entire Irish area outlined by the Irish Exclusive Economic Zone (EEZ).

This is considered tier 1/2 methodologies.


#### 7.3.14 International navigation (GNFR P\_IntShipping)

Emissions from international navigation will largely occur outside the Irish territory, but for the purposes of the submission, emissions are allocated evenly across the sea area outlined by the Irish Exclusive Economic Zone (EEZ).


This approach is a tier 1 method.


#### 7.4 National total emission mapped by the EMEP 0.1 degree x 0.1 degree grid

Figure 7.3 a-e present the mapped national total emissions for NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, NH<sub>3</sub>, and PM<sub>2.5</sub> in Ireland by EMEP 0.1 x 0.1 degree grid.

















*Figure 7.3 National Total Emissions in 2023 for a) NO<sub>x</sub>, b) SO<sub>2</sub>, c) NMVOC, d) NH<sub>3</sub>, and e) PM<sub>2.5</sub>*

## Chapter Eight - Projections

### 8.1 Overview of Emissions Projections

Ireland's Environmental Protection Agency (EPA) is responsible for developing, preparing and reporting periodic projections of air pollutants. The EPA is also the responsible body for preparing greenhouse gas emissions projections that are submitted under the Governance of the Energy Union and Climate Action Regulation (EU) 2018/1999. In this section an overview of the emission projections is discussed.

Section 8.2 describes emission commitments for 2025 and 2030. Section 8.3 provides a short explanation of how energy projections are generated for both "*With Existing Measures*" and "*With Additional Measures*" scenarios. Information on key assumptions and underlying data are also provided.

Section 8.4 presents the emission projections for each of the pollutants covered and considers the key trends across the time series. Sections 8.5 to 8.13 present the key input assumptions and methodologies for the main sectors which include the impact of national policies and measures aimed at reducing greenhouse gas emissions and how these affect air pollutant emission levels.

Article 8 of Directive (EU) 2016/2284 requires biennial reporting from 2017 of projected emissions for SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, NMVOC, PM<sub>2.5</sub> and, if available, Black Carbon (BC) covering projection years 2020, 2025, 2030 and, where available, 2040 and 2050. 2025 is a mandatory reporting year.

The CLRTAP guidelines for reporting emissions projections state that parties to the Gothenburg Protocol within the geographical scope of the EMEP shall regularly update their projections and report every four years, from 2015 onwards, their updated projections for the years 2020, 2025 and 2030 and, where available, also for 2040 and 2050. Parties to the other protocols are encouraged to regularly update their projections and report every four years from 2015. In addition, parties should provide a "*With Existing Measures (WEM)*" and where relevant a "*With Additional Measures*" (WAM) projection estimate.

Projected emission estimates and supporting quantitative information are reported for SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, NMVOC, PM<sub>2.5</sub> and BC under Directive (EU) 2016/2284 utilising the updated 2023 reporting template contained within Annex IV of the Guidelines for Reporting Emissions and Projections Data under the CLRTAP<sup>5</sup>.

This chapter details emission projections under both the *With Existing Measures* scenario and *With Additional Measures* scenario for the following pollutants that are subject to emission reduction commitments for the period 2021-2030: SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, NMVOC and PM<sub>2.5</sub>.

---

<sup>5</sup> [Reporting instructions \(ceip.at\)](https://ceip.at)

## 8.2 Emission Reduction Commitments for 2020 and 2030

The National Emission Ceilings Directive (NECD, 2001/81/EC) was reviewed as part of the Clean Air Policy Package and a new Directive came into effect in December 2016<sup>6</sup>, Directive (EU) 2016/2284. Emission reduction commitments have been set for Ireland for 2020 and 2030 for SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, NMVOC and PM<sub>2.5</sub>. Table 8.1 details the emission reduction commitments in place for each pollutant for 2020 and 2030 for Ireland under Directive (EU) 2016/2284.

**Table 8.1. Emission Reduction Commitments for 2020 and 2030 (expressed as a percentage reduction on 2005 levels)**

Pollutant	SO <sub>2</sub>	NO <sub>x</sub>	NH <sub>3</sub>	NMVOC	PM <sub>2.5</sub>
2020	65%	49%	1%	25%	18%
2030	85%	69%	5%	32%	41%

## 8.3 With Existing Measures and With Additional Measures Scenarios - Approach

This year's projections take into account updated projected activity data provided by a number of key data providers including:

- Energy projections underpinning the 2025 emissions projections prepared by the Sustainable Energy Authority of Ireland (SEAI) in conjunction with the Economic and Social Research Institute (ESRI). The ESRI produce energy demand projections using the I3E model (Ireland Environment, Energy and Economy model)<sup>7</sup>.
- Agricultural projections provided by Teagasc (Agriculture and Food Development Authority) in February 2025 which incorporate the impact of Food Wise 2025<sup>8</sup>, Food Vision 2030<sup>9</sup> and the impacts of the Brexit related Trade and Cooperation Agreement (TCA) reached between the EU for the agriculture sector.

National emission projections utilise two energy projections scenarios compiled by the SEAI:

- The *Baseline or With Existing Measures (WEM)* energy projections project forward Ireland's energy demand, incorporating the expected impacts of policies and measures that were in place (e.g. legislatively provided for) by the end of 2023, the latest inventory year. It includes progress in the implementation of policies and measures as set out in Ireland's National Renewable Energy Action Plan (NREAP), National Energy Efficiency Action Plan (NEEAP) and the National Development Plan 2018-2027 and the 2019, 2021, 2023 and 2024 Climate Action Plans (CAP)<sup>10</sup>. Only those CAP measures that are considered to be "implemented" or "adopted" by the end of the latest inventory year (2023), using the official terminology, are included in the WEM scenario.

<sup>6</sup> DIRECTIVE (EU) 2016/2284 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC

<sup>7</sup> <https://www.esri.ie/current-research/the-i3e-model>

<sup>8</sup> <http://www.agriculture.gov.ie/foodwise2025/>

<sup>9</sup> [gov.ie](http://www.gov.ie) - Food Vision 2030 - A World Leader in Sustainable Food Systems ([www.gov.ie](http://www.gov.ie))

<sup>10</sup> [gov](http://www.gov.ie) - Climate Action Plan 2024 ([www.gov.ie](http://www.gov.ie))

- The *Advanced or With Additional Measures* (WAM) energy projections present an alternative view of future energy demand that accounts for further implementation of the *With Existing Measures* scenario in addition to the implementation of planned government policies and measures in Ireland's 2024 Climate Action Plan<sup>6</sup>. Only planned measures in CAP 2024 with a pathway for achievement are included in the WAM scenario.

For the energy demand projections, the Economic and Social Research Institute (ESRI) used the I3E model<sup>11</sup>. The I3E model is an intertemporal computable general equilibrium (CGE) model which reproduces the structure of the Irish economy in its entirety, including productive sectors, households, and the government, among others.

#### *Main scenario and sensitivity scenarios*

Typically, SEAI request the development of two projections from the I3E reflecting high and low-price scenarios. One price scenario is used for the main scenario and the other is usually used as a sensitivity analysis. The 2024 I3E High-price scenario is used as the basis for the main scenario for the 2024 National Energy Projections.

#### *Baseline Assumptions*

In CGE modelling, a change in a policy variable/parameter or a set of policy variables/parameters is defined as a shock process or experiment. The effects of the experiment are given as the paths of variables that are solved within the model and are reported in terms of a percentage change with respect to the baseline. The baseline is a path reflecting no change in any policy variables/parameters but an economy evolves according to its internal dynamics. Since there are no external interventions in the economy, the base path is also called the business-as-usual path. This pathway now includes a carbon tax that increases to €100 per tonne by 2030 and is constant thereafter.

#### *Fossil Fuel Prices*

Selection of the fossil fuel price projections for the 2025 submissions took place in August 2024. The fuel price projections are selected from available sources and are compared with other price sources to ensure they are within a credible range.

The European Commissions "Central" datasets were selected as the 2025 National Energy Projections (NEP) "High" price scenario. For gas and oil the updated EC "Central" projections are lower than the high price or 'baseline' scenario used in 2024 projections (provided by the European Commission in 2022). These projections were especially high due to Russia's invasion of Ukraine. The Coal projection from the EC is comparable to that used in the 2024 projections.

The International Energy Agency's (IEA) price projections as published each year in the World Energy Outlook (WEO) document were selected for the "Low" price sensitivity for gas, oil and coal.

#### *Carbon prices*

The recommended ETS carbon prices are based on the EU Reference Scenario. The I3E model implements two carbon prices, namely the Irish carbon tax and the EU ETS price. The EU ETS price

---

<sup>11</sup> <https://www.esri.ie/current-research/the-i3e-model>



projections are taken from the EU reference scenario. The Irish carbon tax currently stands at €48.50 a tonne in 2023. Both WEM and WAM scenarios include a varying Irish carbon tax that reaches €100 per tonne by 2030 and a varying EU ETS price that increases annually to €95 per tonne by 2030 and €190 per tonne by 2050.

The software used to model the Irish Electricity Market is PLEXOS which is a power systems modelling tool used for electricity market modelling and planning.

To produce the finalised *Baseline* or WEM energy projections, SEAI amends the output of the energy demand produced by ESRI to take account of the expected impact of energy efficiency measures put in place before the end of 2023 but which are considered too recent to be detectable in any time-series analysis. The *Advanced* or WAM energy projections builds on the *Baseline* projections with adjustments made to account for implementation of additional policies and measures outlined in Ireland's 2024 Climate Action Plan<sup>6</sup>.

Assumptions were finalised in quarter 4 of 2024. Determination of anticipated progress in the implementation of policies and measures was coordinated by the EPA in discussion with the relevant Government Departments.

Further details on the models used for preparing the energy projections (i.e. COSMO, I3E, Plexos Integrated Energy Model, SEAI's National Energy Modelling Framework, SEAI BioHeat Model and the FAPRI Ireland model) are included in the 2023 submission made under Article 18 (4) of the Governance Regulation of the Energy Union and Climate Action (EU) 2018/1999. This is available in the relevant 2024 submission folder at the following link: <https://reportnet.europa.eu/public/country/IE>. Under Name, filter for 'National projections of anthropogenic greenhouse gas emissions [2024]' and download Table 4 under Files.


## 8.4 Key Trends

In the following sections, both the *With Existing Measures (WEM)* and *With Additional Measures (WAM)* scenarios are presented in the context of the emission commitment set for 2030. The year 2025 is also discussed to give an indication of trends to 2030. The WAM scenario is then considered in more detail. The 1990-2023 inventory was used as the baseline historic inventory for the emissions projections. Figures in the following sections include historic and projected air pollutant emission estimates from 1990-2030 for both the WEM and WAM scenarios. A linear reduction pathway (Article 4 (2) Directive (EU) 2016/2284) between the 2020 and 2030 emission reduction commitment (ERC) for each pollutant is also presented. Tables in the following sections present projected emission estimates up to and including 2030 under both the WEM and WAM scenarios. The 2005 emission levels are also provided for context in terms of the base year under the emission reduction commitments.

### 8.4.1 Sulphur Dioxide (SO<sub>2</sub>)


Emission projections for SO<sub>2</sub> for the WEM and WAM scenarios are presented in Figure 8.1. The emission reduction pathway between the ERCs for 2020 and 2030 is also presented. The emission projections predict compliance throughout the projected period. Key sources of SO<sub>2</sub> emissions in the projections include emissions from residential, commercial and industry sectors.

Total SO<sub>2</sub> emissions under the WEM scenario are projected to be 7.29 kt (90.2 per cent below 2005 levels) in 2030. Total SO<sub>2</sub> emissions are projected to be 7.19 kt (90.3 per cent below 2005 levels) in 2030 under the WAM scenario. The ERC for SO<sub>2</sub> for 2030 is an 85.0 per cent reduction on 2005 emission levels.



**Figure 8.1 SO<sub>2</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios**

SO<sub>2</sub> emissions under the WAM scenario by source sector are presented in Figure 8.2. The combined Residential and Commercial sectors account for 66.7 per cent and 64.7 per cent of emissions in 2023 and 2030, respectively, followed by Manufacturing Industries and Construction with 18.7 per cent and 20.8 per cent in 2023 and 2030, respectively.



**Figure 8.2 SO<sub>2</sub> Emission Projections for the With Additional Measures Scenario by Source Sector**

Table 8.2 presents the projected Emissions of SO<sub>2</sub> under the WEM and WAM by sector over the period 2005 to 2030.

**Table 8.2. Projected Emissions of SO<sub>2</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)**

<b>With Existing Measures Scenario</b>				
	<b>2005</b>	<b>2023</b>	<b>2025</b>	<b>2030</b>
Public Electricity and Heat Production	42.50	0.71	0.88	0.63
Manufacturing Industries and Construction	14.15	1.37	1.41	1.50
Residential & Commercial/Institutional	14.32	4.91	4.86	4.83
Agriculture/Forestry/Fishing	0.86	0.03	0.03	0.04
Transport	1.35	0.25	0.25	0.26
Other NFR Sectors	1.07	0.09	0.04	0.04
<b>TOTAL</b>	<b>74.25</b>	<b>7.36</b>	<b>7.48</b>	<b>7.29</b>
<b>With Additional Measures Scenario</b>				
Public Electricity and Heat Production			0.98	0.73
Manufacturing Industries and Construction			1.41	1.50
Residential & Commercial/Institutional			4.86	4.65
Agriculture/Forestry/Fishing			0.03	0.04
Transport			0.25	0.24
Other NFR Sectors			0.04	0.04
<b>TOTAL</b>			<b>7.57</b>	<b>7.19</b>

#### 8.4.2 Nitrogen Oxides (NO<sub>x</sub>)


Emission projections for NO<sub>x</sub> for the WEM and WAM scenarios are presented in Figure 8.3. The emission reduction pathway between the ERCs for 2020 and 2030 is also presented.

Article 4 (3) of the National Emission Ceiling Directive provides that emissions of nitrogen oxides from categories 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with the ERCs for 2020 and 2030. The WEM emissions and WAM emissions (in addition to 2020 and 2030 emission reduction pathway) displayed in Figure 8.3 excludes emissions from these categories (3B and 3D).

Emission projections predict compliance with the ERC for 2030 under the WEM and WAM scenarios. Emissions from transport is the largest contributor to NO<sub>x</sub> emissions.


Total NO<sub>x</sub> (excluding categories 3B and 3D) emissions under the WEM scenario are projected to be 35.18 kt (73.7 per cent below 2005 levels) in 2030. Total NO<sub>x</sub> (excluding categories 3B and 3D) emissions are projected to be 32.80 kt (75.5 per cent below 2005 levels) in 2030 under the WAM scenario. The ERC for NO<sub>x</sub> for 2030 is a 69 per cent reduction on 2005 emission levels.





**Figure 8.3 NO<sub>x</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios**

Figure 8.4 presents the total NO<sub>x</sub> emissions under the WAM scenario by source sector. The largest sectoral contribution is from agriculture and transport. In terms of compliance assessment (i.e. excluding agricultural categories 3B and 3D) transport accounts for 58.5 per cent and 39.8 per cent share of emissions in 2023 and 2030 respectively.



**Figure 8.4 NO<sub>x</sub> Emission Projections for the With Additional Measures Scenario by Source Sector**

Table 8.3 summarises the projected emissions of NO<sub>x</sub> under the WEM and WAM by sector over the period 2005 to 2030 (including categories 3B and 3D).


**Table 8.3. Projected Emissions of NO<sub>x</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)**

<b>With Existing Measures Scenario</b>				
	<b>2005</b>	<b>2023</b>	<b>2025</b>	<b>2030</b>
Public Electricity and Heat Production	32.38	4.33	4.85	3.49
Manufacturing Industries and Construction	16.45	8.15	7.91	8.06
Residential & Commercial	9.31	6.57	6.85	6.98
Agriculture/Forestry/Fishing	12.47	3.39	3.20	2.43
Transport	61.83	32.18	20.73	13.90
Agriculture	31.92	29.44	31.50	31.09
Other NFR Sectors	1.27	0.36	0.31	0.31
<b>TOTAL</b>	<b>165.64</b>	<b>84.42</b>	<b>75.35</b>	<b>66.27</b>
<b>With Additional Measures Scenario</b>				
Public Electricity and Heat Production			4.73	3.01
Manufacturing Industries and Construction			7.91	7.76
Residential & Commercial			6.83	6.22
Agriculture/Forestry/Fishing			3.20	2.43
Transport			20.75	13.06
Agriculture <sup>10</sup>			30.88	29.92
Other NFR Sectors			0.31	0.31
<b>TOTAL</b>			<b>74.59</b>	<b>62.72</b>

### 8.4.3 Ammonia (NH<sub>3</sub>)

Figure 8.5 presents the emission projections for NH<sub>3</sub> for the WEM and WAM scenarios. The emission reduction pathway between the ERCs for 2020 and 2030 is also presented. Overall, the NH<sub>3</sub> emission projections predict compliance with the 2030 ERC under both the WEM and the WAM scenario. Ireland is not compliant with the ERC for 2020 and 2021 but is in compliance from 2022- 2030 under both the WEM and the WAM scenario.

Total NH<sub>3</sub> emissions under the WEM scenario are projected to be 116.44 kt (5.9 per cent below 2005 levels) in 2030. Total NH<sub>3</sub> emissions are projected to be 111.05 kt (10.3 per cent below 2005 levels) in 2030 under the WAM scenario. The ERC for NH<sub>3</sub> for 2030 is a 5 per cent reduction on 2005 emission levels.




**Figure 8.5 NH<sub>3</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios**

The reduction in the WAM scenario compared to the WEM scenario over the period 2024-2030 is attributable to the additional measures (over and above those in WEM scenario) that are included in the most recent Teagasc Marginal Abatement Cost Curve<sup>12</sup>, and the Department of Agriculture, Food and the Marine National Climate and Air Roadmap for the Agriculture Sector “AgClimatise”<sup>13</sup> and various iterations of the Irish Climate Action Plans over recent years. Key measures are discussed in section 8.13.

<sup>12</sup> [Marginal Abatement Cost Curve 2023 - Teagasc | Agriculture and Food Development Authority](#)

<sup>13</sup> <https://www.gov.ie/en/publication/07fbe-ag-climatise-a-roadmap-towards-climate-neutrality/>



**Figure 8.6 NH<sub>3</sub> Emission Projections for the With Additional Measures Scenario by Source Sector**

Ammonia emissions by source sector under the WAM scenario are presented in Figure 8.6. Table 8.4 presents the projected ammonia emissions under the WEM and WAM by sector over the period 2005 to 2030.

**Table 8.4. Projected Emissions of NH<sub>3</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)**

<b>With Existing Measures Scenario</b>				
	<b>2005</b>	<b>2023</b>	<b>2025</b>	<b>2030</b>
Dairy Cattle	10.72	15.64	15.71	16.95
Other Cattle	34.77	30.22	29.46	28.46
Other Livestock	9.99	9.92	10.06	11.24
Direct Soils	53.39	47.15	47.94	46.13
N-excreted on Pasture	12.25	12.61	12.43	12.64
Residential & Commercial	0.12	0.08	0.08	0.13
Road Transport	2.31	0.52	0.52	0.59
Other	0.244	0.21	0.23	0.28
<b>Total</b>	<b>123.79</b>	<b>116.36</b>	<b>116.45</b>	<b>116.44</b>
<b>With Additional Measures Scenario</b>				
Dairy Cattle			15.31	16.10
Other Cattle			29.29	27.68
Other Livestock			9.50	9.38
Direct Soils			42.99	44.81
N-excreted on Pasture			12.18	12.18
Residential & Commercial			0.08	0.12
Road Transport			0.52	0.49
Other			0.23	0.28
<b>Total</b>			<b>110.10</b>	<b>111.05</b>


#### 8.4.4 Non-Methane Volatile Organic Compounds (NMVOCs)

Emission projections for NMVOC under WEM and WAM scenarios are presented in Figure 8.7. The emission reduction pathway between the ERCs for the 2020 and 2030 is also provided.

Article 4 (3) of the Directive (EU) 2016/2284 provides that emissions of NMVOCs from categories 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with the ERCs for 2020 and 2030. The WEM and WAM emission estimates (in addition to 2020 and 2030 emission reduction pathway) displayed in Figure 8.7 excludes emissions from these categories (3B and 3D).

Emission projections predict non-compliance for the ERC for 2030 under both the WEM and WAM scenarios. Additionally, as outlined in chapter 9, emissions were also non-compliant with the ERC for 2020. However, Ireland has applied for an adjustment (in relation to 2H2 Spirit production) as allowed under Article 5 and Part 4 of Annex IV of the Directive (EU) 2016/2284 for 2020. When this adjustment is taken into account the ERC for 2020 is complied with. Figure 8.8 presents NMVOC emissions to 2030 if the same adjustment process was applied for 2030.


Total NMVOC (excluding categories 3B and 3D) emissions under the WEM scenario are projected to be 74.17 kt (3.5 per cent below 2005) in 2030. Total NMVOC (excluding categories 3B and 3D) emissions are projected to be 73.07 kt (4.9 per cent below 2005) in 2030 under the WAM scenario. The ERC for NMVOC for 2030 is a 32 per cent reduction on 2005 emission levels.



**Figure 8.7 NMVOC Emission Projections for the With Existing Measures and With Additional Measures Scenarios<sup>14</sup>**

<sup>14</sup> Article 4 (3) of the National Emission Ceiling Directive provides that emissions of non-methane volatile organic compounds from categories 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with 2020 and 2030 targets. The *With Existing Measures* emissions and *With Additional Measures* emissions in addition to 2020 and 2030 targets displayed in this graph exclude emissions from these categories (3B and 3D).

With respect to Figure 8.8, total NMVOC emissions (excluding categories 3B and 3D and category 2H2 spirit production) under the WEM scenario are projected to be 44.02 kt (35.7 per cent below 2005) in 2030. Total NMVOC emissions (excluding categories 3B and 3D and category 2H2 spirit production) under the WAM scenario are projected to be 42.92 kt (37.3 per cent below 2005). The ERC for NMVOC for 2030 is a 32 per cent reduction on 2005 emission levels. It is therefore projected that in this circumstance the ERC for 2030 will be complied with under both the WEM and WAM scenarios.




**Figure 8.8 NMVOC Emission Projections for the With Existing Measures and With Additional Measures Scenarios where an adjustment is applied for category 2H2 Spirit production<sup>15</sup>**

Projected total NMVOC emissions by source sector under the WAM scenario are presented in Figure 8.9. In terms of compliance assessment (which excludes agricultural categories 3B and 3D) projected emissions from Solvents and Fugitive emissions account for 29.9 per cent and 29.9 per cent of the total under the WEM and WAM scenarios in 2030, respectively. Food and Beverage production accounts for 49.6 per cent and 50.3 per cent of emissions in 2030 under the WEM and WAM scenarios, respectively.

Projected emissions across the time series show an increase in both scenarios to 2030 mostly due to increase in emissions from the Food and Beverages Industry. All other sectors are projected to be relatively stable with no significant increases nor reductions.

Table 8.5 presents emissions under the WEM and WAM by sector over the period 2005 to 2030.

<sup>15</sup> The *With Existing Measures* emissions and *With Additional Measures* emissions in addition to 2020 and 2030 targets displayed in this graph exclude emissions from NFR categories 3B (manure management) and 3D (agricultural soils), and category 2H2 (Spirits production).



**Figure 8.9 NMVOC Emission Projections for the With Additional Measures Scenario by Source Sector**

**Table 8.5. Projected Emissions of NMVOC under the With Existing Measures and With Additional Measures Scenarios (kt)**


<b>With Existing Measures Scenario</b>				
	<b>2005</b>	<b>2023</b>	<b>2025</b>	<b>2030</b>
Public Electricity and Heat Production	0.37	0.27	0.26	0.17
Manufacturing Industries and Construction	2.70	2.85	2.97	3.73
Residential & Commercial	14.05	6.74	6.80	7.20
Agriculture/Forestry/Fishing	1.04	0.20	0.19	0.17
Transport	14.99	3.11	3.67	3.35
Agriculture <sup>11</sup>	43.08	42.09	42.07	42.69
Fugitive and Solvents	28.93	20.27	20.93	22.11
Food and beverages Industry	13.53	29.30	31.26	36.77
Other NFR Sectors	1.24	0.85	0.79	0.67
<b>Total</b>	<b>119.92</b>	<b>105.69</b>	<b>108.94</b>	<b>116.86</b>
<b>With Additional Measures Scenario</b>				
Public Electricity and Heat Production			0.26	0.13
Manufacturing Industries and Construction			2.97	3.61
Residential & Commercial			6.79	6.76
Agriculture/Forestry/Fishing			0.19	0.16
Transport			3.66	3.14
Agriculture <sup>11</sup>			41.74	42.00
Fugitive and Solvents			20.93	21.89
Food and beverages Industry <sup>12</sup>			31.26	36.77
Other NFR Sectors			0.80	0.61
<b>Total</b>			<b>108.59</b>	<b>115.07</b>

#### 8.4.5 Particulate Matter < 2.5 µm in diameter (PM<sub>2.5</sub>)

Emission projections for PM<sub>2.5</sub> for the WEM and WAM scenarios are presented in Figure 8.10. The emission reduction pathway between the ERCs for the 2020 and 2030 is also presented.

Total PM<sub>2.5</sub> emissions under the WEM scenario are projected to be 10.42 kt (44.1 per cent below 2005 levels) in 2030. Total PM<sub>2.5</sub> emissions are projected to be 10.01 kt (46.3 per cent below 2005 levels) in 2030 under the WAM scenario. The ERC for PM<sub>2.5</sub> for 2030 is a 41 per cent reduction on 2005 emission levels. It is therefore projected that the ERC for 2030 will be complied with under both the WEM and WAM scenarios.

Table 8.6 presents the projected PM<sub>2.5</sub> emissions under the WEM and WAM by sector over the period 2005 to 2030.




**Figure 8.10 PM<sub>2.5</sub> Emission Projections for the With Existing Measures and With Additional Measures Scenarios**

Projected emissions across the time series show an increase for both the WEM and WAM scenarios up to 2030 (Figure 8.10). This trend is largely the result of projected increases in emissions due to increase use of Biomass in the Residential and Commercial and the Industry sectors.

Projected total PM<sub>2.5</sub> emissions by source sector under the WAM scenario are presented in Figure 8.11. Emissions from Residential and Commercial combined are the largest contributor to the projected emissions and account for 48.3 per cent and 47.9 per cent of the total under the WEM and WAM scenarios in 2030, respectively.





**Figure 8.11 PM<sub>2.5</sub> Emission Projections for the With Additional Measures Scenario by Source Sector**

**Table 8.6. Projected Emissions of PM<sub>2.5</sub> under the With Existing Measures and With Additional Measures Scenarios (kt)**

<b>With Existing Measures Scenario</b>				
	<b>2005</b>	<b>2023</b>	<b>2025</b>	<b>2030</b>
Public Electricity and Heat Production	1.07	0.21	0.19	0.07
Manufacturing Industries and Construction	2.19	1.18	1.25	1.63
Residential & Commercial/Institutional	8.84	4.44	4.47	5.04
Agriculture/Forestry/Fishing	0.58	0.10	0.10	0.10
Transport	2.94	1.39	1.27	1.27
Agriculture	0.74	0.84	0.83	0.84
Other NFR Sectors	2.28	1.15	1.43	1.47
<b>TOTAL</b>	<b>18.65</b>	<b>9.31</b>	<b>9.54</b>	<b>10.42</b>
<b>With Additional Measures Scenario</b>				
Public Electricity and Heat Production			0.19	0.06
Manufacturing Industries and Construction			1.25	1.62
Residential & Commercial/Institutional			4.47	4.80
Agriculture/Forestry/Fishing			0.10	0.10
Transport			1.27	1.13
Agriculture			0.83	0.84
Other NFR Sectors			1.43	1.47
<b>TOTAL</b>			<b>9.53</b>	<b>10.01</b>

## 8.5 Energy Industries (NFR 1A1)

Public Electricity and Heat Production (1A1a) covers all electricity generation including electricity generated from renewable sources. The Plexos\_Ireland model was used to project future electricity generation. As an electrical systems model, the core input data comprises technical details of generators, transmission lines and loads, as well as fuel costs, operational costs and emission reduction rates and costs.

In 2023, renewable generation accounted for 40.7% of electricity, an increase from 38.6% in 2022. Renewable electricity generation capacity is dominated by wind. In 2030 it is estimated that renewable energy generation increases to approximately 68 per cent of electricity consumption.

The use of peat for electricity generation in Ireland ceased at the end of 2023. One former peat station has planning permission to operate one unit on 100% biomass from 2024 to 2030. The remaining two units will switch from distillate oil to natural gas in 2026.

One coal plant will run to 2025, after which it will be run on heavy fuel oil as a 'unit of last resort' for security of supply reasons only.

In the *Advanced* energy projection (and therefore the *With Additional Measures* emissions scenario) for 2030 it is estimated that renewable energy generation increases to approximately 68 per cent of electricity consumption. This is lower than the estimated 80 per cent stated in the 2024 Climate Action Plan. It is mainly a result of additional expansion in wind energy and Solar PV.

In terms of inter-connection, both scenarios assume that the Greenlink 500MW interconnector to the UK will come on stream in January 2025 and the Celtic 700MW interconnector to France comes on stream in 2027. The North-South Tie Line provides 400 MW by the end of 2025 with 1350 MW being provided from 2027.

Emission factors for NO<sub>x</sub> and SO<sub>2</sub> predominantly are based on those used for the year 2023 in the national inventory (1990-2023). Based on available data, the projections take into account emission limit values provided in the transitional national plan for large combustion plants (e.g. coal fired electricity generation plant).

There has been significant reduction in the use of oil in electricity generation due to the closure and decommissioning of oil fuelled generation plants. Oil is also used as a start-up fuel in coal fired generation stations. The 2023 SO<sub>2</sub> inventory emission factor for oil is assumed for all future years.

NMVOC and PM<sub>2.5</sub> emission factors for most of the other relevant fuel types (i.e. coal, biomass and non-renewable wastes) are assumed to remain constant at the 2023 value utilised in the national inventory (1990-2023).

### 8.5.1 Oil and Gas Refining and Solid Fuel Manufacturing (NFR 1A1b and 1A1c)

Projected SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and PM<sub>2.5</sub> for oil refining and solid fuel manufacture are based on data provided by the relevant operators as energy demand from these sectors is not included in SEAI's energy projections. The oil refining sector (1A1b) in Ireland consists of a single installation. Projections are based on the growth rate of projected greenhouse gas emissions, which are provided to the EPA by the relevant installation operators.

## 8.6 Manufacturing Industries and Construction (NFR 1A2)

Projected SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emissions are based on SEAI's *Baseline* and *Advanced* energy projections and are therefore estimated for both the WEM and WAM scenarios.

Energy projections are provided to the EPA at an aggregated level only (i.e. 1A2) and underpin the air pollutant projections for this sector. The projected emission factors for SO<sub>2</sub> and NO<sub>x</sub> from the combustion of coal, natural gas and petroleum coke are based on the weighted average emission factor for all fuels across the sub sectors 1A2a – 1A2g from the values used in the inventory.

The following policies and measures relevant to the industry sector are included in the *Baseline* (WEM) Projection:

- SEAI Large Industry Energy Network (LIEN) Programme
- Accelerated Capital Allowances - Industry
- SEAI Combined Heat and Power (CHP) Programme
- Excellence in Energy Efficiency Design (EXEED) Industry

The WAM scenario includes an extension of the above policies and measures.

## 8.7 Transport (NFR 1A3)

Transport emissions cover Aviation (1A3a), Road Transportation (1A3b), Rail (1A3c), Navigation (1A3d) and Other transportation (Natural gas pipeline compressors, 1A3e). Projected emissions of NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> from road transport are based on SEAI's *Baseline* and *Advanced* energy projections and are therefore estimated for both the WEM and WAM scenarios.

It is assumed that fuel use in rail and navigation will remain constant at 2023 levels out to 2030.

Other Transportation (1A3e) refers to the use of natural gas for combustion in natural gas pipeline compressor stations. Emissions from this sector are inferred from projected gas demand in the residential, industrial and commercial and institutional services sectors from the energy projections provided by SEAI and are calculated for both the WEM and WAM scenarios.

### 8.7.1 Domestic aviation (NFR 1A3a)

SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and PM<sub>2.5</sub> emission projections from aviation are estimated using the 2023 inventory also in combination with projected data, where available, related to aircraft movements as provided to the EPA by the relevant airport authorities. Emissions associated with all LTO (landing and take-off) cycles are calculated.

### 8.7.2 Road transportation (NFR 1A3b)

The transport sector is a large energy using sector and is reliant on imported fossil fuels in the form of petroleum products and biofuels. Freight transport energy demand is strongly influenced by the level of commercial activity in the economy. Personal transport energy demand is influenced by both the level of employment as well as the oil price. Biofuel uptake in Ireland is driven primarily through the

Biofuel Obligation Scheme, an obligation on fuel suppliers to blend an increasing percentage of biofuel with their fuel.

The Climate Action Plan biofuels target will continue to be delivered through annual increases in the statutory renewable transport fuel obligation (RTFO) on fuel suppliers requiring a minimum proportion of renewable transport fuel supply. The trajectory of these annual increase in the RTFO rate for the next two years, and indicative rates to 2030, has been set out in the updated Renewable Fuels for Transport Policy Statement 2023-2025 published in June 2023. 2023 blend rates are E6.4, B8.6 with planned increases to E10 & B12 in 2025 in both WEM & WAM scenarios. Bioethanol will remain at this level thereafter with Biodiesel increasing to B20 in 2030 in WEM and WAM.

In terms of Electric Vehicles, the baseline energy projection assumes approximately 579,702 Electric Vehicles on the road by 2030 (includes 270,142 Passenger Battery Electric Vehicles, 279,410 Plug in Hybrid Electric Vehicles, 15,000 electric vans & 150 HGVs). The Advanced Scenario assumes 30% of the vehicle fleet or approximately 670,751 Electric Vehicles on the road by 2030 (includes 330,740 Passenger Battery Electric Vehicles, 279,410 Plug in Hybrid Electric Vehicles, 30,000 electric vans & 600 HGVs).

### 8.7.3 Rail (NFR 1A3c)

SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission projections from rail transport are estimated. It is assumed that fuel use in the sector will remain constant at 2023 levels out to 2030. It is also assumed that SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission factors remain the same as for 2023.

### 8.7.4 Navigation (NFR 1A3d)

SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission projections from navigation are estimated. Gasoil/diesel consumption in in-land navigation is assumed to remain constant at the 2023 level out to 2030. SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission factors are assumed to remain constant at the 2023 level.

### 8.7.5 Gas Transmission (NFR 1A3e)

Emission projections for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, and PM<sub>2.5</sub> from natural gas transmission in Ireland's natural gas pipeline network are estimated. Future gas demand for "own use and distribution" is inferred based on projected gas demand in the residential, commercial services and industrial sectors provided by SEAI. Subtracting the amount of gas estimated to be lost from the distribution network allows "own use" gas demand and associated emissions to be estimated. It is assumed that SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and PM<sub>2.5</sub> emission factors remain constant at the 2023 level.

## 8.8 Residential and Commercial/Institutional (NFR 1A4)

The WEM projections include the following policies and measures for Residential and Commercial/Institutional sectors:

- SEAI heat pump grants continuing at current level until 2030;
- 1 TWh Biomethane by 2030;
- 0.07 TWh District Heating by 2030;
- SEAI energy efficiency grants continue at their current level until 2030;
- Current levels of Community Energy and EXEED grants assumed to continue until 2030.

The WAM projections include the following policies and measures for Residential and Commercial/Institutional sectors:

- 4.30 TWh Biomethane by 2030;
- 0.21 TWh District Heating by 2030;
- SEAI energy efficiency grants increase to support target heat pump delivery;
- Current levels of Community Energy and EXEED grants assumed to continue until 2039.

#### 8.8.1 Commercial/Institutional (NFR 1A4a)

Projected SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emissions are based on SEAI's *Baseline* and *Advanced* energy projections and are therefore estimated for both the WEM and WAM scenarios. Oil and gas account for the majority of non-electricity energy demand in this sector. These fuels are used predominantly for space-heating purposes. Projected emissions of NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> are estimated. It is assumed that SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission factors remain constant at the 2023 level.

#### 8.8.2 Residential (1A4b)

Projected SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emissions are based on SEAI's *Baseline* and *Advanced* energy projections and are therefore estimated for both the WEM and WAM scenarios.

Projected emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> are estimated. It is assumed that SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission factors remain constant at 2023 levels.

## 8.9 Combustion in Agriculture and Fishing (NFR 1A4c)

Projected emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> are estimated for the combustion of diesel in the Agriculture sector and in the Fishing sector. Projected fuel use in the Agriculture sector is included in the energy projections by SEAI. For mobile combustion the same modelling approach is used as in the national inventory which takes into account the relevant Tier legislation as it affects the age structure of mobile agricultural machinery. For stationary combustion in agriculture and combustion in fishing, SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> emission factors remain constant at 2023 levels.

## 8.10 Cross cutting measures impacting key energy sectors

There are cross cutting measures that have a significant impact across a number of sectors in terms of energy consumption and a reduction in emissions. These included carbon tax and supports for renewable heat which are described below.

- Carbon tax is a cross-cutting measure that applies to industry, residential, commercial services, transport and agriculture fuel. Under the Baseline and Advanced energy scenarios (and therefore WEM and WAM scenarios), there is a varying carbon tax that increases by €7.50 per annum, reaching €100 per tonne by 2030.
- In terms of renewable heat, the WEM and WAM scenarios assume that current funding under the Support Scheme for Renewable Heat will be extended beyond 2030.
- It is also assumed that there are over 121, 000 domestic buildings with heat pumps by 2030 under the WEM scenario. Under the WAM scenario it is assumed that over 540,000 heat pumps will be in place in domestic buildings, 400,000 of these are in new homes. Both WEM and WAM scenarios assume that there will be no new gas boilers from 2025 in new dwellings, based on building regulations. District heating is also assumed to be deployed (growth to 0.07 and 0.21 TWh by 2030 in the WEM and WAM scenario, respectively) to be split across public, commercial/services and residential buildings.
- Under the WAM scenario, a total of 4.30 TWh of biomethane use across the heat sector (residential and commercial/services buildings) and transport sectors by 2030 is assumed.

## 8.11 Fugitive emissions (NFR 1B)

Projected NMVOCs emissions from category 1B are estimated from three sources: Refining of oil products, Distribution of Oil Products, and transmission and distribution of natural gas. Refinery NMVOC emissions are projected using the latest inventory year refinery NMVOC emissions and using the projected trend in CO<sub>2</sub> emissions from the refinery over the projected period as a proxy. Emissions from the distribution of oil products are estimated by calculating the latest inventory year's ratio of imported petrol/gasoline to total petrol/gasoline consumed and multiplying it by each projected year's total transport petrol/gasoline that is available from the energy forecasts for each of the scenarios (WEM and WAM), providing the future trend in imported petrol/gasoline for each year. The latest inventory emission factors and abatement efficiencies are also used to produce the projected emission estimates. NMVOC from the transmission and distribution of natural gas is projected using

the estimated total gas leaked for each year calculated from the projected primary demand of natural gas. The percentage NMVOC in natural gas is taken from gas analysis reports for the current inventory year and flatlined for the projected years. PM<sub>2.5</sub> emissions from category 1B are projected for coal storage and coal handling. Projections for coal storage are flatlined from the latest inventory year which is based on the area of coal storage and relevant emission factor that is used for the latest inventory year. Projections for coal handling is based on the projected coal used for electricity generation and applying the emission factor that is used in the latest inventory.

## 8.12 Mineral Products (NFR 2A), Chemical industry (NFR 2B), Metal Industry (NFR 2C)

Emission of air pollutants (e.g. SO<sub>2</sub> and NO<sub>x</sub> emissions) from Industrial Processes cannot usually be separated from the emissions from fuel combustion in industry. Emissions from industrial processes are therefore assumed to be included in projected estimates for the Manufacturing Industries and Construction (1A2) sector. For 2A1 Cement Production, emissions are generally reported under NFR Category 1A2f (Non-metallic minerals) and notation keys IE and NA are reported under 2A1 for the pollutants SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and NH<sub>3</sub>. Therefore, only PM<sub>2.5</sub> are projected for 2A1. PM<sub>2.5</sub> are projected using projected clinker production provided by industry experts. A similar approach is undertaken for 2A2 Lime Production.

For 2A3 Glass Production - the only container glass plant closed in 2002, one of the lead crystal plants closed in early 2006, the glass wool plant closed in 2008 and the last one, (second of the two) lead crystal plant closed in 2009. Therefore, after this period, emissions of all pollutants are projected as 'NO' (not occurring).

For 2A5a Quarrying and mining of minerals other than coal, 2A5b Construction and demolition, 2A5c Storage, handling and transport of mineral products and 2A6 Other mineral products (Bricks and ceramics and Asphalt production) do not have SO<sub>2</sub>, NO<sub>x</sub> or NH<sub>3</sub> emissions associated with them and, only particulate emissions are projected for these subcategories. PM<sub>2.5</sub> emissions are projected for 2A5a and 2A5c using estimated GDP growth rates received by ESRI. Projected emissions for 2A5b are estimated based on forecasted building completions.

The chemical industry NFR category 2B is not a dominant industry in Ireland in relation to industrial processes and is not an important source of emissions. The only source of emissions for which estimates are collated are NO<sub>x</sub> emissions from Nitric Acid Production for the years 1990-2002 after which the plant was closed therefore no emissions are reported for projected years.

For 2C Metal production the following subcategories are reported; 2C1 Iron and Steel Production in Ireland has been limited to a single large electric arc furnace installation, which closed in 2001 but was operational throughout the period 1990-2001; 2C2 Ferroalloys Production – Ireland has a single operating facility from 2014 onwards and no SO<sub>2</sub>, NO<sub>x</sub>, or NH<sub>3</sub> emissions occur from this facility; 2C3 Aluminium Production – Ireland is an important producer of alumina at one large plant using the Bayer process (extraction of AlO<sub>3</sub> using NaOH). The production of alumina using the Bayer process does not give rise to significant emissions and therefore process emissions are not estimated for this source; 2C5 Lead production – A significant quantity of Lead is mined in Ireland, but such mining is assumed not to be a significant source of emissions to air. Estimates at facility level of Lead emissions have been

obtained from AERs and are reported as NO since 2009 therefore emissions from these categories are not projected.

### 8.13 Non-Energy Products from Fuels and Non-Energy Products from Fuels and Solvent Use (NFR 2D) and NFR 2H, I, J, K, L

Emissions projections of NMVOC from solvent use and other products are estimated. NMVOC emissions for NFR 2D3b, 2D3d, 2D3e, 2D3f, 2D3g, 2D3h, 2D3i are all included in NFR 2D3a, population growth is used to project emissions out to 2030 with the 2023 inventory used as a starting point.

Projected emissions for NFR category 2H; The 6 subcategories for which emission estimates are made for Ireland are as follows with the relevant SNAP code in parentheses, where applicable: Bread (SNAP 040605), Beer (SNAP 040607), Spirits (SNAP 040608), Meat fish etc. frying/curing (SNAP 040627), Coffee Roasting, Feedstock. These activities do not have SO<sub>2</sub>, NO<sub>x</sub> or NH<sub>3</sub> emissions associated with them. Emissions projections of NMVOC are reported for this category. For subcategory Spirits emissions are projected using the whiskey industry projected growth figures out to 2032 and information from CSO on the proportion of spirit production for whiskey. The global whiskey market is projected to expand steadily during the forecast period 2024-2032 and this is reflected in a 5 per cent annual increase in spirit production for the whiskey market.

Projected emissions for NFR category 2I Wood processing, as stated in the EMEP/EEA 2019 guidebook this source category is only important for particulate emissions.

Projected emissions for NFR category 2J Production of POPs as stated in the EMEP/EEA 2019 guidebook this source category is not significant, since the contribution to the total national emissions is less than 1 per cent of the national emissions of any pollutant, these are not occurring in Ireland and projected emissions are not reported for this category.

Projected emissions for NFR category 2K Consumption of POPs and Heavy metals no emissions occur in this subcategory in Ireland and projected emissions are not reported.

Projected emissions for NFR category 2L Other production, consumption, storage, transportation or handling of bulk products category in Ireland's air pollutant inventory includes emissions of PCDD/F and PCBs from leakage from electrical equipment and emissions of PCBs from fragmentisers and shredders, therefore no SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> or PM<sub>2.5</sub> projected emissions are associated with these activities.

### 8.14 Agriculture (NFR 3)

The agriculture sector is the largest source of NH<sub>3</sub> emissions in Ireland. Projected estimates of NH<sub>3</sub> from the Agriculture sector are undertaken using the same methodological approach as the current national inventory.

Two scenarios were developed for agricultural emission projections, a WEM scenario and a WAM scenario. Projected activity data (animal numbers, crop areas and fertiliser use) are provided by Teagasc (The Irish Agriculture and Food Development Authority) to the EPA in Q1 2025 in order to prepare agricultural emission projections. This includes forecasted livestock populations, crop areas and fertilizer use. It assumes that agricultural policy continues as currently agreed and that the Brexit



related Trade and Cooperation Agreement (TCA) reached between the EU and the UK governs UK-EU trade over the projection period. Additionally, it assumes no new bilateral trade agreements are agreed and implemented between the EU, UK, and other countries. Notably the political agreement between the European Union and four Mercosur countries (Argentina, Brazil Paraguay and Uruguay) that was reached in December 2024 is not assumed to be implemented due to its, as yet, unratified status.

The majority of the data supplied to the EPA is disaggregated at the level of that used in inventory estimates.

Under WEM and WAM scenario dairy cow numbers are projected to increase reflecting the continuing profitability of dairy production in Ireland. Dairy cow numbers in 2030 reach 1.67 million. This represents a 5.9 per cent increase relative to 2023. In contrast, the continuing low levels of profitability of beef cow production systems is reflected in a projected contraction of beef cow population. Beef cow numbers in 2030 are projected to decline to 0.71 million. This represents a 15.4 per cent decrease relative to 2023. The total cattle herd in 2030 are projected to be 7.03 million. This represents a 2.5 per cent decrease relative to 2023.

Projected growth in dairy cow numbers and contraction in beef cow numbers leads to a change in the composition of the Irish bovine population and in the intensity of grassland use. Dairy production systems operate at a higher stocking rate than beef production systems and this higher stocking rate is reflected in higher projected use of nitrogen fertiliser per hectare and in total aggregate nitrogen fertiliser use by the Irish agricultural sector.

Total nitrogen fertiliser use in 2030 is projected to be 333,745 tonnes (in the absence of measures included in the WEM and WAM scenarios) an increase of 19.0 per cent on fertiliser use in 2023.

Irish ewe and total sheep numbers are projected to increase over the period to 2030. By 2030 total Irish sheep numbers are projected to increase to 6.03 million. This represents a 10.4 per cent increase relative to 2023. The projected increase in sheep production reflects the projected evolution of sheep prices on world and EU markets as global growth in the supply of meat (beef, sheep, pig and poultry meat). The total volume of pig and poultry output is projected to increase. The total crop land area is projected to contract due to the higher level of profits per hectare in grassland farming (dairying) compared to tillage. By 2030, the total cereal area harvested (barley, oats and wheat) in Ireland is projected to decrease to 229,000 hectares. This represents a 15.0 per cent decrease relative to 2023.

The WEM scenario assumes mitigation over the period 2024-2030 attributable to some of the measures that are included in the Teagasc Marginal Abatement Cost Curve<sup>16</sup>, and the DAFM Roadmap towards Climate Neutrality “AgClimate”<sup>17</sup>. Legislation levers (namely Nitrates Directive Action Plan) are in place to deliver the following key measures under the WEM scenario: low emission slurry spreading for cattle and pig slurry, and reduced nitrogen fertiliser use (due to increased liming of soils to correct soil fertility issues). The WAM scenario on the other hand includes a reduction in the crude protein content of dairy cow and pig diets, covering of uncovered slurry stores for both cattle and pigs, slurry acidification in store, reduced slaughter age and reduced age of first calving, drying of poultry

---

<sup>16</sup> <https://www.teagasc.ie/media/website/environment/climate-action/climate-centre/MACC-2023.pdf>

<sup>17</sup> <https://www.gov.ie/en/publication/07fbc-ag-climate-a-roadmap-towards-climate-neutrality/>

manure, a limit on straight urea fertiliser sales, replacement of straight urea and part replacement of CAN/NPK type fertiliser products with inhibited urea and a total fertiliser sales target of 300,000 tonnes in 2030.

Projected emissions of NMVOC and PM<sub>2.5</sub> from manure management are estimated using the same approaches and methodologies as the national inventory. Projected activity data that is utilized for NH<sub>3</sub> emission projections is also used to estimate projected emissions of NMVOC and PM<sub>2.5</sub> for the agriculture sector.

## 8.15 Waste (NFR 5)

Air pollutant emission projections in the form of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub> are estimated for the waste sector. NMVOC are estimated from landfill gas production (5A), incineration of industrial waste (5C1bi), cremation (5C1bv), and domestic wastewater handling (5D1). NO<sub>x</sub> and SO<sub>2</sub> emissions are estimated from incineration of industrial waste and cremation. NH<sub>3</sub> emissions are estimated from composting (5B1) and anaerobic digestion at biogas facilities (5B2). PM<sub>2.5</sub> emissions are estimated from incineration of industrial waste, cremation, and landfill gas production.

Solid waste disposal to landfill produces significant quantities of landfill gas. Projected landfill gas production is based on greenhouse gas emission estimates for the sector undertaken by the EPA and submitted to the European Commission under Regulation 525/2013. The emission factor utilized in the national inventory of 5.65 g NMVOC/m<sup>3</sup> landfill gas is used in projected emission estimates. Ireland has met all Landfill Directive<sup>18</sup> targets for diversion of biodegradable municipal waste from landfill to date.

The incineration of Industrial waste (5Cbi) is highly regulated in Ireland. There are currently only a small number of facilities based in the pharmaceutical and chemical sectors that operate incinerators for the treatment of industrial waste. It is assumed that the quantity of industrial waste incinerated and the emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and PM<sub>2.5</sub> at these facilities will remain constant at the 2023 level for each projected year to 2030.

The practice of Cremation (5C1bv) is less popular in Ireland than in other countries. However, due to the decrease in the number of burial plots available, particularly in larger cities and towns, the number of cremations in Ireland has increased. There are currently eight crematoria operating in Ireland. It is assumed that the number of cremations will increase for each projected year to 2030 based on historic growth levels.

---

<sup>18</sup> Council Directive 1999/31/EC on the landfill of waste

## Chapter Nine - Adjusted annual national emission inventories

### 9.1 Meeting the Requirements for an Adjustment in the period 2020 to 2030

#### 9.1.1 Emission Reduction Commitments

The amended Gothenburg Protocol and National Emission reduction Commitments Directive (Directive (EU) 2016/2284) requires parties to demonstrate compliance with emission reduction commitments (ERCs) for 2020 onwards using 2005 as a baseline on which ERCs are based. This is a different approach to the concept of emission ceilings for the period 2010 to 2019 as outlined in the Gothenburg Protocol and Directive 2001/81/EC. As a result, different considerations apply to the demonstration of compliance with emission reduction commitments.

Article 5(1) Directive (EU) 2016/2284 allows Member States to establish adjusted annual national emission inventories where non-compliance with emission ceilings or reduction commitments occur due to applying improved emission inventory methods in accordance with best science. The information provided in this chapter follows the reporting requirements of the adjustment process presented in Article 5 and Part 4 of Annex IV of the Directive (EU) 2016/2284.

Parties who wish to use adjustments to demonstrate compliance with ERCs cannot use existing inventory adjustments accepted with respect to emissions ceilings (for the period 2010-2019) to demonstrate compliance with ERCs. This is primarily for the following reasons:

- The reference version of the EMEP/EEA Guidebook (that is used to determine the scientific knowledge and understanding when the ERCs were set) is 2009 and as a result, it is likely that the validity and quantification of most adjustments will have changed.
- The calculations that are required for an adjustment with respect to ERCs require consideration, and reporting, of information relating to emissions in 2005 as well as 2020 onwards. Previously accepted adjustments do not provide this information.
- A valid adjustment under ERC might involve revising down the emissions in the compliance year in question, or it might involve revising up the emissions in 2005 to change non-compliance into compliance. However, in many cases it is likely that both years would need to be revised to capture relevant changes that impact across the entire time series, and the magnitude and direction of the revisions will determine whether the net effect would bring a Party into compliance and is therefore considered eligible for an adjustment application.
- Article 4 of Directive (EU) 2016/2284 states that emissions of NMVOCs and NO<sub>x</sub> from activities falling under the NFR sectors 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with ERCs as set out in the directive.

Table 9.1 provides details of the emission values in this submission for the years 2005 and 2023, the required emission reduction commitments under Directive (EU) 2016/2284 and the percentage reduction in emission levels for all five pollutants covered (NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, NH<sub>3</sub> and PM<sub>2.5</sub>). For the pollutants NO<sub>x</sub> and NMVOC, emission from the NFR sectors 3B and 3D are not included in the compliance assessment.

*Table 9.1 Summary of National Emissions in 2005 and 2023, the ERCs for 2020-2030 and actual emission reduction*

	2005	2023
<b>NO<sub>x</sub> (kt )*</b>	133.719	54.983
ERC (%)		-49%
Actual reduction in emissions (%)		-55.9%
<b>SO<sub>2</sub> (kt )</b>	74.253	7.360
ERC (%)		-65%
Actual reduction in emissions (%)		-90.1%
<b>NMVOC (kt)*</b>	68.467	63.595
ERC (%)		-25%
Actual reduction in emissions (%)		-17.2%
<b>NH<sub>3</sub> (kt)</b>	123.790	116.361
ERC (%)		-1%
Actual reduction in emissions (%)		-6.0%
<b>PM<sub>2.5</sub> (kt)</b>	18.646	9.310
ERC (%)		-18%
Actual reduction in emissions (%)		-50.1%

\* Emissions of NMVOCs and NO<sub>x</sub> from activities falling under the NFR sectors 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with ERCs

For NMVOC, an adjustment application accompanies this submission and is detailed in the following section.

### 9.1.2 NMVOC Adjustment Application

Ireland considers that non-compliance with the ERC for 2020 for NMVOC is as a result of the inclusion of a new source of emissions, which was not included in the national inventory when ERC targets were set.

NMVOC emissions from the food and beverage industry were not included in Ireland's national emissions inventory submission in 2012, when the ceilings were set. A methodology is presented in the 2023 EMEP/EEA Guidebook, and this is currently used to estimate emissions that are included in the Irish national emissions inventory.

For a new source, the Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria were not met, however paragraph 2 bis (a) (i) of Decision 2012/12 (as amended by ECE.EB.AIR/127/Add.1 (2014/1)), indicates that a new source is eligible for an adjustment for a new source if a methodology existed in the relevant version of the EMEP/EEA Guidebook, but the "Party can demonstrate that it was unable to apply this methodology due to a lack of relevant national statistical data...". The national data for spirit manufacture was confidential and was therefore not available to be provided to the inventory agency when the ceilings were set. It was therefore concluded that NMVOC emissions from the food and beverage industry are a new source and are eligible for an adjustment.

## Quantification

Emissions have been calculated by using the Tier 2 methodologies presented in the Inventory Guidebook (EMEP/EEA, 2023), with EFs taken from Chapter 2H2, Table 3-28.

As a new source, quantification of the adjustment is achieved by subtracting the emissions from the national inventory total as shown in Table 9.2. On the basis of the data presented in Table 9.2, NMVOC emissions are in compliance with the emission reduction commitment when the adjustment is taken into account.

**Table 9.2. Adjustments for emissions from the Food and Beverage Industry (NMVOC)**

	<b>2005</b>	<b>2023</b>
<b>NMVOC (kt)</b>	76.843	63.595
ERC (%)		-25%
Actual reduction in emissions (%)		-17.2%
Adjustment (kt)	-8.376	--23.194
<b>Adjusted NMVOC (kt)</b>	68.467	40.401
Adjusted reduction in emissions (%)		-41.0%

As described in section 8.4.4 under both the With Measures and With Additional scenarios, the emission reduction commitments for 2020-2029 and 2030 for NMVOC are not achieved (Figure 8.7) and will only be complied with if the above adjustment is applied (Figure 8.8).



## Glossary

<b>ADDF</b>	Annual Average Daily Flow
<b>AIM</b>	Animal Identification and Movement
<b>As</b>	Arsenic
<b>B[a]P</b>	Benzo[a]pyrene
<b>B[b]F</b>	Benzo[b]fluoranthene
<b>B[k]F</b>	Benzo[k]fluoranthene
<b>BCF</b>	British Coatings Federations
<b>CAP</b>	Common Agricultural Policy
<b>Cd</b>	Cadmium
<b>CEPE</b>	European Council of Producers and Importers of Paints, Printing Inks and Artists
<b>CEPMEIP</b>	Co-ordinated European Programme on Particulate Matter Emission Inventories,
<b>CLEEN</b>	Chemical Legislation European Enforcement Network
<b>CLRTAP</b>	Convention on Long Long-Range Transboundary Air Pollution
<b>CO</b>	Carbon monoxide
<b>CORINAIR</b>	Co-ordinated Information on the environment in the European Community-AIR.
<b>CMMS</b>	Cattle Movement and Monitoring Scheme
<b>Cr</b>	Chromium
<b>CSO</b>	Central Statistics Office
<b>Cu</b>	Copper
<b>DEHLG</b>	Department of Environment, Heritage and Local Government
<b>DM</b>	Dry matter
<b>DTTAS</b>	Department of Transport, Tourism and Sport
<b>DQO</b>	Data quality objective
<b>EAPA</b>	European Asphalt Pavement Association
<b>ELV</b>	End-of-Life Vehicle
<b>EMEP</b>	European Monitoring and Evaluation Programme, a co-operative programme for
<b>EPA</b>	Environmental Protection Agency
<b>ESB</b>	Electricity Supply Board
<b>ESP</b>	Electrostatic precipitators
<b>ETS</b>	Emissions Trading Scheme
<b>EUROSTAT</b>	Statistical Agency of the European Union
<b>FFS</b>	Farm Facilities Survey
<b>Fossil Fuel</b>	Peat, coal, oil and natural gas and associated derivatives
<b>FUS</b>	Fertiliser Use Survey
<b>GHG</b>	Greenhouse gas
<b>Gg</b>	Gigagram ( $10^9$ g) = kilotonne = 1,000 tonnes
<b>HCB</b>	Hexachlorobenzene
<b>HFO</b>	Heavy fuel oil
<b>Hg</b>	Mercury
<b>IBEC</b>	Irish Business and Employers' Confederation
<b>IEA</b>	International Energy Agency
<b>IEF</b>	Implied Emission Factor
<b>IFFPG</b>	Irish Farm Film Producers Group
<b>IIR</b>	Informative Inventory Report
<b>I[123-cd]P</b>	Indeno[1,2,3-cd]pyrene
<b>IPC</b>	Integrated Pollution Control

<b>IPCC</b>	Intergovernmental Panel on Climate Change
<b>IPPC</b>	Integrated Pollution Prevention and Control
<b>KDP</b>	Key Data Provider
<b>ktoe</b>	Kilotonnes of oil equivalent
<b>LCP</b>	Large Combustion Plant Directive
<b>LFG</b>	Landfill gas
<b>LTO</b>	Landing and take-off
<b>MoU</b>	Memorandum of Understanding
<b>MSW</b>	Municipal solid waste
<b>NAIS</b>	National Atmospheric Inventory System
<b>NAEI</b>	National Atmospheric Emissions Inventory
<b>NCT</b>	National Car Testing
<b>NETCEN</b>	National Environmental Technology Centre
<b>NEC</b>	National Emission Ceilings
<b>NFR</b>	Nomenclature for Reporting Codes
<b>NH<sub>3</sub></b>	Ammonia
<b>Ni</b>	Nickel
<b>NMVOC</b>	Non-methane volatile organic compound
<b>NO<sub>x</sub></b>	Nitrogen oxides
<b>NRA</b>	National Roads Authority
<b>OCLR</b>	Office of Climate, Licensing, Research and Resource Use
<b>OLG</b>	Office of Licensing and Guidance
<b>PAH</b>	Polycyclic aromatic hydrocarbon
<b>Pb</b>	Lead
<b>PCB</b>	Polychlorinated biphenyl
<b>PER</b>	Pollution Emissions Register
<b>PM</b>	Particulate matter
<b>PM<sub>10</sub></b>	Particulate matter <10 µm in diameter
<b>PM<sub>2.5</sub></b>	Particulate matter <2.5 µm in diameter
<b>POP</b>	Persistent organic pollutant
<b>PVC</b>	Polyvinyl chloride
<b>QA/QC</b>	Quality assurance/quality control
<b>S.I.</b>	Statutory Instrument
<b>Se</b>	Selenium
<b>SEAI</b>	Sustainable Energy Authority of Ireland
<b>SNAP</b>	Selected Nomenclature for Air Pollution
<b>SO<sub>2</sub></b>	Sulphur dioxide
<b>SO<sub>x</sub></b>	Sulphur oxides
<b>TAN</b>	Total ammoniacal nitrogen
<b>Teagasc</b>	Irish Agriculture and Food Development Authority
<b>TPM</b>	Total particulate matter
<b>TSP</b>	Total suspended particulates
<b>UAN</b>	Uric acid nitrogen
<b>UNECE</b>	United Nations Economic Commission for Europe
<b>VOC</b>	Volatile organic compounds
<b>WEEE</b>	Waste Electrical and Electronic Equipment Regulation
<b>WtE</b>	Waste to Energy
<b>Zn</b>	Zinc



## References

- ABFI, 2013 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2013, [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Home/\\$File/Beer+Market+Report+2013.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Home/$File/Beer+Market+Report+2013.pdf)
- ABFI, 2014 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2014, [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesABFI/Media~Newsroom~the-grape-depression-tax-take-on-a-standard-bottle-of-wine-is-now-over-50/\\$File/Irish+Beer+Market-2014.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesABFI/Media~Newsroom~the-grape-depression-tax-take-on-a-standard-bottle-of-wine-is-now-over-50/$File/Irish+Beer+Market-2014.pdf)
- ABFI, 2015 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2015, [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Publications~2015-beer-market-report/\\$File/IBA+Report-2015+web+version.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Publications~2015-beer-market-report/$File/IBA+Report-2015+web+version.pdf)
- ABFI, 2020 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2020, <https://www.ibec.ie/drinksireland/-/media/documents/drinks-ireland-new-website/beer/beer-market-report-2020.pdf>
- AEA, 2012. Provision of Technical Assistance to support the National Inventory Agency of the Office of Climate, Licensing, Research and Resource Use in the EPA. Mapping Methods for Gridded Data (AEA/R3274/ED57379).
- AEA/CTC, 2008. Inventories of Persistent Organic Pollutants in Ireland 1990 and 1995-2006. A report produced for Ireland's Environmental Protection Agency. 152 pp.
- Barry S. and O'Regan, B. (2014). FS-2 Activity data for emissions of Non-methane VOC Fellowship 2006-2012. State of Knowledge Report. Submitted to the Environmental Protection Agency, Johnstown Castle, Co. Wexford.
- Basu, M.K., Wilson, H.J. and Krishnan, G., 1991. Mercury risk from teeth. *Nature* 349, 109.
- Benezon, N., 1999. HCB emissions for Ontario 1988, 1998 and 2000. Draft report for Environment Canada, Canada.
- BiPRO (Beratungsgesellschaft für integrierte Problemlösungen), 2005. Study to facilitate the implementation of certain waste related provisions of the Regulation on Persistent Organic Pollutants (POPs). BiPRO GmbH, Munich, Germany. European Commission, Brussels. [http://ec.europa.eu/environment/waste/studies/pdf/pops\\_waste\\_full\\_report.pdf](http://ec.europa.eu/environment/waste/studies/pdf/pops_waste_full_report.pdf)
- Bus-Eireann-Annual-Report-2020 [ONLINE] Available at: <https://www.cie.ie/CIECorporate/media/reports/Bus-Eireann-Annual-Report-2020.pdf>.
- Carey, P. et al., 1996. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- CEPMEIP, 2001. CEPMEIP emission factors for particulate matter. Berdowski, J., Visschedijk, Creemers, E. and Pulles, T.
- CEU (Council of the European Union), 1999. Council Directive 1999/13/EC of 11 March on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations. OJ L85, 29 March 1999.

- Collins, C., Meaney, B., Nolan, K., Maher, H. and Murphy, D. 2004a. National Waste Database Interim Report 2002. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Collins, C., Meaney, B., Nolan, K., Maher, H. and Corish, C. 2004b. National Waste Database Interim Report 2003. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Collins, C., Le Bolloch, O. and Meaney, B. 2005. National Waste Report 2004. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Corinair, Technical annexes Volume 2, Default emission factors handbook. European Commission 1994.
- Crowe, M. et al., 2000. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- CTC/AEA, 2005. NMVOC Inventory for Ireland (SNAP Sector 6 Solvent and Other Product Use). Report to the Environmental Protection Agency
- DAF, 2003. Pesticide Usage Survey (No. 1) Grassland and Fodder Crops. Department of Agriculture and Food, Dublin, Ireland.
- DAF, 2004. Pesticide Usage Survey (No. 2) Arable Crops. Department of Agriculture and Food, Dublin, Ireland.
- DAFM, 2011. Pesticide Usage in Ireland. Vegetable Crops Survey Report, 2011. <http://www.pcs.agriculture.gov.ie/media/pesticides/content/sud/pesticidestatistics/Pesticide%20Usage%20-%202011%20Vegetable%20Survey%20Report.pdf>
- DAFM, 2012. Pesticide Usage in Ireland. Arable Crops Survey Report, 2012. <http://www.pcs.agriculture.gov.ie/media/pesticides/content/sud/pesticidestatistics/Pesticide%20Usage%20-%202012%20Arable%20Survey%20Report.pdf>
- DAFM, 2013. Pesticide Usage in Ireland. Grassland and Fodder Crops Survey Report, 2013. <http://www.pcs.agriculture.gov.ie/media/pesticides/content/sud/pesticidestatistics/Pesticide%20Usage%20-%202013%20Grassland%20&%20Fodder%20Crops%20Survey%20Report.pdf>
- DAFM, 2015. Explanatory Handbook for cross compliance requirements. <https://www.agriculture.gov.ie/media/migration/farmingschemesandpayments/crosscompliance/CrossComplianceHandbook130916.pdf>
- DAFM, 2016. Pesticide Usage in Ireland. Arable Survey Report, 2016. <https://www.pcs.agriculture.gov.ie/sud/pesticidestatistics/>
- DAFM, 2017. Pesticide Usage in Ireland. Grassland and Fodder Crops Survey Report, 2017. <https://www.pcs.agriculture.gov.ie/sud/pesticidestatistics/>
- Dammingen, U, Luttich, M., Haenel, H-D, Dohler, H., Eurich-Menden, B and Osterburg, B. (2007). Calculations of emissions from German agriculture – National Emission Inventory Report 2008 for 2006.
- Directive 2004/42/CE of the European Parliament and of the Council of 21 April 2004 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC
- DOE (Department of the Environment), 1992. The Environmental Protection Agency Act, 1992. S.I. No. 7 of 1992. Department of the Environment, Custom House, Dublin, Ireland.

- DoT (Department of Transport) 2021, Irish Bulletin of Vehicle and Driver Statistics 2020, <https://www.gov.ie/en/publication/aa05b-irish-bulletin-of-vehicle-and-driver-statistics-2020/>
- Drinks Ireland (2021), Irish Spirits Market Report 2020, Drinks Ireland Spirits, <http://drinksireland.ie/spirits>
- Duffy, P., Black, K., O'Brien, P., Hyde, B., Ryan, A.M., Ponzi, J. and Alam, S.. 2017. Ireland National Inventory Report 2017. Greenhouse Gas Emissions 1990-2015 Reported to the United Nations Framework Convention on Climate Change.
- Duffy, P., Black, K., Hyde, B., Kehoe, A., MacFarlane, B., Monaghan, S., Ponzi J., Ryan, A.M., 2024. Ireland National Inventory Report 2024. Greenhouse Gas Emissions 1990-2022 Reported to the United Nations Framework Convention on Climate Change.
- Dyke, P.H., Foan, C., Wenbom, M. and Coleman, P.I., 1997. A review of Dioxin releases to land and water in UK. Science of the Total Environment 127, 119-131.
- EAPA, European Asphalt Pavement Association (2001). Asphalt in Figures 2001
- EAPA, European Asphalt Pavement Association (2007). Asphalt in Figures 2007
- EAPA, European Asphalt Pavement Association (2012). Asphalt in Figures 2012
- EAPA, European Asphalt Pavement Association (2016). Asphalt in Figures 2016
- EAPA, European Asphalt Pavement Association (2018). Asphalt in Figures 2018, [https://eapa.org/wp-content/uploads/2020/02/Asphalt-in-figures\\_2018.pdf](https://eapa.org/wp-content/uploads/2020/02/Asphalt-in-figures_2018.pdf)
- EMEP/CORINAIR (2000). Atmospheric Emission Inventory Guidebook – Second edition 1999
- EMEP/CORINAIR (2006). Emission Inventory Guidebook
- EMEP/EEA (2009). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 9, 2009.
- EMEP/EEA (2013). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 10. 2013.
- EMEP/EEA (2016). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 21. 2016.
- EMEP/EEA (2019). Air Pollutant Emission Inventory Guidebook. EEA technical Report No.13. 2019.
- EMEP/EEA (2023). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 06. 2023.
- EP and CEU (European Parliament and Council of the European Union), 2001. Directive 2001/81/EC of 23 October 2001 on national emissions ceilings for certain atmospheric pollutants. O.J. L309, 27 November 2001.
- EP and CEU (European Parliament and Council of the European Union), 2003. Directive 2003/87/EC establishing a scheme for greenhouse gas emission allowance trading within the Community and amending Council Directive 96/61/EC. O.J. L275, 25 October 2003.
- EP and CEU (European Parliament and Council of the European Union), 2004a. Decision 280/2004/EC of the European Parliament and of the Council concerning a mechanism for monitoring Community greenhouse gas emissions and for implementing the Kyoto Protocol. O.J. L49, 19 February 2004.

- EP and CEU (European Parliament and Council of the European Union), 2004b. Directive 2004/42/CE on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC. O.J. L143, 30 April 2004.
- ERU Corinair 1990 Emissions Inventory for Ireland, July 1993.
- Feeney, B. 2015. Craft Beer and Microbreweries in Ireland, 2016, A Report for the Independent Craft Brewers of Ireland and Bord Bia,  
<http://www.bordbia.ie/industry/manufacturers/insight/publications/bbreports/RecentMarketingReports/Craft%20Beer%20and%20Microbreweries%20in%20Ireland%202016.pdf>
- Finn, J. et al., 2001. Estimation of Emissions of Non-Methane Volatile Organic Compounds from SNAP Sector 6: Solvent and Other Product Use for Ireland in 1998. Prepared for the Environmental Protection Agency.
- Gkatzoflias, D., Kourdis, C, Ntziachristos, L. and Samaras, Z., 2012. COPERT 4. Computer programme to calculate emissions from road transport. Laboratory of Applied Thermodynamics, Mechanical Engineering Department, Aristotle University of Thessaloniki, Greece.  
<http://emisias.com/products/copert/documentation>
- Hyde, B.P., Carton, O.T., O'Toole, P. and Misselbrook, T., 2003. A new inventory of ammonia emissions from Irish Agriculture. Atmospheric Environment 37, 55-62.
- Hyde, B., Carton, O.T. and Murphy, W.E., 2008. Farm Facilities Survey – Ireland 2003. Report prepared for the Department of Agriculture by Teagasc, Johnstown Castle, Wexford, Ireland. 150 p.
- IAPA, Irish Asphalt Pavement Producers Association 2019, 2020, 2021, 2022 Email correspondence from IAPA.
- IPCC (Intergovernmental Panel on Climate Change), 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories.
- Irish Spirits Association (ISA) 2015, ISA homepage of the website 2017,  
<http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesSpirits/Home!OpenDocument>
- Irish Whiskey Association 2015, Vision for Irish whiskey 2015,  
[http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesSpirits/Home/\\$File/Vision+for+Irish+Whiskey+May+2015.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesSpirits/Home/$File/Vision+for+Irish+Whiskey+May+2015.pdf)
- Kirchmann, H., and Winter, E. 1989. Ammonia volatilisation during aerobic and anaerobic manure decomposition. Plant and Soil 115. Pp. 35-41.
- Le Bolloch, O., Meaney, B., Cope, J. and Doyle, M., 2006. National Waste Report 2005: Data Update. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Le Bolloch, O., Cope, J., Meaney, B. and Kurz, I., 2007. National Waste Report 2006. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Le Bolloch, O., Cope, J., Kurz, I., Meaney, B. and Higgins, T., 2009. National Waste Report 2007. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

- Lorenz, W., Wichmann, H. and Bahadir, M., 1996. Bilanzierung der Freisetzung von polychlorierten Dibenzo-p-dioxinen und Dibenzofuranen bei Brandunfällen – ein Diskussionsbeitrag, Gefahrstoffe – Reinhaltung der Luft 56.
- McCoole, F., Derham, J., Kurz, I. and Higgins, T. 2009. National Waste Report 2008. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McCoole, F., Derham, J., Kurz, I. and McDonagh, M. 2011. National Waste Report 2009. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McCoole, F., Kurz, I., McDonagh, M., Derham, J., O'Neill, D. 2012. National Waste Report 2010. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McCoole, F., Kurz, I., McDonagh, M., Derham, J., O'Neill, D. 2013. National Waste Report 201. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Meaney, B., Collins, C., Nolan, K., Cahill, E., Delaney, J., Murray, B., Healy, J. and Carty, G. 2003. National Waste Database Report 2001, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Ministry of Health and Environmental Protection, The Netherlands. Handbook of Emission Factors, Non-Industrial Sources, 1980.
- Misselbrook, T.H. and Gilhespy, S.L. (2020). Inventory of Ammonia Emissions from UK Agriculture 2018. Rothamsted Research, North Wyke, 36 pp (unpublished).
- Misselbrook, T.H., Pain, B.F., Headon, D.M., 1998. Estimates of ammonia emission from dairy cow collecting yards. Journal of Agricultural Engineering Research 71, 127-135.
- Misselbrook, T.H., Chadwick, D.R., Chambers, B.J., Smith, K., Webb, J., Demmers, T. & Sneath, R.W. (2004). Inventory of Ammonia Emissions from UK Agriculture – 2003. Annual project report to Defra for contract AM0127
- Misselbrook, T.H., Chadwick, D.R., Gilhespy, S.L., Chambers, B.J., Smith, K.A., Williams, J. and Dragosits, U. 2010. Inventory of ammonia emissions from UK agriculture 2009. North Wyke Research, 34 pp. (Unpublished)
- Misselbrook, T.H., Gilhespy, S.L., Cardenas, L.M., Williams, J. and Dragosits, U. (2016). Inventory of Ammonia Emissions from UK Agriculture 2015. Rothamsted Research, North Wyke, 36 pp (unpublished).
- Misselbrook, T.H., Gilhespy, S.L., Carswell, A.M. and Cardenas, L.M. (2023) Inventory of Ammonia Emissions from UK Agriculture 2021, Defra Contract SCF0107, Inventory Submission Report, April 2023.
- Netcen/CTC, 2006. Ireland heavy Metal Inventories 1990, 1995 to 2004. A report Produced for Ireland's Environmental Protection Agency by Netcen, an operating division of AEA Technology PLC and Clean Technology Centre, Cork. 89 pp.
- O'Mara, F., 2007. Development of Emission Factors for the Irish Cattle Herd. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

- Owusu-Twum, M.Y., Kelleghan, D., Gleasure, G., Forrestal, P., Lanigan, G.J., Richards, K.J. and Krol, D.J. 2024. Ammonia emission factors from cattle production systems in Ireland – a review. Irish Journal of Agricultural and Food Research. Vol. 62(1):75-95. DOI: 10.15212/ijafr-2023-0108.
- Office of the Revenue Commissioners, 2019 Revenue Statistics and Economic Branch. Illegal Tobacco Products Research Surveys 2019, <https://www.revenue.ie/en/corporate/documents/research/tobacco-surveys-2019.pdf>
- Pastramas, N.; Samaras, Ch.; Mellios, G.; Ntziachristos, L. (2014). Update of the Air Emissions Inventory Guidebook - Road Transport 2014 Update.
- Plejdrup, M.S., Nielsen, O.-K. & Bruun, H.G., 2017: MapElre – A high resolution spatial emission model for Ireland.
- SEAI (Sustainable Energy Authority Ireland), 2020. Energy in Ireland 2019 – SEAI; <https://www.seai.ie/publications/Energy-in-Ireland-2020.pdf>
- SEAI (Sustainable Energy Authority Ireland), 2023. Energy in Ireland 2022 – SEAI; [Energy-in-Ireland-2023.pdf \(seai.ie\)](https://www.seai.ie/publications/Energy-in-Ireland-2023.pdf)
- Thistlethwaite, G., Baggott, S.L. and Goodwin, J., 2005. National Inventory System Scoping Report for the Greenhouse Gas Emission Inventory in Ireland. AEAT/ENV/R/1990 Issue 1, NETCEN, UK.
- UK NAEI, Crown 2018 copyright Defra & BEIS via [naei.defra.gov.uk](http://naei.defra.gov.uk), licensed under the [Open Government Licence](#) (OGL).
- UNECE, 2014. Guidance document on preventing and abating ammonia emissions from agricultural sources. United Nations Economic and Social Council Economic Commission for Europe, Executive Body for the Convention on Long-range Transboundary Air Pollution, 100 pp. [Guidance document on preventing and abating ammonia emissions from agricultural sources](#)
- UNEP Toolkit, 2013. Standardized Toolkit for Identification and Quantification of Dioxin and Furan Releases.
- URS Dames & Moore, 2000. Inventory of Dioxin and Furan Emissions to Air, Land and Water in Ireland for 2000 and 2010. Environmental Protection Agency, Johnstown Castle Estate, Wexford, Ireland.
- Winther, M, and Nielsen, Ole-Kenneth. Fuel use and emissions from non-road machinery in Denmark from 1985-2004 - and projections from 2005-2030. Denmark: N. p., 2006
- Xu, Y., 2008. Review and update of the HCB inventory of emissions to air, land and water for the UK. AEA. Group, Oxfordshire, UK.
- Zimmerman, J., 2014. A review of crop residue burning. MODIS fire detection archive for Ireland.

## Annex A to H

(click on hyperlinks below to view relevant Annex or navigate to <https://www.epa.ie/publications/monitoring--assessment/climate-change/air-emissions/irelands-unece-submissions-2025.php>)

<a href="#">Annex A</a>	A.2 Key Category Analysis 2023
<a href="#">Annex B</a>	A.3 Fuel Tourism in Road Transport and Nitrogen Oxides Emissions Based on Fuels Used
<a href="#">Annex C</a>	Expanded Energy Balance Sheet for 2023
<a href="#">Annex D</a>	Emission Factors for Energy (NFR 1)
<a href="#">Annex E</a>	Emission Factors for Industrial Processes (NFR 2)
<a href="#">Annex F</a>	Emission Factors for Agricultural Activity Data and Emission Factors (NFR 3)
<a href="#">Annex G</a>	Emission Factors for Waste (NFR 5)
<a href="#">Annex H</a>	Uncertainty Analysis
	2023 NECD Review findings and responses

## Annex H - 2023 NECD Review findings and responses

Sector/ID	NFR, Pollutant(s), Year(s)	Recommendation	Revised Estimate or Potential Technical Correction	Party response	IIR Section
IE-2D3a-2023-0001	2D3a Domestic solvent use including fungicides, NMVOC, 1990-2022	For 2D3a Domestic Solvent Use Including fungicides, NMVOC, all years the TERT notes Ireland uses the Tier 2b methodology from the 2019 EMEP/EEA Guidebook, with population statistics and NMVOC emission factors from Table 3.5 of the 2023 EMEP/EEA Guidebook. The 2023 EMEP/EEA Guidebook recommends that these emission factors be used in only specific cases, for instance if the product statistics for the use of the Tier 2b approach are not complete in terms of the product types covered by domestic solvent use. This was raised during the 2023 NECD inventory review. The TERT notes that it is preferred to use emission factors from table 3-3 or 3-4 of the 2023 EMEP/EEA Guidebook, and that the per capita emission factors from table 3-5 should only be used if product statistics for a product category are not complete. In response to a question raised during the review, Ireland explains that product statistics are not readily available in Ireland. Ireland also explained that further investigation is planned to source activity data for solvent use in Ireland and a research project is currently going through the proposal stage and if successful an update on progress will be included in the next IIR. <b>The TERT recommends that Ireland move to a higher tier method to calculate the NMVOC emissions, and that the per capita emission factors from table 3-5 of the 2023 EMEP/EEA Guidebook are only used for gap filling.</b>		ESIG data was requested in 2024 and the team is working to assess the appropriateness of using this dataset for the 2026 submission.	Section 4.5
IE-3Dc-2024-0001	3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products, PM10, 2005-2021	For the category 3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products, PM10 and years 2005-2021 the TERT notes that there are recalculations without justification. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Ireland explained that a correction was made to the Tier 2 PM10 EF for agricultural crop operations (wet climate conditions) for Harvesting Oats, which was changed from 2.3 to 3.4 kg ha <sup>-1</sup> (see table 3-6 2023 EMEP/EEA Guidebook chapter 3D). <b>The TERT recommends that Ireland include justifications for all recalculations, even when they are minor, in the next IIR submission.</b>		Justification for recalculations are presented in all IIR chapters as required	All chapters



Sector/ID	NFR, Pollutant(s), Year(s)	Recommendation	Revised Estimate or Potential Technical Correction	Party response	IIR Section
IE-5A-2022-0001	5A Biological treatment of waste - Solid waste disposal on land, PM2.5, PM10, 1990-2022	<p>The TERT notes with reference to PM2.5 and PM10 emissions from 5A Biological treatment of waste – Solid waste disposal on land for all years that there may be an underestimate of emissions. The TERT notes that this issue may be because the activity data (AD) doesn't include all amounts of inert mineral waste handled/landfilled (e.g. Industrial waste, building and demolition wastes, sand, clay and misc. fill materials). The activity data should be the total amount of waste handled and not only waste disposed in managed landfills. This was raised during the 2022 and 2023 NECD inventory review. The TERT notes that the issue is below the threshold of significance for a technical correction. In response to a question in the 2023 review, Ireland stated that the issue would be revisited in the 2024 submission. The TERT notes that no results from this revisit are found in the 2024 IIR. In response to a question, Ireland states that it hasn't been possible to get a robust dataset to implement this work during the 2024 reporting round.</p> <p><b>The TERT reiterates the recommendation that Ireland estimate PM emissions including inert waste and report the results and conclusion of its investigations in the 2025 submission.</b></p>		It hasn't been possible to get a robust dataset to implement this work during the 2025 reporting round. The issue is below the threshold of significance for a technical correction.	Table 4
IE-5B2-2019-0001	5B2 Biological treatment of waste - Anaerobic digestion at biogas facilities, NH3, 2010-2022	<p>For 5B2 Biological treatment of waste - Anaerobic digestion at biogas facilities, NH3 and years 2010-2022 the TERT notes that Ireland reports NH3 emissions from anaerobic digestion of wastes in this NFR category. However, it's unclear which processes these emissions respond to, as the IIR lacks an explanation on activity data and methodology included. This was raised during the 2019, 2020, 2021, 2022 and 2023 NECD review. The TERT notes that the issue is below the threshold of significance for a technical correction. In response to a question Ireland provided a table with the activity data used for both 5B1 Biological Treatment of Waste - Composting and 5B2 Biological treatment of waste - Anaerobic digestion at biogas facilities and stated that the IIR will be updated to better explain the approach.</p> <p><b>The TERT reiterates the recommendation that Ireland estimate and report emissions from digestion of all types of waste by collecting data on inputs to digesters not covered by EWC codes and update the explanation of the approach in the 2025 submission.</b></p>		Text has been included in Chapter 6 of the IIR	Table 4

Sector/ID	NFR, Pollutant(s), Year(s)	Recommendation	Revised Estimate or Potential Technical Correction	Party response	IIR Section
IE-5C2-2021-0001	5C2 Open burning of waste, PM10, CO, BC, 1990-2022	<p>For 5C2 Open burning of waste, PM10, CO and BC (but in fact all pollutants where Tier 1 emission factors are available) and years 1990-2022 the TERT notes that there is a lack of transparency regarding the use of 'NE' (not estimated) where a Tier 1 methodology is provided in the 2023 EMEP/EEA Guidebook. This was raised during the 2021, 2022 and 2023 NECD inventory review. In response to questions raised during the reviews, Ireland explained that there are laws and regulations that prohibit the burning of small-scale agricultural wastes and that it's illegal to burn waste or have bonfires. Ireland further explained that still some agricultural plastics are assumed to be burned, a small amount of municipal solid wastes is assumed to be burned (part of the uncollected waste), bonfires take place and some building and demolition materials are burned. However, Ireland does not report all these source emissions under 5C2 Open burning of waste, namely bonfires, domestic burning of MSW and burning of construction wastes are reported under 5E Other waste and Ireland stated that this will not be changed. The TERT notes that this is not in line with the 2023 EMEP/EEA Guidebook and relates to issues of comparability between inventories. Additionally, the TERT notes that no activity data is provided regarding the burning of plastics, bonfires, domestic burning of MSW and burning of construction (building and demolition) wastes in the IIR. The TERT notes that the issue is below the threshold of significance for a technical correction.</p> <p><b>The TERT recommends that Ireland improve the transparency of this sector by including the activity data for burning of plastics, bonfires, domestic burning of MSW and burning of construction (building and demolition) wastes in the IIR and improve the comparability by re-evaluating the allocation of these sources to 5C2 Open burning of waste in the next submission.</b></p>		<p>5C2 is the open burning of agricultural wastes.</p> <p>Therefore it is appropriate to continue to report other open burning under 5E as stated in the IIR</p>	Section 6.6
IE-5D1-2022-0002	5D1 Domestic wastewater handling, NH3, 1990-2022	<p>For 5D1 Domestic wastewater handling, NH3 and years 1990-2022 the TERT notes that there is a lack of transparency regarding the use of notation keys. This does not relate to an over- or under-estimate of emissions. This was raised during the 2022 and 2023 NECD inventory review. In response to a question raised during the review, Ireland explained that there are no latrine and/or dry toilets used during the complete time series 1990-2022 and that the notation keys will be changed to 'NO' (not occurring) in the 2025 submission of the NFR tables.</p> <p><b>The TERT reiterates the recommendation that Ireland change the notation key for NH3 from this category to 'NO'.</b></p>		The notation key used will be changed for the next submission	Section 6.5

Sector/ID	NFR, Pollutant(s), Year(s)	Recommendation	Revised Estimate or Potential Technical Correction	Party response	IIR Section
IE-5D1-2022-0001	5D1 Domestic wastewater handling, NMVOC, 1990-2022	<p>For 5D1 Domestic wastewater handling and NMVOC for all years, the TERT noted that there is a lack of transparency regarding activity data used and reported emissions. In the IIR, Ireland still reports that emissions from this source are negligible and that 'NE' (not estimated) is reported for this source. However, the NFR tables include NMVOC emission data for this subcategory (e.g 5.4 t in 2021) but there is no information about activity data and emission factor used in the IIR. This does not relate to an over- or under-estimate of emissions. This was raised during the 2022 and 2023 NECD inventory review. In response to a question raised during the 2022 review, Ireland provided the activity data, explained that the activity for 5D2 Industrial Wastewater Handling was included in 5D1 Domestic wastewater handling and stated that the text would be updated in section 6.5. The TERT notes that the text isn't updated in the 2023 IIR and no activity data was added to the 2023 NFR tables. In response to a question Ireland in the 2023 review stated that the issue would be completed in the 2024 submission. The TERT notes in the 2024 review that Ireland explains in the IIR that a Tier 1 methodology is used, but again no activity data is added to the NFR tables. In response to a question, Ireland stated that this will be completed in the next submission.</p> <p><b>The TERT reiterates the recommendation that Ireland include activity data in the next submission.</b></p>		Additional text added to IIR	Section 6.5



Headquarters  
PO Box 3000, Johnstown Castle Estate  
County Wexford, Y35 W821, Ireland  
Bosca Poist 3000, Eastát Chaisleán Bhaile Sheáin  
Contae Loch Garman, Y35 W821, Éire

T: +353 53 9160600  
F: +353 53 9160699  
E: [info@epa.ie](mailto:info@epa.ie)  
W: [www.epa.ie](http://www.epa.ie)  
Lo Call: 1890 33 55 99

EPA Regional Inspectorate Dublin  
McCumiskey House  
Richview  
Clonskeagh Road  
Dublin 14  
D14 YR62  
Tel: 01-268 0100  
Fax: 01-268 0199

EPA Regional Inspectorate Cork  
Inniscarra  
Co. Cork  
P31 VX59  
Tel: 021-4875540  
Fax: 021-4875545

EPA Regional Inspectorate Castlebar  
John Moore Road  
Castlebar  
Co. Mayo  
F23 KT91  
Tel: 094-9048400  
Fax: 094-9021934

EPA Regional Inspectorate Kilkenny  
Seville Lodge  
Callan Road  
Kilkenny  
R95 ED28  
Tel: 056-7796700  
Fax: 056-7796798

EPA Regional Inspectorate Monaghan  
The Glen  
Monaghan  
H18 YT02  
Tel: 047-77600  
Fax: 047-84987

E: [info@epa.ie](mailto:info@epa.ie)  
W: [www.epa.ie](http://www.epa.ie)  
LoCall: 1890 33 55 99



Printed on an environmentally friendly paper stock