

The Environmental Protection Agency

Ireland's Informative Inventory Report 2018

Air Pollutant Emissions
in Ireland 1990- 2016



ENVIRONMENTAL PROTECTION AGENCY

The Environmental Protection Agency (EPA) is responsible for protecting and improving the environment as a valuable asset for the people of Ireland. We are committed to protecting people and the environment from the harmful effects of radiation and pollution.

The work of the EPA can be divided into three main areas:

Regulation: *We implement effective regulation and environmental compliance systems to deliver good environmental outcomes and target those who don't comply.*

Knowledge: *We provide high quality, targeted and timely environmental data, information and assessment to inform decision making at all levels.*

Advocacy: *We work with others to advocate for a clean, productive and well protected environment and for sustainable environmental behaviour.*

Our Responsibilities

Licensing

We regulate the following activities so that they do not endanger human health or harm the environment:

- waste facilities (*e.g. landfills, incinerators, waste transfer stations*);
- large scale industrial activities (*e.g. pharmaceutical, cement manufacturing, power plants*);
- intensive agriculture (*e.g. pigs, poultry*);
- the contained use and controlled release of Genetically Modified Organisms (*GMOs*);
- sources of ionising radiation (*e.g. x-ray and radiotherapy equipment, industrial sources*);
- large petrol storage facilities;
- waste water discharges;
- dumping at sea activities.

National Environmental Enforcement

- Conducting an annual programme of audits and inspections of EPA licensed facilities.
- Overseeing local authorities' environmental protection responsibilities.
- Supervising the supply of drinking water by public water suppliers.
- Working with local authorities and other agencies to tackle environmental crime by co-ordinating a national enforcement network, targeting offenders and overseeing remediation.
- Enforcing Regulations such as Waste Electrical and Electronic Equipment (WEEE), Restriction of Hazardous Substances (RoHS) and substances that deplete the ozone layer.
- Prosecuting those who flout environmental law and damage the environment.

Water Management

- Monitoring and reporting on the quality of rivers, lakes, transitional and coastal waters of Ireland and groundwaters; measuring water levels and river flows.
- National coordination and oversight of the Water Framework Directive.
- Monitoring and reporting on Bathing Water Quality.

Monitoring, Analysing and Reporting on the Environment

- Monitoring air quality and implementing the EU Clean Air for Europe (CAFÉ) Directive.
- Independent reporting to inform decision making by national and local government (*e.g. periodic reporting on the State of Ireland's Environment and Indicator Reports*).

Regulating Ireland's Greenhouse Gas Emissions

- Preparing Ireland's greenhouse gas inventories and projections.
- Implementing the Emissions Trading Directive, for over 100 of the largest producers of carbon dioxide in Ireland.

Environmental Research and Development

- Funding environmental research to identify pressures, inform policy and provide solutions in the areas of climate, water and sustainability.

Strategic Environmental Assessment

- Assessing the impact of proposed plans and programmes on the Irish environment (*e.g. major development plans*).

Radiological Protection

- Monitoring radiation levels, assessing exposure of people in Ireland to ionising radiation.
- Assisting in developing national plans for emergencies arising from nuclear accidents.
- Monitoring developments abroad relating to nuclear installations and radiological safety.
- Providing, or overseeing the provision of, specialist radiation protection services.

Guidance, Accessible Information and Education

- Providing advice and guidance to industry and the public on environmental and radiological protection topics.
- Providing timely and easily accessible environmental information to encourage public participation in environmental decision-making (*e.g. My Local Environment, Radon Maps*).
- Advising Government on matters relating to radiological safety and emergency response.
- Developing a National Hazardous Waste Management Plan to prevent and manage hazardous waste.

Awareness Raising and Behavioural Change

- Generating greater environmental awareness and influencing positive behavioural change by supporting businesses, communities and householders to become more resource efficient.
- Promoting radon testing in homes and workplaces and encouraging remediation where necessary.

Management and structure of the EPA

The EPA is managed by a full time Board, consisting of a Director General and five Directors. The work is carried out across five Offices:

- Office of Environmental Sustainability
- Office of Environmental Enforcement
- Office of Evidence and Assessment
- Office of Radiation Protection and Environmental Monitoring
- Office of Communications and Corporate Services

The EPA is assisted by an Advisory Committee of twelve members who meet regularly to discuss issues of concern and provide advice to the Board.



IRELAND

INFORMATIVE INVENTORY REPORT 2018

AIR POLLUTANT EMISSIONS IN IRELAND 1990–2016 REPORTED TO THE SECRETARIAT OF THE UN/ECE CONVENTION ON LONG-RANGE TRANSBOUNDARY AIR POLLUTION AND TO THE EUROPEAN UNION

P. Duffy, B. Hyde, A.M. Ryan and M.S. Alam

Environmental Protection Agency
An Ghníomhaireacht um Chaomhnú Comhshaoil
PO Box 3000, Johnstown Castle, Co. Wexford. Ireland

Telephone: +353 53 9160600 Fax: +353 53 9160699
Email: info@epa.ie Web site: www.epa.ie

LoCall 1890 33 55 99

CONTENTS

List of Tables	viii
List of Figures	xi
Acknowledgements	xiii
EXECUTIVE SUMMARY	1
Chapter One Introduction	3
1.1 Convention on Long-Range Transboundary Air Pollution	3
1.2 Inventory Reporting and Review under the Convention on Long-Range Transboundary Air Pollution (CLRTAP)	3
1.3 Inventory Reporting and Review under Directive (EU) 2016/2284	4
1.4 Inventory Reporting by the Environmental Protection Agency	5
1.4.1 National Atmospheric Inventory System	5
1.4.2 Scope of Inventories under the Convention on Long-Range Transboundary Air Pollution	6
1.4.3 Inventory Preparation	9
1.4.4 Data Acquisition	10
1.4.5 Quality Assurance and Quality Control (QA/QC)	10
1.4.6 Inventory Compilation	13
1.5 Key Category Analysis	14
1.6 Uncertainty Assessment	16
Chapter Two Analysis of Key Emission Trends	18
2.1 Introduction	18
2.2 Main Pollutants	18
2.2.1 Sulphur Dioxide (SO ₂)	18
2.2.2 Nitrogen Oxides (NO _x)	19
2.2.3 Ammonia (NH ₃)	20
2.2.4 Non-Methane Volatile Organic Compounds (NMVOCs)	22
2.2.5 Carbon Monoxide (CO)	22
2.3 Particulate Matter	23
2.3.1 Particulate Matter <10 µm in Diameter (PM ₁₀)	23
2.3.2 Particulate Matter <2.5 µm in Diameter (PM _{2.5})	25
2.3.3 Total Suspended Particulates (TSP)	26
2.3.4 Black Carbon	27
2.4 Priority Metals	28
2.4.1 Lead (Pb)	28
2.4.2 Cadmium (Cd)	29
2.4.3 Mercury (Hg)	30
2.5 Other Metals	31
2.5.1 Arsenic (As)	31
2.5.2 Chromium (Cr)	32
2.5.3 Copper (Cu)	33
2.5.4 Nickel (Ni)	34
2.5.5 Selenium (Se)	35
2.5.6 Zinc (Zn)	36
2.6 Persistent Organic Pollutants (POPs)	37
2.6.1 Dioxins and Furans (PCDD/F)	37
2.6.2 Hexachlorobenzene (HCB)	38
2.6.3 Polychlorinated Biphenyls (PCBs)	39
2.6.4 Polycyclic Aromatic Hydrocarbons (PAHs)	40

Chapter Three ENERGY	42
3.1 Overview of the Energy (NFR 1) Sector	42
3.2 Public Electricity and Heat Production (NFR 1A1a)	43
3.2.1 Emissions of Sulphur Dioxide and Nitrogen Oxides	43
3.2.2 Emissions of Other Substances	45
3.3 Petroleum Refining (NFR 1A1b)	45
3.4 Manufacture of Solid Fuels and Other Energy Industries (NFR 1A1c)	45
3.5 Manufacturing Industries and Construction (NFR 1A2)	45
3.6 Transport (NFR 1A3)	46
3.6.1 Domestic and International LTOs (NFR 1A3aii (i) and 1A3ai (i))	47
3.6.2 Road Transport (NFR 1A3b)	48
3.6.3 Railways (NFR 1A3c), National Navigation (NFR 1A3d(ii)) and Other Transportation (NFR 1A3e)	50
3.7 Other Sectors (NFR 1A4)	50
3.7.1 Agriculture/Forestry/Fishing (NFR 1A4ci and ii)	50
3.7.2 Agriculture/Forestry/Fishing (NFR 1A4cii)	51
3.8 Fugitive emissions from Solid Fuels (NFR 1B1)	51
3.8.1 Coal mining and handling (1B1a)	51
3.9 Fugitive emissions from Oil and Natural Gas (NFR 1B2)	52
3.9.1 Oil (NFR 1B2a)	52
3.9.2 Venting and flaring (oil, gas, combined oil and gas) (NFR 1B2c)	53
3.10 Recalculations in the Energy Sector	53
3.11 Quality Assurance/Quality Control	63
3.12 Planned Improvements	63
3.13 Memo Items	63
Chapter Four INDUSTRIAL PROCESSES AND PRODUCT USE	65
4.1 Overview of the Industrial Processes and Product Use (NFR 2) Sector	65
4.2 Mineral Products (NFR 2A)	65
4.2.1 Cement Production (NFR 2A1)	65
4.2.2 Lime Production (NFR 2A2)	66
4.2.3 Glass Production (NFR 2A3)	67
4.2.4 Quarrying and Mining of Minerals Other than Coal (NFR 2A5a)	68
4.2.5 Construction and Demolition (NFR 2A5b)	69
4.2.6 Storage, Handling and Transport of Mineral Products (NFR 2A5c)	69
4.2.7 Other (NFR 2A6)	70
4.2.7.1 Bricks and Ceramics Production	70
4.2.7.2 Asphalt Production	70
4.3 Chemical Industry (NFR 2B)	71
4.3.1 Nitric Acid Production (NFR 2B2)	71
4.4 Metal Production (NFR 2C)	71
4.4.1 Iron and Steel Production (NFR 2C1)	72
4.4.2 Ferroalloys Production (NFR 2C2)	72
4.4.3 Aluminium Production (NFR 2C3)	74
4.4.4 Lead Production (NFR 2C5)	74
4.4.5 Other Metal Production (NFR 2C7)	74
4.5 Overview of NMVOC emissions from Solvent and Other Product Use (NFR 2D-2L) Sector	75
4.5.1 Domestic Solvent Use Including fungicides (NFR2D3a)	79
4.5.2 Road Paving with Asphalt (NFR 2D3b)	79
4.5.3 Coating Application (NFR 2D3d)	80
4.5.4 Degreasing and Dry Cleaning (NFR 2D3e and 2D3f)	83
4.5.5 Chemical Products Manufacture and Processing (NFR 2D3g)	84
4.5.6 Printing (NFR 2D3h)	88

4.5.7	Other Use of Solvents and Related Activities (NFR 2D3i).....	89
4.5.8	Food and Beverage Industry (NFR 2H2).....	91
4.6	Other production, consumption, storage, transportation or handling of bulk product (NFR 2L)	94
4.5.1	Leakage from Electrical Equipment (NFR 2L).....	94
4.5.2	Fragmentisers and Shredders (NFR 2L)	95
4.7	Recalculations in the Industrial Processes Sector	96
4.8	Quality Assurance/Quality Control	97
4.9	Overview of Notation Keys	97
4.10	Planned Improvements	98
Chapter Five AGRICULTURE.....		103
5.1	Overview of the Agriculture (NFR 3) Sector.....	103
5.2	Manure Management (NFR 3B)	106
5.2.1	Cattle (NFR 3B1).....	107
5.2.2	Sheep (NFR 3B2).....	108
5.2.3	Swine (NFR 3B3)	109
5.2.4	Poultry (NFR 3B4g).....	110
5.2.5	Other livestock – Goats (NFR 3B4d), Horses (3B4e), Mules and Asses (NFR 3B4f) and Other Animals (3B4h).....	111
5.2.6	Uncertainties	111
5.3	Agricultural Soils (NFR 3D)	112
5.3.1	Direct Soil Emissions - Inorganic N-fertilizers (NFR 3Da1)	112
5.3.2	Direct Soil Emissions -Livestock manure applied to soils (NFR 3Da2a)	112
5.3.3	Direct Soil Emissions -Sewage sludge applied to soils (NFR 3Da2b).....	112
5.3.4	Direct Soil Emissions -Other organic fertilizers applied to soils (NFR 3Da2c)	112
5.3.5	Direct Soil Emissions -Urine and dung deposited during grazing (NFR 3Da3)	112
5.3.6	Farm-level agricultural operations (NFR 3Dc).....	113
5.3.7	Off farm storage, handling and transport (NFR 3.D.d)	113
5.3.8	Cultivated crops (NFR 3De)	113
5.3.9	Use of Pesticides (NFR 3Df)	113
5.4	Field Burning of Agricultural residues (NFR 3F)	114
5.5	Uncertainties associated with Synthetic Fertilizer (3Da1), Organic fertilizers (3Da2c) and urine and dung deposited by grazing animals (3Da3)	114
5.6	Recalculations in the Agriculture Sector	114
5.7	Quality Assurance/Quality Control	115
5.8	Planned Improvements.....	115
Chapter Six WASTE.....		122
6.1	Overview of the Waste (NFR 5) Sector	122
6.2	Biological treatment of waste - Solid waste disposal on land (NFR 5A)	122
6.2.1	Main Pollutants.....	122
6.2.2	Heavy Metals	123
6.2.3	Landfill Gas and Persistent Organic Pollutants	123
6.3	Biological treatment of waste -Composting (NFR5B1)	124
6.4	Waste Incineration (NFR 5C)	124
6.4.1	Clinical Waste (5C1biii)	125
6.4.2	Industrial Waste (5C1bi), Hazardous Waste (5C1bii) and Sewage Sludge (5C1biv)	126
6.4.3	Crematoria (5C1bv).....	127
6.4.4	Open Burning of Agricultural Wastes (5C2)	128
6.5	Wastewater handling (NFR 5D)	129
6.6	Other Waste (NFR 5E)	129
6.6.1	Accidental Fires.....	129

6.6.2	Other Burning.....	130
6.7	Recalculations in the Waste Sector.....	132
6.8	Quality Assurance/Quality Control.....	132
6.9	Planned Improvements.....	132
Chapter Seven GRIDDED AND LPS DATA		136
7.1	Overview of Gridded and LPS data reporting.....	136
7.2	Mapping methodology.....	140
7.3	Mapping methods for each GNFR	144
7.3.1	Public Power (GNFR A_PublicPower)	144
7.3.2	Industry (GNFR B_ Industry).....	144
7.3.3	Other stationary Combustion (GNFR C_ OtherStationaryComb).....	144
7.3.4	Fugitive Emissions (GNFR D_ Fugitive).....	145
7.3.5	Solvents (GNFR E_Solvents).....	145
7.3.6	Road transport (GNFR F_RoadTransport).....	145
7.3.7	National navigation (GNFR G_Shipping)	145
7.3.8	Aviation (LTO) (GNFR H_Aviation)	145
7.3.9	Off Road mobile sources (GNFR I_OffRoad).....	145
7.3.10	Waste handling and treatment (GNFR J_Waste).....	146
7.3.11	Agricultural livestock (GNFR K_AgriLivestock)	146
7.3.12	Agricultural soils (Other emissions) (GNFR L_AgriOther)	146
7.3.13	Aviation (Cruise) (GNFR O_AviCruise).....	146
7.3.14	International navigation (GNFR P_IntlShipping)	147
7.4	National total emissions mapped by EMEP 0.1 degree x 0.1 degree grid emission	147
Chapter Eight PROJECTIONS		153
8.1	Overview of Emissions Projections	153
8.2	Emission Reduction Targets for 2020 and 2030.....	154
8.3	With Existing Measures and With Additional Measures Scenarios.....	154
8.4	Key Trends.....	156
8.4.1	Sulphur Dioxide (SO ₂)	156
8.4.2	Nitrogen Oxides (NO _x)	158
8.4.3	Ammonia (NH ₃)	160
8.4.4	Non-Methane Volatile Organic Compounds (NMVOCs)	162
8.4.5	Particulate Matter < 2.5 µm in diameter (PM _{2.5})	164
8.5	Energy Industries (1A1).....	166
8.5.1	Oil Refining and Solid Fuel Manufacturing (1A1b, 1A1c)	167
8.6	Manufacturing Industries and Construction (1A2).....	168
8.7	Transport (1A3)	168
8.7.1	Domestic aviation (1A3a)	169
8.7.2	Road transportation (1A3b)	169
8.7.3	Rail (1A3c)	169
8.7.4	Navigation (1A3d)	169
8.7.5	Gas Transmission (1A3e).....	170
8.8	Residential and Commercial/Institutional (1A4).....	170
8.8.1	Commercial/Institutional (1A4a)	170
8.8.2	Residential (1A4b).....	170
8.9	Combustion in Agriculture and Fishing (1A4c).....	171
8.10	Mineral products (NFR 2A).....	171
8.11	Non-Energy Products from Fuels and Solvent Use (NFR 2D).....	171
8.12	Agriculture (NFR 3)	172
8.13	Waste (NFR 5)	172

Chapter Nine ADJUSTED ANNUAL NATIONAL EMISSION INVENTORIES.....	174
9.1 Introduction	174
9.2 Meeting the requirements for an Adjustment.....	178
9.2.1 New Emission Sources.....	178
9.2.2 Significantly Different Emission Factors	178
9.3 NOx Adjustment Applications.....	179
9.3.1 1A3b Road Transport (NO _x)	179
9.3.2 1A3c Railways (NO _x).....	186
9.3.3 1A3dii National Navigation (NO _x)	186
9.3.4 1A4ai Commercial/Institutional Stationary Combustion (NO _x).....	187
9.3.5 1A4bi Residential Stationary Combustion (NO _x)	188
9.3.6 1A4ciii National Fishing (NO _x)	191
9.3.7 3B1a-3B4h Manure Management (NO _x)	192
9.3.8 3D Fertilizers applied to soils (NO _x)	192
9.4 NMVOC Adjustment Applications.....	193
9.4.1 Food and Beverage industry, 2H2; Spirits (NMVOC)	193
9.4.2 Manure Management, 3B (NMVOC)	194
9.4.3 Inorganic N Fertilisers, 3D1a (NMVOC)	195
9.5 Impact of flexibility on compliance.....	195
References	197
Glossary	202
Annex A	205
Annex A.1 <i>Annex 1 Table 2016.....</i>	206
Annex A.2 <i>Key Category Analysis 2016.....</i>	210
Annex A.3 <i>Fuel Tourism in Road Transport and Nitrogen Oxides Emissions Based on Fuels Used</i>	227
Annex B <i>Expanded Energy Balance Sheets for 2016</i>	231
Annex C <i>Emission Factors for Energy (NFR 1)</i>	233
Annex D <i>Emission Factors for Industrial Processes (NFR 2)</i>	271
Annex E <i>Agricultural Activity Data and Emission Factors (NFR 3).....</i>	280
Annex F <i>Emission Factors for Waste (NFR 5)</i>	294
Annex G <i>Uncertainty Analysis</i>	298

List of Tables

Table	Page
1.1 Key Data Providers and Information Covered by Memoranda of Understanding	12
1.2 Key Category Analysis of Ireland's Air Pollutant Inventory 2016	14
1.3 Emissions Uncertainties	15
3.1 Fugitive Emissions from Coal mining and handling	48
3.2 Fugitive Non-Methane Volatile Organic Compound Emissions	48
3.3 Recalculations in Energy NFR 1A1	50
3.4 Recalculations in Energy NFR 1A2	52
3.5 Recalculations in Energy NFR 1A3 (a & b)	53
3.6 Recalculations in Energy NFR 1A3 (c, d & e)	54
3.7 Recalculations in Energy NFR 1A4	55
4.1 Emission Time Series for Dioxins and Furans and Polychlorinated Biphenyls from Cement Production	60
4.2 Emission Time Series from Glass Production	62
4.3 Quarrying and mining of minerals other than coal (2A5a)	62
4.4 Construction and Demolition emissions (2A5b)	
4.5 Storage, handling and transport of mineral products emissions (2A5c)	
4.6 Dioxin and Furan Emission Time Series from Bricks and Ceramics Production	63
4.7 Dioxin and Furan Emission Time Series from Asphalt Production	63
4.8 Nitrogen Oxides Emission Time Series from Nitric Acid Production	64
4.9 Emission Time Series from Iron and Steel Production	65
4.10 Emission Time Series from Ferroalloys Production	66
4.11 Emission Time Series for Zinc from Aluminium Production	66
4.12 Emission Time Series for Lead Production	67
4.13 Emission Time Series for Non-Ferrous Metal Production	67
4.14 NMVOC Emissions from Domestic solvent use	68
4.15 Emissions from Road Paving with Asphalt (NFR 2D3b)	69
4.16 NMVOC emissions from Paint Application; Car Repairing	70
4.17 NMVOC emissions from Paint Application; Construction and Building and Domestic Use	71
4.18 NMVOC Emissions from Paint application; Boat Building	
4.19 NMVOC Emissions from Paint application; Wood	
4.20 NMVOC Emissions from Paint application; Other industrial sources	73
4.21 NMVOC Emissions from Paint Application; Other Non-Industrial sources	74
4.22 NMVOC Emissions from Metal Degreasing, Electronics Manufacture and other Industrial cleaning	75
4.23 NMVOC emissions from Dry Cleaning	
4.24 NMVOC Emissions from Polyurethane Processing	75

4.25	NMVOC Emissions from Rubber Processing	75
4.26	NMVOC Emissions from Pharmaceutical Products Manufacturing	76
4.27	NMVOC Emissions from Paint Manufacture	77
4.28	NMVOC Emissions from Inks Manufacturing	77
4.29	NMVOC Emissions from Adhesives Manufacturing	78
4.30	NMVOC Emissions from Textile Finishing	78
4.31	NMVOC Emissions from Printing	79
4.32	NMVOC Emissions from Glass Wool Blowing/Enduction	80
4.33	NMVOC Emissions from Fat, edible and non-edible oil extraction	81
4.34	NMVOC Emissions from Application of Glues and Adhesives	82
4.35	NMVOC Emissions from Preservation of Wood	83
4.36	NMVOC Emissions from Underseal Treatment and Conservation of Vehicles	84
4.37	NMVOC Emissions from Vehicle Dewaxing	84
4.38	Emissions from Use of Tobacco	85
4.39	Emissions from Use of Fireworks	86
4.40	NMVOC Emissions from Food and Beverage industry; Bread	87
4.41	NMVOC Emissions from Food and Beverage industry; Beer	88
4.42	NMVOC Emissions from Food and Beverage industry; Spirits	89
4.43	NMVOC Emissions from Food and Beverage industry; Meat, fish etc. frying/curing	89
4.44	NMVOC Emissions from Food and Beverage industry; Coffee roasting	90
4.45	NMVOC Emissions from Food and Beverage industry; Feedstock	90
4.46	Time Series of Polychlorinated Biphenyls and Dioxin and Furan Emissions from Leakage from Electrical Equipment	69
4.47	Emission Time Series for Polychlorinated Biphenyls from Fragmentisers and Shredders	70
4.48	Notation keys used in the 2016 NMVOC inventory	99
4.49	Recalculations for Industrial Processes 1990–2015 (Heavy Metals)	86
4.50	Recalculations for Industrial Processes 1990–2015 (Particulates)	88
4.51	Recalculations for Industrial Processes 1990–2015 (NMVOC)	110
5.1	Emission Estimates for Hexachlorobenzene Emissions from Pesticide Use	99
5.2	NH ₃ Recalculations for Agriculture 1990–2015	101
5.3	NMVOC Recalculations for Agriculture 1990–2015	102
5.4	TSP Recalculations for Agriculture 1990–2015	103
5.5	PM ₁₀ Recalculations for Agriculture 1990–2015	104
5.6	PM _{2.5} Recalculations for Agriculture 1990–2015	105
6.1	Pollutant Emissions by Waste Source Category	107
6.2	Emission Time Series for NMVOC, TSP, PM ₁₀ and PM _{2.5} from Solid Waste Disposal on Land	108
6.3	Emission Time Series for Mercury from Solid Waste Disposal on Land	108
6.4	Time Series of Activity Data and Emissions of Dioxins and Furans and Polychlorinated Biphenyls from	

	Landfill Gas	108
6.5	Time Series of Activity Data and Emissions of NH ₃ and CO from Composting	109
6.6	Time Series of Emissions from the Incineration of Clinical Waste	110
6.7	Time Series of Emissions from the Incineration of Industrial (incl. Hazardous & Sludge) Waste	111
6.8	Time Series of Emissions from Crematoria	113
6.9	Time Series of Emissions from the Open Burning of Farm Plastics	114
6.10	Time Series of Emissions from Accidental Fires from Vehicles and Buildings	115
6.11	Time Series of Emissions from Other Waste Burning	116
6.12	Recalculations for Waste 1990–2015	118
7.1	List of GNFR categories	122
7.2	Correspondence list for GNFR and NFR categories	122
8.1	Emission Reduction Commitments for 2020 and proposed reduction commitments for 2030 (expressed as a percentage reduction of 2005 levels)	138
8.2	Key assumptions underpinning the energy forecasts	139
8.3	Projected Emissions of SO ₂ under the With Measures and With Additional Measures Scenarios (kt)	141
8.4	Projected Emissions of NO _x under the With Measures and With Additional Measures Scenarios (kt)	143
8.5	Projected Emissions of NH ₃ under the With Measures and With Additional Measures Scenarios (kt)	145
8.6	Projected Emissions of NMVOC under the With Measures and With Additional Measures Scenarios (kt)	147
8.7	Projected Emissions of PM _{2.5} under the With Measures and With Additional Measures Scenarios (kt)	149
9.1	Summary of National Emissions and the NECD Emission Ceilings	157
9.2a	Summary of NO _x Adjustments	158
9.2b	Summary of NMVOC Adjustments	159
9.3	Adjustment Quantification for Road Transport-1A3bi Passenger Cars (NO _x)	160
9.4	Adjustment Quantification for Road Transport-1A3bii Light Duty Vehicles (NO _x)	160
9.5	Adjustment Quantification for Road Transport-1A3biii Heavy Duty Vehicles (NO _x)	160
9.6	Adjustment Quantification for Road Transport-1A3biii Heavy Duty Vehicles, Buses (NO _x)	161
9.7	Adjustment Quantification for Road Transport-1A3biv Mopeds and Motorcycles (NO _x)	161
9.8	Adjustment Quantification for Road Transport-1A3b Road Transport (NO _x)	161
9.9	Adjustment Quantification for Railways (NO _x)	162
9.10	Adjustment Quantification for National Navigation (NO _x)	163
9.11	Adjustment Quantification for Commercial/Institutional Stationary Combustion (NO _x)	163
9.12	Adjustment Quantification for Residential Stationary Combustion (NO _x)	164
9.13	Adjustment Quantification for National Fishing (NO _x)	165
9.14	Adjustment Quantification for Manure management (NO _x)	166
9.15	Adjustment Quantification for Fertilizer applied to soils (NO _x)	166
9.16	Adjustments for emissions from Food and Beverage Industry (NMVOC)	167
9.17	Adjustments from Manure Management (NMVOC)	167
9.18	Adjustment from Inorganic N Fertilisers (NMVOC)	167

List of Figures

Figure	Page
1.1 National Atmospheric Inventory System Overview	7
2.1 Emission Trend for Sulphur Dioxide 1990–2016	18
2.2 Emission Trend for Nitrogen Oxides 1990–2016	19
2.3 Emission Trend for Ammonia 1990–2016	20
2.4 Emission Trend for Non-Methane Volatile Organic Compounds 1990–2016	21
2.5 Emission Trend for Carbon Monoxide 1990–2016	22
2.6 Emission Trend for Particulate Matter <10 µm in Diameter 1990–2016	23
2.7 Emission Trend for Particulate Matter <2.5 µm in Diameter 1990–2016	24
2.8 Emission Trend for Total Suspended Particulates 1990–2016	25
2.9 Emission Trend for Black Carbon 1990–2016	26
2.10 Emission Trend for Lead 1990–2016	26
2.11 Emission Trend for Cadmium 1990–2016	27
2.12 Emission Trend for Mercury 1990–2016	28
2.13 Emission Trend for Arsenic 1990–2016	29
2.14 Emission Trend for Chromium 1990–2016	30
2.15 Emission Trend for Copper 1990–2016	31
2.16 Emission Trend for Nickel 1990–2016	32
2.17 Emission Trend for Selenium 1990–2016	33
2.18 Emission Trend for Zinc 1990–2016	34
2.19 Emission Trend for Dioxins and Furans 1990–2016	35
2.20 Emission Trend for Hexachlorobenzene 1990–2016	36
2.21 Emission Trend for Polychlorinated Biphenyls 1990–2016	37
2.22 Emission Trend for Polycyclic Aromatic Hydrocarbons 1990–2016	38
3.1 Nitrogen Oxide Implied Emission Factors for Category 1A1a	41
3.2 Sulphur Dioxide Implied Emission Factors for Category 1A1a	41
3.3 Number of Domestic LTOs from Irish airports 1990–2016	44
3.4 Number of International LTOs from Irish airports	45
4.1 Non-Methane Volatile Organic Compounds Emission Trend for 2D-2H Other Solvent and Product Use 1990–2016	75
5.1 Emission Trend for Ammonia from Agriculture 1990–2016	92
5.2 Emission Trend for NMVOC from Agriculture 1990–2016	92
7.1 General decision tree for emissions mapping	126
7.2 Example of Geokey preparation (a,b,c,d)	127
7.4(a) National total Emissions NO _x 2015	132
7.4(b) National Total Emissions SO ₂ 2015	133

7.4(c.)	National Total Emissions NMVOC 2015	134
7.4(d)	National Total Emissions NH ₃ 2015	135
7.4(e.)	National Total Emissions PM _{2.5} 2015	136
8.1	SO ₂ Emissions Projections for the With Measures and With Additional Measures Scenarios	140
8.2	SO ₂ Emissions Projections for the With Additional Measures Scenario by Source Sector	141
8.3	NO _x Emissions Projections for the With Measures and With Additional Measures Scenarios	142
8.4	NO _x Emissions Projections for the With Additional Measures Scenario by Source Sector	143
8.5	NH ₃ Emissions Projections for the With Measures and With Additional Measures Scenarios	144
8.6	NH ₃ Emissions Projections for the With Additional Measures Scenario by Source Sector	145
8.7	NMVOC Emissions Projections for the With Measures and With Additional Measures Scenarios	146
8.8	NMVOC Emission Projections for the WAM Scenario by Source Sector	147
8.9	PM _{2.5} Emissions Projections for the With Measures and With Additional Measures Scenarios	148
8.10	PM _{2.5} Emission Projections for the WAM Scenario by Source Sector	149

Acknowledgements

The authors wish to express their appreciation to the various Government Departments, the Sustainable Energy Authority of Ireland, the Central Statistics Office, Aether (UK), the Electricity Supply Board and the many other data suppliers and individuals who have contributed to the subject matter of this inventory report.

EXECUTIVE SUMMARY

As a Party to the United Nations Economic Commission for Europe (UN/ECE) Convention on Long-Range Transboundary Air Pollution (CLRTAP), Ireland is required to annually report emission data for a wide range of air pollutants and other substances released into the atmosphere. The data are needed to support the work of the Convention in addressing well-known environmental problems such as urban pollution, acidification and tropospheric ozone formation arising from classic pollutants, such as nitrogen oxides (NO_x), sulphur dioxide (SO₂), non-methane volatile organic compounds (NMVOCs), ammonia (NH₃), carbon monoxide (CO) and particulate matter (PM), and for the implementation of its Protocols on Heavy Metals and Persistent Organic Pollutants.

The UN/ECE revised 2014 Reporting Guidelines, Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution¹, describe the scope and reporting of the emission inventories and projections under the Convention. They specify the methodologies and procedures to be followed for submitting consistent and comparable data on an annual basis in a timely, efficient and transparent manner to meet the needs of the Convention. Under the Guidelines, Parties are strongly encouraged to submit an Informative Inventory Report (IIR) to support the evaluation of their up-to-date annual inventories and projections. The objective of the IIR is to describe the methodologies, input data, background information and the entire process of inventory compilation for transboundary air pollutant emissions and to give explanations for any improvements and recalculations of the inventories reported in previous submissions. The report is needed by expert review teams to assess the transparency, completeness and overall quality of the inventories as part of the review process being developed for the submissions from Parties to the Convention.

Member States of the European Union are required to report an Informative Inventory Report annually under Article 8(3) of Directive [\(EU\) 2016/2284](#). This Directive sets out reduction commitments of certain atmospheric pollutants and repeals Directive [2001/81/EC](#). This report to the European Union fulfils this reporting obligation.

The Environmental Protection Agency (EPA) in Ireland has overall responsibility for national air emission inventories and projections pursuant to the establishment of the National Atmospheric Inventory System (NAIS) in 2007. The EPA Office of Environmental Sustainability (OES) performs the role of inventory agency in Ireland and undertakes all aspects of inventory preparation and management and is responsible for the submission of the results to CLRTAP. The present report constitutes Ireland's ninth IIR submitted under the Convention, covering annual inventories for the period 1990–2016. The report aims to provide a comprehensive description of the procedures, methodologies and activity data used for the compilation of Ireland's air emission inventories and projections as presented in Ireland's 2018 submission under CLRTAP and to the European Union under Directive (EU) 2016/2284. The report shows how Ireland follows the guidelines for estimating and reporting of emission data to ensure the transparency, accuracy, consistency, comparability and completeness of the reported emissions. In addition to complying with reporting requirements in this regard, the 2018 IIR is intended to inform the Government Departments and institutions involved, as well as other stakeholders in Ireland, of the level of emissions

¹ [Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution](#)

and the state of the art of Ireland's emission inventories and projections as they address the challenges to comply with commitments already established for air pollutants and to control emissions in general. An attempt has been made to give adequate descriptions of all methodological approaches and to provide pertinent information to facilitate the assessment of the emission estimates and the understanding of emission trends. The IIR is published on the web site of the EPA (<http://erc.epa.ie/clrtap>). It will be further developed for future submissions and updated annually in accordance with the UN/ECE Reporting Guidelines and Directive (EU) 2016/2284.

Chapter One

Introduction

1.1 Convention on Long-Range Transboundary Air Pollution

The Convention on Long-Range Transboundary Air Pollution (CLRTAP) came into being in 1979 following the recognition that co-operation at international level was necessary to address environmental problems such as acidification associated with the transboundary transport and deposition of acidifying gases emitted into the atmosphere. The Convention was the first international legally binding instrument to deal with problems of air pollution on a broad regional basis. Besides laying down the general principles of international co-operation for air pollution abatement, the Convention sets up an institutional framework bringing together research and policy. The Executive Secretary of the United Nations Economic Commission for Europe (UN/ECE) acts as Secretariat to CLRTAP, and the Convention entered into force in 1983.

The aim of the Convention is that Parties shall endeavour to limit and, as far as possible, gradually reduce and prevent air pollution, including long-range transboundary air pollution. This objective is pursued under eight protocols that identify specific measures to be taken by Parties to cut their emissions of a wide range of air pollutants. The extent to which Parties to the Convention have ratified the various protocols varies. Of the eight protocols to date, Ireland has ratified the 1994 Oslo Protocol on Further Reduction of Sulphur Emissions and the 1988 Sofia Protocol concerning the Control of Emissions of Nitrogen Oxides.

1.2 Inventory Reporting and Review under the Convention on Long-Range Transboundary Air Pollution (CLRTAP)

The Guidelines for Reporting Emissions and Projections Data under the Convention on Long-Range Transboundary Air Pollution (ECE/EB.AIR/125)² specify the scope, methodologies, formats and deadlines for annual inventory submissions by Parties to the Convention. These Guidelines were adopted by the Executive Body in December 2013 (ECE/EB.AIR/122/Add.1, decisions 2013/3 and 2013/4) and published in 2014. They are a revised version of the 2009 Guidelines for Reporting Emission data under the Convention (ECE/EB.AIR/97), which were approved by the Executive Body in 2008 (ECE/EB.AIR/96, para. 83 (b)). While the Guidelines make it clear that Parties are required to report only on the substances and for the years set forth in the protocols that they have ratified and that have entered into force, Ireland endeavours to estimate and report emissions for the full range of substances set down in Annex I of the Guidelines. These substances are nitrogen oxides (NO_x), sulphur oxides (SO_x), non-methane volatile organic compounds (NMVOCs), carbon monoxide (CO), ammonia (NH₃), particulate matter (PM), black carbon (BC), heavy metals (HM) and persistent organic pollutants (POPs).

² The Guidelines for Reporting Emissions and Projections Data under the Convention on Long-Range Transboundary Air Pollution ([ECE/EB.AIR/125](https://www.ece.org/filearchive/doc/02/02_125.pdf))

The Guidelines state that an Informative Inventory Report (IIR) should be prepared for inclusion in the annual submission and is required under Article 8(3) of Directive (EU) 2016/2284. The objective of the IIR is to describe the methodologies, input data, background information and the entire process of inventory compilation for air pollutants, as well as any improvements and recalculations of the inventories reported in previous submissions. The report is needed to support the evaluation of emission trends and may be used by expert review teams to assess the transparency, completeness and overall quality of the inventories as part of the review process established for submissions by the Parties to CLRTAP and the review process to be established under the NECD.

The present report constitutes Ireland's IIR for 2018, the ninth such report. It contains specific information on the national inventory for the years 1990–2016 as submitted to CLRTAP in February 2018, including descriptions of methods, data sources, quality assurance/quality control (QA/QC) activities and trend analysis. The structure of the report follows the structure proposed in Annex II to the reporting guidelines.

The IIR focuses on the year 2016 and the status of the inventories achieved for the time series up to 2016 and is the basis for methodological description for the purposes of facilitating technical review and general assessment of Ireland's emission inventories. The IIR is designed to capture the cyclical nature of the reporting process and to clarify the chronology of changes and revisions that are part of normal inventory development. In this way, the report provides the basis for technical assessment and expert review of Ireland's air pollutant inventories. An attempt has been made to give adequate descriptions of all methodological approaches and to provide all the pertinent inventory information to facilitate the assessment of the emission estimates and the understanding of emission trends.

The IIR will be further developed and updated annually in accordance with the UN/ECE guidelines and is published on the web site of the EPA (<http://erc.epa.ie/clrtap>). Such updating is necessary to keep the UN/ECE Secretariat and other interested parties informed of the status of Ireland's air pollutant inventories and to document on-going improvements, recalculations and other developments affecting the estimates of emissions. Ireland is contributing to the Stage 3 in-depth review process for transboundary emission inventories by not only providing this IIR, but also providing expert reviewers to evaluate the submissions from other Parties to the Convention.

1.3 Inventory Reporting and Review under Directive (EU) 2016/2284

Directive (EU) 2016/2284³ came into force on the 31st of December 2016. This inventory report and data submission fulfils Ireland's reporting requirements as set out in Article 10(2) of the directive. The inventory submission is fully consistent with the submission under the LRTAP Convention and will be subject to review in accordance with Article 10(3) of the directive.

Ireland's national total emissions of nitrogen oxides (NO_x), ammonia (NH₃) and non-methane volatile organic compounds (NMVOCs) are not in compliance with the ceilings in Article 4(1) and Annex I to Directive 2001/81/EC⁴. Ireland has established adjusted annual national emission inventories for NO_x and NMVOCs as allowed for under the flexibilities outlined in Articles 21(2) and 5(1) of the Directive 2016/2284. The adjusted inventories are established

³ [DIRECTIVE \(EU\) 2016/2284 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC](#)

⁴ [DIRECTIVE 2001/81/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 23 October 2001 on national emission ceilings for certain atmospheric pollutants](#)

in accordance with Part 4 of Annex IV of the directive and are explained in detail in chapter 9 of this report.

1.4 Inventory Reporting by the Environmental Protection Agency

Under Section 52 of the Environmental Protection Agency Act, 1992 (DOE, 1992), the Agency is required to establish and maintain databases of information on the environment and to disseminate such information to interested parties. Section 55 of the Act states that the Agency must provide, of its own volition or upon request, information and advice to Ministers of the Government in the performance of their duties. This includes making available such data and materials as are necessary to comply with Ireland's reporting obligations and commitments within the framework of international agreements. These requirements are the regulatory basis on which the EPA prepares annual inventories of air pollutants in Ireland. The activities related to the compilation and reporting of air pollutant emissions constitute one specific on-going project in the Agency's work programme.

The Department of Communications, Climate Action and Environment (DCCAE) has designated the EPA as the agency with responsibility for the submission of emission data to international bodies, including the Secretariat for CLRTAP and the European Union. The Agency's Office of Environmental Sustainability (OES) currently compiles the national air pollutant emission inventories on behalf of the DCCAE for submission under CLRTAP and NECD.

1.4.1 National Atmospheric Inventory System

In 2005, UK consultants from the National Environmental Technology Centre (NETCEN) carried out a scoping study to identify the essential elements and structure of a national inventory system for Ireland to meet the needs of Decision 280/2004/EC (EP and CEU, 2004a) and to comply with obligations under Articles 5 and 7 of the Kyoto Protocol. The report (Thistlethwaite et al., 2005) describes how institutional arrangements among the EPA, its parent government department and other stakeholders may be reorganised, extended and legally consolidated across all participating institutions to strengthen inventory capacity within the EPA and to ensure that more formal and comprehensive mechanisms of data collection and processing are established for long-term implementation. The report sets out the extent of institutional participation, resource requirements and the form of legal arrangements necessary to perform the functions prescribed in the guidelines for national systems and enable Ireland to meet the objectives specified in those guidelines. The scoping study developed a QA/QC system as an integral part of the national system and the report made recommendations on internal inventory review and proposed a database system to facilitate more efficient data management and reporting. Whilst developed to meet the needs of Decision 280/2004/EC and the Kyoto Protocol, Ireland's national system is also implemented to achieve emission inventories for transboundary gases for submission under CLRTAP and NECD.

The National Atmospheric Inventory System (NAIS) for Ireland was adopted by Government decision in April 2007. It establishes the necessary institutional, legal and procedural arrangements for the compilation of robust inventories of emissions of greenhouse gases (GHGs) and air pollutants to the atmosphere. It sets out formal procedures for the planning, preparation and management of the national atmospheric inventory and identifies clearly the roles and responsibilities of all the organisations involved in inventory compilation, reporting and review. A schematic overview of the national system is presented in Figure 1.1.

The principal objective of the NAIS is to ensure that Ireland can compile robust and verifiable annual inventories of emissions and report its emission estimates in accordance with relevant international obligations. The NAIS also facilitates the formal review of information submitted under international obligations, including the Kyoto Protocol, protocols under

CLRTAP and the NECD. Implementation of the national system ensures the transparency, consistency, comparability, completeness and accuracy of the national inventory in accordance with the established reporting guidelines, which incorporate methodological guidance and good practice.

Within the NAIS, the EPA's OES is designated as the single national entity with overall responsibility for the national emission inventory in Ireland. The OES also performs the role of inventory agency, i.e. the Office compiles the annual inventory and delivers Ireland's submissions to the various international organisations (European Commission (EC), European Environment Agency (EEA), United Nations Framework Convention on Climate Change (UNFCCC) and UN/ECE in accordance with agreed deadlines and reporting formats. In addition to the primary data received from the Key Data Providers (KDPs), the inventory team obtains considerable supplementary information from other teams in the OES, the Office of Evidence and Assessment and the Office of Environmental Enforcement within the EPA. These sources include Annual Environmental Reports (AERs) submitted by licensed facilities and the National Waste Database. The inventory team also draws on national research related to air pollutant emissions and special studies undertaken from time to time to acquire the information needed to improve the estimates for particular categories and gases. The approval of the completed annual inventory involves sign-off by the QA/QC manager and the inventory manager before it is transmitted to the Board of Directors of the EPA via the Programme Manager of OES. Any issues arising from the Board of Directors' examination of the estimates are communicated to the inventory experts for resolution before final adoption of the inventory for submission and publication.

1.4.2 Scope of Inventories under the LRTAP Convention and Directive (EU) 2016/2284

The scope of Ireland's emission inventories under LRTAP Convention and Directive (EU) 2016/2284 is provided in Annex A1. It covers a wide range of air pollutants and other substances, which are reported in a standard electronic format for a predefined nomenclature of source categories set down in the UN/ECE Reporting Guidelines. The air pollutants are referred to in seven groups as follows:

1. Main pollutants (NO_x, NMVOCs, SO_x, NH₃, CO);
2. Particulate matter (PM_{2.5}, PM₁₀, total suspended particulates (TSP), black carbon (BC));
3. Priority heavy metals (lead (Pb), cadmium (Cd), mercury (Hg));
4. Other heavy metals (arsenic (As), chromium (Cr), copper (Cu), nickel (Ni), selenium (Se), zinc (Zn));
5. Annex I Persistent Organic Pollutants: the POPs listed in Annex I to the Protocol on POPs which are substances scheduled for elimination;
6. Annex II Persistent Organic Pollutants: the POPs listed in Annex II to the Protocol on POPs which are substances scheduled for restrictions on use;
7. Annex III Persistent Organic Pollutants: the POPs listed in Annex III to the Protocol on POPs which are substances referred to in Article 3, Para. 5 (a), of the Protocol. Polycyclic aromatic hydrocarbons (PAHs): for the purpose of the emission inventories, the following four indicator compounds should be used: benzo[b]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene and indeno [1,2,3-cd] pyrene. Hexachlorobenzene (HCB) is also included in Annex I to the Protocol as a substance for elimination.

The list of source categories for inventory purposes is known as the NFR (Nomenclature for Reporting). It comprises coded activities across all socio-economic sectors identified as sources of one or more of the substances listed above and provides for the inclusion of other activities that may be specific to individual countries. Many of the NFR categories are split into a varying number of subcategories, which are designed to reflect their importance as sources of one or more pollutants and to provide an adequate level of transparency. In the compilation of annual inventories, significant subdivision of the given NFR categories is normally applied for the process of calculating the relevant emissions. The NFR facilitates the comparison of emissions among reporting countries and the synthesis and assessment of submissions at the UN/ECE level. The current version of the NFR, NFR 14, is given in Annex I of the Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution (ECE/EB.AIR/125), which were adopted by the Executive Body in December 2013.

The reporting format also includes a number of *Memo Item* entries. These items refer to sources of emissions whose contributions are not included in a Party's national total but which are to be reported because of their importance in relation to the overall assessment of emissions and for comparisons among Parties. The notable emission sources excluded from the reported national total for transboundary gases and included as Memo Items are emissions from international and domestic aviation during the cruise phase of a flight, and international shipping.



A set of notation keys has been adopted for use in completing the NFR templates to provide explanation and transparency where a numerical value does not appear for a particular pollutant and/or category combination. The notation keys are as follows:

- (a) NO (not occurring) for activities that do not occur in the country;
- (b) NE (not estimated) where emissions do occur but are not estimated, usually because they are considered negligible or the necessary data cannot be obtained;
- (c) NA (not applicable) for activities that do not generate emissions of a particular pollutant;
- (d) IE (included elsewhere) for emissions relating to a subcategory that are reported in another subcategory, usually at the next highest level;
- (e) C (confidential) for emissions that could lead to the disclosure of confidential information;
- (f) NR (not relevant) for emissions that are not required by the ratified protocols.

At four-year intervals from 2017 onwards, the inventory submissions under CLRTAP should include compilations of emissions for a list of defined large point sources and aggregated sectoral gridded data for the European Monitoring and Evaluation Programme (EMEP) grid cells overlying the national territory. This information is used in EMEP models for evaluating long-range transport of air pollutants and for assessing emission deposition relationships in Europe. Ireland's 2017 submission contained data for large point sources and sectoral gridded data of emissions for 2015 on the EMEP grid. This information is provided in chapter 7 of this report.

Parties to the Gothenburg Protocol shall report national projections every four years from 2015 onwards, for the years 2020, 2025 and 2030 and where available, also for 2040 and 2050, by 15th March for the pollutants: SO_x (as SO₂), NO_x, NMVOCs, NH₃ and PM_{2.5}, with voluntary reporting for black carbon. Other Parties are encouraged to provide projections for these pollutants. The Directive (EU) 2016/2284 requires emission projections every two years from 2017 onwards from European Member States. Information on projections is provided in chapter 8 of this report.

1.4.3 Inventory Preparation

The air pollutant emission inventory database normally contains information on measured emission quantities, activity statistics (populations, fuel consumption, vehicle/kilometres of travel, industrial production and land areas), emission factors and the associated emission estimates for the NFR list of source categories. In practice, very few measured emission data are available for the range of gases covered and, consequently, the emissions from most activities are estimated by applying emission factors for each source/gas combination to appropriate activity data for the activity concerned. Virtually all emissions may be ultimately derived on the basis of the product of activity data and emission factor. Even in the case where emission estimates for particular categories are reported directly to the inventory agency they will normally have been derived in this manner.

The reporting guidelines provide the general guidance for the preparation and reporting of annual inventories by Parties. They incorporate the methodologies given in the EMEP/EEA (European Environment Agency) Emission Inventory Guidebook, hereafter referred to as the Inventory Guidebook. The inventory preparation process involves the acquisition of the required statistical data for the inventory year concerned and the application of emission factors that characterise the rate of emission of the gases concerned. Some data analysis and preparatory calculations are generally needed in order to make available suitable combinations of activity data and emission factors at the level of disaggregation that gives the best estimate of emissions in the individual emission source categories. In the case of some source/gas combinations, it may be necessary to apply sophisticated models to

generate the activity data, the emission factors or the emissions. The methods recommended by the Inventory Guidebook use a tiered system. This provides methodologies at different levels of detail and sophistication, which take account of these issues and other factors, such as data availability, technical expertise, inventory capacity and other circumstances, which may vary considerably across countries.

1.4.4 Data Acquisition

In its capacity as the inventory agency, the OES of the EPA acquires the principal items of activity data from identified Key Data Providers (KDP) relevant to each of the NFR sectors. Most KDPs provide data directly to the OES, but some secondary KDPs provide their input to one of the primary KDPs for processing and incorporation into the information subsequently transmitted to the OES. Some KDPs may also deliver estimates of emissions or removals for their particular area of coverage or expertise. Table 1.1 lists the KDPs and the data they supply for use in transboundary gas emission inventories.

The NAIS provides for a formal Memorandum of Understanding (MoU) between each KDP and the inventory agency regarding the scope, quality and submission date of the data to be provided for the purposes of the national emission inventory. In the majority of cases, the data concerned are already routinely collected and published by the KDPs under existing mandates and established reporting programmes. Additional MoUs may be developed under the NAIS in cases where new or supplementary data sources need to be targeted. Under Section 69 of the EPA Act 1992, formal legal powers are assigned to the EPA, whereby the Agency may require any public body to provide information related to environmental quality and may make arrangements with other bodies for the provision of similar information. This provision can also be invoked by the OES to acquire specific information for inventory purposes as the need arises.

The Emissions Trading Unit was established under the EPA Office of Licensing and Guidance (OLG) in late 2003 to implement Directive 2003/87/EC (EP and CEU, 2003) in Ireland. The Emissions Trading Unit currently forms part of the OES and is another key component of the national system. Information compiled for participants in the Emissions Trading Scheme (ETS) under Directive 2003/87/EC is an important source of activity-specific and company-specific data on GHG emissions for approximately 100 installations in Ireland. The inventories for transboundary gases draw on relevant information regarding fuel quantities and fuel properties available under the ETS for these installations and fuel data are used for reconciliation with the national energy balance for major categories and matching of activity data for GHG inventories. The inventory agency in the OES obtains useful support and activity data from other EPA offices and programme areas, including the Environmental Licensing Programme, Office of Environmental Enforcement, and the Environmental Research Programme. These programmes and offices make various contributions that are used to determine or substantiate the activity data or emission factors for particular categories or individual activities, which ensures that country-specific information is exploited to the maximum extent possible. In all cases, consistency is maintained with data application for GHG inventories and vice versa.

1.4.5 Quality Assurance and Quality Control (QA/QC)

Quality Assurance (QA) activities include a planned system of review procedures conducted by personnel not directly involved in the inventory compilation and development process. Reviews, preferably by independent third parties, should be performed upon a finalised inventory following the implementation of QC procedures. Reviews verify that data quality objectives were met, ensure that the inventory represents the best possible estimates

of emissions and sinks, given the current state of scientific knowledge and data available, and support the effectiveness of the QC programme.

A QA/QC spreadsheet tool has been developed to manage and maintain Ireland's QA/QC system. This tool consists of several spreadsheets that provide procedures, guidance, forms and templates as required for the general QA/QC functions. The supporting manual (Thistlethwaite et al., 2005) provides a general overview to the QA/QC system and guidance on the application of the plan and procedures. The QA/QC plan identifies the specific data quality objectives related to the principles of transparency, consistency, completeness, comparability and accuracy required for Ireland's national inventory and provides specific guidance and documentation forms and templates for the practical implementation of QA/QC procedures. The spreadsheets include a brief introduction and a statement of the data quality objectives (DQOs) and how they will be met through the QA/QC system with reference to the relevant spreadsheet tool template sheets and forms. The Introduction sheet links to the QA/QC plan which provides the schedules and procedures for the QA/QC system and lists all of the QA/QC activities that exist or are planned to make up Ireland's QA/QC system. The Plan sheet consists of tables that contain three different categories of QA/QC activity:

1. General activities covering the planning and management practices and procedures;
2. Activities that should be undertaken on an annual basis for management and preparation of the inventory;
3. Periodic activities that should be undertaken in response to specific events in the inventory and for periodic peer review or verification.

The inventory agency has implemented this approach to QA/QC for twelve annual reporting cycles. This involved the allocation of responsibilities linked to the national system and the use of the template spreadsheet system to record the establishment and maintenance of general inventory checking and management activities covering the overall compilation process, as well as the undertaking of specific annual activities and any necessary periodic activities in response to specific events or outcomes in inventory reporting and review. The system facilitates record keeping related to the chain of activities from data capture, through emission calculations and checking, to archiving and the identification of improvements.

Ireland's calculation spreadsheets in all sectors are structured on a time-series basis. This organisation is designed to facilitate the QA/QC process as well as more efficient trend analysis and to ensure ease of transfer of the outputs to the NFR tables. The inventory compilation is directly linked to the primary statistical inputs, which facilitates rapid year-on-year extension of the time series and efficient updating and recalculation, where appropriate, in the annual reporting cycle. Internal aggregation to various levels corresponding to the NFR tables provides immediate and complete checks of the results.

Quality Control (QC) is a system of routine technical activities, to measure and control the quality of the inventory as it is being developed. The QC system is designed to:

- (i) Provide routine and consistent checks to ensure data integrity, correctness, and completeness;
- (ii) Identify and address errors and omissions;
- (iii) Document and archive inventory material and record all QC activities.

Quality control activities include general methods such as accuracy checks on data acquisition and calculations and the use of approved standardised procedures for emission calculations, measurements, estimating uncertainties, archiving information and reporting.

Higher-tier QC activities include technical reviews of source categories, activity and emission factor data, and methods.

In recent years, the inventory agency has created and implemented a number of QA/QC tools. These spreadsheet-based tools are used to better inform the QA/QC process whilst providing transparent descriptions of the outcomes of checks. In each spreadsheet-based tool comments are added to explain anything highlighted by the checking process. The text from these tools (such as the recalculations assessment) then helps inform the update of the IIR. The tools used include:

1. Recalculations assessment – this spreadsheet tool calculates the percentage change of emission estimates between the current and previous inventory submissions using conditional formatting to highlight the significant changes. There is a separate table for each pollutant in which the entire time series is evaluated for all NFR codes. This check highlights the recalculations that have been made in the inventory so that they can be verified and justified by the inventory agency.
2. Trend assessment – this spreadsheet tool calculates the percentage change between the most recent year and the preceding year of the current inventory submission using conditional formatting to highlight the significant changes. There is a separate table for each pollutant in which the entire time series is evaluated for all NFR codes. This check helps identify any time series inconsistency with the newly reported data of the most recent year. Provided this is run annually and alongside the recalculations check, time series consistency should be maintained.
3. Pollutant specific assessments – there are two tools that check the following rules that should be maintained in an inventory: $TSP \geq PM_{10} \geq PM_{2.5}$; Total PAHs (1-4) = $B(a)p + B(b)F + B(k)P + I(123)-cd$. These simple checks help maintain the accuracy of the inventory.
4. Data value assessment – two tools check the entire time series for all pollutants to ensure that none of the following are reported: zero values, errors, negative values. These simple checks help maintain the accuracy and transparency of the inventory.
5. Annex I reporting template assessment – this tool evaluates whether all compiled Annex I reporting template files are comparable in structure and content to the template. This helps maintain the comparability of the inventory.
6. Notation keys assessment – this tool summarises the use of the different notation keys within the inventory. This tool has been implemented for all NEC pollutants across the entire time series. The tool helps the inventory agency evaluate, justify and document the use of notation keys in the inventory. By increasing the accuracy of the notation keys, the transparency of the inventory is improved and this allows the inventory agency to clearly identify areas where potential improvements could be made (e.g. the use of IE or NE).

The online tool provided by the Centre on Emissions Inventories and Projections (CEIP), RepDab, is used by the inventory agency to check the format, completeness and internal consistency of the Annex I reporting template submission files. Further details regarding the checks that are carried out can be found on the [CEIP](#) website.

In the 2018 reporting cycle, the inventory agency updated some of the default emission factors in accordance with the recently published EMEP/EEA Air Pollutant Emission Inventory Guidebook (2016) and implemented suggested changes/requests for further information as presented in the Final Review Report of the 2017 Comprehensive Technical Review of National Emission Inventories. These are outlined in the sectoral chapters of this report.

Table 1.1 Key Data Providers and Information Covered by Memoranda of Understanding

Key Data Provider	Data Supplied	Deadline	Sector in which Data are Used
Sustainable Energy Authority of Ireland (SEAI)	National Energy Balance; Detailed national energy consumption disaggregated by economic sector and fuel	30 September	Energy, Waste
Department of Agriculture Food and the Marine (DAFM)	Nitrogen fertiliser sales, cattle populations from the AIM (Animal Identification and Movement) database, sheep statistics, poultry statistics	30 September	Agriculture
Central Statistics Office	Annual population, livestock populations, crop statistics, housing survey data	30 September	Agriculture, Industrial Processes, Waste
Gas Networks Ireland	Analysis results for indigenous and imported natural gas	30 September	Energy
Marine Institute	Annual report on discharges, spills and emissions from offshore gas production installations	30 October	Energy
Emissions Trading Unit (OES, EPA)	Verified CO ₂ estimates and related fuel and production data for installations covered by the EU ETS	30 April	Energy, Industrial Processes
Department of Communications, Climate Action and Environment (DCCA)	National Oil Balance (as a component of the energy balance)	30 September	Energy
Road Safety Authority*	Road transport statistics from the National Car Testing (NCT) Service	30 April	Energy

*These bodies have MoUs with the SEAI rather than with the OES.

1.4.6 Inventory Compilation

The source data, calculation workbooks and outputs for all emissions to air are held on the server in the Monaghan Regional Inspectorate of the EPA. The annual inventory compilation for transboundary gases is undertaken in separate *Data Processing* folders for each sector, which are linked to the *Source Data* folders for the respective sectors at the same level. The *Outputs* folder and the *QA/QC* folder are also at this level. The *Outputs* folder contains the files used for the official submissions to the EU and the UN/ECE and for preparing summary reports and relevant media statements at national level. All calculation workbooks for the individual sectors contain a *QA/QC* worksheet, which are compiled collectively in the *QA/QC* folder. Data processing to compute the emission estimates is carried out at the most detailed level of aggregation possible, consistent with data availability and the outputs needed to populate the reporting template format for the category concerned. These outputs are primarily the estimates of emissions and the corresponding activity data for each category.

Quality control procedures are an integral part of the inventory preparation and reporting cycle. Within the inventories team, quality control for each sector is undertaken by an inventory compiler who has not produced the emission estimates for that sector. Quality control involves a series of checks covering the data inputs and any necessary pre-processing, the calculation of emissions, and the generation of the output records that are subsequently compiled in the NFR

templates. The checks cover such items such as the comparison of inputs with those of previous years, the identification of errors and omissions, validating internal linking and calculation algorithms, replicating the aggregation of subcategories and ensuring an adequate level of completeness in NFR files to achieve transparency for external review purposes. A colour code system is used to distinguish between such elements as data taken from another spread sheet, calculated values, extrapolated or interpolated values, outputs for the NFR, and checks and annotations.

1.5 Key Category Analysis

Key category analysis for transboundary air emissions is explained in the Inventory Guidebook, and is the same concept as that presented in the Intergovernmental Panel on Climate Change (IPCC) 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006). This defines a key emission category as one that is prioritised within the national inventory system because its emission estimate has a significant influence on the Party's total inventory in terms of the absolute level of emissions, the trend in emissions, or both. The Inventory Guidebook provides several methods for undertaking the analysis of key categories that can be applied at any appropriate level of source aggregation, depending on the information available. The simplest approach (Approach 1) – identification based on contribution to emission level – is used here for the inventories of all substances to highlight which sources of emissions are the most important in Ireland.

In level assessment, key categories are those categories whose combined contribution to the total emission level, determined from the ranking of all categories on the basis of their individual contributions to the level, is 80 per cent. Information about key categories is considered to be crucial to the choice of methodology for individual sources and to the management and reduction of overall inventory uncertainty. The identification of such categories is recommended in order that inventory agencies can give them priority in the preparation of annual inventories, especially in cases where resources may be limited. Information on key categories is clearly also vital for the development of policies and measures for emissions reduction.

It is well established that fuel combustion in a small number of economic sectors is the major source of most air pollutants. This is true for classic pollutants such as SO_x and NO_x, which are reasonably well quantified in emission inventories and for which emissions have decreased considerably, and also for other substances (PM, POPs) for which inventories have much higher levels of uncertainty. The relative contributions of key categories are clearly shown by the results of the simple key category analysis shown in Table 1.2 and Annex A.2, Tables 1-15, which summarises NFR Level 2 key categories by pollutant. The dominance of categories under 1A (Energy: Combustion) highlights the importance of combustion sources. The four key categories (1A1a, 1A2f, 1A3b and 1A4b) dominate the results of the key category analysis presented in Table 1.2. Agriculture sources (3B, 3D) and are the main sources of emissions for NH₃, NMVOC's and particulate matter and the additional categories 2D and 2H for NMVOCs. Waste sector sources are the main driver of emissions for As, Cr, dioxins and PCBs.

Table 1.2. Key Category Analysis of Ireland's Air Pollutant Inventory 2016

Pollutant	Key Categories									Total (%)
NO_x	1A3bi 15.16%	1A3biii 11.66%	3Da1 11.62%	3Da3 10.62%	1A3bii 8.18%	1A1a 7.13%	1A2f 6.28%	1A3dii 5.81%	3Da2a 5.29%	81.75%
CO	1A3bi 41.59%	1A4bi 19.93%	1A1a 17.80%	1A2f 4.63%						83.95%
NM VOC	3B1b 24.67%	2H2 19.61%	2D3a 9.93%	3B1a 8.57%	1A4bi 7.43%	3De 3.53%	2D3d 2.69%	1B2aiv 2.52%	1A3bi 2.36%	81.31%
SO_x	1A4bi 48.39%	1A1a 27.00%	1A2f 9.73%							85.12%
NH₃	3Da2a 29.65%	3B1b 26.80%	3Da3 12.16%	3B1a 11.51%						80.12%
TSP	2D3b 44.79%	1A4bi 11.91%	3Da1 10.90%	2A5a 5.84%	3B1b 3.85%	3B1a 3.16%				80.44%
PM₁₀	3Da1 23.88%	1A4bi 23.82%	2D3b 13.08%	2A5a 6.27%	3B1b 3.80%	1A3bvi 3.67%	3B1a 3.15%	1A2f 2.88%		80.55%
PM_{2.5}	1A4bi 44.06%	2D3b 12.28%	1A2f 4.92%	3B1b 4.78%	3B1a 3.87%	1A3bvi 3.74%	1A3bi 3.54%	1A1a 2.84%		80.04%
Pb	1A3bi 59.80%	1A4bi 15.53%	1A3bvi 10.44%							85.77%
Cd	1A1a 22.98%	1A2gviii 21.93%	1A4bi 14.42%	1A3bi 10.77%	1A2f 10.71%					80.81%
Hg	1A1a 33.93%	1A4bi 26.83%	1A2f 9.62%	1A3bi 7.18%	5A 6.36%					83.91%
As	5C1bi 51.33%	1A1a 40.27%								91.60%
Cr	5C1bi 25.19%	1A3bvi 20.76%	1A1a 12.61%	1A4bi 8.63%	1A2f 7.44%	1A4ai 6.15%				80.77%
Cu	1A3bvi 56.09%	1A3bi 25.76%								81.85%
Ni	1A4ai 30.65%	1A2f 23.79%	1A2e 12.79%	1A1a 10.55%	1A2gviii 8.05%					85.82%
Se	1A4bi 53.08%	1A1a 44.01%								97.09%
Zn	1A3bvi 22.65%	1A4bi 20.59%	1A3bi 15.13%	1A2gviii 12.14%	1A2f 8.19%	1A1a 7.34%				86.04%
PCDD/F	1A4bi 62.67%	5E 10.61%	2A1 6.96%							80.24%
PCBs	2A1 41.64%	1A4bi 21.51%	5E 20.55%							83.69%
HCb	3Df 68.63%	1A1a 27.35%								95.98%
PAHs	1A4bi 86.28%									86.28%
	1 Energy		2 IPPU		3 Agriculture		5 Waste			

1.6 Uncertainty Assessment

Undertaking a quantitative estimate of emissions uncertainty requires a substantial amount of detailed data on the uncertainty of both activity data and emissions factors for a diverse range of source types. It has not been possible to collect these data in full. However, it has been possible to characterise the uncertainties associated with sources in a more approximate way.

A semi-quantitative uncertainty analysis has been used to determine the overall emissions uncertainty for a number of pollutants for 2016 data. This uses a Tier 1 propagation of errors to obtain an uncertainty for the total emission. However, the uncertainty assigned to the activity data and emission factor for each individual source is obtained from a combination of expert judgement and ranges of uncertainty obtained from the EMEP/EEA emission inventory guidebook. The results provide a good indication as to which sources are contributing the most to the overall uncertainty, and therefore where improvement effort should be targeted.

The methodology and results of the Tier 1 uncertainty analysis are presented in detail in Annex G, tables G.1 to G.6. The results can be summarised as follows:

Table 1.3 Emissions Uncertainties

Pollutant	Emission (kilotonnes, 2016)	Uncertainty in 2016 (%)	Trend Uncertainty 1990-2016 (%)
NO _x	112.28	40.0	2.7
SO ₂	13.77	10.7	0.6
NM VOC	108.41	81.9	25.8
NH ₃	116.72	88.0	9.6
PM _{2.5}	15.47	99.5	24.0

The total uncertainty in the NO_x emission in 2016 is dominated by the contribution inorganic N-fertilisers (3.D.a.1), animal manure applied to soils (3.D.a.2.a) and urine and dung deposited by grazing animals (3.D.a.3), ± 200 per cent for each sub category of 3.D. These three categories combined account for 97.6 per cent of the total NO_x emissions uncertainty. The next largest contributor to uncertainty for NO_x emissions is fuel used by off road mobile machinery in agriculture (1.A.4.c.iii) with an uncertainty of over 100 per cent.

Emissions of SO₂ are well characterised when compared to NO_x for most emissions because they are combustion related, with emission factors for SO₂ (i.e. the sulphur content of the fuel) more readily determined than for NO_x. Solid fuel combustion, coal, biomass and peat in the residential sector combined account for 89.7 per cent of the overall uncertainty.

Emissions of NM VOC from non-combustion sources are typically high in uncertainty because they are difficult to characterise by measurement. Manure management from cattle (3.B.1.a & 3.B.1.b) contributes to 91.5 per cent of the overall uncertainty because of the magnitude of the emission of these categories (33.2 per cent in 2016) and, as with all manure management categories, has poorly characterised emission factors (± 300 per cent). Fugitive NM VOC losses from fuel extraction and distribution (1.B.2) and domestic solvent use (2.D.3.a) are large contributors because they have poorly characterised emission factors and activity data, respectively.

The uncertainty associated in NH₃ emissions are driven by the emission factors, with the activity data; number of livestock and nitrogen amounts, typically being well characterised by comparison. The sources making the largest contributions to the overall uncertainty are ammonia losses from inorganic N-fertiliser use (3.D.a.1), animal manure applied to soils

(3.D.a.2.a) and urine and dung deposited by grazing animals (3.D.a.3). The emission factors for these sources are currently assigned an uncertainty of ± 200 per cent, and they contribute to 96.9 per cent of the overall uncertainty.

Emissions of $PM_{2.5}$ are generally high in uncertainty because many combustion sources are either not well characterised, or are variable in emission, with small changes to combustion conditions having very large impacts on $PM_{2.5}$ emissions. Residential coal, biomass and peat combustion (1.A.4.b) and road paving with asphalt (2.D.3.b) are the largest contributors to the overall uncertainty, contributing 61.3 per cent and 38.1 per cent respectively. These are relatively large sources with very uncertain emission factors.

It is interesting to note that electricity generation and industrial combustion do not feature as major contributors to the overall uncertainty for any pollutants. This is because the use of point specific data allows the emission estimates to be particularly well characterised.

Chapter Two

Analysis of Key Emission Trends

2.1 Introduction

Ireland's 2018 submission under the CLRTAP and the Directive (EU) 2016/2284 includes emission estimates for the period 1990–2016 in respect of all substances listed in Section 1.4.2 above. The primary emission time series is prepared on the basis of Ireland's published national energy balances, which record the amounts of fuels sold in the country. In recognition of the significant cross-border movement that occurs with respect to automotive fuels in some parts of Europe, the reporting guidelines allow for the reporting of emissions from road transport on the basis of fuels used within the country. This may result in a significant decrease in the national total emissions for some pollutants and the adjusted total is considered more appropriate for the assessment of performance in relation to certain protocols. This issue is relevant to Ireland in the case of the Sofia Protocol on NO_x emissions and, to facilitate the assessment, Ireland has also submitted inventories in which the estimates for road transport are based on fuels used in the country. Emission inventories based on fuel sold and fuel used are provided in this submission for all pollutants for the period 1990–2016 and for the year 1987, the base year for the Sofia Protocol.

This chapter provides an overview of the emission trends for the period 1990–2016 for all substances included in Ireland's 2018 submission under the CLRTAP and NECD. The general analysis of trends is performed only in respect of emissions estimated on the basis of fuels sold in Ireland.

2.2 Main Pollutants

2.2.1 Sulphur Dioxide (SO₂)

Total sulphur dioxide emissions decreased by 92.5 per cent, from 183.57 kt in 1990 to 13.77 kt in 2016 (Figure 2.1). The Commercial/Institutional and Residential (1A4a and 1A4b) sectors combined account for 50.1 per cent of the total in 2016, and decreased by 82.0 per cent between 1990 (38.28 kt) and 2016 (6.90 kt). The Public Electricity and Heat Production (1A1a) sector remains one of the main sources of SO₂ emissions, contributing 27.0 per cent of the total in 2016, and decreased by 96.4 per cent between 1990 (103.04 kt) and 2016 (3.72 kt). An increase in consumption of coal, peat and oil in this sector in 1994 followed by decreased peat and oil consumption the following year caused a peak in emissions in 1994.

In 1998 an increase in consumption of coal followed by a decrease the following year caused another peak in emissions in 1998. Combustion sources in the Manufacturing Industries and Construction (1A2) sector largely account for the remainder of the emissions, with contribution of 18.8 per cent in 2016. Emissions in this sector peaked in 1994 due to an

increase in SO₂ from oil combustion in the installation which is the main contributor to emissions in Non-ferrous metals (1A2b). Emissions in Manufacturing Industries and Construction (1A2) have decreased in the 1990-2016 time series by 92.0 per cent. Combustion in Agriculture/Forestry/Fishing (1A4c) sector accounts for 0.3 per cent and Transport (1A3) combustion sources account for 1.9 per cent of national total emissions of SO₂ in 2016. The remainder of the SO₂ emissions arise from combustion sources in the Petroleum Refining (1A1b) and Manufacture of Solid Fuels and Other Energy Industries (1A1c) sectors, Other Product Use (2G) Waste Incineration (5C1), which combined account for 1.9 per cent of the total in 2016 and are presented in Other NFR sectors in Figure 2.1. In 1990, coal combustion accounted for 51.5 per cent of SO₂ emissions and fuel oil contributed 30.3 per cent. By 2016, the share of SO₂ emissions from coal had decreased to 48.9 per cent and that from fuel oil had decreased to 6.6 per cent.




Figure 2.1 Emission Trend for Sulphur Dioxide 1990–2016

2.2.2 Nitrogen Oxides (NO_x)

Total nitrogen oxides emissions have decreased by 38.7 per cent, from 175.05 kt in 1990 to 107.30 kt in 2016 (Figure 2.2). Road Transport (1A3b) is the principal source of NO_x emissions, contributing 36.7 per cent (and 39.33 kt) of the total in 2016, with the transport sector as a whole accounting for 41.0 per cent (and 43.98 kt) of the national total. The Manufacturing Industries and Construction (1A2) sector accounts for an increasing percentage of the national total. The contribution of the sector in 1990 to the national total was 5.1 per cent (8.93 kt), which increased to 12.1 per cent share in 2007 (and 18.42 kt) as a result of the increases in cement production for construction during the economic boom in Ireland over the previous decade then reduced to 8.3 per cent share (8.46 kt) of the national total in 2011 due to the economic crisis impacting upon the sector. In 2016 the sector contribution increased to a 10.5 per cent share (and 11.31 kt) of the national total largely driven by an increase in cement production.

The Public Electricity and Heat Production (1A1a) sector is another main source of NO_x emissions, accounting for 7.5 per cent of emissions in 2016. Emissions from this sector have decreased by 82.7 per cent between 1990 (46.37 kt) and 2016 (8.00 kt). Commercial/Institutional and Residential (1A4a and 1A4b) sectors combined account for 7.00 per cent of the total and combustion sources in Agriculture/Forestry/Fishing (1A4c) sector account for 3.8 per cent in 2016. The remainder of the combustion sources of NO_x

arise in the Petroleum Refining (1A1b) and Manufacture of Solid Fuels and Other Energy Industries (1A1c) sectors, as well as combustion sources in Other product Use (2G) and the Waste sector (5C1), which are presented in Other NFR sectors that together account for 0.6 per cent of the total in 2016. Agricultural sources of NO_x, reported for the first time in this submission accounted for 29.6 per cent of emissions in 2016 (31.77 kt), having reduced by 3.4 per cent since 1990 (32.90 kt).

The largest sources of NO_x emissions within agriculture are associated Inorganic fertilizer application (3Da1) and Urine and Dung Deposited by grazing animals (3Da3).

The reductions in NO_x emissions arising from the use of catalytic converters in cars and heavy-duty vehicles have only become apparent in recent years, as the technology has been offset by large increases in vehicle numbers in the past 10 years. This effect is exaggerated in latter years by so-called fuel tourism, whereby a significant proportion of the automotive fuel sold in Ireland – the basis for the emission time series given in Figure 2.2 – is used by vehicles in other countries. The estimated level of fuel tourism is given in Annex A.3, together with the adjusted annual NO_x emissions based on fuels used in Ireland, which is relevant to the assessment of obligations in relation to the Sofia Protocol on NO_x emissions.




Figure 2.2 Emission Trend for Nitrogen Oxides 1990–2016

2.2.3 Ammonia (NH₃)

Total ammonia emissions have increased by 6.3 per cent, from 109.8 kt in 1990 to 116.70 kt in 2016 (Figure 2.3). Livestock production has historically accounted for the bulk of national total ammonia emissions in Ireland and, in 2016, Manure Management (3B) and Organic Fertilisers (3Da2a) applied to soils combined accounted for 89.4 per cent of the national total. In 2016, Manure Management at 55.09 kt and 47.2 per cent share of the total in 2016 showed an increase by 11.1 per cent from 49.60 kt in 1990. Organic Fertilisers cover emissions from the two sectors: Animal Manure applied to soils (3Da2a) and Sewage Sludge applied to soils (3Da2b), which combined at 35.05 kt accounted for 30.0 per cent of the total ammonia emissions in 2016 (indicating an 8.6 per cent increase from 32.26 kt in 1990). Urine and Dung deposited by grazing animals (3Da3) accounted for 12.2 per cent of total

emissions in 2016 (14.19 kt), having increased by 1.9 per cent since 1990. Inorganic N-fertilizers applied to soils (sector 3Da1) decreased by 18.3 per cent from 1990 (13.69 kt) and at 11.19 kt in 2016 accounted for 9.6 per cent of the national total. The small contribution by Transport (1A3) sources peaked in 2005, the main driver of which has been the increased use of cars with early generation three way catalysts in Road Transport (1A3b). Transport emissions have increased from 0.05 kt in 1990 to 0.76 kt (and 0.7 per cent share of the total) in 2016. The remainder of the ammonia emissions arise from Commercial/Institutional and Residential (1A4a and 1A4b) sectors combined (0.1 per cent share) and Other NFR sectors (Combustion in Manufacturing Industries and Construction (1A2), Combustion in off road Agricultural machinery (1A4cii), Other Product Use (2G), Biological Treatment of Waste (5B1)) that together account for 0.4 per cent of the total in 2016.

Within livestock production, Manure Management (3B) at 55.09 kt in 2016 is the largest source of NH₃. In Ireland, approximately two-thirds (61.0 per cent) of animal manure is excreted at pasture annually, reflecting the relatively short period that cattle are housed.

Dairy Cattle (3B1a) and Non-Dairy Cattle (3B1b) account for the major part (38.7 per cent) of Agriculture sector ammonia emissions in 2016. Other livestock, which includes Sheep (3B2), Swine (3B3), Goats (3B4d), Horses (3B4e), Mules and asses (3B4f), Poultry (3B4g) and Other animals (manure management; 3B4h), combined account for 9.0 per cent of total Agriculture and 8.9 per cent of the national total ammonia emissions in 2016 (Figure 2.3). Throughout the 1990s, the cattle herd increased to reach a peak in 1998 of 7.6 million head, which along with associated increases in fertiliser nitrogen consumption increased ammonia emission totals from the whole agriculture sector, from 109.48 kt in 1990 to 122.20 kt in 1998. As a result of reforms to the Common Agricultural Policy (CAP), animal numbers and associated fertiliser nitrogen use have reduced, and ammonia emissions in agricultural sectors had fallen to 102.70 kt in 2011. However, in response to growth plans for the sector emissions have increased in recent years with a total of 115.53 kt in 2016 largely as a result of increased dairy cattle and other cattle populations, increased fertilizer use and increased use of urea as an inorganic fertilizer.




Figure 2.3 Emission Trend for Ammonia 1990–2016

2.2.4 Non-Methane Volatile Organic Compounds (NMVOCs)

Total non-methane volatile organic compound emissions have decreased by 25.8 per cent, from 145.81 kt in 1990 to 108.25 kt in 2016 (Figure 2.4). The NMVOC emissions are determined largely by the Agriculture sectors: 3B Manure Management and 3Da1 Inorganic N-fertilizers and emissions from solvents and other product use and the food and beverages industry. The Agriculture categories combined accounted for 42.4 per cent of the national total and showed an increase by 15.2 per cent between 1990 (39.85 kt) and 2015 (45.90 kt). The combined solvents use (2D and 2G) and fugitive emissions from oil sectors emissions (1.B.2.a) produced 21.4 per cent of the 2016 total of NMVOC emissions in Ireland, having decreased by 13.7 per cent between 1990 (26.80 kt) and 2015 (23.13 kt). The Food and Beverage Industry (2H2) contributed to 19.6 per cent of total emissions in 2016, having increased by 121.1 per cent from 9.62 kt in 1990 to 21.26 kt in 2016.

Combustion sources in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors are also important sources, accounting for 8.0 per cent of national total NMVOC emissions in 2016, a reduction of 70.2 per cent between 1990 (28.99 kt) and 2016 (8.63 kt). Technological controls for volatile organic compounds (VOCs) emitted by motor vehicles have been more successful than in the case of NO_x, and have contributed to a significant reduction in emissions from Road Transport (1A3b), with the total transport sector's contribution having decreased by 85.5 per cent between 1990 (36.28 kt) and 2016 (5.26 kt). This equates to contributions to the national total of 24.9 per cent in 1990, falling to 4.9 per cent in 2016. A total of five NFR source categories make up the source Other NFR sectors (Figure 2.4) contributing 0.5 per cent of national total NMVOC emissions in 2016. The largest source of emissions within this categorisation is Solid waste disposal on land sector (5A).




Figure 2.4 Emission Trend for Non-Methane Volatile Organic Compounds 1990–2016

2.2.5 Carbon Monoxide (CO)

Carbon monoxide emissions continue to decline, driven by major reductions due to three way catalysts in gasoline vehicles in Road Transport (1A3b), which is the principal source of CO, and a large decrease in the use of solid fuels for space heating in the Residential

(1A4b) sector (Figure 2.5). National total CO emissions have reduced from 367.90 kt in 1990 to 101.27 kt in 2016, a reduction of 72.5 per cent.

The Transport sector accounted for 49.4 per cent of national total emissions in 2016, a major reduction (81.5 per cent) from 270.97 kt in 1990 to 50.07 kt in 2016. Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined are another important source, accounting for 21.5 per cent of national total CO emissions in 2016, a reduction of 65.7 per cent between 1990 (63.47 kt) and 2016 (21.76 kt). Public Electricity and Heat Production (1A1a) sector reached its peak in 2001 (23.82 kt) and has decreased by 23.9 per cent to reach 18.25 kt in 2016, a reduction of 0.6 per cent on 1990 levels (18.14 kt). Combustion sources from Manufacturing Industries and Construction (1A2) account for 9.5 per cent of the national total in 2016 and at 9.58 kt showed a 13.3 per cent decrease on their 1990 levels (11.06 kt). Agriculture/Forestry/Fishing (1A4c) combustion sources account for 1.2 per cent of the total CO emissions in 2016. Petroleum refining (1A1b), Manufacture of soil fuels and other energy industries (1A1c), Other Product Use (2G), Biological Treatment of waste (5B1) and Waste incineration (5C) emissions form the Other NFR sectors category and combined account for the remainder of CO emissions (0.4 per cent of the total) in 2016.




Figure 2.5 Emission Trend for Carbon Monoxide 1990–2016

2.3 Particulate Matter

Particulate matter emission estimates include PM with diameter less than 10 μm (PM_{10}), PM with diameter less than 2.5 μm ($\text{PM}_{2.5}$) and total suspended particulates (TSP).

2.3.1 Particulate Matter <10 μm Diameter (PM_{10})

Emissions of particulate matter <10 μm diameter amounted to 28.21 kt in 2016, a 42.5 per cent reduction from 49.08 kt in 1990 (Figure 2.6). The main determinant of the trend in PM_{10} emissions is Agriculture sector with 35.8 per cent share of the national total, and combustion in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined with 25.5 per cent share of the total in 2016.

Emissions from Agriculture arise from Inorganic N-fertilizers (3Da1), Off-farm storage, handling and transport of bulk agricultural products (3Dd) and Manure Management (3B)

categories that together in 2016 at 10.11 kt indicated a 0.1 per cent increase on their 1990 levels. Part replacement of coal and peat in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors are the second largest contributor to the total PM₁₀ emissions, with 71.2 per cent reduction in emissions from these sectors, from being the largest contributor to the PM₁₀ emissions (50.9 per cent of national total emissions) at 24.99 kt in 1990, emissions have fallen to 7.19 kt in 2016 (and 25.5 per cent of the national total emissions). Emissions from Transport (1A3), at 7.2 per cent share of the total in 2016 increased throughout the 1990s with increased total vehicle kilometre travel particularly of diesel vehicles. However, the effect of the increase in vehicle numbers seen over the last decade has been offset somewhat by changes in the age structure of the national fleet and developments in diesel fuel technology resulting in decreased Transport emissions by 35.2 per cent (from 3.12 kt in 1990 to 2.02 kt in 2016).

Manufacturing Industries and the Construction sector (1A2) used to account for an increasing percentage of the national total PM₁₀, until reaching its peak of 2.58 kt in 2005. This is also evident with some other pollutants, and is due to the increase in cement production post-2000 following the entry into the market of two new plants. Emissions for this sector have decreased since 2005, and accounted for 5.7 per cent of the national total in 2016 (1.62 kt), representing a decrease across the 1990-2016 time series of 10.4 per cent. Public Electricity and Heat Production (1A1a) sector emissions accounted for 2.4 per cent of the national total in 2016 and at 0.67 kt reduced by 30.0 per cent from 1990 levels (0.96 kt). The decrease was due to the increased use of natural gas and wind for electricity generation, in proportion to coal and peat which still account for a large share of the fuel mix used. Petroleum refining (1A1b), Manufacture of solid fuels and other energy industries (1A1c), Quarrying and mining of minerals other than coal (2A5a), Construction and demolition (2A5b), Road paving with asphalt (2D3b), Other solvent use (2D3i), Other product use (2G), Biological treatment of waste – solid waste (5A), Incineration (5C) and Other waste (5E) are included under the other NFR sectors heading, which combined accounted for 22.5 per cent of national emission in 2016. The largest contributors are within the other NFR grouping are Quarrying and mineral of minerals other than coal (2A5a) and Road paving with asphalt (2D3b).




Figure 2.6 Emission Trend for Particulate Matter <10 µm in Diameter 1990–2016

2.3.2 Particulate Matter <2.5 µm Diameter (PM_{2.5})

National total emissions of particulate matter <2.5 µm diameter amounted to 14.94 kt in 2016, a 57.9 per cent reduction on 35.46 kt in 1990 (Figure 2.7). Emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined are the main determinant of the trend with their 47.1 per cent share of the national total PM_{2.5} emissions in 2016. There has, however, been a reduction of 71.3 per cent in emissions from these sectors between 1990 (24.48 kt and 69.0 per cent share) and 2016 (7.03 kt). Reduced use of coal and peat, with increased use of gasoil, kerosene and natural gas in the two sectors has resulted in lower emissions and a reduction in the contribution to the national total.

Emissions from Agriculture arise from Manure Management (3B) and Inorganic N-fertilizers (3Da1) sectors that together in 2016 at 1.80 kt accounted for 12.1 per cent of the national total and indicated a 1.1 per cent decrease on their 1990 levels (1.82 kt). Transport (1A3) contributed 1.57 kt (10.5 per cent share) to the national total in 2016. Emissions from Transport sector (1A3), dominated by Road Transport (1A3b) increased from 1990 (2.90 kt) to a peak in 1996 (3.30 kt), but have been decreasing since 2005, with a 45.7 per cent reduction between 1990 and 2016 which is largely due to technological advances and the age structure of the national fleet which in turn have been balanced by the increases in vehicle numbers over the time series (see comments in section 2.3.1 on the trends of PM₁₀ emissions). Emissions from Manufacturing Industries and Construction (1A2) have decreased from 1.59 kt in 1990 to 1.48 kt in 2016 and 9.9 per cent share of national total emissions (a decrease of 6.6 per cent in the trend), the result of decreased fuel use of petroleum coke in cement production (Non-metallic minerals, 1A2f).

Electricity and Heat Production (1A1a) sector accounts for 2.9 per cent of the national total emissions in 2016, a reduction of 31.9 per cent from 0.65 kt in 1990 and 0.44 kt in 2016. Combustion from the Agriculture/Forestry/Fishing sector accounted for 1.0 per cent of national total PM_{2.5} emissions in 2016 and has reduced by 84.5 per cent from 1990 when emissions were 0.93 kt, compared to 0.14 kt in 2016. The remainder of the PM_{2.5} emissions arise from Petroleum refining (1A1b), Manufacture of soil fuels and other energy industries (1A1c), Fugitive emissions from solid fuels: Coal mining and handling (1B1a), Quarrying and mining of minerals other than coal (2A5a), Construction and demolition (2A5b), Storage, handling and transport of mineral products (2A5c), Road paving with asphalt (2D3b), Other solvent use (2D3i), Other product use (2G), Biological treatment of waste – solid waste (5A), Incineration (5C) and Other waste (5E) are included under the other NFR sectors heading, which combined accounted for 16.5 per cent of national emission in 2016. Road paving with asphalt (2D3b) contributes over 77 per cent of this Other category.




Figure 2.7 Emission Trend for Particulate Matter <2.5 µm in Diameter 1990–2016

2.3.3 Total Suspended Particulates (TSP)

Total suspended particulate emissions have decreased by 29.7 per cent, from 87.92 kt in 1990 to 61.85 kt in 2015 (Figure 2.8). The main driver of the TSP trend is emissions from Other NFR sectors which includes a wide range of source categories namely Petroleum refining (1A1b), Manufacture of soil fuels and other energy industries (1A1c), Fugitive emissions from solid fuels: Coal mining and handling (1B1a), Quarrying and mining of minerals other than coal (2A5a), Construction and demolition (2A5b), Storage, handling and transport of mineral products (2A5c), Road paving with asphalt (2D3b), Other solvent use (2D3i), Other product use (2G), Biological treatment of waste – solid waste (5A), Incineration (5C) and Other waste (5E) which combined accounted for 55.2 per cent of the national total in 2016. Road paving with asphalt (2D3b) accounts for the majority of emissions within this grouping (83.5 per cent). Emissions from the Other NFR sectors have reduced by 11.2 per cent from 38.42 kt in 1990 to 34.14 kt in 2016. Emissions from the agriculture (Manure Management (3B) and Inorganic Fertiliser (3Da1)) sector were 15.04 kt (and 24.3 per cent share of the total) in 2016, a 2.4 per cent increase from 1990 levels (14.69 kt). Combined emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors were the third largest contributor to the total TSP emission in 2016. There has been, similar to emissions from both PM₁₀ and PM_{2.5}, a reduction of 71.2 per cent in emissions from these sectors between 1990 (27.49 kt) and 2016 (7.90 kt). In the time series, the part replacement of coal and peat with natural gas, gasoil and kerosene has resulted in the contribution of these sectors falling from 27.5 per cent of the national total in 1990 to 12.8 per cent in 2016.

Emissions from Transport (1A3) decreased by 35.2 per cent to 2016 (2.02 kt and 3.3 per cent share of national total) compared to 1990 (3.12 kt and 3.6 per cent share). Manufacturing Industries and Construction (1A2) account for an increasing proportion of emissions post-2000 as a result of the entry into the market of two new cement production plants (Non-metallic minerals, 1A2f). Emissions from the 1A2 sector had their peak in 2005 (2.93 kt) and although have been generally declining since, emissions increased 20.0 per cent between 2013 and 2016. At 1.78 kt in 2016 (2.9 per cent of the total) emissions have decreased from 2005 by 39.3 per cent and by 12.7 per cent from 1990 (2.04 kt). Public Electricity and Heat Production sector (1A1a) emissions have decreased by 33.5 per cent over the time series, from 1.23 kt in 1990 to 0.82 kt (and 1.3 per cent share of the total emissions) in 2016. Emissions from Agriculture/Forestry/Fishing in 2016 (0.15 kt, 0.2 per cent of the national total) have decreased by 83.7 per cent since 1990 (0.94 kt).




Figure 2.8 Emission Trend for Total Suspended Particulates 1990–2016

2.3.4 Black Carbon (BC)

Black Carbon emissions have decreased by 56.0 per cent, from 4.16 kt in 1990 to 1.83 kt in 2016 (Figure 2.9). The main driver of the BC trend is emissions from Transport (1A3). Emissions from the sector have reduced by 44.6 per cent between 1990 (1.38 kt) and 2016 (0.76 kt). Combined emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors were the second largest contributor to total BC emissions in 2016. In 2016, combined emissions from the sectors were 0.57 kt (31.4 per cent share) and have reduced by 68.4 per cent since 1990 (1.82 kt). Manufacturing Industry and Construction (1A2) emissions accounted for 19.4 per cent of the total in 2016 (0.36 kt) having increased by 5.0 per cent since 1990 (0.34 kt). Emissions from Agriculture/Forestry/Fishing in 2016 were 0.07 kt, which equates to an 85.4 per cent reduction on 1990 (0.51 kt). Public Electricity and Heat Production (1A1a) emissions have decreased by 54.4 per cent over the time series, from 0.02 kt in 1990 to 0.01 kt in 2016. The remaining 2.7 per cent in 2016 accounts for emissions from Petroleum refining (1A1b), Manufacture of solid fuels and other energy industries (1A1c), Road paving with asphalt (2D3b), Other product use (2G), and Industrial waste incineration (5C1bi).




Figure 2.9 Emission Trend for Black Carbon 1990–2016

2.4 Priority Metals

2.4.1 Lead (Pb)

Over the 1990–2016 time series, total national Pb emissions have decreased by 92.3 per cent, from 171.18 t in 1990 to 13.13 t in 2016 (Figure 2.10). The Pb emissions trend is largely determined by Road Transport (1A3b). Emissions of Pb have decreased considerably since 1990. There was a marked decrease between 1999 and 2000 when the lead content of petrol was reduced. In addition there was an increase in the use of unleaded gasoline in road transport throughout the 1990s and the subsequent phasing out of leaded gasoline in 2000/2001. The contribution of Transport (1A3) to the much-reduced national total emissions has decreased by 94.1 per cent, from 155.50 t (90.8 per cent share) in 1990 to 9.22 t (70.2 per cent share) in 2016.

The second largest contributors to Pb emissions at 17.3 per cent share of national total in 2016 were the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined. The use of coal and peat in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors is being part replaced with natural gas, gasoil or kerosene. Emissions from these sources have fallen from 7.98 t in 1990 to 2.27 t in 2016, a reduction of 71.6 per cent. Combustion in Manufacturing Industries and Construction (1A2) accounted for 8.1 per cent share of the total in 2016, and emissions from Public Electricity and Heat Production sector (1A1a) were responsible for 4.0 per cent share in 2016. Emissions from Metal Production (2C) have decreased in recent years due to the closure of a number of foundries and were negligible in 2016. Similarly, emissions from Waste Incineration (5C) have also decreased to almost zero as a result of an outright ban on the incineration of clinical wastes in the mid-1990s. Incineration in 5C is now solely in relation to the destruction of vapours.




Figure 2.10 Emission Trend for Lead 1990–2016

2.4.2 Cadmium (Cd)

Total national emissions of Cd have decreased from 0.56 t in 1990 to 0.28 t in 2016 (Figure 2.11), a reduction of 50.1 per cent. Emissions of cadmium are largely determined by the Manufacturing Industries and Construction (1A2) sector (38.1 per cent share of national total in 2016), specifically combustion sources in Non-Ferrous Metals (1A2b). Across the time series, the contribution of combustion sources within the Manufacturing Industries and Construction (1A2) sector reached its peak in 2005 (0.12 t) subsequently decreasing to 0.08 t in 2012 and increasing back up to 2005 levels in 2014, 2015 and 2016. Public Electricity and Heat Production (1A1a) decreased from 1990 by 22.9 per cent and accounted for 24.6 per cent of the national total in 2016. Emissions from this source increased throughout the 1990s as a result of the combustion of increasing quantities of coal and peat for electricity generation. The use of coal has reduced across the time series.

Residential (1A4b) and Commercial/Institutional (1A4a) combustion is also an important source of Cd emissions, with combined emissions from these sectors accounting for 20.8 per cent of the national total in 2016 due to the continued use of the fossil fuels (coal, peat and oil); however, emissions from the sector have fallen by 51.2 per cent, from 0.12 t in 1990 to 0.06 t in 2016. Transport (1A3) sector emissions have been increasing in the time series (by 106.6 per cent) and in 2016 it accounted for 15.6 per cent of the national total cadmium emissions. Combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 0.6 per cent of the total Cd emissions in 2016. As a result of the closure of the Irish Steel plant in late 2001, emissions from Metal Production (2C) are no longer a driver of the trend (0.0 per cent share in 2016) in national total emissions of Cd compared to a 38.5 per cent share (and main contributor to the total) in 1990.




Figure 2.11 Emission Trend for Cadmium 1990–2016

2.4.3 Mercury (Hg)

Total national emissions of Hg have decreased from 0.82 t in 1990 to 0.35 t in 2016, a reduction of 57.1 per cent (Figure 2.12). Emissions from Public Electricity and Heat Production (1A1a) decreased by 3.7 per cent in the trend and accounted for 32.3 per cent (the largest source) of national total mercury emissions in 2016 (0.11 t). Emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined account for the second largest share of national total mercury emissions (26.2 per cent in 2016), having decreased by 69.9 per cent between 1990 and 2016 as a result of an increase in the use of natural gas, kerosene and gasoil and a decrease in the use of coal and peat. Combustion sources in Manufacturing Industries and Construction (1A2) accounted for 19.2 per cent of total emissions in 2016, a 22.2 per cent reduction since 1990. Emissions from this sector are largely dependent on the increased use of petroleum coke and coal as a fuel source in the cement industry after the entry of a number of additional cement producers into the Irish market post-2000 to reach their peak in 2005. The sectoral emissions decreased since 2005 and especially after 2008 when the Irish economy was impacted by the recession.

Biological treatment of waste – Solid waste disposal on land (5A) and Waste Incineration (5C) combined accounted for a 8.2 per cent share of total emissions in 2016. The Transport (1A3) sector accounted for 13.8 per cent and combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 0.2 per cent of the total Hg emissions in 2016. Glass Production (2A3) and Metal Production (2C) are no longer occurring due to plant closures (glass since 2010 and metal since 2002).




Figure 2.12 Emission Trend for Mercury 1990–2016

2.5 Other Metals

2.5.1 Arsenic (As)

Emissions of arsenic have decreased by 23.0 per cent from 1.64 t in 1990 to 1.27 t in 2016 (Figure 2.13). These emissions are largely dominated by incineration of hazardous and clinical wastes and crematoria in Waste Incineration sector (5C). Waste Incineration accounted for 51.3 per cent of national total arsenic emissions, having increased by 17.6 per cent from 0.55 t in 1990 to 0.65 t in 2016. The absolute and percentage contributions of this sector are increasing largely due to the increase in the number of cremations undertaken in Ireland which has increased by 237 per cent since 1990 and incineration of wood products historically treated with a preservative containing arsenic. Continued use of coal, peat and fuel oil as part of the fuel mix contributes largely (40.9 per cent share of the total in 2016) to the trend in emissions from Public Electricity and Heat Production (1A1a); however, arsenic emissions from the sector have decreased by 16.2 per cent over the time series due to the replacement of less-efficient peat plants with new plant, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.

The continued use of fossil fuels in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined results in the sectors accounting for 4.3 per cent of national total emissions in 2016, however, they have decreased by 68.4 per cent in the trend. Within the Manufacturing Industries and Construction (1A2) sector, the sub-sector Non-metallic minerals (1A2f) is responsible for the majority of total sector's emissions due to the increase in cement production and associated fuel use in the sector, in particular petroleum coke post-2000 with the entry into the market of new cement plants and reflecting more recent post-recession decrease in production and consequential lower emissions from the sector. The sector's As emissions contributed 3.0 per cent the total in 2016, a decrease of 42.7 per cent on the 1990 level. Metal Production (2C) is no longer an important source of As emissions, following the closure of the Irish Steel plant in 2001 and a reduction in emissions from Integrated Pollution Prevention and Control (IPPC)-licensed facilities. Emissions from this sector accounted for 13.6 per cent in 1990 (third largest contributor) and are reported as not occurring in 2016 due to the closure of the foundry which was a responsible for emissions from this sector between 2001 and 2013.




Figure 2.13 Emission Trend for Arsenic 1990–2016

2.5.2 Chromium (Cr)

Emissions of Cr have decreased by 46.6 per cent from 4.53 t in 1990 to 2.42 t in 2016 (Figure 2.14). Transport (1A3) and Waste Incineration (5C) sectors are the main two drivers of the chromium trend. The Transport (1A3) sector accounted for 26.9 per cent of estimated national total emissions in 2016. Emissions from this source category have increased by 103.1 per cent over the time series due to the large increase in vehicle numbers (150.4 per cent) in Ireland between 1990 (0.32 t) and 2015 (0.65 t). The incineration of hazardous and clinical wastes and crematoria contribute to emissions from Waste Incineration (5C). The sector is the second largest source of chromium emissions and accounted for 26.0 per cent of 2016 national chromium emissions, an increase of 17.4 per cent in the trend. The absolute and percentage contributions of this sector are increasing largely due to an increase in industrial waste incineration (5C1bi) which includes the incineration of wood that has historically been treated with preservative containing chromium.

Continued use of coal, peat and fuel oil as part of the fuel mix for Public Electricity and Heat Production (1A1a) means that the sector contributed 13.6 per cent to the emissions total in 2016. However, there has been a reduction in emission levels of 16.2 per cent between 1990 and 2016 as a result of the replacement of less-efficient peat plants with new plant, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy. Emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined accounted for 15.2 per cent of the national total in 2016, having decreased, by 63.8 per cent over the time series, reflecting the part replacement of coal and peat with natural gas, gasoil and kerosene. Within the Manufacturing Industries and Construction (1A2) sector, the sub-sector Other (1A2f) is responsible for the majority of emissions largely due to the cement industry, as is evident with other heavy metal estimates. Emissions in 1A2 accounted for 17.7 per cent of national total chromium emissions and have decreased by 18.3 per cent in the whole trend. Similar to other heavy metals estimates, the closure of the Irish Steel plant in 2001 has significantly reduced the effect of Iron and Steel Production (2C1) on emission trends, from 31.5 per cent share of the total (and the main contributor to chromium emissions) in 1990. Emissions from the sector are reported as not occurring in 2014, 2015 and 2016 due to the closure of the foundry which was a responsible for emissions from this sector in 2013.




Figure 2.14 Emission Trend for Chromium 1990–2016

2.5.3 Copper (Cu)

Total copper emissions in Ireland were steadily increasing over the 1990–2008 period and have been steadily decreasing since (Figure 2.15). Total emissions in 2016 (18.31 t) were 12.2 per cent lower than in their peak in 2008 (21.27 t) but they are 66.0 per cent higher than in 1990 (11.03 t). This trend is determined mostly by the Transport (1A3) sector that accounted for 89.9 per cent of estimated national total copper emissions in 2016. Emissions from this source category have more than doubled over the time series due to a large increase in vehicle numbers in Ireland; an increase of 110.8 per cent between 1990 (7.81 t) and 2016 (16.45 t).

Continued use of coal, peat and fuel oil as part of the fuel mix results in a contribution of 3.1 per cent to the total emissions from Public Electricity and Heat Production (1A1a) in 2016. However, a reduction in emission levels of 18.8 per cent is evident between 1990 and 2016 as a result of the replacement of older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy. The Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined accounted for 2.4 per cent of emissions in 2016, showing a 70.0 per cent decrease in the trend. Waste Incineration (5C) emissions accounted for 1.9 per cent of the national total in 2016, having increased by 16.8 since 1990. Combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 1.4 per cent of the total Cu emissions in 2016 (26.4 per cent decrease since 1990). The Manufacturing Industries and Construction (1A2) sector has decreased by 35.8 per cent between 1990 and 2016 (accounting for 1.3 per cent of the total in 2016). Emissions from the sector were increasing proportionately since 1990 reaching their peak in 2005 as a result of the entry into the market of new operators in the cement production sub-sector post-2000. The use of fossil fuels for combustion decreasing since 2005 as a result of economic circumstances. The closure of the Irish Steel plant in 2001 meant that copper emissions from Metal Production (2C) have made an insignificant contribution to the national total since 2001.




Figure 2.15 Emission Trend for Copper 1990–2016

2.5.4 Nickel (Ni)

National total emission estimates of nickel were steadily increasing over the 1990–1999 period and have been generally decreasing from 1999 onwards (Figure 2.16). Total emissions in 2016 (8.45 t) were 76.8 per cent lower than in their peak in 1999 (36.48 t) and 62.2 per cent lower than in 1990 (22.33 t). The main contributor to the trend is Manufacturing Industries and Construction (1A2) with a 48.7 per cent share in 2016 (4.11 kt) having reduced by 41.4 per cent since 1990 (7.02 kt). The second largest contributor to total Ni emissions in 2016 is emissions from the combined Residential (1A4b) and Commercial/Institutional (1A4a) sectors accounting for 33.2 per cent. A switch within these sectors from solid fuels (coal and peat) to less carbon intensive liquid fuels and natural gas has resulted in a 61.2 per cent in emissions since 1990. Public Electricity and Heat Production (1A1a) sector accounted for a 10.8 per cent share of the total nickel emissions in 2016. However, a reduction in emission levels of 77.7 per cent is evident between 1990 and 2016 as a result of the replacement of older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.

Combustion sources from Agriculture/Forestry/Fishing (1A4c) sector accounted for 2.4 per in 2016. Similar to other heavy metals estimates, the closure of the Irish Steel plant in 2001 has significantly reduced the effect of Metal Production (2C) on emission trends, from 13.0 per cent share of the total (and fourth largest contributor to nickel emissions) in 1990. Emissions in 2016 are reported as not occurring due to the closure of a foundry in the sector which was the last remaining source of emissions in the sector.




Figure 2.16 Emission Trend for Nickel 1990–2016

2.5.5 Selenium (Se)

National total emission estimates of Se have decreased by 59.5 per cent, from 8.76 t in 1990 to 3.55 t in 2016 (Figure 2.17). The main contributor to the trend has been fuel combustion in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors which combined accounted for 53.2 per cent of selenium emissions in 2016, having decreased by 72.2 per cent since 1990. The second largest contributor to the trend is the Public Electricity and Heat Production (1.A.1.a) sector and in 2016 it accounted for 44.7 per cent share of total selenium emissions. Emissions from this sector have decreased by 12.7 per cent from their 1990 level of 1.82 t to 1.59 t in 2016 due to the replacement of older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.

Transport (1A3) sector has been increasing in the time series and with its 1.6 per cent share of the national total in 2016 it has more than doubled in the trend compared to 1990. Emissions from Manufacturing Industries and Construction accounted for 0.4 per cent of the total, having decreased by 31.5 per cent from 1990. The remainder of the selenium emissions arise from combustion in Agriculture/Forestry/Fishing (1A4c) sector with its 0.1 per cent share of the national total in 2015. Glass production (2A3) under Industrial Processes and Product Use (IPPU) sector used to be an important contributor to the selenium emissions trend throughout the 1990s and up to 2002, accounting for an average of 1.4 per cent of the national total. This is no longer a contributor to the trend following closure of all glass production plants between 2002 and 2009. Metal Production (2C) accounted on average for 0.1 per cent of selenium emissions up to 2001 when closure of Ireland's only steel plant reduced emissions to negligible levels.




Figure 2.17 Emission Trend for Selenium 1990–2016

2.5.6 Zinc (Zn)

National total emissions of Zn amounted to 53.66 t in 1990 and have fallen by 63.9 per cent to 19.34 t in 2016. In the 1990–2001 period, the main determinant of the trend in zinc emissions was Metal Production (2C), accounting on average for 55.2 per cent of national total emissions throughout that period (Figure 2.18). However, following the closure of Ireland's only steel plant in 2001, emissions from this source sector are now almost negligible and limited to relatively small IPPC-licensed foundries and galvanising plants. The main determinant for the trend since 2002 has been the Transport (1A3) sector (41.4 per cent of national total zinc emissions in 2016) specifically Road Transport (1A3b) sub-sector has increased significantly since 1990 as a result of the increase in the number of vehicles on Irish roads. As a result, emissions from Transport sector have increased substantially (108.3 per cent), from 3.85 t in 1990 to 8.02 t in 2016.

The second largest source is emissions from combustion in the combined sectors of Residential (1A4b) and Commercial/Institutional (1A4a) accounted for 24.9 per cent of the national total in 2016. However, reduced use of coal and peat through part replacement with natural gas and gasoil has resulted in the reduction of zinc emissions from these sectors by 65.3 per cent, from 13.88 t in 1990 to 4.81 t in 2016. Emissions in Manufacturing Industries and Construction (1A2) have increased by 40.8 per cent from 3.37 t in 1990 to 4.74 t and contributed 24.5 per cent to national total zinc emissions in 2016. Emissions from combustion in Public Electricity and Heat Production (1A1a) emissions have decreased by 42.9 per cent since 1990 and accounted for a 8.3 per cent share of national total in 2016. Similar to other pollutants the reduction is due the replacement of older less-efficient generation plants, reductions in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy. The remainder of the zinc emissions arise from combustion in Agriculture/Forestry/Fishing (1A4c) sector with its 0.9 per cent share of the national total in 2016. Clinical Waste Incineration (5C) accounted for 0.1 per cent of the national total in 1990 however, following an outright ban on incineration of clinical waste in hospitals in 1997, emission from 5C are no longer a source.




Figure 2.18 Emission Trend for Zinc 1990–2016

2.6 Persistent Organic Pollutants (POPs)

As part of Ireland's emission inventory improvement programme, the inventory agency tendered a project in 2007 to develop an inventory of persistent organic pollutants in Ireland. The project report (AEA/CTC, 2008) provides detailed information in relation to the methodological choice and activity data for the diverse range of sources that give rise to emissions of POPs within Ireland. For the purposes of identifying the major contributors to the trend in PCDD/F, PCBs, HCB and PAHs, some of this information is provided in the following sections. The approach was updated for the 2018 submission using the EFs provided in the EMEP/EEA Inventory guidebook (2016).

2.6.1 Dioxins and Furans (PCDD/F)

Dioxin and furan emission levels decreased from 62.43 g I-TEQ in 1990 to 21.04 g I-TEQ in 2016 (66.3 per cent reduction on 1990 levels). The main contributors to the trend are the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined with a 64.1 per cent share of total emissions in 2016; however, emissions from these sectors have reduced (by 71.2 per cent) from 46.79 g I-TEQ in 1990 to 13.48 g I-TEQ in 2016 (Figure 2.19).

The second largest contributor to the trend is the Waste sector which is dominated by emissions from the Other Waste (5E) sector, where building and vehicle fires and residential burning of waste are the emission sources. Total emissions from the waste sector accounted for 13.3 per cent (2.80 g I-TEQ) of national total emissions in 2016, a reduction of 68.0 per cent on 1990 levels (8.75 g I-TEQ). Emissions from the sector peaked in 2003 (8.92 g I-TEQ) due to an increase in the assumed combustion of household waste that remains unaccounted for in national statistics. The introduction of unleaded petrol and technological improvements in road vehicles partly offset increased numbers of vehicles in the national fleet, however there is a 30.2 per cent increase in emissions in Transport (1A3), from 1.04 g I-TEQ in 1990 to 1.36 I-TEQ (6.4 per cent of total emissions) in 2016.

Process emissions from the manufacture of cement (2A1) in Industrial Processes and Product Use sector continue to be an important source of PCDD/F emissions in Ireland, accounting for 7.1 per cent of national emissions in 2016 (1.49 g I-TEQ), a 54.8 per cent reduction on 1990 levels (3.30 g I-TEQ). Combustion in Agriculture/Forestry/Fishing (1A4c)

sector accounted for 6.6 per cent of emissions in 2016 (1.38 g I-TEQ) a decrease of 29.7 per cent on the 1990 emissions (1.96 g I-TEQ).

Public Electricity and Heat Production (1A1a) emissions have decreased by 9.3 per cent from their 1990 level of 0.58 g I-TEQ to 0.53 g I-TEQ and are responsible for a 2.5 per cent share of national total in 2016 due to the replacement of older less-efficient generation plants, reduction in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy.




Figure 2.19 Emission Trend for Dioxins and Furans 1990–2016

2.6.2 Hexachlorobenzene (HCB)

Figure 2.20 outlines the trend in hexachlorobenzene emissions in Ireland across the 1990–2016 time series. The graph indicates that HCB emissions from Secondary Aluminium Processing (2C) which, for the period up to and including 1996, dominated the inventory with a contribution of 40 kg per year and is no longer a source of HCB emissions within Ireland due to the banning of hexachloroethane (HCE)-based cover gas use (HCB was present as a contaminant in such cover gases). Emissions since 1997 are more relevant to trend analysis up to 2016.

There is very limited information on the release of HCB to air for most source sectors in Ireland. The main source is the use of contaminated pesticide in agricultural practices significantly increasing up to 2003 (1.27 kg), reaching its peak in 2004 (1.69 kg) and decreasing to 1.15 kg and 68.6 per cent share of total emissions in 2016, a 6.2-fold increase since 1990 (0.19 kg). The Public Electricity and Heat Production (1A1a) sector was the second largest source in 2016 accounting for 0.46 kg and a 27.7 per cent share of the national total, having decreased by 5.8 per cent between 1990 and 2016.

Incineration of hazardous and clinical wastes and crematoria included in the waste incineration (5C) accounted for 0.3 per cent share of national emissions in 2016, a decrease of 92.7 per cent since 1990. The remainder of the HCB emissions arise mainly from combustion in the Residential (1A4b) and Commercial/Institutional (1A4a), Manufacturing Industries and Construction (1A2) and Agriculture/Forestry/Fishing (1A4c) sectors, which

combined are presented as “Other” NFR sectors in Figure 2.20 and account for 2.9 per cent of the total in 2016.




Figure 2.20 Emission Trend for Hexachlorobenzene 1990–2016

2.6.3 Polychlorinated Biphenyls (PCBs)

Estimated national total emissions of polychlorinated biphenyls have decreased by 69.6 per cent from 40.73 kg in 1990 to 12.39 kg in 2016. Emissions peaked in 2003 (71.62 kg) (Figure 2.21). Industrial Processes (which includes Cement Production (2A1) and Other Production, consumption, storage, transportation or handling of bulk products (2L)) is the main contributor to the trend in PCB emissions in Ireland and were responsible for 41.8 per cent (5.18 kg) of the national total in 2016. Other Waste (5E) is the second largest contributor to the trend in PCB emissions in Ireland. In 2016, the emissions from the Waste sector were 3.39 kg (27.4 per cent of national total emissions), a decrease of 78.1 per cent compared to 1990 (15.50 kg). Of particular note for PCB emissions is the contribution of the NFR Sector 2L (Other Production, consumption, storage, transportation or handling of bulk products), which in Ireland’s inventory covers PCB use as dielectric fluid in electrical equipment such as transformers and capacitors. However, through the introduction of Hazardous Waste Management Plans and the Waste Electrical and Electronic Equipment (WEEE) Regulations, emissions since 2006 have been decreasing in general. Emissions from the Waste sector peaked in 2003 (59.93 kg) due to an estimated increase in the quantity of household waste that remains unaccounted for in national statistics and which is assumed to be burned. Combined emissions from the Residential (1A4b) and Commercial/Institutional (1A4a) sectors accounted for 21.6 per cent of the national total in 2016 (2.67 kg), a 72.8 per cent decrease from those estimated for 1990 (9.85 kg).

Public Electricity and Heat Production (1A1a) emissions were almost negligible in 2016. Emissions in this sector have reduced by 97.7 per cent since 1990 due to the replacement of older less-efficient generation plants, a reduction in the quantities of coal combusted and fuel switching from oil to natural gas and wind energy. Increases in cement production have led to increases in fuel combustion based emissions in Manufacturing Industries and Construction (1A2) up until 2005. This source category’s emissions have decreased by 49.9 per cent between 1990 (1.57 kg) and 2016 (0.78 kg), with a contribution of 6.3 per cent to the national total in 2016.




Figure 2.21 Emission Trend for Polychlorinated Biphenyls 1990–2016

2.6.4 Polycyclic Aromatic Hydrocarbons (PAHs)

For the purposes of this report, total PAHs in the form of the sum of emissions of benzo[a]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene, and indeno[1,2,3-cd]pyrene are presented in Figure 2.22. All together the emissions from these combined four pollutants decreased by 69.6 per cent between 1990 (48.84 t) and 2016 (14.86 t). For all four PAHs, the main source sectors are the same. The inventories are dominated by emissions from combustion in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors. In the Residential (1A4b) sector (main driver of the trend), the lack of combustion controls or abatement, together with relatively low-temperature combustion conditions, leads to high emissions of PAHs. Even though national total emissions of PAHs have generally declined across the time series, the Residential (1A4b) and Commercial/Institutional (1A4a) sectors combined accounted for 88.9 per cent of the national total in 2016, having decreased by 71.7 per cent from 46.62 t in 1990 to 13.22 t in 2016. The decline in emission levels is due primarily to the decline in the use of coal and sod peat for residential space heating, as reported in the National Energy Balance.

Combustion emissions in Manufacturing Industries and Construction (1A2) sector accounted for 8.5 per cent of the total in 2016 (1.26 t), a decrease of 36.9 per cent since 1990 (2.00 t). Emissions from Transport (1A3) have more than doubled (an increase of 160.2 per cent) to 0.30 t representing a 2.0 per cent share of the national total in 2016 for all four PAHs as compared to 0.2 per cent 1990 (0.12 t). Other Waste (5E) is the main driver in the Waste sector resulting in the Waste sector accounting for 0.2 per cent in the total PAHs emissions in 2016, a 22.0 per cent reduction since 1990.




Figure 2.22 Emission Trend for Polycyclic Aromatic Hydrocarbons 1990–2016

Chapter Three

Energy

3.1 Overview of the Energy (NFR 1) Sector

The Energy sector covers combustion and fugitive sources of emissions associated with the production, transport, conversion and use of fossil fuels. Emissions from combustion in this sector account for the bulk of total national emissions for the majority of substances covered in this IIR. Estimates of the various pollutants are included for all emission sources that occur in the country and the required level of disaggregation is achieved for sufficiently detailed completion of the NFR tables.

Annual energy balance sheets published by Sustainable Energy Authority Ireland (SEAI) are the principal source of activity data for computing the emissions in the Energy sector. Ireland's energy statistics are compiled using a combination of top-down and bottom-up methods and the annual energy balances have undergone major improvement over recent years to take account of emission inventory requirements and more harmonised reporting to Eurostat and the International Energy Agency (IEA). The annual submission of up-to-date energy balances from SEAI to the inventory agency is one of the primary data inputs covered by a MoU in Ireland's national system (Chapter One). A fully consistent set of energy balance sheets for the years 1990–2016 underlies the time-series estimates of emissions for *Energy* in this submission. The 2016 energy balance is provided in Annex B.

Substantial plant-level fuel-use data are also available for many important categories in the Energy sector, especially for more recent years, which allows bottom-up estimates to be derived for some pollutants using Tier 3 methods. These data are obtained through direct arrangements with the operators of certain plants through their returns under the Large Combustion Plant (LCP) Directive and the EU ETS and under environmental reporting related to their IPPC permits.

The emissions of SO₂ from fuel combustion are determined from the fuel properties, and fully representative emission factors are readily obtained for the fossil fuels used in most emission categories in Ireland. In general, other pollutants emitted from combustion sources are heavily reliant on emission factors from non-national sources. For all other pollutants, the emission factors are taken from the inventory guidebook (EMEP/EEA, 2009, 2016). The Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP), which is aimed at supporting national experts in reporting PM emission inventories, serves as the reference for emission factors in the category 1A3a for the different forms of Particulate Matter (PM). PM emission factors for other sectors are derived from the Inventory Guidebook (EMEP/EEA, 2016). In the past, emission inventories for heavy metals and POPs received little attention in Ireland and special studies were necessary to compile national emission estimates for reporting purposes. The separate detailed studies on emissions of heavy metals and POPs in Ireland were undertaken by consultants (AEA/CTC, 2008; Netcen/CTC, 2006) and they made use of the best available emission factors for many emission sources, with strong dependence on UK information sources. The results from these studies were the basis for developing time-series emissions of heavy metals and POPs in the Energy sector in previous submissions. In this submission,

many of the EFs for heavy metals and POPs are from the Inventory Guidebook (EMEP/EEA, 2016).

3.2 Public Electricity and Heat Production (NFR 1A1a)

The production of electricity and heat from fossil fuels has traditionally been the most important source of key pollutants such as SO₂ and NO_x in most countries. Approximately 76.1 per cent of electricity production in Ireland (SEAI, 2016) is dependent on fossil fuels and Category 1A1a therefore remains one of the major emission categories. Emissions of SO₂, NO_x, CO, three particulate matter pollutants, BC, all nine heavy metals and POPs have decreased more or less significantly since 1990. NMVOCs were the only emissions that have increased in the time series due to the increase use of natural gas. The level of emissions in Sector 1A1a depends heavily on the mix of fossil fuels and renewables used for electricity production. In 1990, heavy fuel oil (HFO), coal, peat and natural gas were the principal fuels used. The use of HFO, coal and peat declined as natural gas became the preferred fuel during latter years. Incineration emissions from Ireland's only waste-to-energy plant, however small, have been increasing since its operation started in 2011.

3.2.1 Emissions of Sulphur Dioxide and Nitrogen Oxides

Until 2000, the Electricity Supply Board (ESB) operated all public electricity power plants in Ireland. After 2000, several new gas-fired plants and one peat-fired plant were built by other operators, while the ESB replaced old peat-burning stations with new stations also burning peat and has been engaged in a major retrofit and improvement programme for plants in general. The shift to natural gas and the use of low-sulphur coal, combined with a decline in the sulphur content of fuel oil, have reduced SO₂ emissions by 95.0 per cent from 103.04 kt in 1990 to 3.72 kt in 2016. The sector's contribution to the overall SO₂ emissions in 2016 was 27.0 per cent as opposed to 56.1 per cent share in 1990. Incineration of non-renewable waste in MSW accounted for 0.7 per cent of SO_x emissions from power generation in 2016.

At the same time, the changed fuel mix, together with the application of extensive NO_x emission control technology and the more modern plants, has decreased NO_x emissions by 82.7 per cent from 46.37 kt (and 26.0 per cent share of total NO_x emissions) in 1990 to 8.00 kt (and 7.5 per cent share of the total) in 2016. In comparison to the year 2015, NO_x emissions decreased by 16.1 per cent in 2016, and SO₂ emissions continued to decline in 2016 by 28.8 per cent due to increase use of natural gas. The ESB has supplied estimates of SO₂ and NO_x on a plant-by-plant basis to the inventory agency for all years since 1990, mainly for the purpose of compiling SO₂ and NO_x inventories under the LCP Directive. The emissions for power plants operated by other companies are obtained either directly from their LCP, AER or PRTR submissions or they can be estimated by the inventory agency from fuel data made available under the ETS. Incineration of non-renewable waste in MSW accounted for 1.3 per cent of NO_x emissions from power generation in 2016.




Figure 3.1 Nitrogen Oxide Implied Emission Factors for Category 1A1a

The weighted average emission factors of SO₂ and NO_x per fuel type (coal, peat, oil, natural gas, biomass and waste (MSW)) in Category 1A1a are given in Figures 3.1 and 3.2 as implied emission factors (IEFs) to illustrate the level of decrease due to the factors mentioned above. The SO₂ emission factors reflect the sulphur content and net calorific value of the fuels used in the particular year and they account for sulphur retention levels in the fuel ash of 5 per cent and 10 per cent for coal and peat, respectively. The NO_x emission factors (apart from landfill gas) are compiled from plant-level estimates that are determined from measurement, unit load factor and plant performance. Emission factors for landfill gas are default values as per the revised Inventory Guidebook (EMEP/EEA, 2016).




Figure 3.2 Sulphur Dioxide Implied Emission Factors for Category 1A1a

3.2.2 Emissions of Other Substances

The emissions of all substances other than SO₂ and NO_x in Category 1A1a are estimated by the inventory agency using the fuel-use energy data given by the national energy balance and appropriate emission factors taken from the Inventory Guidebook (EMEP/EEA, 2016) and plant-specific factors for non-renewable wastes from waste incineration. The emission factors for 2016 (split by pollutants and fuel type) and their sources are listed in Table C.1 of Annex C.

3.3 Petroleum Refining (NFR 1A1b)

Emissions from fuel combustion at one small oil refinery in Ireland are estimated in this source category. Detailed information on the fuels used in different parts of the refinery in recent years is available through the company's AER, PRTR and ETS submissions. This allows for the selection of appropriate emission factors from national data and from international literature sources recognised as being fit for purpose by the emissions inventory community. In the case of heavy metals and POPs, the estimates are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2016). The emission factors for 2015 and their sources are listed in Table C.2 of Annex C.

3.4 Manufacture of Solid Fuels and Other Energy Industries (NFR 1A1c)

Emissions from this source category refer to combustion emissions from the production of peat briquettes from milled peat at two plants, one natural gas production platform and one new natural gas refinery in Ireland. As in the case of the oil refinery, the energy balance fuel-use data are supplemented by information reported by the plants under the ETS, which again allows for the selection of appropriate emission factors using national data and high quality international sources. The estimates for heavy metals and POPs are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2016). The emission factors for 2016 and their sources are listed in Table C.3 of Annex C.

3.5 Manufacturing Industries and Construction (NFR 1A2)

This category covers emissions from combustion in manufacturing industries and construction activities. Category 1A2 is split into the following seven subcategories:

- | | |
|---------|--|
| 1.A.2.a | Iron and Steel |
| 1.A.2.b | Non-Ferrous Metals |
| 1.A.2.c | Chemicals |
| 1.A.2.d | Pulp, Paper and Print |
| 1.A.2.e | Food Processing, Beverages and Tobacco |
| 1.A.2.f | Non-metallic minerals |
| 1.A.2.g | Other Industry |

Where it is possible to separate process emissions from those associated with fuel use, the process emissions associated with these industrial groups are reported in the Industrial Processes (NFR 2) sector (Chapter Four). The relevant process emissions in Ireland are largely those related to cement manufacture and some metal industries.

Comparison of the Sustainable Energy Authority of Ireland (SEAI) Energy Balance data with ETS fuel-use data indicates that the combustion activities within the 1.A.2 category are dominated by a limited number of large industrial processing plants. It is assumed that all biomass reported as fuel use within the SEAI Energy Balance is clean untreated wood and wood-processing waste. The ETS data for major wood-processing facilities indicate the use of large-scale biomass boilers, fired using wood biomass, chippings, pulp and wood dust.

EPA contacts indicate that none of these wood-based fuels are pre-treated and hence emission factors applicable to clean wood use in large-scale boilers have been used to estimate POP emissions from these sources.

The Iron and Steel (1A2a) sector was dominated in the 1990s by fuel use and emissions from one electric arc furnace but, since its closure in 2001, the fuel use reported in 2002 in this sector related to a small number of iron foundries, which in total used only a very small amount of gasoil and LPG and since 2003 combustion emissions are not occurring. The process emissions from this industrial activity are described in chapter 4, section 4.4.1. The Non-Ferrous Metal (1A2b) subcategory is dominated by the very significant fuel use (mainly natural gas replacing popular in the 1990s fuel oil) reported at a single large alumina plant, whilst the Chemicals (1A2c) subcategory includes natural gas, kerosene, fuel oil, gasoil and LPG use at large chemical plants. The Food Processing, Beverages and Tobacco (1A2e) sector covers a diverse range of industrial plants, much of which is related to agriculture, with just over 30 installations reporting to the EU ETS and using predominantly natural gas and liquid fuels. The bulk of the fuel use reported in the energy balance under the 1A2f subcategory is accounted for by major cement works, lime producers, a small number of brickworks and fuel use at boiler plant within industries such as the pharmaceutical, glass and tile manufacturing sectors. All other industrial fuel use is reported under subcategory 1A2g. This sector covers a diverse range of manufacturing branches ranging from textile and leather, through machinery, transport equipment, wood products, mining (excluding fuel mining) and quarrying to other manufacturing businesses. Ireland's national Energy statistics do not provide an estimate of fuel used in mobile construction. All emissions associated with the category 1A2g are reported in 1A2gviii and mobile emissions, 1A2gvii, are reported as "included elsewhere" (IE) in 1A2gviii.

The revised and expanded energy balance sheets developed by SEAI incorporate a mapping of industrial fuel use in combustion into the NFR subcategories 1A2a through 1A2g under sector 1A2 Manufacturing Industries and Construction. This facilitates the complete disaggregation of emissions at subcategory level. In addition, information on fuel consumption in 2016 was obtained in respect of a small number of energy-intensive industries (e.g. alumina production and cement manufacture) from their ETS returns, allowing their respective energy use amounts to be reconciled with the breakdown given in the national energy balance. Emissions in subcategories 1A2a through 1A2g are estimated on a top-down basis using disaggregated fuel use from the energy balance and the mix of country-specific and default emission factors as shown in Tables C.4 through C.10 of Annex C. The estimates for heavy metals and POPs are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2016). Emission factors for heavy metals and POPs (including references) for NFR 1A2 (a–g) are shown in Table C.11 of Annex C.

Regarding NMVOC emissions in 1A2gviii, raised in the review, a new methodology was applied in 2017 reporting. In previous submissions, an EF of 300kg/TJ was used for NMVOC, EMEP/EEA 2013, 1.A.2 Combustion in manufacturing, Table 3-5, 'biomass'. In the 2017 and this submission, the fuel used for 3 large plants for all years where data is available (2005-2016, EU ETS), was subtracted from the total fuel use and a Tier 2 EF was used, 12kg/TJ for NMVOC and all other pollutants, EMEP/EEA 2016, 1.A.4 Small Combustion, Table 3-48, 'wood'. The residual amount of biomass fuel still uses the Tier 1 EF.

3.6 Transport (NFR 1A3)

As abatement measures continue to reduce emissions of key pollutants from major stationary combustion sources, transport in general, and road transport in particular, has become more important as a source of atmospheric emissions in many countries. The effects of technological emission controls for passenger cars and other vehicles in Ireland have, to a large extent, been offset by the substantial increases in vehicle numbers, with the

result that major reductions in the emissions of pollutants such as NO_x did not occur until 2007 when the economic situation caused an overall decrease in most emissions evident from 2008. In 2016 NO_x emissions from this sector reached 43.98 kt (41.0 per cent share of total) reflecting a decrease by 33.4 per cent on 66.07 kt emissions (and 37.7 per cent share) in 1990.

Road transport in Ireland is a larger source of NO_x than electricity production, and road traffic also continues to be the major source of CO, BC, Pb, Cr, Cu and Zn, although BC, Pb, Cr, Cu and Zn emissions are very small in absolute terms.

3.6.1 Domestic and International LTOs (NFRs 1A3aii(i) and 1A3ai(i))

As a relatively small island state, aviation emissions are dominated by the international component. Under the LRTAP Convention, only the landing and take-off (LTO) component of emissions for both domestic and international flights is reported in the national total. The cruise component, domestic and international, is reported as a memo item. After the motorway network to Dublin from Galway and Cork was completed around 2008/09 and the upgrades of the Cork to Dublin rail line, domestic air travel was no longer competitive, leading to cessation of specific routes. Almost all flights ceased from Galway airport post 2011.

The fuel consumption associated domestic and international LTOs is estimated using a Tier 3a approach (Table 3.6.2, 2006 IPCC guidelines) based on origin and destination data for domestic air travel provided by the Irish Aviation Authority (IAA), the fuel consumption rates given by the Inventory Guidebook (EMEP/EEA, 2009,2013) appropriate to the type of aircraft concerned and the length of the flights within Ireland. This approach is used for all years from 2004 to 2016 where airport pair data is available.

The inventory agency received 2016 flight data for all Irish airports from the IAA in 2017 completing the time series of data for 2004 to 2016. This data included all flights, domestic and international, on an origin and destination basis and by aircraft type for over 25 different Irish origin airports. For the years 1990 to 2003, the number of flights for each airport was estimated based on domestic passenger and aircraft movement statistics as well as the relationship between all Irish airports and Dublin airport which is the principal destination of all civil flights. Figures 3.3 and 3.4 and Table C.13 of Annex C present the number of LTOs, domestic and international, from Irish airports for all years from 1990 to 2016.




Figure 3.3 Number of Domestic LTOs from Irish airports 1990-2016

For data handling purposes, the inventory agency aggregated approximately 15 small regional airport/aerodrome pairs to “Other” which account for approximately 3.2 per cent of all domestic flights along with nine Irish airports which account for the remaining 96.8 per cent of all domestic flights in 2016. Table C.12 of Annex C outlines the distance between the airport pairs in nautical miles (nm) used in estimating fuel used in the cruise phase.




Figure 3.4 Number of International LTOs from Irish airports 1990-2016

The tier 3a methodology estimates both LTO and cruise emissions for domestic flights and LTOs for international flights based on origin and destination, flight distances and by aircraft type. The fuel used in the cruise element of international flights is derived by the following equation;

$$Int\ Cruise_{(fuel)} = Total_{(fuel)} - (Int\ LTO_{(fuel)} + Domestic\ LTO_{(fuel)} + Domestic\ Cruise_{(fuel)})$$

The inventory agency estimated fuel consumption for the LTO and cruise phases of each domestic flight based on 37 aircraft types using fuel consumption emission factors from the Inventory Guidebook (EMEP/EEA, 2009). The fuel consumption for international LTOs is based on 136 aircraft types using fuel consumption emission factors from the Inventory Guidebook (EMEP/EEA, 2009). Tables C.14 and C.15 of Annex C outline the emission factors used for domestic and international LTOs for fuel, NO_x, HC and CO by aircraft type. Tables C.16 and C.17 of Annex C present implied emission factors (IEF) for fuel consumption, NO_x, HC and CO used in the LTO cruise phase of flights weighted by number of flights per airport.

3.6.2 Road Transport (NFR 1A3b)

The emissions of nine well-known pollutants (SO₂, NO_x, NMVOCs, NH₃, CO, TSP, PM₁₀, PM_{2.5}, BC) as well as seven heavy metals (Pb, Cd, Cu, Cr, Ni, Se, Zn) and POPs (Dioxins and Furans, PAHs) reported under sub-category 1.A.3.b Road Transport are estimated directly from the COPERT 5 model (Pastramas N. et al., 2014) developed within the CORINAIR programme for estimating a range of emissions from this important source.

The national total emissions in all cases are determined by the quantity of fuel sold in the country, as given by the energy balance. Approximately 65.82 petajoule (PJ) equivalent fuel energy from petrol, diesel, liquefied petroleum gas (LPG) and biofuels was consumed by road transport in Ireland in 1990. Consumption went up to 190.86 PJ in 2007 followed by a sharp decline to 145.88 PJ in 2012 and subsequently increasing to 165.46 PJ in 2016. The energy share in biofuel has continued to increase since of its introduction in 2005 and was 5.36 PJ in 2015 but decreased to 4.96 PJ in 2016. It is known that significant proportions, e.g. 1.0 per cent of petrol and 13.2 per cent of diesel in 2016 of sold in Ireland are consumed outside the country (referred to as fuel tourism) and therefore separate estimates of emissions are produced based on fuel amounts used within Ireland. The use of compressed natural gas (CNG) in road transport is recorded in national energy statistics for 2014 and 2015, however, the emissions were not estimated as the fuel amounts were considered negligible. For some countries, including Ireland, the national totals determined by fuels used are the basis for assessment of their performance in relation to relevant Protocols under CLRTAP. Annex A.3 outlines the methodology used to estimate the quantities of automotive fuels used in Ireland and includes the adjusted annual emissions for 1987 and the years 1990–2016.

The emissions of SO₂ for road transportation are computed from the amounts of petrol and diesel used by motor vehicles, as reported in the energy balance, and the sulphur content of the fuels. For the other substances, the COPERT 5 model estimates emissions based on distance travelled using a detailed bottom-up approach (Tier 3) that accounts for such factors as fuel type, fuel consumption, engine capacity, driving speed and a wide range of applicable technological emission controls that may be applied across the different vehicle categories according to the age of vehicles.

The architecture of the COPERT 5 has changed in comparison to the COPERT 4 v11 methodology used in the previous submission; for feeding input data, estimation of emission and output file structure. The emissions are now estimated based on the energy rather than the fuel use, and thus several input types were required to adjust to adopt the change. However, the emission factors and methodology remain the same in general between the latest COPERT IV and COPERT 5 versions.

A total of 180 vehicle categories are determined by these variables in COPERT 5. Figures C.1 through C.4 of Annex C show the trend in emission factors for NO_x, SO₂, NH₃ and

NMVOCs per fuel type for road transportation over the period 1990–2016 determined by the COPERT variables for the vehicle fleet in Ireland. The methodologies for estimation of PM_{2.5}, PM₁₀ and TSP are from the COPERT output of non-exhaust emission that includes brake and tyre wear emissions and road abrasion emissions for PM_{2.5}, PM₁₀ and TSP were calculated and reported using emission factors from the Inventory Guidebook (EMEP/EEA 2016) and fuel adjusted mileage data from COPERT 5 at the vehicle category level.

The primary model inputs for each year are the populations of vehicles in the relevant categories, their annual kilometres of travel in three selected speed classes, total fuel amounts and the fuel specifications. The numbers of vehicles are taken from annual bulletins of vehicle and driver statistics (DTTAS, 2016) and these are allocated to the different control technologies (Euro I, Euro II, etc.) based on their age and the application dates of the controls in Ireland. Information to assign values of annual kilometres of travel for the three speed classes (corresponding to urban, rural and highway driving) used for the individual vehicle categories is taken from the National Roads Authority (NRA) and odometer records from the National Car Testing (NCT) and Commercial Vehicle Roadworthiness Test (CVRT) Service. Before the emissions are estimated in COPERT 5, fuel balancing is undertaken in the model using inbuilt consumption rates for the different vehicle categories to ensure that the total fuel amounts calculated from annual kilometres and consumption rates for all vehicle categories match the input fuel quantities (the fuel sold as given in the energy balance or the fuel used as determined by Annex A.3) and therefore that emissions relate to the relevant total fuel amount. This fuel balancing may involve some adjustment to kilometres travelled or to the shares of total kilometres in the different road classes.

Lubricant emissions from the road transport sector were estimated using COPERT 5 where the lubricant used was considered as an input in the model. Thus, emissions from lubricant for all engines including 2 and 4 stroke engines were included in COPERT output and reported under relevant categories in the NFR.

3.6.3 Railways (NFR 1A3c), National Navigation (NFR 1A3dii) and Other Transportation (NFR 1A3e)

The emissions under sub-categories 1A3c Railways and 1A3d National Navigation are calculated from the amounts of fuel used by these activities and the country-specific SO_x emission factors and default Inventory Guidebook (EMEP/EEA, 2016) emission factors for oil. No solid fuels have been used in railways since 1970. All emission factors for railways and navigation are given in Table C.20 of Annex C. These are minor sources of emissions in Ireland.

The emissions reported in sub-category 1A3e Other Transportation refer to the use of natural gas in pipeline compressor stations and emission factors (country-specific and default values) for this sub-category are given in Table C.21 of Annex C. The fuel use is estimated as the difference between the value given for natural gas under own use/losses in the energy balance sheets (Annex B) and the amount of gas estimated to be lost from the distribution network.

3.7 Other Sectors (NFR 1A4)

The NFR Subcategory 1A4 Other Sectors covers combustion sources in the Commercial/Institutional (1A4a), the Residential (1A4b), and Agriculture/Forestry/Fishing (1A4c) sectors. The Residential sector remains the most important source of emissions for most substances in this subcategory in Ireland, while Agriculture/Forestry/Fishing is a major contributor to NO_x emissions, largely because of the influence of agricultural (off-road)

machinery. The activity data for all 1A4 subcategories are taken directly from the energy balance. As there are no fuels allocated to the sector 1A4aⁱⁱ Commercial/Institutional: Mobile or 1A4bⁱⁱ Residential: Household and Gardening (Mobile) in national energy statistics, it is accounted for in category 1A3b Road Transport where gasoline fuel is fully accounted. The emission factors for sub-category 1A4 Other Sectors are given in Tables C.22 through C.28 of Annex C.

3.7.1 Agriculture/Forestry/Fishing (NFR 1A4ci and ii)

This subcategory covers both stationary combustion (1A4ci) and mobile combustion (1A4cii) in Agriculture. The energy balance does not currently provide information on the end use of gasoil in agriculture. However, based on information received from agricultural experts, 90 per cent of total gasoil in the sector is assigned to mobile machinery with the remainder assigned to stationary combustion. For both stationary and mobile combustion, the emission factors used are presented in Table C.26 of Annex C.

Ireland utilises the Tier 2 approach outlined in the Inventory Guidebook (EMEP/EEA 2013) to estimate emissions of NO_x, NMVOC, CO, PM₁₀, PM_{2.5} and TSP from mobile combustion in this sector. The Tier 2 emission factors are based on data from the Danish Inventory (Winther and Nielsen, 2006). The emission factors are grouped according to the EU emission legislation stages, and three additional layers are added to cover the emissions from engines prior to the first EU legislation stages. The country-specific SO_x emission factors and default Inventory Guidebook (EMEP/EEA, 2016) emission factors for NH₃ and BC are used for this sub-category. The implied emission factors are presented in Table C.27 of Annex C. Emission Factors for Heavy Metals and POPs are based on emission factors from the Inventory Guidebook (EMEP/EEA, 2009, 2016). The category 1A4ci Agriculture/Forestry/Fishing: Stationary, the NO_x Tier 1 liquid fuel emission factors from the Inventory Guidebook (EMEP/EEA, 2016) are an average of Tier 2 EFs and are not considered representative for Ireland considering one of the liquid fuel values for NO_x is 942g/GJ based on reciprocating engines which are not occurring in Ireland. The Tier 1 value of 100g/GJ (EMEP/EEA, 2009) used by Ireland is still valid, as all the guidebook editions refer to Chapter B216 originally. A generic emission factor of 100 g/GJ was applied for all liquid fuels for sectors 1A2a to 1A2g, which was obtained from “Table 3-4: Tier 1 emission factors for 1.A.2 combustion in industry using liquid fuels”, for “Other Liquid Fuels” in page 16, 1.A.2 Manufacturing industries and construction (combustion), EMEP/EEA emission inventory guidebook 2009. Reference EMEP/CORINAIR B216. Table 1-1 of the Inventory Guidebook (EMEP/EEA, 2016) refers to the appropriate chapters for emission factors, either 1A1, 1A2 or 1A4. Very few emission factors were used from Chapter 1A2 as Ireland does not have these types of industry.

3.7.2 Agriculture/Forestry/Fishing: National Fishing (NFR 1A4ciii)

Emissions from this sub-category were reported for the first time in the 2013 submission. The national energy balance now includes marine diesel used in national fishing for the all years from 1990 to 2015. The emission factors used for this sub-category are country-specific (SO_x) and Inventory Guidebook (EMEP/EEA, 2013) default values for all other pollutants and are presented in Table C.28 of Annex C.

3.8 Fugitive emissions from Solid Fuels (NFR 1B1)

3.8.1 Coal mining and handling (1B1a)

Emissions from the NFR subcategory 1B1a Coal mining were reported for the first time in the 2015 submission. The national energy balance now includes coal mined in the years 1990 to 1995 when the last commercial coal mine was closed in Ireland. The emission factors used and the resulting time series of NMVOC, TSP, PM₁₀ and PM_{2.5} emissions are based on the Inventory Guidebook (EMEP/EEA, 2013) default values and are presented in

Table 3.1. Emissions of PM_{2.5} from the handling and storage of coal other than the domestic production has been estimated for the first time in 2018 using emission factors from Inventory Guidebook (EMEP/EEA, 2016).

Table 3.1. Fugitive Emissions from Coal mining and handling

Activity Data		1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Coal mining	kt	25.00	1.00	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Area of coal storage	ha	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20	31.20
Coal handling	kt	1,867	2,248	2,145	2,130	1,762	1,491	1,165	1,305	1,373	1,745	1,459	1,417	1,695	1,656
Emission Factors															
Coal mining	kg NMVOC/t	0.80	0.80	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal mining	kg TSP/t	0.09	0.09	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal mining	kg PM ₁₀ /t	0.04	0.04	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal mining	kg PM _{2.5} /t	0.01	0.01	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Coal storage	t PM _{2.5} /ha	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
Coal handling	g PM _{2.5} /t	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Emissions															
Coal mining	kt NMVOC	0.020	0.001	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal mining	kt TSP	0.002	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal mining	kt PM ₁₀	0.001	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal mining	kt PM _{2.5}	0.000	0.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Coal storage	kt PM _{2.5}	0.001	0.001	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013
Coal handling	kt PM _{2.5}	0.001	0.001	0.0006	0.0006	0.0005	0.0004	0.0003	0.0004	0.0004	0.0005	0.0004	0.0004	0.0005	0.0005

3.9 Fugitive emissions from Oil and Natural Gas (NFR 1B2)

3.9.1 Oil (NFR 1B2a)

The NFR subcategory 1B2a is an important source of fugitive NMVOC emissions. Emissions of NMVOCs are estimated from two sources, Refining/Storage (1B2aiv) and Distribution of Oil Products (1B2av). Emissions from Refining/Storage are estimated using a Tier 3 methodology using plant specific data for years 2007-2016 from Ireland's only refinery. An implied emission factor of 0.8818 kg/t was used for years before 2007 using an average of 2007-2016 implied emission factors and the crude oil throughput for the refinery. This revision to the methodology has resulted in increased NMVOC emissions in this category.

The emissions from 1B2av, which is a key category, were estimated using a Tier 2 approach for the first time in 2018 using emission factors and abatement efficiencies from the Inventory Guidebook (2016). The Stage I and Stage II abatement controls were applied as appropriate to the implementation of legislation in Ireland. The resulting time series of NMVOC emissions are presented in Table 3.2. Emissions of NMVOCs from these subcategories nearly doubled in the period from 1990 to 2007 in line with the increases in crude oil throughput for the refinery and petrol distribution for the transport sector, but have since decreased due to the economic situation and the implementation of abatement controls in petrol distribution causing an overall decrease in most emissions. A slight increase in emissions in the Refining/Storage source can be seen from 2014 to 2016 due to increased crude oil throughput.

Table 3.2. Fugitive Non-Methane Volatile Organic Compound Emissions

Activity Data		1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Crude to refineries	kt	1,804	2,229	3,278	3,309	3,389	3,272	2,812	2,905	2,949	3,068	2,838	2,752	3,340	3,200
Petrol distribution	kt	885	1,037	1,493	1,711	1,771	1,688	1,537	1,387	1,314	1,195	1,124	1,064	1,009	942
Emission Factors															
Crude to refineries	kg NMVOC/t	0.88	0.88	0.88	0.88	0.87	0.88	0.99	0.92	0.84	0.81	0.86	0.95	0.84	0.85
Petrol distribution	kg NMVOC/t	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
Emissions															
Crude to refineries	kt NMVOC	1.59	1.97	2.89	2.92	2.95	2.87	2.80	2.66	2.49	2.48	2.45	2.62	2.80	2.73
Petrol distribution	kt NMVOC	1.79	2.10	3.30	3.81	3.88	3.69	3.39	3.15	2.97	2.60	2.22	2.12	1.95	1.76
Total emissions	kt NMVOC	3.38	4.06	6.19	6.73	6.83	6.56	6.19	5.80	5.46	5.08	4.67	4.74	4.75	4.50

3.9.2 Venting and flaring (oil, gas, combined oil and gas) (NFR 1B2c)

Regarding the flares at refineries, Ireland explained that the emissions had been included within the category 1A1b Petroleum Refining, as site emissions were calculated based on total metered fuel use.

3.10 Recalculations in the Energy Sector

The results of recalculations for the combustion categories 1A1, 1A2, 1A3 and 1A4 are given in Tables 3.3 through 3.7 below.

Recalculations have been undertaken for the years 1990–2015 in the Energy sector (combustion and fugitive) to account for the following changes. All references to EFs from the EMEP/EAA, 2013 have been updated to EMEP/EEA, 2016 where relevant:

1A1 Energy Industries

- Revised emissions for the oil refinery, 1A1b, for 2014 and 2015 from PRTR data. Revised historical EFs for SO₂ based on reported data from 2000 to 2016.
- Revised emissions data for NO_x and SO₂ as recommended during review, data is now plant specific. Revised the EF for Peat for NMVOC as recommended in the review, 0.80 kg NMVOC/TJ, Table 5.1, Inventory Guidebook (2016).
- Also, included emissions from a new natural gas refinery for 2015 and 2016 and reported emissions from a natural gas offshore platform to 1A1c, previously reported in 1A3e. See Table 3.3.

1A2 Manufacturing Industries and Construction

- Updated EFs from EMEP/EEA 2013 to 2016 except for liquid fuels which are from EMEP/EEA 2009 to be consistent with other air pollutants. See Table 3.4.

1A3 Transport

- Revised methodology used in road transport (1A3b) from COPERT 4v11.3 to COPERT 5. The most significant changes are for HMs for the earlier time series relating to leaded gasoline. See Table 3.5.
- Updated the country specific SO₂ EF for 2015 for Gasoil used in railways (1A3c). See Table 3.6.

1A4 Other Sectors

- Changed liquid fuel EFs for HMs and POPs in commercial/institutional combustion (1A4a) from EMEP/EEA, 2013 to EMEP/EEA, 2009 to be fully consistent with other

air pollutants.

- Changed the NO_x EF for Biomass in Residential combustion (1A4b) from 80 g/GJ to 50 g/GJ, EMEP/EEA, 2013 to 2016.
- Updated the country specific SO₂ EF for 2015 for Gasoil used in agriculture/forestry/fishing (1A4c).
- Changed liquid fuel EFs for HMs and POPs in agriculture/forestry/fishing (1A4c) from EMEP/EEA, 2013 to EMEP/EEA, 2009 to be fully consistent with other air pollutants.
- Revised Tier 2 EFs for air pollutants for agriculture/forestry/fishing (1A4cii) mobile combustion. See Annex C Table C.27.

Table 3.3. Recalculations in Energy NFR 1A1

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
1A1a Public Electricity and Heat Production	NO _x	kt	46.37	41.39	39.72	32.38	29.87	27.33	22.02	13.32	11.52	8.00	10.30	8.57	7.54	9.48
	SO ₂	kt	103.04	91.63	79.87	42.50	36.80	30.84	25.19	15.69	9.46	9.38	9.58	8.69	5.95	5.20
	NM VOC	kt	0.19	0.25	0.36	0.37	0.38	0.38	0.38	0.35	0.37	0.31	0.30	0.27	0.26	0.26
	Pb	t	0.63	0.75	0.78	0.73	0.64	0.57	0.54	0.45	0.44	0.44	0.54	0.47	0.48	0.54
	Cd	t	0.09	0.11	0.12	0.11	0.10	0.08	0.08	0.06	0.06	0.06	0.07	0.06	0.06	0.07
	Hg	t	0.12	0.13	0.13	0.13	0.12	0.11	0.11	0.09	0.09	0.09	0.11	0.09	0.10	0.11
1A1b Petroleum Refining	NO _x	kt	0.47	0.52	0.77	0.94	0.81	0.89	0.77	0.75	0.80	0.63	0.68	0.58	0.54	0.54
	SO ₂	kt	0.75	0.58	0.78	0.91	1.03	1.03	0.98	0.88	0.67	0.90	0.43	0.57	0.57	0.57
	NM VOC	kt	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02
	Pb	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1c Manufacture of Solid Fuels and Other Energy Industries	NO _x	kt	0.08	0.06	0.07	0.10	0.11	0.10	0.11	0.13	0.11	0.08	0.09	0.11	0.09	0.07
	SO ₂	kt	0.24	0.18	0.22	0.29	0.33	0.31	0.33	0.39	0.32	0.25	0.27	0.32	0.26	0.20
	NM VOC	kt	0.08	0.06	0.07	0.10	0.11	0.10	0.11	0.13	0.11	0.08	0.09	0.11	0.09	0.07
	Pb	t	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Submission 2018																
1A1a Public Electricity and Heat Production	NO _x	kt	46.37	41.39	39.72	32.38	29.87	27.33	22.02	13.32	11.52	8.00	10.30	8.57	7.58	9.53
	SO ₂	kt	103.04	91.63	79.87	42.50	36.80	30.84	25.19	15.69	9.46	9.38	9.58	8.69	5.96	5.22
	NM VOC	kt	0.19	0.25	0.36	0.37	0.38	0.38	0.38	0.35	0.37	0.31	0.30	0.27	0.26	0.26
	Pb	t	0.63	0.75	0.78	0.73	0.64	0.57	0.54	0.45	0.44	0.44	0.54	0.47	0.48	0.54
	Cd	t	0.09	0.11	0.12	0.11	0.10	0.08	0.08	0.06	0.06	0.06	0.07	0.06	0.06	0.07
	Hg	t	0.12	0.13	0.13	0.13	0.12	0.11	0.11	0.09	0.09	0.09	0.11	0.09	0.10	0.11
1A1b Petroleum Refining	NO _x	kt	0.47	0.52	0.77	0.94	0.81	0.89	0.77	0.75	0.80	0.63	0.68	0.58	0.54	0.38
	SO ₂	kt	0.48	0.51	0.78	0.91	1.03	1.03	0.98	0.88	0.67	0.90	0.43	0.57	0.01	0.03
	NM VOC	kt	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02
	Pb	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1c Manufacture of Solid Fuels and Other Energy Industries	NO _x	kt	0.16	0.12	0.15	0.29	0.30	0.22	0.37	0.25	0.31	0.21	0.18	0.22	0.22	0.24
	SO ₂	kt	0.13	0.10	0.12	0.16	0.18	0.12	0.04	0.14	0.20	0.25	0.24	0.19	0.03	0.09
	NM VOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Pb	t	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 3.3. Recalculations in Energy NFR 1A1 (continued)

% Change in Emissions			1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A1a Public Electricity and Heat Production	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%	0.6%
	SO ₂	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.3%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
1A1b Petroleum Refining	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-29.4%
	SO ₂	%	-36.8%	-12.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.1%	-0.2%	-0.1%	-98.0%	-95.6%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
1A1c Manufacture of Solid Fuels and Other Energy Industries	NO _x	%	99.1%	99.1%	99.1%	201.0%	175.1%	112.2%	235.0%	95.4%	190.3%	157.6%	99.2%	107.4%	152.9%	266.6%
	SO ₂	%	-45.2%	-45.2%	-45.2%	-45.0%	-45.1%	-62.0%	-86.8%	-63.4%	-38.9%	3.5%	-14.1%	-41.8%	-87.7%	-55.4%
	NM VOC	%	-99.2%	-99.2%	-99.2%	-96.2%	-97.0%	-96.9%	-96.8%	-97.7%	-97.3%	-96.9%	-97.2%	-97.6%	-97.4%	-96.3%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Table 3.4. Recalculations in Energy NFR 1A2

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
1A2 Manufacturing Industries and Construction	NO _x	kt	8.93	8.05	10.53	17.31	16.59	18.45	15.77	10.37	9.96	8.48	10.43	10.55	10.73	10.95
	SO ₂	kt	32.56	38.06	37.56	11.98	8.50	8.96	7.87	5.11	5.86	5.24	4.49	4.87	3.34	2.30
	NM VOC	kt	1.67	1.27	1.77	2.78	2.95	2.91	2.76	2.35	2.47	2.26	2.23	2.28	2.76	2.87
	Pb	t	1.31	0.49	0.77	1.41	1.21	1.22	1.08	0.79	0.80	0.71	0.69	0.62	0.79	0.79
	Cd	t	0.05	0.04	0.06	0.10	0.10	0.10	0.09	0.08	0.09	0.08	0.08	0.08	0.10	0.10
	Hg	t	0.09	0.04	0.06	0.10	0.09	0.09	0.08	0.06	0.06	0.06	0.06	0.06	0.06	0.07
Submission 2018																
1A2 Manufacturing Industries and Construction	NO _x	kt	8.93	8.05	10.53	17.31	16.59	18.42	15.73	10.34	9.93	8.46	10.41	10.53	10.71	10.85
	SO ₂	kt	32.56	38.06	37.56	11.98	8.50	8.96	7.87	5.11	5.86	5.24	4.49	4.87	3.34	2.44
	NM VOC	kt	1.67	1.27	1.77	2.78	2.95	2.90	2.75	2.34	2.46	2.25	2.23	2.28	2.76	2.88
	Pb	t	1.73	1.10	1.51	2.09	1.83	1.82	1.67	1.24	1.24	1.03	0.97	0.90	1.04	1.03
	Cd	t	0.06	0.05	0.08	0.12	0.11	0.11	0.10	0.09	0.10	0.09	0.08	0.09	0.10	0.11
	Hg	t	0.09	0.04	0.06	0.09	0.09	0.09	0.08	0.06	0.06	0.06	0.06	0.05	0.06	0.07
% Change in Emissions																
1A2 Manufacturing Industries and Construction	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.2%	-0.2%	-0.3%	-0.3%	-0.3%	-0.2%	-0.2%	-0.1%	-0.9%
	SO ₂	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	6.4%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.3%	-0.4%	-0.4%	-0.3%	-0.3%	-0.3%	-0.3%	-0.2%	0.6%
	Pb	%	32.4%	123.3%	96.8%	49.0%	50.7%	49.5%	54.3%	57.9%	54.3%	44.9%	40.2%	46.2%	31.4%	30.9%
	Cd	%	15.6%	29.4%	22.3%	12.3%	11.1%	11.5%	12.4%	10.2%	9.0%	7.1%	6.6%	6.5%	4.6%	6.7%
	Hg	%	-0.7%	-2.3%	-1.9%	-1.0%	-1.0%	-1.3%	-1.3%	-1.4%	-1.3%	-1.1%	-1.0%	-1.1%	-0.8%	-1.7%

Table 3.5. Recalculations in Energy NFR 1.A.3 (a & b)*

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
1A3a(i) & (ii) International and Domestic Aviation	NO _x	kt	2.34	2.37	3.61	4.50	5.15	5.42	5.07	4.05	4.01	3.60	3.10	3.47	3.79	4.22
	SO ₂	kt	0.35	0.38	0.59	0.81	0.93	0.98	0.92	0.72	0.74	0.66	0.55	0.64	0.71	0.80
	NM VOC	kt	0.22	0.20	0.28	0.30	0.32	0.35	0.32	0.26	0.24	0.24	0.20	0.21	0.22	0.24
1A3b Road Transportation	NO _x	kt	54.00	54.92	59.53	54.45	54.37	54.36	48.74	42.42	38.23	36.74	35.03	36.03	36.59	37.51
	SO ₂	kt	5.39	5.14	1.66	0.54	0.54	0.35	0.14	0.05	0.04	0.04	0.04	0.04	0.04	0.05
	NM VOC	kt	34.66	33.04	23.16	15.25	13.83	12.60	11.31	9.81	8.51	7.69	6.84	6.28	5.74	5.36
	NH ₃	kt	0.04	0.41	1.66	2.47	2.29	2.14	2.07	1.79	1.53	1.36	1.18	1.06	0.96	0.87
	PM _{2.5}	kt	2.40	2.71	3.71	3.18	3.20	3.23	2.92	2.61	2.39	2.24	2.09	2.06	1.96	1.97
	Pb	t	109.16	67.17	11.37	12.85	13.13	13.58	13.71	12.82	12.10	11.53	10.66	10.21	9.73	9.62
	Cd	t	0.02	0.02	0.04	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04
	Hg	t	0.01	0.01	0.02	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02
Submission 2018																
1A3a(i) & (ii) International and Domestic Aviation	NO _x	kt	2.33	2.36	3.60	4.50	5.15	5.38	5.07	4.05	4.01	3.60	3.10	3.47	3.79	4.22
	SO ₂	kt	0.35	0.38	0.59	0.81	0.93	0.98	0.92	0.72	0.74	0.66	0.55	0.64	0.71	0.80
	NM VOC	kt	0.22	0.20	0.28	0.30	0.32	0.35	0.32	0.26	0.24	0.24	0.20	0.21	0.22	0.24
1A3b Road Transportation	NO _x	kt	54.70	55.41	60.24	55.19	55.14	55.06	49.21	42.87	38.53	37.24	35.68	36.83	38.25	39.36
	SO ₂	kt	5.37	5.13	1.66	0.54	0.54	0.35	0.14	0.05	0.04	0.04	0.04	0.04	0.04	0.05
	NM VOC	kt	33.15	31.57	23.01	15.39	13.98	12.72	11.65	10.09	8.72	7.88	7.00	6.45	5.91	5.42
	NH ₃	kt	0.04	0.41	1.69	2.51	2.34	2.17	2.09	1.81	1.54	1.36	1.18	1.06	0.96	0.87
	PM _{2.5}	kt	2.43	2.74	3.74	3.28	3.30	3.32	2.99	2.67	2.44	2.29	2.13	2.12	2.03	1.99
	Pb	t	143.95	87.44	11.53	12.86	13.14	13.60	13.76	12.88	12.18	11.61	10.74	10.30	9.83	9.71
	Cd	t	0.02	0.02	0.04	0.05	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.05
	Hg	t	0.01	0.01	0.02	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02
% Change in Emissions																
1A3a(i) & (ii) International and Domestic Aviation	NO _x	%	-0.3%	-0.2%	-0.1%	0.0%	0.0%	-0.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO ₂	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NM VOC	%	0.1%	0.2%	0.1%	0.0%	0.0%	0.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
1A3b Road Transportation	NO _x	%	1.3%	0.9%	1.2%	1.4%	1.4%	1.3%	1.0%	1.1%	0.8%	1.4%	1.8%	2.2%	4.5%	5.0%
	SO ₂	%	-0.2%	-0.2%	-0.1%	-0.1%	-0.1%	0.0%	0.0%	-0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.2%
	NM VOC	%	-4.3%	-4.5%	-0.7%	0.9%	1.1%	1.0%	3.0%	2.9%	2.5%	2.5%	2.4%	2.6%	2.9%	1.2%
	NH ₃	%	1.3%	1.5%	1.9%	1.8%	1.8%	1.7%	1.1%	0.9%	0.4%	0.4%	0.3%	0.1%	-0.1%	0.3%
	PM _{2.5}	%	1.5%	1.1%	0.8%	3.2%	3.1%	2.8%	2.3%	2.3%	1.9%	1.9%	1.7%	2.7%	3.8%	0.9%
	Pb	%	31.9%	30.2%	1.3%	0.1%	0.1%	0.1%	0.4%	0.5%	0.7%	0.7%	0.7%	0.9%	1.0%	1.0%
	Cd	%	5.1%	5.5%	3.5%	4.0%	3.4%	3.0%	5.3%	7.5%	8.7%	9.0%	9.7%	9.5%	9.5%	8.4%
	Hg	%	-0.5%	-0.4%	-0.1%	0.0%	0.0%	0.5%	1.7%	2.5%	3.4%	3.6%	3.4%	3.8%	4.1%	4.3%

* Emissions from both LTO and cruise are included here to capture all possible recalculations. However, the cruise component is reported as a memo item under the LRTAP Convention.

Table 3.6. Recalculations in Energy NFR 1A3 (c, d & e)

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
1A3c Railways	NO _x	kt	2.20	1.84	2.03	2.02	2.02	2.18	2.31	2.03	2.01	2.03	1.95	1.94	1.78	1.81
	SO ₂	kt	0.25	0.14	0.12	0.11	0.11	0.11	0.05	0.06	0.06	0.06	0.06	0.05	0.02	0.02
	NMVOC	kt	0.20	0.16	0.18	0.18	0.18	0.19	0.21	0.18	0.18	0.18	0.17	0.17	0.16	0.16
	PM _{2.5}	kt	0.06	0.05	0.05	0.05	0.05	0.06	0.06	0.05	0.05	0.05	0.05	0.05	0.05	0.05
	Pb	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1A3d(ii) National Navigation	NO _x	kt	2.14	2.29	3.78	5.20	6.12	4.83	5.01	4.88	4.90	4.25	4.49	4.39	5.50	5.43
	SO ₂	kt	1.16	1.26	1.22	0.49	0.21	0.16	0.08	0.10	0.10	0.07	0.10	0.10	0.10	0.10
	NMVOC	kt	0.07	0.08	0.13	0.18	0.22	0.17	0.18	0.17	0.17	0.15	0.16	0.16	0.20	0.19
	PM _{2.5}	kt	0.12	0.13	0.17	0.17	0.11	0.09	0.09	0.09	0.09	0.08	0.08	0.08	0.10	0.10
	Pb	t	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3e(i) Pipeline Compressors	NO _x	kt	0.05	0.10	0.05	0.13	0.13	0.11	0.12	0.13	0.14	0.13	0.12	0.13	0.13	0.12
Submission 2018																
1A3c Railways	NO _x	kt	2.20	1.84	2.03	2.02	2.02	2.18	2.31	2.03	2.01	2.03	1.95	1.94	1.78	1.81
	SO ₂	kt	0.25	0.14	0.12	0.11	0.11	0.11	0.05	0.06	0.06	0.06	0.06	0.05	0.02	0.02
	NMVOC	kt	0.20	0.16	0.18	0.18	0.18	0.19	0.21	0.18	0.18	0.18	0.17	0.17	0.16	0.16
	PM _{2.5}	kt	0.06	0.05	0.05	0.05	0.05	0.06	0.06	0.05	0.05	0.05	0.05	0.05	0.05	0.05
	Pb	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1A3d(ii) National Navigation	NO _x	kt	2.14	2.29	3.78	5.20	6.12	4.83	5.01	4.88	4.90	4.25	4.49	4.39	5.50	5.43
	SO ₂	kt	1.16	1.26	1.22	0.49	0.21	0.16	0.08	0.10	0.10	0.07	0.10	0.10	0.10	0.10
	NMVOC	kt	0.07	0.08	0.13	0.18	0.22	0.17	0.18	0.17	0.17	0.15	0.16	0.16	0.20	0.19
	PM _{2.5}	kt	0.12	0.13	0.17	0.17	0.11	0.09	0.09	0.09	0.09	0.08	0.08	0.08	0.10	0.10
	Pb	t	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3e(i) Pipeline Compressors	NO _x	kt	0.05	0.10	0.05	0.13	0.13	0.11	0.12	0.13	0.14	0.13	0.12	0.13	0.13	0.12

Table 3.6. Recalculations in Energy NFR 1A3 (c, d & e) (continued)

% Change in Emissions																
1A3c Railways	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO ₂	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-26.5%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM _{2.5}	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1A3d(ii) National Navigation	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	SO ₂	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM _{2.5}	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
1A3e(i) Pipeline Compressors	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Table 3.7. Recalculations in Energy NFR 1A4

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
1A4a Commercial/Institutional	NO _x	kt	3.06	2.84	3.16	3.30	3.13	3.24	3.56	3.11	3.12	2.86	2.88	2.68	2.45	2.37
	SO ₂	kt	11.60	5.56	2.39	1.93	1.80	1.70	1.16	0.74	0.71	0.68	0.65	0.55	0.26	0.25
	NMVOC	kt	0.47	0.42	0.53	0.62	0.61	0.65	0.71	0.60	0.62	0.54	0.57	0.56	0.53	0.52
	NH ₃	kt	0.00	0.00	0.00	0.00	0.00	0.01	0.02	0.02	0.02	0.02	0.03	0.04	0.04	0.03
	PM _{2.5}	kt	0.56	0.40	0.39	0.48	0.45	0.45	0.49	0.32	0.30	0.30	0.28	0.25	0.22	0.20
	Pb	t	0.18	0.03	0.02	0.15	0.15	0.15	0.16	0.02	0.02	0.02	0.02	0.03	0.03	0.02
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01
	Hg	t	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01
1A4b Residential	NO _x	kt	7.38	5.82	5.45	5.94	5.84	5.66	6.13	6.16	6.36	5.44	5.18	5.37	4.77	4.99
	SO ₂	kt	26.76	17.32	15.07	12.22	10.78	10.54	9.21	9.15	8.80	8.05	7.62	8.29	7.00	6.74
	NMVOC	kt	28.52	18.05	12.35	10.99	10.65	10.18	10.78	11.38	10.75	9.97	9.73	10.46	8.93	8.68
	NH ₃	kt	0.15	0.10	0.06	0.05	0.05	0.06	0.05	0.06	0.06	0.05	0.06	0.07	0.06	0.06
	PM _{2.5}	kt	23.92	15.17	10.35	9.23	8.94	8.56	9.05	9.56	9.03	8.37	8.19	8.81	7.53	7.32
	Pb	t	7.40	4.67	3.20	2.84	2.75	2.64	2.80	2.96	2.79	2.59	2.52	2.70	2.31	2.25
	Cd	t	0.11	0.07	0.05	0.04	0.04	0.04	0.04	0.05	0.05	0.04	0.04	0.05	0.04	0.04
	Hg	t	0.30	0.19	0.14	0.14	0.13	0.13	0.14	0.14	0.14	0.12	0.12	0.13	0.11	0.11
1A4c Agriculture/Forestry/Fishing	NO _x	kt	8.70	14.28	12.99	12.88	11.69	10.63	10.40	8.69	7.55	6.69	6.35	5.71	5.01	4.43
	SO ₂	kt	1.41	1.34	0.94	0.86	0.82	0.74	0.37	0.42	0.37	0.09	0.09	0.08	0.05	0.05
	NMVOC	kt	1.28	1.58	1.26	1.04	0.93	0.83	0.81	0.65	0.56	0.49	0.44	0.37	0.31	0.27
	NH ₃	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	PM _{2.5}	kt	0.85	0.94	0.68	0.54	0.49	0.44	0.43	0.35	0.30	0.27	0.24	0.20	0.17	0.14
	Pb	t	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Submission 2018																
1A4a Commercial/Institutional	NO _x	kt	3.06	2.84	3.16	3.30	3.13	3.24	3.56	3.11	3.12	2.86	2.88	2.68	2.45	2.47
	SO ₂	kt	11.52	5.54	2.38	1.93	1.80	1.70	1.16	0.74	0.71	0.68	0.65	0.55	0.26	0.26
	NMVOC	kt	0.47	0.42	0.53	0.62	0.61	0.65	0.71	0.60	0.62	0.54	0.57	0.56	0.53	0.55
	NH ₃	kt	0.00	0.00	0.00	0.00	0.00	0.01	0.02	0.02	0.02	0.02	0.03	0.04	0.04	0.03
	PM _{2.5}	kt	0.56	0.40	0.39	0.48	0.45	0.45	0.49	0.32	0.30	0.30	0.28	0.25	0.22	0.20
	Pb	t	0.58	0.38	0.37	0.49	0.45	0.45	0.49	0.29	0.27	0.27	0.26	0.23	0.19	0.18
	Cd	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.01
	Hg	t	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4b Residential	NO _x	kt	7.33	5.79	5.43	5.92	5.83	5.64	6.11	6.14	6.34	5.43	5.16	5.35	4.76	4.97
	SO ₂	kt	26.76	17.32	15.07	12.22	10.78	10.54	9.21	9.15	8.80	8.05	7.62	8.29	7.00	6.69
	NMVOC	kt	28.52	18.05	12.35	10.99	10.65	10.18	10.78	11.38	10.75	9.97	9.73	10.46	8.93	8.68
	NH ₃	kt	0.15	0.10	0.06	0.05	0.05	0.06	0.05	0.06	0.06	0.05	0.06	0.07	0.06	0.06
	PM _{2.5}	kt	23.91	15.16	10.34	9.21	8.93	8.55	9.04	9.55	9.03	8.37	8.19	8.81	7.53	7.33
	Pb	t	7.40	4.67	3.20	2.84	2.75	2.64	2.80	2.96	2.79	2.59	2.52	2.70	2.31	2.25
	Cd	t	0.11	0.07	0.05	0.04	0.04	0.04	0.04	0.05	0.05	0.04	0.04	0.05	0.04	0.04

1A4c Agriculture/Forestry/Fishing	Hg	t	0.29	0.19	0.13	0.12	0.12	0.11	0.12	0.12	0.12	0.11	0.11	0.11	0.10	0.09
	NO _x	kt	8.76	14.31	12.91	12.76	11.59	10.53	10.27	8.57	7.43	6.56	6.19	5.55	4.84	4.27
	SO ₂	kt	1.41	1.34	0.94	0.86	0.82	0.74	0.37	0.42	0.37	0.09	0.09	0.08	0.05	0.04
	NM VOC	kt	1.38	1.71	1.35	1.09	0.98	0.87	0.86	0.69	0.60	0.53	0.48	0.40	0.34	0.30
	NH ₃	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	PM _{2.5}	kt	0.93	1.05	0.76	0.60	0.53	0.47	0.47	0.37	0.32	0.28	0.25	0.21	0.17	0.15
	Pb	t	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01
	Cd	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Hg	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 3.7. Recalculations in Energy NFR 1A4 (continued)

% Change in Emissions			Pollutant	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A4a Commercial/Institutional	NO _x	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	4.4%
	SO ₂	%		-0.7%	-0.3%	-0.4%	-0.1%	-0.2%	-0.1%	-0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.7%
	NM VOC	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	6.2%
	NH ₃	%					0.0%	0.0%	0.0%	0.0%	-0.1%	-0.1%	0.1%	0.0%	0.0%	0.0%	0.0%
	PM _{2.5}	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%
	Pb	%		214%	1058%	1425%	222%	199%	192%	204%	1507%	1704%	1313%	1023%	666%	500%	790%
	Cd	%		281%	1157%	1442%	277%	194%	104%	73%	62%	71%	54%	41%	26%	20%	32%
	Hg	%		-15%	-45%	-56%	-34%	-35%	-37%	-38%	-68%	-70%	-67%	-69%	-69%	-70%	-70%
1A4b Residential	NO _x	%		-0.8%	-0.6%	-0.4%	-0.3%	-0.3%	-0.3%	-0.3%	-0.3%	-0.3%	-0.3%	-0.3%	-0.4%	-0.3%	-0.3%
	SO ₂	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.7%
	NM VOC	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	NH ₃	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM _{2.5}	%		0.0%	-0.1%	-0.1%	-0.2%	-0.2%	-0.1%	-0.1%	0.0%	-0.1%	-0.1%	0.0%	0.0%	0.0%	0.1%
	Pb	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Cd	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Hg	%		-1.5%	-3.9%	-8.4%	-11.8%	-12.5%	-12.3%	-12.7%	-11.3%	-13.3%	-11.9%	-12.8%	-12.2%	-12.6%	-13.2%
1A4c Agriculture/Forestry/Fishing	NO _x	%		0.7%	0.2%	-0.6%	-0.9%	-0.9%	-1.0%	-1.2%	-1.4%	-1.6%	-2.0%	-2.4%	-2.7%	-3.3%	-3.5%
	SO ₂	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-13.8%
	NM VOC	%		7.6%	8.2%	6.7%	4.7%	5.1%	5.7%	6.2%	6.8%	7.7%	8.5%	9.3%	9.9%	10.7%	11.9%
	NH ₃	%		0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	PM _{2.5}	%		8.4%	11.0%	11.7%	9.6%	9.2%	8.6%	7.9%	7.0%	6.3%	5.8%	5.6%	5.3%	5.1%	5.6%
	Pb	%		395%	303%	382%	314%	336%	342%	429%	390%	468%	539%	466%	365%	344%	376%
	Cd	%		12.3%	11.8%	12.3%	11.9%	12.0%	12.1%	12.4%	12.3%	12.6%	12.8%	12.6%	12.2%	12.1%	12.2%
	Hg	%		-1.9%	-1.5%	-1.9%	-1.6%	-1.7%	-1.7%	-2.1%	-1.9%	-2.3%	-2.6%	-2.2%	-1.8%	-1.7%	-1.9%

3.11 Quality Assurance/Quality Control

The time series spreadsheet system developed for individual categories, together with direct linking to the energy balance, allows for simple and efficient checking of activity data, emission factors, annual emissions and aggregated totals. Year-on-year changes immediately highlight any omissions, anomalies or internal errors. Initial checks are conducted by the inventory compiler as part of the calculation process, which is followed by a second check and completion of the QA/QC sheets which are integral to the calculation workbooks by another member of the inventories team. Cross-checks are performed for fuel data against the available supplementary sources for particular categories, such as 1A1a Public Electricity and Heat Production and some industrial processes, while maintaining consistency with fuel-use application in the estimation of GHG emissions. When new versions of the COPERT model are introduced for calculations in 1A3b, the previous year's activity data is run in the new model to compare with the current year, and the current year's activity data is run in the old version of the model to compare with the previous year. This allows firstly for identification of changes due to the model, and secondly for identification of changes due to the activity data.

3.12 Planned Improvements

The nature of the substances covered in transboundary emission inventories and the diverse range of sources involved, many of which are unintentional or uncontrolled releases, inevitably result in estimates that can be highly uncertain. There is heavy reliance on default emission factors made available through the Inventory Guidebook or other references, which are often based on limited information or which can only be applied in a limited or aggregated way because the necessary detailed activity data are not available. For some substances and categories, the emissions must be estimated in an indirect way, such as on a per-capita basis, which adds further to level of uncertainty.

Notwithstanding these difficulties, there is scope for improvement in Ireland's transboundary emission inventories by updating emission factors using the information in the latest version of the Inventory Guidebook (EMEP/EEA, 2016) and by accounting more completely for technological improvements over time, which should be reflected in reduced emissions for the time series. This submission included many changes to emission factors based on the latest Inventory Guidebook and this work will continue in future submissions. The inventory agency plans to address these needs by attempting to apply Tier 2 emission factors that refer to different types of stationary combustion appliance for the fuels in common usage, especially in sub-categories 1A4a Commercial/Institutional combustion and 1A4b(i) Residential combustion. The inventory agency will also endeavour to reduce the many remaining instances of the use of notation key "NE" by providing an estimate or concluding that the source category does not occur "NO" in future submissions.

3.13 Memo Items

The memo items of the NFR reporting format refer to activities for which the emissions are excluded from national totals. The use of fuels in domestic and international aviation (cruise phase) and marine bunkers are the most important of these activities. Some of the associated emissions, particularly from international aviation, are increasing very rapidly and it is therefore important that they are closely monitored for comparison with other sources and for the benefit of the international organisations that will have to develop control strategies for them in the future. The estimation of emissions for memo items is described here because they are calculated as part of the general estimation procedures for the Energy sector.

The national energy balance sheets include marine bunkers as a specific item and the emissions may be calculated directly. The approach used to for the cruise element of aviation is explained in Section 3.6.1 and this data is provided to SEAI for inclusion in the national energy balance.

Emission factors for international cruise aviation and navigation are documented in Tables C.19 and C.29 of Annex C.

Chapter Four

Industrial Processes and Product Use

4.1 Overview of the Industrial Processes and Product Use (NFR 2) Sector

The Industrial Processes sector has historically not been a large source of emissions in Ireland. Indeed, major industrial processes within the chemical sector and metal production that are common to many other developed countries have never been part of the economy in Ireland. Hence, many of the production processes within this sector are not relevant to the inventories of air pollutants in Ireland. Also of note is the fact that for a number of pollutants, it has not been possible to separate emissions from the combustion of fuel within industry and those associated with production processes. For all industries, fuel-based estimates of emissions have been collated and are reported under NFR Sector 1A2 (Manufacturing Industries and Construction). Where specific information is available in relation to process emissions as distinct from those associated with fuel combustion, they are reported under NFR Sector 2 (Industrial Processes and Product Use) and are discussed in this chapter. In most these cases, process-specific information is sourced from Annual Environmental Reports, which form part of the reporting obligations under IPPC permits in Ireland. In some cases, production data (estimated and/or calculated) are also used where available.

Relevant subcategories under Mineral Products (2A), Chemical Industry (2B), Metal Production (2C), Solvent and other product use (2D-2H) and Other Production Processes (2L) for which process emissions of various pollutants are reported under Industrial Processes and Product Use in Ireland are described in the following sections. Among these subcategories, Cement Production (2A1) is a key category for emissions of PCBs and Dioxins, accounting for 41.8 and 7.0 per cent, respectively, of the pollutants' national total emissions in 2016.

4.2 Mineral Products (NFR 2A)

The industrial processes for which estimates are included in Ireland's air pollution inventory under NFR 2.A are as follows:

- 2.A.1 Cement Production
- 2.A.2 Lime Production
- 2.A.3 Glass Production
- 2.A.5.a Quarrying and mining of minerals other than coal
- 2.A.5.b Construction and demolition
- 2.A.5.c Storage, handling and transport of mineral products
- 2.A.6 Other mineral products

4.2.1 Cement Production (NFR 2A1)

Cement manufacture is a major mineral industry. During cement manufacture, raw materials, such as limestone, are finely ground and then transformed in a kiln at high temperatures

(calcination) to produce clinker. Gypsum is then blended with clinker to produce cement. The combustion process in the cement kiln is an integral part of the production process, where the fuel ash becomes part of the cement clinker. It is therefore not possible in most cases to distinguish the process and combustion emissions from one another. As a result, because most of the pollutants originate from the fuels used, all emissions are generally reported under NFR Category 1A2f (Non-metallic minerals) and notation key IE is reported under 2A1 for these pollutants. The above is certainly true of all pollutants in Ireland's air pollutant inventory, with the exception of PCDD/F and PCBs, which are reported under Category 2A1 and are discussed in the following paragraphs.

There are at present four cement plants in Ireland, all of which use the dry kiln process, and they are currently fuelled by coal, petroleum coke and fuel oil, while a small amount of meat and bone-meal is used at one plant. Literature sources, in particular, the UNEP Toolkit (2013) and the EMEP/EEA Guidebook (2006), provide some POP emission factors on an overall "per unit production basis". These emission factors have been used to determine the total emissions from cement plants. Fuel-use data are available from plant operators as part of their reporting requirements under the EU ETS (Directive 2003/87/EC). Emissions from fuel use are calculated using combustion emission factors, while process emissions are calculated as the difference between the fuel-based estimates and the "per unit production" emission estimates.

Emission factors from the UNEP Toolkit (2013) for releases of PCDD/F to air range from 0.05 to 5 µg I-TEQ/t of cement produced, depending on the operating conditions and standards of abatement at production plants. As a result, plant-specific PCDD/F emission factors per unit production have been used to estimate total PCDD/F emissions from the industry in Ireland. The fuel-use data have been used to estimate the PCDD/F emissions from fuel combustion reported in subcategory 1A2f. The differences between fuel-based and per-unit-production-based values are reported under Sector 2A1 and these estimates range from 0.9 to 2.5 g I-TEQ across the time series. In 2016 at 1.48 g I-TEQ this sector accounted for 7.0 per cent share of the total PCDD/F emissions, having decreased by 21.1 per cent since 1990 (0.39 I-TEQ). Emissions for the time series are presented in Table 4.1.

There are limited data available on emission factors for PCBs from cement production, and uncertainties associated with the data are large, although cement production is unlikely to create significant emissions of PCBs under steady operation. Nevertheless, the Inventory Guidebook (EMEP/EEA, 2006) provides an emission factor of 1 µg/t of cement produced, and this factor has been adopted for the inventory across all years. Electrostatic precipitators (ESPs) are typically used to abate emissions from cement kilns due to the high running temperatures of the kilns (in excess of 1,500°C). While this form of abatement works well for PCDD/F, this is not the case with PCBs or PAHs. Thus, it has been decided that the Inventory Guidebook emission factor should be used for all years without amendment. As for PCDD/F, fuel-use data have been used to estimate the fuel-combustion-derived PCB emissions, which are reported in subcategory 1A2f. The differences in emissions between fuel-based estimates and per unit production estimates are reported under Category 2.A.1. In 2016 at 5.18 kg this sector accounted for 41.6 per cent share of the total PCB emissions, meaning these emissions have more than trebled since 1990 (1.07 kg). These emissions are presented in Table 4.1 (non-fuel emissions).

Table 4.1. Emission Time Series for Dioxins and Furans and Polychlorinated Biphenyls from Cement Production

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCDD/F (g I-TEQ)	1.871	1.862	1.258	0.925	0.914	0.910	1.024	1.421	1.507	1.577	1.564	1.650	1.521	1.499	1.476
PCBs (g)	1.070	1.064	2.259	3.363	3.365	3.399	2.962	1.958	1.743	1.546	1.861	2.907	4.074	4.736	5.177

4.2.2 Lime Production (NFR 2A2)

The lime production process involves the grinding and “burning” of limestone (CaCO_3) to produce what is commonly termed “quicklime” (CaO). It can then be further treated by the addition of water, a process called slaking, to produce slaked lime (CaOH), which generates large amounts of heat and steam. The finished product can then be packaged and distributed for use. Currently, there are two lime plants in Ireland and a third that operated until 1999. It is understood that all three utilised limestone quarries and kilns to burn the limestone raw material. The nature of the fuel used and the abatement in place varies from plant to plant.

Process emissions from lime production are derived following the approach taken for the estimation of emissions from Cement Production (2A1) described in section 4.2.1, whereby process emission estimates are obtained as the difference between total emissions on a “per unit production” basis and those estimated from fuel combustion. In this case, it was found that fuel combustion estimates of emissions were generally larger than those estimated on a “per unit production” basis and therefore all emissions from lime production are assumed to be included in those reported within Sector 1A2f.

4.2.3 Glass Production (NFR 2A3)

The manufacture of glass was not a predominant industry in Ireland, being limited to three sectors: lead crystal, container glass, and glass wool. The only container glass plant closed in 2002, one of the lead crystal plants closed in early 2006, the glass wool plant closed in 2008 and the last one, (second of the two) lead crystal plant closed in 2009. The pollutants for which process emission estimates have been made are particulate matter (TSP, PM_{10} , $\text{PM}_{2.5}$, BC) priority metals (Pb, Cd and Hg), other metals (As, Cr, Cu, Ni, Se and Zn), and PCDD/F. In addition, fuel-derived emissions from the glass industry are already accounted for within the Manufacturing Industries and Construction (1A2) sector.

Metal emissions can occur from glass processes from the metals contained in fossil fuels burnt to melt the glass and from metal additives to the glass. Metal emissions from industry fossil fuel use are accounted for within NFR Sector 1A2g (Other Manufacturing Industries). The metals emitted from glass production processes depend on the type of glass produced. Lead oxide and sometimes arsenic trioxide are used in the production of lead crystal glass and both metals can be emitted to air. Selenium and chromium compounds are used as colouring agents for container glass. Metal compounds are not believed to be used to any great extent in the production of glass wool. Of the glass processes outlined, all the plants were regulated under Integrated Pollution Prevention Control IPPC licences. Therefore, there is some information available from their AERs until they closed. Other licence information includes some details of plant design and operation, including capacities, fuel types and operating hours. In addition, confidential information in relation to production statistics has also been supplied to the inventory team.

Emission data for individual metals reported in AERs are limited to Pb emissions from the two lead crystal installations. These data have been used to estimate emissions of Pb from lead crystal production, while emission factors are used for the other glass processes. Literature emission factors are used from the Inventory Guidebook (EMEP/EEA, 2016) and, where deemed more appropriate, emission factors from the UK National Atmospheric Emissions Inventory (NAEI) database. The emission factors used are presented in Table D.1, Annex D. Total emissions for each metal from glass production are presented in Table 4.2.

Emission estimates from particulate matter were included in the inventory for the first time in the 2017 submission. The methodology uses confidential production data and emission

factors from the Inventory Guidebook (EMEP/EEA 2016). The emission factors used are presented in Table D.1, Annex D.

The potential for PCDD/F emissions from glass production is generally low because of the long residence times in high-temperature conditions, although chlorine can be introduced via fuels and raw materials, and therefore there is some potential for PCDD/F emissions. However, in the plants in Ireland, the main energy sources used were gas and electricity, and therefore PCDD/F emissions from fuel combustion were likely to be low. The information on abatement technology is uncertain for the glass manufacturing plants in Ireland. The URS Dames & Moore (2000) PCDD/F inventory report implies that one furnace would be fitted with abatement by 2005, but it is assumed that this is the large facility that closed in 2002. IPPC licence information implies that the environmental performance at the glass wool plant was improved in 1999, but the nature of the improvements is not clear.

Emission factors for PCDD/F are provided in the UNEP Toolkit (2013) for two different classes of facility: 0.2 µg I-TEQ/t of glass produced for a facility with no dust control, and 0.015 µg I-TEQ/t of glass produced for a facility with abatement. However, there have been improvements in environmental performance; therefore, the emission factor of 0.2 µg I-TEQ/t is used to estimate emissions in 1990, with a linear decrease to 0.11 µg I-TEQ/t in 2000 (URS Dames & Moore, 2000) and with a subsequent decrease to 0.015 µg I-TEQ/t by 2003 (when the container plant had closed) and remaining at this level up to 2009 when the last plant ceased its operation. Dioxin and furan emission estimates for glass production are presented in Table 4.2.

Table 4.2. Emission Time Series from Glass Production

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
TSP (t)	22.56	22.56	21.88	6.01	6.21	6.07	5.08	0.01	NO	NO	NO	NO	NO	NO	NO
PM10 (t)	20.08	20.08	19.48	5.30	5.47	5.35	4.47	0.01	NO	NO	NO	NO	NO	NO	NO
PM2.5 (t)	17.68	17.68	17.15	4.67	4.82	4.71	3.94	0.01	NO	NO	NO	NO	NO	NO	NO
BC (t)	0.10	0.10	0.09	0.09	0.10	0.09	0.08	0.00	NO	NO	NO	NO	NO	NO	NO
As (kg)	17.57	17.57	17.55	0.18	0.18	0.18	0.15	NE	NO	NO	NO	NO	NO	NO	NO
Cd (kg)	7.44	7.44	7.41	0.25	0.26	0.25	0.21	NE	NO	NO	NO	NO	NO	NO	NO
Cr (kg)	23.90	23.90	23.69	1.77	1.85	1.80	1.50	NE	NO	NO	NO	NO	NO	NO	NO
Cu (kg)	1.70	1.70	1.49	1.77	1.85	1.80	1.50	NE	NO	NO	NO	NO	NO	NO	NO
Pb (t)	310.77	310.77	310.55	128.90	88.52	96.78	96.46	7.91	NO	NO	NO	NO	NO	NO	NO
Hg (kg)	0.58	0.58	0.51	0.60	0.63	0.61	0.51	NE	NO	NO	NO	NO	NO	NO	NO
Ni (kg)	18.46	18.46	18.25	4.00	3.36	3.46	3.16	0.14	NO	NO	NO	NO	NO	NO	NO
Se (kg)	90.11	90.11	90.10	0.12	0.12	0.12	0.10	NE	NO	NO	NO	NO	NO	NO	NO
Zn (kg)	33.47	33.47	31.07	33.57	30.32	30.63	27.13	0.80	NO	NO	NO	NO	NO	NO	NO
PCDD/F (g I-TEQ)	0.54	0.39	0.26	0.01	0.01	0.01	0.01	0.00	NO	NO	NO	NO	NO	NO	NO

4.2.4 Quarrying and Mining of Minerals Other than Coal (NFR 2A5a)

Emissions of PM_{2.5}, TSP and PM₁₀ have been estimated for the first time in the 2018 submission for all years of the time series 1990- 2016. Activity data was taken from national statistics from the Central statistics office and emission factors from Chapter 2.A.5.a, Table 3.3 of the guidebook (EMEP/EEA, 2016). Activity data and emissions are presented in Table 4.3.

Table 4.3. Quarrying and mining of minerals other than coal (2.A.5.a)

Activity Data		1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Metallic minerals	Mt	2.7	2.6	3.5	4.7	4.6	4.4	4.2	4.3	4.2	4.2	3.8	3.7	3.7	3.7
Non-metallic minerals	Mt	28.8	26.8	63.1	94.5	111.9	92.0	59.1	39.3	35.9	31.4	33.0	32.8	32.8	32.8
Emissions factors															
TSP	g/t	102	102	102	102	102	102	102	102	102	102	102	102	102	102
PM10	g/t	50	50	50	50	50	50	50	50	50	50	50	50	50	50
PM2.5	g/t	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Emissions															
TSP	kt	3.2	3.0	6.8	10.1	11.9	9.8	6.5	4.4	4.1	3.6	3.8	3.7	3.7	3.7
PM10	kt	1.6	1.5	3.3	5.0	5.8	4.8	3.2	2.2	2.0	1.8	1.8	1.8	1.8	1.8
PM2.5	kt	0.2	0.1	0.3	0.5	0.6	0.5	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2

4.2.5 Construction and Demolition (NFR 2A5b)

Emissions of PM_{2.5}, TSP and PM10 have been estimated for the first time in the 2018 submission for all years of the time series 1990- 2016. Activity data was taken from national statistics from the Central statistics office and the US EPA Tier 1 methodology and emission factors from Chapter 2.A.5.b section 3.2 of the guidebook (EMEP/EEA, 2016). The emissions are presented in Table 4.4.

Table 4.4. Construction and Demolition (2.A.5.b)

Emissions		1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
TSP	kt	1.39	1.05	1.56	6.12	6.60	8.30	7.27	14.42	0.55	0.62	1.30	0.79	1.51	1.54
PM10	kt	0.42	0.32	0.47	1.83	1.98	2.48	2.18	4.31	0.17	0.19	0.39	0.24	0.45	0.46
PM2.5	kt	0.042	0.032	0.047	0.183	0.198	0.248	0.218	0.431	0.017	0.019	0.039	0.024	0.045	0.046

4.2.6 Storage, Handling and Transport of Mineral Products (NFR 2A5c)

Emissions of PM_{2.5} have been estimated for the first time in the 2018 submission for all years of the time series 1990- 2016. Activity data was taken from national statistics from the Central statistics office and emission factors from Table 3.4 Chapter 2.A.5.a of the guidebook (EMEP/EEA, 2016). Activity data and emissions are presented in Table 4.5.

Table 4.5. Storage, handling and transport of mineral products (2.A.5.c)

Activity Data		1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Metallic minerals	Mt	2.68	2.60	3.49	4.65	4.61	4.43	4.23	4.28	4.18	4.21	3.82	3.66	3.66	3.66
Non-metallic minerals	Mt	28.82	26.81	63.10	94.46	111.88	92.03	59.05	39.27	35.92	31.37	32.96	32.78	32.78	32.78
Emissions factors															
PM2.5-handling	g/t	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60
Emissions															
PM2.5-handling	kt	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

4.2.7 Other (NFR 2A6)

The industrial processes included within NFR Sector 2A6 are Bricks and Ceramics Production and Asphalt Production. Each of these subcategories is described in the following sections in terms of the pollutants for which emission estimates are made.

4.2.7.1 Bricks and Ceramics Production

The production of bricks and ceramics is a small sector in Ireland with a total of four IPPC-licensed facilities in operation. Emission estimates are only made for PCDD/F as there are no data available in relation to process emissions of other pollutants and, furthermore, they are expected to be negligible (AEA/CTC, 2008). Direct production information in relation to the bricks and ceramics sector is not available; however, raw material input data are provided by the companies under the EU ETS. For the purposes of inventory estimates, as a worst-case scenario it is assumed that raw material input equals product output. Emission factors are sourced from the UNEP Toolkit (2013) in which two classes of facility are suggested: 0.2 µg I-TEQ/t of brick produced for a facility with no dust control and 0.02 µg I-TEQ/t of brick produced for a facility with abatement. The URS Dames & Moore (2000) report suggests an emission factor of 0.11 µg I-TEQ/t, which is the average of the two emission factors, and this value, has been used across the time series. The UNEP Toolkit (2013) does not include emission factors for ceramics production and therefore the emission factor for bricks is also applied to ceramics production. Dioxin and furan emission estimates for bricks and ceramics production are presented in Table 4.6.

Table 4.6. Dioxin and Furan Emission Time Series from Bricks and Ceramics Production

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Bricks production														
PCDD/F (mg I-TEQ)	12.14	13.02	15.38	15.34	12.90	7.00	1.80	1.74	2.48	0.08	0.07	NO	1.62	2.84
Ceramics production														
PCDD/F (mg I-TEQ)	NO	NO	NO	0.032	0.049	0.043	NO	NO	NO	NO	NO	NO	NO	NO

4.2.7.2 Asphalt Production

In the context of this inventory, the term “asphalt” is used to describe a bituminous product that may contain varying amounts of aggregate, used to build and maintain roads, whilst “bitumen” is assumed to be a heavy oil tar product which is used at elevated temperatures particularly in roofing materials for some buildings. Currently, only PCDD/F emission estimates from asphalt are included in Ireland’s air pollution inventory.

Information in relation to the production of asphalt in Ireland is sourced from the European Asphalt Pavement Association (EAPA, 2001,2007, 2012 & 2016), which generates an annual report outlining the quantity and end use of asphalt produced in European countries. Production data are available from 1994 onwards, with pre-1994 production estimates assumed to be equal to those in 1994. The production levels until 2006 show an upward year-on-year trend due to increased road building in Ireland, from 2007 the trend has been decreasing.

In Ireland, bag filters were fitted to most asphalt production facilities prior to 2000 and it was suggested that all facilities would have bag filters by 2001 (URS Dames & Moore, 2000). The UNEP Toolkit (2013) gives a range in emission factors of 0.007 to 0.07 µg I-TEQ/t asphalt produced. Given the above information, the emission factor for PCDD/F from asphalt production of 0.07 µg I-TEQ/t is adopted for 1990. A linear decrease in the emission factor is then assumed to 0.039 µg I-TEQ/t by 2000, and a further linear decrease is assumed to 0.007 µg I-TEQ/t by the end of 2002. The emission factor is assumed to be 0.007 µg I-TEQ/t

from 2003 and onwards. Dioxin and furan emission estimates for asphalt production are presented in Table 4.7.

Table 4.7. Dioxin and Furan Emission Time Series from Asphalt Production

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCDD/F (mg I-TEQ)	133.0	92.2	111.7	23.8	23.1	19.6	23.1	16.1	12.6	13.3	16.1	12.6	13.3	13.3

4.3 Chemical Industry (NFR 2B)

The chemical industry is not a dominant industry in Ireland in relation to industrial processes and is not an important source of emissions. The only source of emissions for which estimates are collated are NO_x emissions from Nitric Acid Production for the years 1990-2002.

4.3.1 Nitric Acid Production (NFR 2B2)

Nitric acid is used as a raw material mainly in the manufacture of nitrogen-based fertiliser. It may also be used in the production of adipic acid and explosives, for metal etching, and in the processing of ferrous metals. In the manufacture of nitrogenous fertilisers, the Haber Bosch process is utilised in which NH₃ is made by combining nitrogen from the air with hydrogen from natural gas and water, using the energy from the gas and a catalyst. Nitric acid is produced by burning (oxidising) the NH₃ over a catalyst. The nitric acid is combined with more NH₃ to produce ammonium nitrate, which is solidified into granules or bead-like prills for application to land using a fertiliser spreader. Up to its closure in 2002, there was one such plant in Ireland, which utilised the above process to produce calcium ammonium nitrate and other nitrogenous fertiliser blends. The inventory agency received direct correspondence from the plant in relation to the quantities of nitric acid produced and the measured emissions of NO_x. Emission estimates and associated activity data for NO_x emissions from nitric acid production are presented in Table 4.8. Abatement measures were installed at the plant in the mid-1990s and they are reflected in emission estimates from 1995.

Table 4.8. Nitrogen Oxides Emission Time Series from Nitric Acid Production

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
Nitric acid (kt)	338.8	260	260	260	260	260	260	260	260	260	260	260	130
NO _x (kt)	1.680	1.672	1.823	0.960	0.280	0.280	0.280	0.280	0.280	0.280	0.303	0.374	0.187

4.4 Metal Production (NFR 2C)

This category includes a wide range of processes such as primary and secondary iron and steel production, aluminium production and other non-ferrous production. In this category, emissions are estimated for the following subcategories and pollutants:

- 2C1 Iron and Steel Production – As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, PCDD/F;
- 2C2 Ferroalloys Production – As, Cd, Cr, Ni, Pb, Zn, HCB;
- 2C3 Aluminium Production – Zn;
- 2C5 Lead production – Pb;
- 2C7 Other metal production – Cd, Cr, Cu, Ni, Pb, Zn;

Ireland is a major European producer of Zn and Pb ores. The preparation of Pb and Zn concentrates does not produce emissions and the concentrates are exported for further processing.

4.4.1 Iron and Steel Production (NFR 2C1)

This sector covers the manufacture of iron and steel, an energy-intensive process likely to generate high emissions to air from the use of furnaces and sintering processes, as well as the manual handling of the raw material to finished goods, which can include hot and cold rolling, and turning, temping and cutting of metal to reach a desired end product. Steel production in Ireland has been limited to a single large electric arc furnace installation, which closed in 2001 but was operational throughout the period 1990–2001. One small foundry remained in operation contributing with negligible amount of emissions after the large plant's closure. The main plant produced up to 360 kt of steel per annum mainly from recycled scrap steel. It received an IPPC licence to operate just months before its closure, therefore no AERs were filed by the plant. However, some emission testing was carried out with respect to heavy metal emissions as part of its licence application.

Heavy metal emission estimates have been calculated using the aforementioned emission testing results for Cd, Cr, Pb, Ni and Zn, whilst for the remaining pollutants (i.e. As, Cu, Hg, and Se) Inventory Guidebook (EMEP/EEA, 2016) emission factors have been used, assuming no abatement at the plant. The emission factors used are presented in Table D.3, Annex D. Emission estimates for the 1990–2001 time series are shown in Table 4.9. Metal production data are available from the site for the period 1994–2001, with pre-1994 production assumed to be equal to that in 1994. Emission estimates are calculated by multiplying the production data by the relevant pollutant emission factor and assume that no abatement was in place at the plant.

Table 4.9. Emission Time Series from Iron and Steel Production

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
As (t)	0.130	0.117	0.103	0.130	0.106	0.124	0.136	0.135	0.143	0.134	0.144	0.060	NO
Cd (t)	0.216	0.194	0.170	0.216	0.176	0.205	0.226	0.223	0.237	0.222	0.239	0.099	NO
Cr (t)	1.428	1.284	1.126	1.428	1.165	1.358	1.494	1.476	1.568	1.467	1.577	0.657	NO
Cu (t)	0.023	0.021	0.018	0.023	0.019	0.022	0.024	0.024	0.025	0.023	0.025	0.011	NO
Pb (t)	1.753	1.576	1.382	1.753	1.430	1.667	1.834	1.812	1.925	1.802	1.936	0.807	NO
Hg (kg)	32.60	29.30	25.70	32.60	26.60	31.00	34.10	33.70	35.80	33.50	36.00	15.00	NO
Ni (t)	2.694	2.421	2.124	2.694	2.198	2.562	2.818	2.785	2.958	2.768	2.975	1.240	NO
Se (t)	0.007	0.006	0.005	0.007	0.005	0.006	0.007	0.007	0.007	0.007	0.007	0.003	NO
Zn (t)	27.75	24.94	21.87	27.75	22.64	26.38	29.02	28.68	30.47	28.51	30.64	12.77	NO
PCDD/F (g I-TEQ)	0.743	0.891	0.783	0.990	0.810	0.942	1.035	1.023	1.086	1.017	1.092	0.462	0.012
PCBs (kg)	0.619	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE
B[a]p (t)	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

Electric arc furnaces are significant sources of POPs. The overall approach to report emissions of POPs from the iron and steel category has been to account for emissions from fuel combustion within Sector 1A2a, with the process emissions reported (where possible) under Sector 2C1, which have been estimated using the approach described with respect to Cement Production (2A1). Emission factors based on per unit production were used to calculate initial estimates of total emissions. These factors are sourced from the Inventory Guidebook (2016). The difference between estimates determined on this basis and those reported for fuel combustion sector 1A2a is then reported in sector 2C. Emission estimates for the sector are presented in Table 4.9 for PCDD/F.

4.4.2 Ferroalloys Production (NFR 2C2)

This sector covers several secondary sites engaged in iron and steel manufacture. Two types of installation are distinguished. The first type covers installations involved in the

manufacture of ductile iron for use in street furniture, public benches, waste bins and manhole covers, and the second is the manufacture of cast iron for appliances. The process of creating ductile iron utilises electric arc furnaces to smelt the raw materials, iron and magnesium. In the manufacture of cast iron, ferrous and non-ferrous metals, including scrap metal, are used within the process. Since 1990, there have been three relevant facilities in Ireland. Due to a change in operations, one of these plants reported negligible emission estimates from 2003 onwards; one facility closed in 2014 leaving a single operating facility from 2014 onwards.

A number of the larger metal processing sites are regulated under IPPC. Some metal emission estimates and particulate emission estimates have been reported in AERs; however, not all installations report emissions in all years. In some cases, only production data and emissions of TSP are available. Where production data only were provided by the plant operator, they were used to calculate emissions of TSP using USEPA factors for dust emissions from abated/unabated iron foundry cupola processes. Estimates for metal emissions were then obtained from the TSP estimates based on Inventory Guidebook (EMEP/EEA, 2016) dust composition data for foundry dust. Abatement techniques are also taken into account in emission calculations at a plant-specific level, where this applies. Emission estimates of HCB for the time series 1990-1996 are presented in Table 4.10.

Estimates for TSP, PM₁₀, PM_{2.5} and BC have also been made for this category. TPM data reported in AERs were used to estimate emissions from TSP. Fractionation profiles based on the emission factors within the Inventory Guidebook (EMEP/EEA, 2016) were used to estimate PM₁₀, PM_{2.5} and BC emissions.

The only source of HCB is the secondary manufacture of aluminium, for which the Inventory Guidebook (EMEP/EEA, 2016) indicates a factor of 5 g/t of aluminium. This factor has been used to estimate HCB emissions across the time series until use of the HCE-based cover gas was banned in 1996 and emissions are reported as not occurring for years after 1996.

Where production data are not available, TSP estimates have been used to estimate metal production across the time series using the BiPRO waste report (2005). Other POPs like PCDD/F, PCB and PAHs are reported under fuel combustion sector 1A2b. The HCB emission factor used is presented in Table D.3, Annex D, and process emission estimates for the 1990-1996 time series are presented in Table 4.10.

Table 4.10. Emission Time Series from Ferroalloys Production

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
TSP (Mg)	70.57	70.57	5.57	1.32	1.69	1.03	1.02	1.93	1.28	0.66	0.29	0.64	NO	NO	NO
PM ₁₀ (Mg)	59.99	59.99	4.74	1.12	1.43	0.88	0.86	1.64	1.09	0.56	0.24	0.54	NO	NO	NO
PM _{2.5} (Mg)	42.34	42.34	3.34	0.79	1.01	0.62	0.61	1.16	0.77	0.39	0.17	0.38	NO	NO	NO
Cr (Mg)	0.252	0.252	0.019	0.005	0.005	0.004	0.004	0.007	0.004	0.002	0.001	0.002	NO	NO	NO
As (Mg)	0.069	0.069	0.005	0.001	0.001	0.001	0.001	0.002	0.001	0.001	0.000	0.001	NO	NO	NO
Cd (Mg)	0.033	0.033	0.003	0.001	0.001	0.000	0.001	0.001	0.008	0.000	0.000	0.000	0.000	0.000	NO
Ni (Mg)	0.115	0.115	0.008	0.002	0.002	0.002	0.002	0.003	0.002	0.001	0.000	0.001	NO	NO	NO
Pb (Mg)	1.712	1.712	0.183	0.031	0.035	0.024	0.023	0.047	0.035	0.016	0.007	0.015	0.0001	0.0001	NO
Zn (Mg)	1.407	1.407	0.345	0.022	0.036	0.027	0.022	0.040	0.041	0.011	0.005	0.011	0.0001	0.0001	0.0001
PCDD/F (g I-TEQ)	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE
HCB (kg)	40.000	40.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
PCBs (kg)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
B[a]p (Mg)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

4.4.3 Aluminium Production (NFR 2C3)

Ireland is an important producer of alumina at one large plant using the Bayer process (extraction of Al_2O_3 using NaOH). The production of alumina using the Bayer process gives rise to significant metal emissions and therefore process emissions are not estimated for this source. Ireland has some secondary aluminium processing for which estimates of Zn have been made following reports from the plant involved. The plant closed in late 2006 and therefore estimates are only provided for the 1990–2006 time series as presented in Table 4.11. Production data for the plant are not available and therefore estimates were made using PM as an indicator. The UK NAEI emission factor of 2.725 g/t is then applied (Table D.3, Annex D).

Table 4.11. Emission Time Series for Zinc from Aluminium Production

Year	1990	1995	2000	2001	2002	2003	2004	2005	2006
Zn (t)	0.029	0.029	0.029	0.029	0.043	0.019	0.026	0.011	0.011

4.4.4 Lead Production (NFR 2C5)

A significant quantity of Lead is mined in Ireland, but such mining is assumed not to be a significant source of emissions to air. Estimates at facility level of Lead emissions have been obtained from AERs. Emission estimates for the time series are presented in Table 4.12 and are reported as not occurring since 2009.

Table 4.12. Emission Time Series for Lead Production

Pollutant (unit)	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Pb (kg)	7.845	7.845	7.845	7.845	0.018	0.008	0.013	0.006	0.025	0.034	0.030	NO

4.4.5 Other Metal Production (NFR 2C7)

This category covers all other metal manufacture and manipulation, including any emissions from the mining of raw materials. A significant quantity of Zn is mined in Ireland, but such mining is assumed not to be a significant source of emissions to air. Ireland has a number of small aluminium casting companies in addition to facilities for wire manufacture and the manufacture of refined or secondary Pb and Cu products, as well as a number of Zn galvanising plants.

Estimates at facility level of the heavy metals Cd, Cr, Cu, Ni, Pb and Zn have been obtained from AERs with respect to these secondary metal operations. Emission estimates for the time series are presented in Table 4.13. Emissions arise from very few plants in the latter parts of the time series, and are therefore very sensitive to changes in activity of individual plants.

Table 4.13. Emission Time Series for Non-Ferrous Metal Production

Pollutant	Units	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Cd (kg)	kg	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	NO	0.20	0.20	0.20	0.20	0.10	NO
Cr (kg)	kg	15.00	211.00	211.00	256.33	194.49	0.08	0.06	0.07	NO	NO	NO	NO	NO	NO	NO
Cu (kg)	kg	3.18	3.18	3.18	22.27	7.92	25.51	0.61	NO	NO	NO	NO	NO	NO	NO	NO
Pb (kg)	kg	0.15	0.15	0.15	9.85	0.46	3.75	3.11	0.02	NO	0.30	0.30	0.40	0.41	0.30	0.20
Ni (kg)	kg	82.00	82.00	82.00	4.97	3.98	40.69	0.13	NO	NO	NO	NO	NO	NO	NO	NO
Zn (kg)	kg	233.00	233.00	233.00	121.28	172.26	103.38	18.98	4.34	3.10	0.30	0.30	22.35	1.50	0.30	0.30

Process emissions of POPs (where applicable) are included in combustion emissions and reported as included elsewhere (IE) for category 2C7.

4.5 Overview of NMVOC emissions from Solvent and Other Product Use (NFR 2D-2L) Sector

The emission estimates presented in Solvent and Other Product Use (NFR 2D-2H) include Domestic solvent use including fungicides (2D3a), Road Paving with asphalt (2D3b), Coating Applications (2D3d), Degreasing and surface cleaning (2D3e), Dry Cleaning (2D3f), Chemical Products, Manufacture and Processing (2D3g), Printing (2D3h), Other Solvent Use (2D3i), Other Product use (2G) and Food and Beverages industry (2H2). Emissions are the result of continuing improvement of NMVOC emission estimates for Ireland through the outsourcing of tendered projects by the EPA. Road Paving with Asphalt 2D3b, Other Solvent use (Fat, Edible and non-edible oil extraction 2D3i, Use of Tobacco products and Fireworks 2G and Food and Beverages industry (2H2) have been included in Ireland's submission as a result of ongoing inventory improvements.

In 2012, the inventory agency commissioned a research project to update the NMVOC emission inventory for 2006-2013. This was a follow-on project to CTC/AEA (2005) and Finn et al. (2001) and resulted in a revised dataset where new data and methodologies had become available. This approach was taken in accordance with the Inventory Guidebook (EMEP/EEA 2016) methodology for NMVOC emissions for Solvent and Other Product Use (NFR 2D-2L). The results of this project were provided in the 2016 submission. This project continued into 2016 and resulted in further improvements to the NMVOC emission inventory from Solvent Usage.

Emissions data were gathered using a similar methodology to previous approaches. Bottom-up data was mainly obtained from submissions of Annual Environmental reports (AERs) which detail emissions in a variety of reporting formats ranging from the Solvent Mass Balance Summary, Solvent Management Plan (SMP), Pollution Release and Transfer Register (PRTR), or the Annual Environmental Report returns Workbook. In addition, new data sources were used from legislation designed to limit and report solvent usage (Solvent Directive 1999/13/EC). In conjunction with these data, the number of operators within each category was estimated using NACE codes provided by the Central Statistics Office (CSO) or from expert opinion.

Top-down methods were used for activities not covered by the IPPC licensing system nor under the Solvent Directive (1999/13/EC). The most significant included the use of non-industrial paints, metal degreasing and the use of domestic solvents. Input in the form of activity data, solvent usage or VOC emissions data for each individual activity were collated into spreadsheets. Emissions were estimated by applying the Inventory Guidebook (EMEP/EEA 2016) methods, default emission factors and general guidance as appropriate. Scaling up to national level was applied where necessary. The emission factors used are presented in Table D.4, Annex D

Emissions reported in NFR 2014 format are aggregated from the Selected Nomenclature for Air Pollutants (SNAP) categories. SNAP codes are used in the Inventory Guidebook (EMEP/EEA 2016) where sectoral emission sources and emission factors are provided in this system. Therefore, SNAP codes are adopted in these categories as it ensures that reporting of emissions is consistent with the guidebook and therefore other Parties submissions. Additionally, the use of SNAP codes facilitates a sub-sectoral analysis of drivers and trends.

For a number of sources, it was not possible to obtain reliable country-specific data (SNAP 060107: Paint Application: Wood, SNAP 060109: Non-Industrial Paint Application, SNAP

060408: Domestic Solvent Use). As a consequence of this, UK and other Parties' emission factors, and in some cases activity data (scaled by surrogate data), were used in the estimation methodology.

Obtaining country-specific data has been identified as an important issue in the past (Barry, S. and Regan, B., 2014). While new activity data were obtained for sectors that previously relied upon proxy sources, a number of sectors are still estimated using proxy information sources. Further reducing the dependence of these would require substantial investment, and the improvement that this would bring over using proxy based data is thought to be relatively small due to the similarity in lifestyle behaviour between the countries operating within the EU and therefore a common market place.

The main drivers associated with trends in implied emission factors relate to reduced solvent content of products, and paints. The trends in activity data reflect the fact that Ireland experienced rapid economic growth from the late 1990s to 2007. As a result, there was a substantial increase in the number of vehicles, growth in the number of individual households, and generally a higher per capita consumption of paints, cosmetics, toiletries, and other solvent containing products. Since 2007, there has been a rapid economic downturn, which has had a marked impact on consumption, and therefore emissions of NMVOC. As economic conditions began to improve emissions have also increased from 2012 to 2016.

Figure 4.1 illustrates the overall trend and shows a 20.8 per cent increase in total emissions between 1990 (33.0 kt of NMVOC) and 2016 (39.9 kt of NMVOC). The main contributor to the trend is sector 2H2 Food and Beverages industry, with 121 per cent increase between 1990 and 2016 and was responsible for 53.3 per cent share of emissions from solvent and other product use in 2016. The second largest contributor is sector 2D3a (Domestic Solvent Use including fungicides) with a contribution of 27.0 per cent of emissions from solvent and other product use (having increased by 35.8 per cent since 1990). Sector 2D3d (Coating Applications) accounted for 7.3 per cent in 2016 showing a decrease of 56.8 per cent between 1990 and 2016.




Figure 4.1 NMVOC Emission Trend for 2D-2H Other Solvent and Product 1990–2016

Sector 2D3a (Domestic Solvent Use including fungicides) is the second largest contributor to NMVOC emissions (Table 4.14). A Tier 2 method was implemented in this submission. The method uses population data obtained from the C.S.O and per capita emission factors for product use from the Inventory guidebook (EMEP/EEA, 2016). This approach was used as the statistics required for the use of Tier 2b approach were not complete in terms of the product types covered by domestic solvent use.

Sector 2D3b (Road paving with asphalt) emissions contribute less than 0.1 per cent of 2016 emissions. Emissions from the sector have decreased by 13.6 per cent since 1990.

Emissions from Coating applications (2D3d) decreased between 2007 and 2012 before increasing in 2013 and 2014 and decreasing again in 2015 and 2016 and reaching their lowest level in 2016 (2.92 kt). The main driver of NMVOC emissions from this emission category is the application of decorative paint (SNAP codes:060103/060104). A number of factors contributed to the ongoing decrease in emissions including; the substantial reduction in the solvent content of paint in recent years to comply with the Deco-Paints Directive (EP and CEU, 2004b), a greater awareness of environmental issues from the general public in addition to the economic downturn in Ireland. From discussions with industry, pressure from some of the larger retailers is noted to be one of the key drivers for the decrease in solvent use in architectural paint. The sales of water-based paints have decreased by 5.2 per cent between 1990 and 2016 whereas solvent-based paint sales have decreased by 43.3 per cent over the same period.

Emissions from 2D3e (Degreasing) decreased by 52.0 per cent between 1990 and 2016. Emissions peaked in 1996 at 2.5 kt. The methodology is based on net consumption of solvents (imports minus exports) provided by the CSO. The analysis showed that the main solvent used in this sector is Dichloromethane. The reductions are assumed to be driven by improved management practices and abatement technologies (open-top tanks have been

phased out in the European Union as a result of the Solvents Emissions Directive 1999/13/EC). Emissions from this emission source accounted for 2.1 per cent of the total emissions from solvent and other product use in 2016.

Data obtained under the reporting requirements of Solvent Directive (1999/13/EC) was used to estimate emissions from 2D3f (Dry Cleaning). Solvent usage, emissions data and national statistics were used to estimate emissions from this emission source. Emissions decreased by 77.9 per cent over the 1990-2016 period. Emissions from 2D3f accounted for 0.2 per cent of the total emissions from solvent and other product use in 2016.

2D3g (Chemical products) accounts for 2.9 per cent of emissions in solvent and other product use in 2016. In 1990, this sector accounted for 9.2 per cent. This emission category consists of fourteen emission sources, however, the majority of the emissions sources contribute very little to the overall emissions from the Chemical products sector. The diversity within these sectors is very large in terms of the type of process, the products made and the scale involved. The main driver of emissions from this emission source is Pharmaceutical Production (SNAP code 060306). Emissions from pharmaceutical production accounted for 68.7 per cent of emissions in the 2D3g in 2016. Emissions from 2D3g decreased by 61.4 per cent between 1990 and 2016. Emissions decreased as a result of the introduction of new management practices or through the use of abatement technology according to the (CTC/AEA, 2005 and Barry S. and O'Regan B., 2014). This indicates that current policy strategies are having an impact on solvent use and emissions. In addition, large reductions in emissions were found in several emission sources between 1990 and 2016. For instance, emissions reduced significantly from SNAP code 060303 (Polyurethane Processing), SNAP code 060305 (Rubber Processing). This was mainly a result of plant closures with the last rubber processing facility closing in 2016.

2D3h (Printing) emissions decreased 49.8 per cent over the 1990 to 2016. In 2016, the sector accounted for 3.7 per cent of total emissions from solvent and other product use. Emissions from this sector have increased for four consecutive years (5.4 per cent in 2013, 9.1 per cent in 2014, 4.4 per cent in 2015 and 0.1 per cent in 2016). The economic downturn in Ireland in 2009 may be responsible for the prior decrease in emissions and a return to better economic conditions may be driving the emission increases in recent years. However, it's important to note that the print industry is included under the Solvent Directive (1999/13/EC) and is subject to IPPC licencing where applicable.

2D3i (Other Solvent use) emissions increased 95.3 per cent over the 1990 to 2016. In 2016, the sector accounted for 3.4 per cent of total emissions from solvent and other product use. This emission category consists of 8 sources which include SNAP codes; 060401 (Glass Wool Enduction), 060402 (Mineral Wool Induction), 060404 (Fat, edible and non-edible oil extraction), 060405 (Application of Adhesives and Glues), 060406 (Preservation of Wood), 060407 (Underseal Treatment and Conservation of Vehicles), 060409 (Vehicle Dewaxing) and 060412 (Other). Application of Glues and Adhesives contribute 77.7 percent of total emissions from solvent and other product use in 2016. Emissions from this category have increased 97.0 per cent between 1990 and 2016 due to increased consumption of glue products. The methodology for reporting 060404 (Fat, edible and non-edible oil extraction) was amended to include emissions from 1990 to 2016, whereas previously values were available from 2008 onwards and were based on emissions from one IPPC licenced facility. The updated methodology is Tier 2 method based on Oilseed rape crop yield data provided by the CSO and emission factor from the Inventory guidebook (EMEP/EEA 2016).

Emission sources from 2G4 include 060602 (Use of tobacco) and 060303 (Use of shoes). Emissions from SNAP code 060602 (Use of tobacco) has been included in this submission using Tier 2 emission factor from Inventory guidebook (EMEP/EEA 2016) and excise volumes data obtained from Revenue. Emissions from category 2G4 accounted for 0.04 per

cent of emissions from solvent and other product use in 2016 and have decreased by 49.7 per cent between 1990 and 2016.

Sector 2H2 (Food and Beverage industry) is the largest contributor (53.3 per cent in 2016) to NMVOC emissions (Table 4.40-4.45). Tier 2 methodologies were applied to SNAP codes 040605 Bread, 040607 Production of Beer and 040608 Production of spirits, 040627 Meat frying and meat rendering, coffee roasting and feedstock, using activity data from C.S.O and Eurostat.

4.5.1 Domestic Solvent Use including fungicides (NFR 2D3a)

This subcategory covers SNAP sector 060408. This category addresses NMVOC emissions from the general use of products containing solvents by members of the public in their homes, but does not include the use of decorative paints. Many domestic products are also used in industry and commerce and in many cases, it is difficult or impossible to separate total sales into domestic and industrial components. Products that contain VOCs can be divided into a number of categories such as Cosmetic and Personal Care Products, Household Products, DIY products, Car Care Products, Varnish remover, Sealant and fillings Agents, Pharmaceutical Products Use and Pesticides. In this submission, a Tier 2 methodology was used with per-capita emission factors. This is the recommended approach to use where product statistics for the use of the Tier 2b approach are not complete in terms of the product types covered by domestic solvent use. Further study is planned to source appropriate product statistics to develop a Tier 2b approach. Estimates of NMVOC emissions from domestic solvent use are provided in Table 4.14.

Table 4.14. Non-Methane Volatile Organic Compound Emissions from Domestic Solvent Use

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
kt	7.93	8.14	8.57	9.35	9.89	10.14	10.25	10.30	10.34	10.37	10.38	10.42	10.48	10.77

4.5.2 Road Paving with asphalt (NFR 2D3b)

This sector covers the use of asphalt for road paving and covers SNAP sector 040611. This source is estimated using a Tier 2 methodology using annual weight of warm and hot mix asphalt used in Ireland for years 1993-2016 and the Tier 2 emission factor from the Inventory guidebook (EMEP/EEA 2016). PM_{2.5}, PM₁₀, TSP and BC were estimated for the first time in the 2018 submission for all years 1990-2016. Estimates of NMVOC, PM₁₀, TSP, PM_{2.5} and BC emissions from Road Paving with asphalt are provided in Table 4.15.

Table 4.15. Emissions from Road Paving with Asphalt (NFR 2D3b)

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.04	0.03	0.05	0.05	0.05	0.04	0.05	0.04	0.03	0.03	0.03	0.03	0.03	0.03
TSP (kt)	33.00	25.50	43.50	51.00	49.50	42.00	49.50	34.50	27.00	28.50	27.00	27.00	28.50	28.50
PM ₁₀ (kt)	4.40	3.40	5.80	6.80	6.60	5.60	6.60	4.60	3.60	3.80	3.60	3.60	3.80	3.80
PM _{2.5} (kt)	2.20	1.70	2.90	3.40	3.30	2.80	3.30	2.30	1.80	1.90	1.80	1.80	1.90	1.90
BC (kt)	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01

4.5.3 Coating Application (NFR 2D3d)

This sector covers the use of paints within the industrial, trade and domestic sectors. The term paint includes pigmented coatings and clear coatings such as lacquers and varnishes,

with the exception of glues, adhesives and inks. Unless captured on release and either recovered or destroyed, the solvent content of paint can be considered to be emitted to the atmosphere. The subcategories covered in this source category are presented below, with the relevant SNAP code in parentheses. SNAP codes not included below are deemed not to occur in Ireland.

Paint Application – Car Repairing (060102)
 Paint Application – Construction and Buildings (060103)
 Paint Application – Domestic Use (060104)
 Paint Application – Boat Building (060106)
 Paint Application – Wood (060107)
 Paint Application – Other Industrial Paint Application (060108)
 Paint Application – Other Non-Industrial Paint Application (060109)

Dependent on the SNAP code of interest, both bottom-up and top-down approaches have been used in emission estimates. Where there is an absence of country-specific data, per-capita emission factors derived from a number of EU member states national inventories were used to estimate emissions in Ireland using population statistics. Further details of the methodological choices for this source category are provided in Barry. and O'Regan. (2014), CTC/AEA (2005) and in Finn et al. (2001).

4.5.3.1 Paint Application: Car Repairing (SNAP 060102)

Activity data was obtained from a number of sources. From 2006-2012, sales data was obtained from a large supplier and data was scaled up based on market share and expert opinion. Data used in 1998 was calculated by Finn et al. (2001), data for 2000 and 2001 was provided by the British Coating Federation. Data was extrapolated and interpolated for the intervening years using passenger car numbers reported by the Department of Transport, Tourism and Sport each year (DTTAS, 2016). Emission factors were obtained using survey data from AEA/CTC (2005) and Barry and O'Regan (2014), default emission factors provided by the Inventory Guidebook (EMEP/EEA 2016) and where necessary, emission factors were calculated based on the average decrease in VOC content in known coating applications. The emission estimate includes thinners (EF 1000-835g/L), body fillers (EF 249-175g/L), top coat (720-420g/L) and primers (720-540g/L). This is considered a Tier 2 method. Emission estimates are provided in Table 4.16.

Table 4.16. Non-Methane Volatile Organic Compound Emissions from Paint Application: Car Repairing

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.19	0.21	0.31	0.43	0.81	0.68	0.44	0.38	0.36	0.26	0.24	0.25	0.22	0.19

4.5.3.2 Paint Application: Construction and Building (SNAP 060103) and Domestic Use (SNAP 060104)

Activity data was obtained from the Irish Decorative Surface Coating Association (IDSCA) for the period 2006-2016. The Irish Business and Employers' Confederation (IBEC) collated the total product sales for both water-based and solvent-based paints and provided the information to the inventory agency for the period 2000-2004. Following the experience in the UK (CTC/AEA, 2005), total product sales are proportioned between trade (Construction and Buildings) and retail (Domestic Use) use, assuming a 44:56 split in 1998, reaching 40:60 in 2003, and 30:70 in 2013. The split in 2016 is assumed to be the same as 2013. Estimates of paint sales prior to 1998 were extrapolated using GDP ($R=0.70$).

A number of emission factors were used to calculate NMVOC emissions from decorative coating applications. A survey of products found in popular retail chain stores was completed to establish a realistic emission factor for decorative surface coating products for recent years:

- Interior matt walls and ceiling paint was found to be 30g of VOC/l for solvent based paints and 22.5g of VOC/l for water based paints.
- Interior glossy walls and ceilings were found to have 76g of VOC/l for solvent based paint and 50g of VOC/l for water based paints.
- Exterior walls of mineral substrate were found to have 126g of VOC/l of solvent based paints and 9g of VOC/l for water based paints.
- Interior/exterior trim and cladding paints for wood and metal have an average solvent content of 324g of VOC/l of solvent based paints and 43g of VOC/l of water based paints.
- Primers were found to have an average solvent based paints of 201g of VOC/l and 45g of VOC/l for water based paints.

These emission factors were used for 2010-2016 while emission factors prior to this were assumed to be similar to 2007 limits outlined in the decorative paints directive. This was considered a Tier 2 method Emission estimates for this category are provided in Table 4.17.

Table 4.17. Non-Methane Volatile Organic Compound Emissions from Paint Application: Construction and Building (060103) and Domestic use (060104)

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
060104 kt	1.510	1.572	1.577	1.440	1.434	1.092	0.817	0.590	0.562	0.500	0.509	0.556	0.403	0.302
060103 kt	1.922	2.000	2.090	2.350	2.549	2.028	1.587	1.197	1.195	1.114	1.188	1.298	0.939	0.704

4.5.3.3 Paint Application: Boat Building (SNAP 060106)

Paint application in the Marine Sector includes a diverse range of products designed to prevent corrosion and protect ships hulls against damage from fouling. The formulation varies depending on the area being coated and application techniques also vary ranging from spraying to brushing and application by roller.

Activity data were obtained from a major marine coating supplier from 2010-2016 and was upscaled based upon the company's market share. Previous annual emissions were assumed to be the same as 2010. The supplier also provided an estimated industry product breakdown. Emission factors between products are relatively similar with Top coats, primers and anti-corrosion products having an estimated VOC content of 400g of VOC per kg of product while anti-fouling products are estimated to contain 440g of VOC per kg of product.

Paint Application in the Marine Sector in Ireland can be divided into domestic sector, cargo or freight sector and fishery sector. Larger vessels which require more product application are unable to dry dock in Ireland due to a lack of facilities to handle larger vessels. Therefore, sales data are adjusted based upon expert opinion to account for this (50 per cent of paint sales are applied elsewhere). This sector is a minor emission source for this reason. The methodology is considered a Tier 1 method. Estimates of NMVOC emissions from this source category are provided in Table 4.18.

Table 4.18. Non-Methane Volatile Organic Compound Emissions from Paint Application: Boat Building

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.090	0.090	0.090	0.090	0.090	0.090	0.090	0.090	0.102	0.089	0.103	0.111	0.115	0.144

4.5.3.4 Paint Application: Wood (SNAP 060107)

This subcategory refers to all paints used for the wood and wooden products sector but excludes the use of wood preservatives and creosote. Some activity data were available; however, no indication of the number of operators or market size was obtainable. Therefore, the emissions estimate was downscaled from UK data where consumption patterns and product range were considered to be comparable to Ireland. This involved using UK emissions data from the UK National Atmospheric Emissions Inventory (NAEI) and calculating per capita emissions (kg/per person) and applying this to Ireland using national population statistics. The methodology is considered a Tier 1 method. Estimates of NMVOC emissions from this source sector are provided in Table 4.19.

Table 4.19. Non-Methane Volatile Organic Compound Emissions from Paint Application: Wood

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	1.115	0.969	0.760	0.596	0.623	0.512	0.427	0.425	0.428	0.426	0.428	0.429	0.431	0.439

4.5.3.5 Paint Application: Other Industrial Sources (SNAP 060108)

The methodology for this source category involves the use of IPPC emissions data and scaling up to account for emissions in the non-IPPC sector based on information obtained from reporting under the Solvent Directive 1999/13/EC. Emissions varied slightly from those previously reported due to inclusion of corrected up to date emissions data. This category covers paints applied in industrial activities other than those already described in previous sections. Products painted include agricultural, construction and earth-moving equipment, aircraft, cans and drums, domestic appliances, electrical components, freight containers, machine tools, military vehicles, motor-vehicle components, office equipment, paper and plastics, and toys.

The scale of operation varies considerably from large operations employing automated roller coating to small-scale spraying painting. Processes may be enclosed or open air, and both air-dried and stove coatings are used. The emission estimate was up-scaled based on information obtained as a result of Solvents Directive 1999/13/EC. Estimates of NMVOC emissions from this source category are provided in Table 4.20. This methodology is considered a Tier 3 method

Table 4.20. Non-Methane Volatile Organic Compound Emissions from Paint Application: Other Industrial Sources

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	1.425	1.321	1.019	0.996	0.675	0.708	0.633	0.461	0.483	0.587	0.671	0.828	0.95	0.771

4.5.3.6 Paint Application: Other Non-Industrial Sources (SNAP 060109)

This category refers to the use of high performance protective and/or anti corrosive paints applied to structural steel, concrete and other substrates and any other non-industrial coatings. The sector includes coatings for offshore drilling rigs, production platforms and similar structures as well as road marking paints and non-decorative floor paints. Finn et al. (2001) obtained the activity data for this category as the difference between total paint sales in Ireland according to CSO data on paint sales and that used in other SNAP sectors under SNAP 0601. However, as no other data are available, emissions have been calculated following the advice of CTC/AEA (2005) using extrapolation from UK per capita estimates.

In order to establish whether the use of UK data is appropriate, per capita emissions were compared to other reporting parties. It was found that per capita emissions from this category range from 0.08 to 0.45 kg/person. The UK's per capita estimate is calculated at 0.14 kg per person. This is considered to be a realistic estimate for Irish emissions and was used to estimate emissions. This involved calculating UK per capita emissions (kg/per person) and applying this to Ireland using national population statistics. Estimates of NMVOC emissions from this source category are provided in Table 4.21. The methodology is considered a Tier 1 method.

Table 4.21. Non-Methane Volatile Organic Compound Emissions from Paint Application: Other Non-Industrial Sources

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.490	0.401	0.466	0.337	0.388	0.321	0.264	0.234	0.251	0.262	0.262	0.345	0.36	0.369

4.5.4 Degreasing and Dry Cleaning (NFR 2D3e and 2D3f)

Degreasing and Dry Cleaning (2D3e and 2D3f) covers the four subcategories that constitute SNAP Sector 0602. The subcategories for which emission estimates have been made are as follows with the relevant SNAP code in parentheses:

Metal Degreasing (060201)
 Dry Cleaning (060202)
 Electronic Components (060203)
 Other Industrial Cleaning (060204)

Activity data were obtained in the form of net consumption statistics (import minus exports) supplied by the CSO. Solvents included in the emissions estimate include perchloroethylene, dichloromethane, trichloroethylene and hydrocarbons from 1992-2016 before which time data are not available. The methodologies outlined in the Inventory Guidebook (EMEP/EEA 2016) and emissions data collected under the Solvents Directive 1999/13/EC are used to derive emission estimates. Further details of the methodological choices for this source category are provided in Barry, S. and O'Regan B. (2014).

4.5.4.1 Metal Degreasing (SNAP 060201), Electronics Manufacture (SNAP 060203) and Other Industrial Cleaning (SNAP 060204)

Degreasing is a process for cleaning water-insoluble substances, such as grease, fats, oils, waxes, carbon deposits, fluxes and tars, primarily from various metal products, but plastic, fibreglass, printed circuit boards and other products may also be treated by the same process. Therefore, a wide range of activities is covered.

The metalworking industries are the major users of solvent degreasing. Many manufacturers of electronic components also employ degreasing, but it is difficult to differentiate between the emissions emanating from degreasing and those from other sources. As a result, for the purposes of inventory estimates, emissions from Other Industrial Cleaning (060204) and Electronic Manufacture (060203) are included with Metal Degreasing (060201) as national statistics do not facilitate disaggregation of individual sectors.

The Inventory Guidebook (EMEP/EEA 2016) Tier 1 methodology is used for inventory estimates, and solvent consumption statistics (import minus exports) are used as the activity data. The default emission factor of 460g of VOC per kg of cleaning product is used. As data are not available for 1990-1991, the annual emission estimates for these years are assumed to be the same as 1992. Estimates of NMVOC emissions from this source category are provided in Table 4.22.

Table 4.22. Non-Methane Volatile Organic Compound Emissions from Metal Degreasing, Electronics Manufacture and Other Industrial Cleaning

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	1.743	2.388	0.847	0.953	1.557	1.531	0.789	0.885	0.925	1.004	0.902	0.922	0.69	0.837

4.5.4.2 Dry Cleaning (SNAP 060202)

Dry cleaning refers to any process to remove contamination from furs, leather, down leathers, textiles or other objects made of fibres, using organic solvents. Dry cleaning can be defined as the use of chlorinated organic solvents, principally perchloroethylene, to clean clothes and other textiles.

Emissions and usage data were obtained from the Solvents Directive 1999/13/EC for the years 2008-2010. In addition, the CSO provides information directly to the inventory agency in relation to perchloroethylene imports and exports. It is assumed that the net consumption (imports minus exports) in any year are used in that year for inventory estimates, even if there is some carryover of stock between years. Data are available from 1992-2016. Based on the percentage of perchloroethylene used in Dry Cleaning compared to national consumption in 2008-2010 and 2012, emissions were calculated for 1990-2007, 2011 and for 2013-2016. Estimates of NMVOC emissions from this source sector are provided in Table 4.23. The methodology is considered a Tier 3 methodology.

Table 4.23. Non-Methane Volatile Organic Compound Emissions from Dry Cleaning

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.282	0.270	0.090	0.085	0.105	0.093	0.063	0.053	0.058	0.075	0.057	0.068	0.04	0.062

4.5.5 Chemical Products, Manufacture and Processing (NFR 2D3g)

The mapping of Chemical Products, Manufacture and Processing (2D3g) to SNAP covers 14 subcategories in SNAP Code 0603. These subcategories are all industrial applications and, similar to Coating Application (2D3d), emission sources not included including Asphalt blowing and Leather tanning activities do not occur in Ireland. The 9 subcategories for which emission estimates are made are as follows with the relevant SNAP code in parentheses:

- PVC Processing (060302)
- Polyurethane Processing (060303)
- Rubber Processing (060305)
- Pharmaceutical Products Manufacturing (060306)
- Paints Manufacturing (060307)
- Inks Manufacturing (060308)
- Adhesives Manufacturing (060309)
- Adhesive and Magnetic Tapes, Films and Photographs Manufacturing (060311)
- Textile Finishing (060312)

Information pertaining to these sectors has been obtained from IPPC licenced companies with the exclusion of PVC processing (060302) which is based upon expert opinion from Finn et al. (2001). Estimates were up-scaled to reflect national emissions using the number of companies for each sector classified under European industrial activity classifications (NACE Rev.2) provided by the CSO. Emissions from Adhesive and Magnetic Tapes, Films and Photographs Manufacturing (060311) are included under SNAP code (060405) Industrial adhesive usage in section 4.5.7.3 Other Use of Solvents and Related Activities (2D3i-2G). Further details as to the exact methodological choices and the use of Inventory

Guidebook (EMEP/EEA 2016) methodologies applied in estimating emissions can be found in Finn et al. (2001), CTC/AEA (2005), and Barry and O'Regan (2014).

4.5.5.1 Polyvinyl Chloride (PVC) Processing (SNAP 060302)

The manufacture of polyvinyl chloride plastic involves an enclosed reaction or polymerisation step using the basic monomer to produce the resin, a drying step, and a final treating and forming step. Plastics are polymerised in completely enclosed vessels. Treatment of the resin after polymerisation varies with the proposed use. The major sources of air emission in plastics manufacture are the raw materials or monomers, solvents, or other volatile liquids emitted during the reaction, sublimed solids such as phthalic anhydride emitted in alkyd production, and solvents lost during storage and handling of thinned resins. Processing of PVC is not significant in Ireland. Emission data have been sourced from the installations involved which suggest an emission of 5 t/annum (Finn et al., 2001). The methodology is considered a Tier 1 method.

4.5.5.2 Polyurethane Processing (SNAP 060303)

This category deals with the application and subsequent discharge of organic compounds as blowing agents for creating polyurethane foams. Emissions are from the release of these blowing agents during foaming, or subsequently by the long-term release over several years. Polyurethane is used in building construction, for heat insulation, and for packaging material. For soft polyurethane foams, water may be used. Hard polyurethane foams utilise organic liquids as blowing agents.

Emission data have been sourced from IPPC-licensed companies involved in the manufacture of polyurethane and other foams. Estimates of NMVOC emissions from this category are provided in Table 4.24. The methodology is considered a Tier 1 method.

Table 4.24. Non-Methane Volatile Organic Compound Emissions from Polyurethane Processing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.162	0.162	0.080	0.044	0.039	0.037	0.035	0.027	0.004	0.004	0.004	0.004	0.004	0.004	0.004

4.5.5.3 Rubber Processing (SNAP 060305)

No detailed information for rubber processing is available within the Inventory Guidebook (EMEP/EEA 2016). Therefore, it is assumed in inventory estimates that this category includes processes such as moulding and mixing of natural and synthetic rubbers. Operations involving trimming and cutting are ignored since NMVOC emissions would not be associated with such operations.

Emission data have been sourced from IPPC-licensed companies involved in rubber processing that utilise organic solvents. Estimates of NMVOC emissions from this category sector are provided in Table 4.25. Emissions from this sector were dominated by the manufacture of tennis balls from two companies both of which have ceased operation and account for the steep decline in emissions. The last rubber processing facility closed in 2016 and emissions in 2016 are reported as NO. The methodology was considered a Tier 2 method.

Table 4.25. Non-Methane Volatile Organic Compound Emissions from Rubber Processing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NM VOC (kt)	0.423	0.424	0.422	0.199	0.199	0.199	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	NO

4.5.5.4 Pharmaceutical Products Manufacturing (SNAP 060306)

Depending on the nature of the pharmaceutical manufacturing facility, organic chemicals are used in the synthesis, extraction, fermentation and purification of Active Pharmaceutical Ingredients. Solvents are also used in the dilution of liquids, granulation, packaging and film coating. Thousands of individual products are categorised as pharmaceuticals. These products are usually produced in modest quantities in relatively small plants using batch processes. A typical pharmaceutical plant will use the same equipment to make several different products at different times.

The pharmaceutical industry is well established in Ireland and subject to IPPC licence requirements. Emission estimates have been made for 1998 and 2004 using an emission factor of 2 per cent of usage data (Finn et al. 2001, CTC/AEA, 2005) and for 2006-2016 using reported fugitive emissions data supplied by IPPC licenced facilities to the EPA. Other years (1990-1997 and 2005 emissions estimates) are interpolated or extrapolated from these estimates. The methodology is considered a Tier 3 method. Estimates of NM VOC emissions from this source category are provided in Table 4.26.

Table 4.26. Non-Methane Volatile Organic Compound Emissions from Pharmaceutical Products Manufacturing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NM VOC (kt)	2.124	2.124	2.072	0.769	0.865	1.119	1.310	1.727	0.873	1.029	1.129	1.121	0.969	0.767	0.802

4.5.5.5 Coating Manufacture: Paint (SNAP 060307)

The manufacture of paint involves the dispersion of coloured oil or pigments in a vehicle, usually an oil or resin, followed by the addition of an organic solvent for viscosity adjustment. Only the physical processes of weighing, mixing, grinding, tinting, thinning and packaging take place. No chemical reactions are involved.

Input and usage data have been sourced from a number of installations for 1998 and 2004 and from 2007 to 2016 emissions data was obtained from AERs. Emissions data were upscaled based on national statistics to reflect national emissions. Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data was available. Other years in the time series are estimated by interpolation and extrapolation. This methodology is considered a Tier 3 method. Emission estimates for NM VOC are provided in Table 4.27

Table 4.27. Non-Methane Volatile Organic Compound Emissions from Paint Manufacture

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NM VOC (kt)	0.162	0.162	0.117	0.083	0.074	0.064	0.069	0.060	0.043	0.027	0.039	0.051	0.052	0.152	0.300

4.5.5.6 Inks Manufacturing (SNAP 060308)

There are four major classes of printing ink: letterpress and lithographic inks, commonly called oil or paste inks, and flexographic and rotogravure inks, which are referred to as solvent inks. These inks vary considerably in physical appearance, composition, method of application, and drying mechanism. Flexographic and rotogravure inks have many elements

in common with the paste inks but differ in that they are of very low viscosity, and they almost always dry by evaporation of highly volatile solvents.

Emissions data were obtained from IPPC licensed facilities for 2008-2016. Where emissions estimates are based on usage data they are calculated based on an assumed emission factor of 2.5 per cent, which is the UK NAEI emission factor for this category (CTC/AEA, 2005). Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data were available. Gaps in the time series were then filled by interpolation and extrapolation. The methodology is considered a Tier 2 method. Estimates of NMVOC emissions from this source category are provided in Table 4.28.

Table 4.28. Non-Methane Volatile Organic Compound Emissions from Inks Manufacturing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.018	0.018	0.006	0.004	0.003	0.003	0.002	0.001	0.001	0.001	0.001	0.001	0.000	0.000	0.000

4.5.5.7 Adhesives Manufacturing (SNAP 060309)

This category includes the manufacture of glues and adhesives as it was difficult to derive separate activity data for glues and adhesives from those obtained from IPPC-licensed installations. Minor changes to emissions previously reported have occurred in this submission due to increased accuracy of reported emission data by facilities.

Emissions and usage data were supplied for a number of years in the time series (1998, 2006-2016). Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data were available. The methodology is considered a Tier 3 method. Estimates of NMVOC emissions from this source category are provided in Table 4.29.

Table 4.29. Non-Methane Volatile Organic Compound Emissions from Adhesives Manufacturing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.051	0.051	0.034	0.051	0.033	0.015	0.034	0.036	0.041	0.051	0.016	0.036	0.022	0.027	0.057

4.5.5.8 Textile Finishing (SNAP 060312)

Textile fabric finishing is part of the textile finishing industry. In fabric printing, a decorative pattern or design is applied to constructed fabric by roller, flat-screen or rotary-screen methods. Pollutants of interest in fabric printing are VOCs from mineral spirit solvents in print pastes or inks. Solvent use in this sector is usually associated with dry processing rather than wet processing of textiles.

Very little information is available for this activity. Two IPPC-regulated companies provided information to allow estimates to be made for a limited number of years (1998, 2004 and 2006-2015). Emissions were assumed to remain at 1998 levels for the period 1990-1997 as no emissions data were available. The remaining years were extrapolated. This was considered a Tier 1 method. The last company closed in 2016 and emissions are reported as NO for 2016. Estimates of NMVOC emissions from textile finishing are provided in Table 4.30.

Table 4.30. Non-Methane Volatile Organic Compound Emissions from Textile Finishing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.078	0.078	0.049	0.033	0.028	0.028	0.030	0.030	0.030	0.030	0.028	0.028	0.028	0.028	NO

4.5.6 Printing (2D3h)

Printing involves the use of various types of inks, which may contain a proportion of organic solvents which may be diluted before use. Different inks have different proportions of organic solvents and require dilution to different extents. Printing can also require the use of cleaning solvents and organic dampeners. The main printing techniques identified include offset, cold-set web offset, heat-set web offset, sheet-fed offset, rotogravure, flexography, letterpress, and screen-printing.

Usage and emission data are sourced from IPPC-regulated companies and scaled for those not regulated by IPPC based on national statistics and average emissions. Estimates of NMVOC emissions from printing are provided in Table 4.31. The large decrease in emissions is due to abatement measures introduced by the companies operating in Ireland due to the Solvent Directive (1999/13/EC) and a general greater awareness of environmental issues by the print industry. While the printing industry was affected by the economic recession from 2008 in Ireland and resulted in a decreasing emission trend, emissions increased in 2009 due to emissions from two IPPC licenced facilities. The methodology is considered a Tier 3 method.

Table 4.31. Non-Methane Volatile Organic Compound Emissions from Printing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	2.912	2.912	1.946	2.126	2.628	2.118	1.393	2.360	2.474	1.788	1.217	1.283	1.400	1.462	1.463

4.5.7 Other Use of Solvents and Related Activities (2D3i-2G)

This sector consists of 11 subcategories, which are a mixture of industrial and non-industrial activities, only 8 of which are applicable to Ireland. In NFR 2D3i (Other solvent use) the categories; Mineral wool enduction (060402) and Other (060412) are not considered to occur in Ireland. In NFR 2G4 Other product use the category Use of shoes (060602) is not estimated for Ireland. The 8 subcategories for which emission estimates are made are as follows with the relevant SNAP code in parentheses:

NFR 2D3i;

Glass Wool Blowing/Enduction (060401)

Fat, edible and non-edible oil extraction (060404)

Application of Glues and Adhesives (060405)

Preservation of Wood (060406)

Underseal Treatment and Conservation of Vehicles (060407)

Vehicle Dewaxing (060409)

NFR 2G4;

Use of Fireworks (060601)

Use of Tobacco (060602)

Both bottom-up and top-down approaches are used in the estimation of emissions from the subcategories outlined, depending on the availability of data for each subcategory. Similar to the other categories, further information in relation to subcategory estimation methodologies can be found in Finn et al. (2001), CTC/AEA (2005) and Barry and O'Regan (2014).

4.5.7.1 Glass Wool Blowing/Enduction (SNAP 060401)

Glass fibre manufacturing is the high-temperature conversion of various raw materials into a homogeneous melt, followed by the fabrication of this melt into glass fibres. The two basic types of glass fibre products, textile and wool, are manufactured by similar processes. Within

the category in Ireland, formaldehyde and phenol are used. Usage and emission data have been sourced from one IPPC-regulated company which ceased operation in 2009. Estimates of NMVOC emissions from this source category are provided in Table 4.32.

Table 4.32. Non-Methane Volatile Organic Compound Emissions from Glass Wool Blowing/Enduction

Year	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008
NMVOC (kt)	0.003	0.003	0.003	0.003	0.003	0.003	0.005	0.005	0.005	0.005	0.001

4.5.7.2 Fat, edible and non-edible oil extraction (SNAP 060404)

This sector covers solvent extraction of edible oils from oilseeds and drying of leftover seeds before resale as animal feed. The extraction of oil from oil seeds is performed either mechanically or through the use of solvents, or both. Where solvent is used, it is generally recovered and cleaned for reuse. The seed may be subjected to solvent treatment many times before all the oil is extracted. The remaining seed residue is then dried and may be used as an animal feed. Emissions of NMVOC, TSP, PM₁₀ and PM_{2.5} for this sector were estimated using statistics obtained from the C.S.O on the national yield of oilseed The Inventory Guidebook (EMEP/EEA, 2016) Tier 2 emission factors from Table 3-4 were applied. Estimates of NMVOC, TSP, PM₁₀ and PM_{2.5} are given in Table 4.33.

Table 4.33. Emissions from Fat, edible and non-edible oil extraction

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.031	0.020	0.014	0.022	0.028	0.050	0.032	0.037	0.044	0.088	0.092	0.077	0.054	0.063	0.054
TSP (kt)	0.022	0.014	0.009	0.016	0.020	0.035	0.022	0.026	0.031	0.061	0.065	0.054	0.038	0.044	0.038
PM ₁₀ (kt)	0.018	0.012	0.008	0.013	0.016	0.029	0.018	0.021	0.025	0.050	0.053	0.044	0.031	0.036	0.031
PM _{2.5} (kt)	0.012	0.008	0.005	0.009	0.011	0.019	0.012	0.014	0.017	0.034	0.035	0.030	0.021	0.024	0.020

4.5.7.3 Application of Glues and Adhesives (SNAP 060405)

This sector covers the use of all adhesives excluding domestic adhesive usage and includes adhesive and magnetic tape production (SNAP 060311). These data include adhesives used for publications and packaging, footwear, construction, transport equipment, rubber and plastic products, abrasives, engineering, laminating and other sectors.

This estimate is based upon net consumption statistics (import minus export data) and the Inventory Guidebook (EMEP/EEA, 2016) default emission factor of 522 g/kg of adhesive. The methodology is considered a Tier 2 method. Estimates of NMVOC emissions from the application of glues and adhesives are provided in Table 4.34.

Table 4.34. Non-Methane Volatile Organic Compound Emissions from Application of Glues and Adhesives

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.540	0.313	0.755	2.986	3.263	4.610	3.138	2.031	1.121	1.293	0.872	0.702	0.972	1.123	1.064

4.5.7.4 Preservation of Wood (SNAP 060406)

This section refers to emissions from the industrial use of wood preservatives. It does not include emissions from the surface coating of timber with paints, varnishes or lacquer (which are covered under SNAP 060107), and it does not cover the use of wood preservatives by the public at large (which is covered under SNAP 060408). Wood preservation is carried out using solvent-based preservatives, water-based preservatives or creosote. Creosote is an oil

product, prepared from coal tar distillation, and contains a high proportion of aromatic compounds such as PAHs. Regulations banning the sale of creosote took effect from June 2003. However, creosote may still be used for industrial applications, e.g. railway sleepers, telegraph poles and fencing, but with tougher restrictions on its composition and how it is applied. Creosote is gradually being replaced by water-borne preservatives. Preservatives based on organic solvents have a wide-ranging content of organic solvent, usually white spirit or other petroleum-based hydrocarbons. Water-borne preservatives consist of solutions of inorganic salts in water, with Cu, Cr and As (CCA)-based preservatives being the most widely used. Water-borne preservatives are not of concern to this inventory, as they do not contain VOCs.

In addition to bottom-up IPPC-licensed data, usage data was provided by the sole Creosote using company in Ireland. The Inventory Guidebook (EMEP/EEA, 2016) emission factor of 105 g/litre creosote applied is used. The methodology is considered a Tier 3 method. Estimates of NMVOC emissions from wood preservation are provided in Table 4.35.

Table 4.35. Non-Methane Volatile Organic Compound Emissions from Preservation of Wood

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.092	0.122	0.216	0.090	0.514	0.491	0.176	0.130	0.292	0.239	0.536	0.342	0.354	0.236	0.252

4.5.7.5 Underseal Treatment and Conservation of Vehicles (SNAP 060407)

The application of coatings to the underside of car bodies is conducted for protection from stone chips and for sound deadening. In the aftermarket sector, coatings are applied to the underside of cars only during repair of damaged bodywork. Finn et al. (2001) stated that sources within the trade suggested that application of underseal in Ireland was zero or minimal. However, further contact with suppliers revealed that a market of 650 l/annum existed at the time (1998). It is assumed that this market existed for all years prior to 1998. However, CTC/AEA (2005) suggested that this market no longer exists in Ireland and that emissions decreased in a linear fashion up to 2003, after which emissions from the activity no longer occur. The approach uses an average solvent content of 20 per cent, a density of 1,000 kg/m³, and assumes that 100 per cent of the solvent is emitted. Estimates of NMVOC emissions from this source category are provided in Table 4.36.

Table 4.36 Non-Methane Volatile Organic Compound Emissions from Underseal Treatment and Conservation of Vehicles

Year	1990	1995	2000	2001	2002	2003
NMVOC (Kg)	130	130	87	65	43	22

4.5.7.6 Vehicle Dewaxing (SNAP 060409)

In the past, some manufacturers of new cars applied a protective covering to parts of the car body after painting to provide protection during transport. Removal of this coating was carried out at the import centres using solvents. However, car manufacturers now invariably use either water-soluble wax that can be removed using hot water or self-adhesive film instead of wax. Consequently, it is assumed that emissions from this activity are now zero. Discussion with car distributors suggested that, historically, 20 per cent of new cars in Ireland were dewaxed and that the practice was discontinued after 2003. An emission factor of 1 kg/car is applied to estimate emissions using vehicle statistics provided by the CSO (Finn et al., 2001). Estimates of NMVOC emissions from vehicle dewaxing are provided in Table 4.37.

Table 4.37. Non-Methane Volatile Organic Compound Emissions from Vehicle Dewaxing

Year	1990	1995	2000	2001	2002	2003
NMVOC (kt)	0.035	0.035	0.023	0.017	0.012	0.006

4.5.7.7 Use of Tobacco (SNAP 060602)

This category comprises NMVOC, NO_x, CO, NH₃, TSP, PM₁₀, PM_{2.5} and BC emissions from the combustion (smoking) of tobacco products. Activity data was obtained from The Office of the Revenue Commissioners regarding the excise volumes of tobacco and includes an estimation of Illegal tobacco imported to Ireland from an illegal products research report produced by the Office of the Revenue Commissioners (Office of the Revenue Commissioners, 2015). The Inventory guidebook (EMEP/EEA, 2016) Tier 2 emission factors in table 3-14 are applied. Estimates of emissions from Use of Tobacco are provided in Table 4.38.

Table 4.38. Emissions from Use of Tobacco

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.034	0.037	0.039	0.032	0.032	0.031	0.028	0.027	0.024	0.025	0.023	0.020	0.019	0.020	0.017
NO _x (kt)	0.013	0.014	0.015	0.012	0.012	0.012	0.011	0.010	0.009	0.009	0.008	0.007	0.007	0.008	0.006
CO (kt)	0.386	0.419	0.449	0.360	0.366	0.353	0.324	0.310	0.278	0.280	0.257	0.224	0.213	0.230	0.194
NH ₃ (kt)	0.029	0.032	0.034	0.027	0.028	0.027	0.024	0.023	0.021	0.021	0.019	0.017	0.016	0.017	0.015
TSP (kt)	0.189	0.205	0.220	0.176	0.179	0.173	0.159	0.152	0.136	0.137	0.126	0.110	0.104	0.113	0.095
PM ₁₀ (kt)	0.189	0.205	0.220	0.176	0.179	0.173	0.159	0.152	0.136	0.137	0.126	0.110	0.104	0.113	0.095
PM _{2.5} (kt)	0.189	0.205	0.220	0.176	0.179	0.173	0.159	0.152	0.136	0.137	0.126	0.110	0.104	0.113	0.095
BC (kt)	0.076	0.082	0.088	0.070	0.072	0.069	0.064	0.061	0.054	0.055	0.050	0.044	0.042	0.045	0.038

4.5.7.8 Use of Fireworks (SNAP 060601)

This category is estimated for the first time in this submission. It comprises NO_x, CO, SO₂, TSP, PM₁₀ and PM_{2.5} emissions from the use of fireworks. Activity data was obtained from The Department of Justice and Equality and relates to the professional use of fireworks in displays. The Inventory guidebook (EMEP/EEA, 2016) Tier 2 emission factors in Table 3-13 are applied. Estimates of emissions from Use of Fireworks are provided in Table 4.39.

Table 4.39. Emissions from Use of Fireworks

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NO _x (t)	0.011	0.012	0.012	0.012	0.012	0.013	0.013	0.012	0.012	0.011	0.011	0.010	0.010	0.008	0.007
CO (t)	0.315	0.331	0.340	0.335	0.338	0.351	0.355	0.344	0.326	0.307	0.293	0.283	0.262	0.208	0.199
SO ₂ (t)	0.133	0.140	0.143	0.141	0.143	0.148	0.150	0.145	0.138	0.130	0.124	0.119	0.111	0.088	0.084
TSP (t)	4.835	5.091	5.218	5.145	5.195	5.397	5.446	5.279	5.011	4.717	4.505	4.340	4.021	3.193	3.059
PM ₁₀ (t)	4.399	4.632	4.747	4.681	4.727	4.910	4.955	4.803	4.559	4.291	4.098	3.949	3.659	2.905	2.783
PM _{2.5} (t)	2.287	2.408	2.468	2.433	2.457	2.552	2.576	2.497	2.370	2.231	2.130	2.053	1.902	1.510	1.447

4.5.8 Food and Beverage Industry (NFR 2H2)

According to the EMEP/EEA Guidebook (EMEP/EEA, 2016) this sector includes emissions from all processes in the food production chain which occur after the slaughtering of animals and the harvesting of crops as well as drink manufacturing including production of alcoholic beverages. For Ireland, Wine production (040606) is not occurring, Sugar production (040625) has not occurred since 2005, when the last sugar factory was closed (no activity

data available), and Flour production (040626) does not occur on a large scale (no activity data available), margarine and fats production is not thought to occur (no activity data available). Emissions include Spirit production, Animal feed production and Bread production which are the most significant source of emissions in the Food and Beverage industry in Ireland. The 6 subcategories for which emission estimates are made for Ireland are as follows with the relevant SNAP code in parentheses, where applicable:

Bread (SNAP 040605)

Beer (SNAP 040607)

Spirits (SNAP 040608)

Meat fish etc. frying/curing (SNAP 040627)

Coffee Roasting

Feedstock

4.5.8.1 Bread (SNAP 040605)

This sector includes bread, cakes and baking products. Activity data on white bread and bread products production was obtained from EUROSTAT for years 1995 to 2016. The data for years 1990-1994 was taken to be the same as 1995 as no data was available for these years. Emissions from cakes was not included in this estimate as no activity data was available. Tier 2 emission factors from the Inventory guidebook (EMEP/EEA, 2016) were used for bread and cakes, biscuits and breakfast cereals i.e. baking goods. The NMVOC emissions from Bread is given in table 4.40.

Table 4.40. Non-Methane Volatile Organic Compound Emissions from Food and Beverage Industry; Bread

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	1.707	1.707	1.006	1.374	1.495	1.512	1.631	1.472	2.577	1.707	1.593	1.838	1.714	1.696	1.643

4.5.8.2 Beer (SNAP 040607)

This includes mainstream beer production and craft beer production which has seen a steady increase since 2005 and a significant increase in Ireland since 2010. Activity data was obtained from a variety of sources including the Irish Brewers Association reports (ABFI, 2013, 2014 & 2015) and Independent Craft Brewers of Ireland and Bord Bia Report (Feeney, 2015). The Inventory Guidebook (EMEP/EEA, 2016) Tier 2 emission factor of 0.035 kg/hL was used. The NMVOC emissions from Beer production is given in table 4.41.

Table 4.41. Non-Methane Volatile Organic Compound Emissions from Food and Beverage Industry; Beer

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.194	0.207	0.225	0.250	0.256	0.266	0.272	0.274	0.274	0.274	0.288	0.282	0.258	0.276	0.276

4.5.8.3 Spirits (SNAP 040608)

Spirit production is a significant source of NMVOC emissions within the Food and Beverage industry due to the growth of the Whiskey production industry in Ireland. In the previous submission activity data was obtained for the years 2008, 2013 and 2014 from the Irish Whiskey Association Report (Irish Whiskey Association, 2015) and Irish Spirits Association data (Irish Spirits Association, 2015). Other years were extrapolated using this data. In the current submission activity data is based on national statistics for the years 1999-2016. Other years were extrapolated using this data. A Tier 2 emission factor of 15kg/hl alcohol

was used from the Inventory Guidebook (EMEP/EEA, 2016). NMVOC emissions from Spirit production in Ireland in given in table 4.42.

Table 4.42. Non-Methane Volatile Organic Compound Emissions from Food and Beverage Industry; Spirits

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	3.966	4.677	4.899	9.294	8.303	9.500	9.368	11.489	12.286	13.308	14.688	16.290	13.560	14.379	15.517

4.5.8.4 Meat, fish etc., frying/curing (SNAP 040627)

Emissions mainly occur from the cooking of meat, fish and poultry, releasing fats and oils and their degradation products. Emissions from fish frying and curing were not estimated due to absence of accurate activity data. AD for fish frying is under investigation and included in the planned improvement section 4.10. Activity data was obtained from the CSO on tonnes of animal slaughterings in Ireland this was taken to be the equivalent of meat rendered in Ireland the Inventory guidebook (EMEP/EEA, 2016) emission factor of 0.33 kg/Mg of meat rendered was used. Activity data on human consumption of meat from the CSO was taken to equate to meat frying and using the Inventory Guidebook (EMEP/EEA, 2016) emission factor of 0.3 kg/Mg product was used this is considered a Tier 2 method. The NMVOC emissions from meat frying/curing is given in table 4.43.

Table 4.43. Non-Methane Volatile Organic Compound Emissions from Food and Beverage Industry; Meat, fish etc. frying/curing

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.398	0.402	0.450	0.438	0.446	0.439	0.424	0.405	0.425	0.426	0.415	0.424	0.457	0.459	0.482

4.5.8.5 Coffee Roasting

The roasting of coffee beans is a source of NMVOC emissions. This activity does not have a relevant SNAP code. Activity data for unroasted coffee imports was obtained from the UN Comtrade Database and the emission factor from the Inventory Guidebook (EMEP/EEA, 2016) Tier 2 emission factor was used to estimate emissions as can be seen in table 4.44.

Table 4.44. Non-Methane Volatile Organic Compound Emissions from Food and Beverage Industry; Coffee roasting

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.001	0.001	0.002	0.003	0.003	0.003	0.002	0.001	0.001	0.003	0.003	0.003	0.003	0.005	0.006

4.5.8.6 Feedstock

The processing of by-products to produce animal feeds is a source of NMVOC emissions in Ireland. The tonnage of animal feed produced was sourced from the CSO and the Inventory Guidebook (EMEP/EEA, 2016) emission factor of 1 kg/Mg feed was used to estimate emissions from this source as can be seen in table 4.45.

Table 4.45. Non-Methane Volatile Organic Compound Emissions from Food and Beverage Industry; Feedstock

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	3.351	3.222	3.100	2.700	3.019	2.583	2.636	2.263	2.490	2.466	3.425	3.655	3.040	3.228	3.337

4.6 Other production, consumption, storage, transportation or handling of bulk product (NFR 2L)

The Other production, consumption, storage, transportation or handling of bulk product category in Ireland's air pollutant inventory includes emissions of PCDD/F and PCBs from leakage from electrical equipment and emissions of PCBs from fragmentisers and shredders. The main use of PCBs since the 1970s, when open uses were banned, has been as dielectric fluids in electrical equipment such as transformers and capacitors. However, the production and use of dielectric fluid containing PCBs has been highly regulated since 1986. Releases to the environment have decreased since 1990 as older PCB-containing equipment is taken out of service and is replaced by PCB-free equipment, which reduces the stocks that may lead to PCB emissions. It is also taken into consideration that, in some cases, trace PCDD/F may be present in PCB dielectric fluid. These arise from the original PCB synthesis process and from oxidation during dielectric breakdown events.

Electrical equipment, including white goods and electronic equipment, is partly recycled by breaking down the products in fragmentisers and shredders. Fragments are separated into ferrous scrap, a fraction containing non-ferrous scrap (which would then be processed separately), and a waste fraction that is typically disposed to landfill. Polychlorinated biphenyls are present in the capacitors of old electrical equipment. Hence, there is potential for PCBs to be released to air during fragmentiser operations.

4.6.1 Leakage from Electrical Equipment (NFR 2L)

The release of PCBs to the environment from electrical equipment is very difficult to estimate with any accuracy due to the large number of components potentially containing PCBs, the range of lifetime and replacement rates for PCB components, and the difficulties for users in identifying such components. Polychlorinated biphenyls have never been manufactured in Ireland. Production ceased in the UK in 1977 and in the rest of Europe and North America in 1986. Manufacturers of electrical equipment were then supplied with alternative dielectric media and replacement products entered the market. However, some countries outside the EU and North America continued to produce these substances until recently. Hence, products from those countries may have continued to contain PCBs until the mid-1990s. Current releases to the environment arise principally from the closed electrical appliances that still exist, as their useful life could be up to 40 years.

Activity data are very difficult to obtain on quantities of PCBs in existing transformers and associated leakage rates. A National Inventory of PCB Holdings for Ireland was originally prepared in 2001. This inventory has been updated a number of times, the most recent data corresponding to 2016. The report for this inventory provides an estimate of the total volume of PCB oil (confirmed and suspected) for 2016 of 12.98 m³. This estimate includes both inventoried (confirmed) large and small holdings and estimated non-inventoried (suspected) holdings. Indications are that this is an overestimate and that many of the suspected holdings do not contain any PCBs. The estimate of holdings for 2016 represents a substantial decrease on the peak value in 2009 (522.06 m³) following a large decrease in 2008 (114.29 m³). This is partly due to methodological changes in the inventory compilation, which has given rise to a step change in the emission estimates. This issue requires further investigation to determine the level of inconsistency that may have been introduced across the current time series. The European Union's Chemical Legislation European Enforcement Network (CLEEN) initiated a project to compare inventories of PCBs in Member States of the EU. The CLEEN project documents summarise a large amount of information held within the EU offices (on PCB stocks) that have been reported by Member States to the EC but have not to date been published or synthesised by the Commission itself. Analysis of the CLEEN data indicates that Ireland has a lower than average PCB per-capita stock when compared with other Member States. All of this qualitative information points towards a lower than

average prevalence of PCB-containing materials within electrical equipment in Ireland and this has been taken into consideration in the estimation method used for category 2L.

The derivation of activity data outlined above provides a time series of estimates of PCB-containing oil stocks in Ireland, based on a worst-case assumption that all of the as yet unreported transformer stocks do contain PCBs. The estimates range from 417,620 dm³ of oil in 1990 to 12,983 dm³ of oil (as reported by the EPA) in 2016. Data from the UK NAEI indicate that annual emissions of PCBs derived from dielectric fluid stocks can be estimated as 0.0005 kg PCBs/kg fluid, of which emissions to air comprise 0.06 g PCBs/kg emitted, with the remainder emitted to land. In the absence of source activity and monitoring data, these factors have been used to estimate Ireland's PCB emission estimates. The time series of PCB emissions from leakage of electrical equipment is presented in Table 4.46. As noted above, the current data gives rise to a sharp increase in emissions for 2009 to decrease again in 2010 and further in each consecutive year, caused by a change in the methodology used for estimating the volume of dielectric fluid containing PCBs. Total quantity of PCB containing oil contained in equipment in-situ at start of year for 2016 is substantially lower than previous years, resulting in decreased emissions.

The data on PCDD/F concentrations in dielectric PCB fluid from Dyke (1997) give a concentration in PCB dielectric fluid of 83.5 µg I-TEQ/kg of PCBs. It is assumed that the evaporation rate is the same for PCBs and PCDD/F so that for every kilogram of PCBs that is emitted to air, 83.5 µg I-TEQ of PCDD/F are emitted. Using this factor, estimates for PCDD/F emissions to air from dielectric fluid stocks in Ireland have been made. The time series of PCDD/F emissions from leakage from electrical equipment is presented in Table 4.46.

Table 4.46. Time Series of Polychlorinated Biphenyls and Dioxin and Furan Emissions from Leakage from Electrical Equipment

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCDD/F (µg I-TEQ)	1.034	1.034	0.842	0.790	0.784	0.752	0.283	1.293	0.625	0.546	0.172	0.061	0.055	0.036	0.032
	0.012	0.012	0.010	0.009	0.009				0.007	0.006	0.002	0.000	0.000	0.000	0.000
PCBs (kg)	4	4	1	5	4	0.009	0.003	0.015	5	5	1	7	7	4	4

4.6.2 Fragmentisers and Shredders (NFR 2L)

The practice of fragmenting or shredding electrical equipment currently occurs in a small number of IPPC-licensed facilities, where any suspected POP-containing components (e.g. capacitors) are removed and the residual material is then exported. White goods are also exported for recovery or treatment. The recycling of electrical and electronic goods has also been improved since the introduction of the WEEE Regulations in 2005. However, prior to the commencement of the WEEE Regulations and the All Island Fridge & Freezer Collection and Export Scheme in 2004, it is possible that white goods may have been shredded within Ireland, although there is little evidence that such practice was widespread. To provide a worst-case estimate for this potential emission, the UK NAEI activity data have been scaled on a per-capita basis to prepare estimates for Ireland for 1990–2005.

The shredding of End-of-Life Vehicles (ELVs) is another operation that may result in the possible emissions of POPs, and it has been found to be a relatively significant source in other European countries. Currently, two companies operate ELV shredders at three locations. The larger company operates two shredders but undertakes no monitoring of POPs on incoming vehicles or auto residue post-shredding. However, due to the ELV regulations all vehicles are “de-polluted” either on-site or prior to receipt from dismantlers, with all suspected contaminated materials being removed. In addition, the de-polluting process is expected to further improve in future years. Approximately 30,000 tonnes of auto residue (de-polluted vehicles) are shredded annually. Prior to the implementation of the ELV

regulations the entire intact vehicle was shredded. Therefore, it can be assumed that for earlier years the shredding of ELVs would have resulted in larger quantities of shredded auto residue, with a higher potential for release of POPs.

Very limited data are available on emissions of POPs to air from fragmentisers, especially for the early part of the time series. Emission estimates for POPs are based on the Inventory Guidebook (2000) factor of 0.004 g/capita/year for PCB emissions from fragmentisers, which is considered to apply in the early part of the time series, around the time of the banning of PCBs (1985). The starting point for the time series of estimates of emissions from fragmentisers in Ireland is the estimated emissions in 1986 using population data and the factor of 0.004 g/capita/year, and this leads to an initial estimate of 14 kg PCBs emitted to air in Ireland in 1986. Assuming a 20-year lifespan of electronic equipment, it is reasonable to assume that 5 per cent of the 1986 emissions are removed each year, as old PCB-containing equipment is disposed to landfill and new PCB-free equipment is used as replacements. This assumption leads to an estimated time series of PCB emissions to air of 11.9 kg in 1990, falling to zero emissions by 2006 and are reported as NO for the period 2006 to 2016. Although this is a very broad “top-down” approach and is subject to significant uncertainty, there are very little additional data available to inform more accurate estimates. Emission estimates for the time series are presented in Table 4.47.

Table 4.47. Emission Time Series for Polychlorinated Biphenyls from Fragmentisers and Shredders

Year	1990	1995	2000	2001	2002	2003	2004	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCBs (kg)	11.92	8.64	5.31	4.62	3.92	3.18	2.43	1.65	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO

4.7 Recalculations in the Industrial Processes Sector

4.7.1 Heavy Metals recalculations

Recalculations occurred to emission estimates for 1990–2015 in sector 2A3 as shown in Table 4.49. This was due to the update of emission factors for Cadmium (Cd) from the UK NAEI emission factors to the 2016 Guidebook (EMEP/EEA, 2016).

4.7.2 Particulate matter recalculations

There were no recalculations to emission estimates for 1990-2015 as can be seen in table 4.50.

4.7.3 Nitrogen Oxides (NOx) recalculations

There were no recalculations to emission estimates for 1990-2015 as can be seen in table 4.50.

4.7.4 Non- Methane Volatile Organic Compounds (NMVOC) recalculations

Recalculations occurred in this reporting round due to new sources of activity data, corrections to activity data and corrected disaggregation of emissions between categories 2D3i and 2G. All recalculations are presented in Table 4.51.

Coating Application (NFR 2D3d)

Revisions were made to two SNAP codes within this sector resulting in recalculations of an average of 2.1 per cent across the time series. The recalculations include revisions made to SNAP 060103 Paint Application; Construction and Buildings and 060104 Paint application; Domestic uses (except 060107). An error was found in the units which were used to

calculate the solvent/ water based paint split. This was corrected and caused a change in the emissions estimates for 1990-2015.

Other solvent use (NFR 2D3i)

No recalculations occurred however 2D3i categories were removed from the category 2G and correctly reported under 2D3i.

Other product use (NFR 2G)

Recalculations occurred in this sector due to changes in Preservation of wood (SNAP 060406). The methodology for this sub category was corrected and resulted in recalculations of an average of 343 per cent reduction across the time series. The removal of 2D3i categories from the 2G category as well as changes to Preservation of wood resulted in an average reduction of 98 per cent across the time series for 2G.

Food and Beverages Industry (NFR 2H2)

Recalculations occurred in this sector in; Spirits (SNAP 040608) where the source of activity data was changed to the national statistics data which had previously been unavailable. This resulted in an average increase of 16.15 per cent across the time series in this sector.

4.8 Quality Assurance/Quality Control

Section 4.7 outlines the recalculations which were undertaken in the Industrial Processes sector in this reporting round. The inventory agency will continue to implement QA/QC procedures with respect to the estimates from the Industrial Processes sector in future submissions. The time series spreadsheet system developed for individual categories as described in Section 1.3.5 now allows for simple and efficient checking of activity data, emission factors, annual emissions and aggregated totals. Year-on-year changes immediately highlight any omissions, anomalies or internal errors. Initial checks are conducted by the inventory compiler as part of the calculation process, which is followed by a second check by another member of the inventories team and completion of the QA/QC sheets in calculation workbooks.

4.9 Overview of Notation Keys

Table 4.48 describes the notation keys used in the 2016 NMVOC emission inventory for Other Solvent and Product Use (2D-2L).

Table 4.48. Notation keys used in 2016 NMVOC inventory

NFR	Snap	Description	Notation Key	Reason
2D3a	060411	Domestic use of pharmaceutical products	IE	Included in 060408
2D3e	060203	Electronic components manufacturing	IE	Included under metal degreasing
2D3e	060204	Other industrial cleaning	IE	Included under metal degreasing
2D3g	060311	Adhesive, magnetic tapes, films and photographs manufacturing	IE	Included under 060405

4.10 Planned Improvements

The inventory team will continue to review emission estimates for this sector in light of any new information that may become available for future submissions. In addition, the inventory team also plans to continue to outsource contracts on a periodic basis to re-examine and extend the inventory time series with respect to emissions of heavy metals and POPs. Main priority will be given to estimate emissions of PCDD/F for all years after 2001 and developing estimates of PCB and HCB for the full time-series from iron and steel production sector 2C1.

Following a recommendation from the Stage 3 Review Report, emission estimates for TSP, PM₁₀ and PM_{2.5} for aluminium production (2C3) will be investigated for inclusion in the next submission.

Following a recommendation from the Stage 3 Review Report, the inventory agency will endeavour to undertake a review of particulate matter estimates in cement production. This will consider whether estimates are already included under 1A2f, or whether calculations should be made in 2A1

The levels of solvent use and the emissions from solvents are changing substantially in response to product replacement and reformulation and emission controls being implemented under IPPC and the Solvents Directive (1999/13/EC). The reduction of solvent content has been captured in the methodologies, but this has relied on a number of assumptions, and the collection of real data is required to determine emissions with improved confidence.

In addition, liaison with industry will allow refinement of the estimates for activities subject to licensed controls and to reflect abatement measures in the time series. However, the per capita approach to estimating NMVOC emissions will remain the only option in several important categories, such as SNAP code 060107 (Paint Application: Wood).

Future improvements include refining the assumptions used in the current emission estimate. In particular, future work will investigate estimating historic values for NMVOC emission factors to more accurately reflect both product formulations and emissions in Ireland. The assumptions used in the current emission estimate will be further investigated and refined. These include further investigating product breakdown used in domestic solvent use (SNAP 060408), vehicle refinishing product sales (SNAP code 060102), decorative paint product sales (SNAP codes 060103 and 060104) and Meat and Fish frying (040627).

Table 4.49. Recalculations for Industrial Processes 1990–2015 (Heavy Metals)

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
2A3 Glass Production	Pb	t	0.311	0.311	0.311	0.129	0.089	0.097	0.096	0.008	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Cd	t	0.00002	0.00011	0.00017	0.00028	0.00031	0.00032	0.00028	0.00000	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Hg	t	0.001	0.001	0.001	0.001	0.001	0.001	0.001	NE	NO	NO	NO	NO	NO	NO
2C Metal Production	Pb	t	3.473	3.387	2.127	0.041	0.036	0.028	0.027	0.047	0.035	0.016	0.007	0.016	0.001	0.000
2C Metal Production	Cd	t	0.249	0.238	0.241	0.001	0.001	0.000	0.001	0.001	0.008	0.001	0.000	0.001	0.000	0.000
2C Metal Production	Hg	t	0.033	0.031	0.036	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Submission 2018																
2A3 Glass Production	Pb	t	0.311	0.311	0.311	0.129	0.089	0.097	0.096	0.008	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Cd	t	0.007	0.007	0.007	0.000	0.000	0.000	0.000	NE	NO	NO	NO	NO	NO	NO
2A3 Glass Production	Hg	t	0.001	0.001	0.001	0.001	0.001	0.001	0.001	NE	NO	NO	NO	NO	NO	NO
2C Metal Production	Pb	t	3.473	3.387	2.127	0.041	0.036	0.028	0.027	0.047	0.035	0.016	0.007	0.016	0.001	0.000
2C Metal Production	Cd	t	0.249	0.238	0.241	0.001	0.001	0.000	0.001	0.001	0.008	0.001	0.000	0.001	0.000	0.000
2C Metal Production	Hg	t	0.033	0.031	0.036	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
% Change in Emissions																
2A3 Glass Production	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%						
2A3 Glass Production	Cd	%	29835.9%	6868.0%	4370.4%	-12.1%	-17.1%	-21.6%	-25.6%							
2A3 Glass Production	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%							
2C Metal Production	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2C Metal Production	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-33.3%
2C Metal Production	Hg	%	0.0%	0.0%	0.0%											

Table 4.50. Recalculations for Industrial Processes 1990–2015 (Particulates)

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
2A3 Glass Production	TSP	kt	0.023	0.023	0.022	0.006	0.006	0.006	0.005	0.000	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM ₁₀	kt	0.020	0.020	0.019	0.005	0.005	0.005	0.004	0.000	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM _{2.5}	kt	0.018	0.018	0.017	0.005	0.005	0.005	0.004	0.000	NO	NO	NO	NO	NO	NO
2A5a Quarrying and mining of minerals other than coal	TSP	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2A5a Quarrying and mining of minerals other than coal	PM ₁₀	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2A5a Quarrying and mining of minerals other than coal	PM _{2.5}	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2A5b Construction and Demolition	TSP	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2A5b Construction and Demolition	PM ₁₀	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2A5b Construction and Demolition	PM _{2.5}	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2A5c Storage and Handling of mineral products	PM _{2.5}	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2B10b Storage, handling and transport of chemical products	TSP	kt	0.179	0.192	0.173	0.148	0.140	0.133	0.124	0.117	0.142	0.120	0.123	0.149	0.140	0.140
2B10b Storage, handling and transport of chemical products	PM ₁₀	kt	0.057	0.061	0.055	0.047	0.045	0.043	0.040	0.038	0.046	0.039	0.039	0.048	0.045	0.045
2B10b Storage, handling and transport of chemical products	PM _{2.5}	kt	0.007	0.008	0.007	0.006	0.006	0.005	0.005	0.005	0.006	0.005	0.005	0.006	0.006	0.006
2C Metal Production	TSP	kt	0.071	0.071	0.006	0.001	0.002	0.001	0.001	0.002	0.001	0.001	0.000	0.001	NO	NO
2C Metal Production	PM ₁₀	kt	0.060	0.060	0.005	0.001	0.001	0.001	0.001	0.002	0.001	0.001	0.000	0.001	NO	NO
2C Metal Production	PM _{2.5}	kt	0.042	0.042	0.003	0.001	0.001	0.001	0.001	0.001	0.001	0.000	0.000	0.000	NO	NO
2D3b Road Paving	TSP	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2D3b Road Paving	PM ₁₀	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2D3b Road Paving	PM _{2.5}	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2D3i Other solvent use	TSP	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
2D3i Other solvent use	PM ₁₀	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
2D3i Other solvent use	PM _{2.5}	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Submission 2018																
2A3 Glass Production	TSP	kt	0.023	0.023	0.022	0.006	0.006	0.006	0.005	0.000	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM ₁₀	kt	0.020	0.020	0.019	0.005	0.005	0.005	0.004	0.000	NO	NO	NO	NO	NO	NO
2A3 Glass Production	PM _{2.5}	kt	0.018	0.018	0.017	0.005	0.005	0.005	0.004	0.000	NO	NO	NO	NO	NO	NO
2A5a Quarrying and mining of minerals other than coal	TSP	kt	3.213	2.999	6.792	10.110	11.508	11.882	9.838	6.455	4.442	4.090	3.629	3.751	3.716	3.716
2A5a Quarrying and mining of minerals other than coal	PM ₁₀	kt	1.575	1.470	3.330	4.956	5.641	5.825	4.823	3.164	2.178	2.005	1.779	1.839	1.822	1.822
2A5a Quarrying and mining of minerals other than coal	PM _{2.5}	kt	0.158	0.147	0.333	0.496	0.564	0.582	0.482	0.316	0.218	0.200	0.178	0.184	0.182	0.182
2A5b Construction and Demolition	TSP	kt	1.388	1.049	1.560	6.124	4.915	6.598	8.295	7.273	14.419	0.552	0.619	1.296	0.791	1.510
2A5b Construction and Demolition	PM ₁₀	kt	0.417	0.315	0.468	1.835	1.475	1.982	2.484	2.176	4.309	0.166	0.186	0.389	0.237	0.453
2A5b Construction and Demolition	PM _{2.5}	kt	0.042	0.032	0.047	0.183	0.147	0.198	0.248	0.218	0.431	0.017	0.019	0.039	0.024	0.045
2A5c Storage and Handling of mineral products	PM _{2.5}	kt	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2B10b Storage, handling and transport of chemical products	TSP	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2B10b Storage, handling and transport of chemical products	PM ₁₀	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2B10b Storage, handling and transport of chemical products	PM _{2.5}	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2C Metal Production	TSP	kt	0.071	0.071	0.006	0.001	0.002	0.001	0.001	0.002	0.001	0.001	0.000	0.001	NO	NO

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2C Metal Production	PM ₁₀	kt	0.060	0.060	0.005	0.001	0.001	0.001	0.001	0.002	0.001	0.001	0.000	0.001	NO	NO
2C Metal Production	PM _{2.5}	kt	0.042	0.042	0.003	0.001	0.001	0.001	0.001	0.001	0.001	0.000	0.000	0.000	NO	NO
2D3b Road Paving	TSP	kt	33.000	25.500	43.500	51.000	52.500	49.500	42.000	49.500	34.500	27.000	28.500	27.000	27.000	28.500
2D3b Road Paving	PM ₁₀	kt	4.400	3.400	5.800	6.800	7.000	6.600	5.600	6.600	4.600	3.600	3.800	3.600	3.600	3.800
2D3b Road Paving	PM _{2.5}	kt	2.200	1.700	2.900	3.400	3.500	3.300	2.800	3.300	2.300	1.800	1.900	1.800	1.800	1.900
2D3i Other solvent use	TSP	kt	0.022	0.014	0.009	0.016	0.020	0.035	0.022	0.026	0.031	0.061	0.065	0.054	0.038	0.044
2D3i Other solvent use	PM ₁₀	kt	0.018	0.012	0.008	0.013	0.016	0.029	0.018	0.021	0.025	0.050	0.053	0.044	0.031	0.036
2D3i Other solvent use	PM _{2.5}	kt	0.012	0.008	0.005	0.009	0.011	0.019	0.012	0.014	0.017	0.034	0.035	0.030	0.021	0.024
% Change in Emissions																
2A3 Glass Production	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%						
2A3 Glass Production	PM ₁₀	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%						
2A3 Glass Production	PM _{2.5}	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%						
2A5a Quarrying and mining of minerals other than coal	TSP	%														
2A5a Quarrying and mining of minerals other than coal	PM ₁₀	%														
2A5a Quarrying and mining of minerals other than coal	PM _{2.5}	%														
2A5b Construction and Demolition	TSP	%														
2A5b Construction and Demolition	PM ₁₀	%														
2A5b Construction and Demolition	PM _{2.5}	%														
2A5c Storage and Handling of mineral products	PM _{2.5}	%														
2B10b Storage, handling and transport of chemical products	TSP	%														
2B10b Storage, handling and transport of chemical products	PM ₁₀	%														
2B10b Storage, handling and transport of chemical products	PM _{2.5}	%														
2C Metal Production	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	
2C Metal Production	PM ₁₀	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	
2C Metal Production	PM _{2.5}	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	
2D3b Road Paving	TSP	%														
2D3b Road Paving	PM ₁₀	%														
2D3b Road Paving	PM _{2.5}	%														
2D3i Other solvent use	TSP	%														
2D3i Other solvent use	PM ₁₀	%														
2D3i Other solvent use	PM _{2.5}	%														
Submission 2017																
2B2 Nitric Acid Production	NOx	kt	0.960	0.280	0.302	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Submission 2018																
2B2 Nitric Acid Production	NOx	kt	0.960	0.280	0.302	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
% Change in Emissions																
2B2 Nitric Acid Production	NOx	%	0.0%	0.0%	0.0%											

Table 4.51. Recalculations for Industrial Processes 1990–2015 (NMVOC)

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
2D3a Domestic solvent use including fungicides	NMVOC	kt	7.927	8.143	8.568	9.347	9.571	9.894	10.141	10.250	10.298	10.344	10.368	10.385	10.422	10.481
2D3b Road Paving	NMVOC	kt	0.035	0.027	0.046	0.054	0.056	0.053	0.045	0.053	0.037	0.029	0.030	0.029	0.029	0.030
2D3c Asphalt Roofing	NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2D3d Coating Applications	NMVOC	kt	6.047	6.004	6.310	6.238	6.406	6.604	5.452	4.273	3.378	3.381	3.234	3.921	4.294	3.697
2D3e Degreasing	NMVOC	kt	1.744	2.386	0.847	0.953	1.674	1.557	1.531	0.789	0.885	0.925	1.004	0.902	0.922	1.501
2D3f Dry Cleaning	NMVOC	kt	0.282	0.272	0.090	0.085	0.124	0.105	0.093	0.063	0.053	0.058	0.075	0.057	0.068	0.044
2D3g Chemical products manufacturing or processing	NMVOC	kt	3.023	3.024	2.786	1.189	1.247	1.470	1.486	1.888	0.999	1.149	1.224	1.248	1.083	1.341
2D3h Printing	NMVOC	kt	2.912	2.912	1.946	2.126	2.628	2.118	1.393	2.360	2.474	1.788	1.217	1.283	1.400	1.462
2D3i Other solvent use	NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2G Other product use	NMVOC	kt	1.794	1.434	1.412	3.190	3.475	4.871	3.512	2.389	1.806	1.741	1.455	1.112	1.367	1.480
2H2 Food and Beverages Industry	NMVOC	kt	9.033	9.632	9.587	10.279	10.875	10.600	10.904	10.841	12.679	12.274	13.608	14.572	15.016	16.441
Submission 2018																
2D3a Domestic solvent use including fungicides	NMVOC	kt	7.927	8.143	8.568	9.347	9.571	9.894	10.141	10.250	10.298	10.344	10.368	10.385	10.422	10.481
2D3b Road Paving	NMVOC	kt	0.035	0.027	0.046	0.054	0.056	0.053	0.045	0.053	0.037	0.029	0.030	0.029	0.029	0.030
2D3c Asphalt Roofing	NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2D3d Coating Applications	NMVOC	kt	6.743	6.560	6.315	6.238	6.379	6.575	5.432	4.259	3.373	3.377	3.241	3.402	3.814	3.414
2D3e Degreasing	NMVOC	kt	1.743	2.388	0.847	0.953	1.674	1.557	1.531	0.789	0.885	0.925	1.004	0.902	0.922	0.687
2D3f Dry Cleaning	NMVOC	kt	0.282	0.270	0.090	0.085	0.124	0.105	0.093	0.063	0.053	0.058	0.075	0.057	0.068	0.044
2D3g Chemical products manufacturing or processing	NMVOC	kt	3.023	3.024	2.786	1.189	1.247	1.470	1.486	1.888	0.999	1.149	1.224	1.248	1.082	0.986
2D3h Printing	NMVOC	kt	2.912	2.912	1.946	2.126	2.628	2.118	1.393	2.360	2.474	1.788	1.217	1.283	1.400	1.462
2D3i Other solvent use	NMVOC	kt	0.701	0.493	1.011	3.103	3.810	5.157	3.346	2.198	1.457	1.620	1.501	1.122	1.380	1.422
2G Other product use	NMVOC	kt	0.034	0.037	0.039	0.032	0.032	0.031	0.028	0.027	0.024	0.025	0.023	0.020	0.019	0.020
2H2 Food and Beverages Industry	NMVOC	kt	9.617	10.216	9.683	14.059	13.522	14.303	14.332	15.905	18.052	18.184	20.412	22.492	19.031	20.043
% Change in Emissions																
2D3a Domestic solvent use including fungicides	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3b Road Paving	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3c Asphalt Roofing	NMVOC	%														
2D3d Coating Applications	NMVOC	%	11.5%	9.3%	0.1%	0.0%	-0.4%	-0.4%	-0.4%	-0.3%	-0.1%	-0.1%	0.2%	-13.2%	-11.2%	-7.7%
2D3e Degreasing	NMVOC	%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-54.3%
2D3f Dry Cleaning	NMVOC	%	0.0%	-0.6%	-0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3g Chemical products manufacturing or processing	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-26.5%
2D3h Printing	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2D3i Other solvent use	NMVOC	%														
2G Other product use	NMVOC	%	-98.1%	-97.4%	-97.2%	-99.0%	-99.1%	-99.4%	-99.2%	-98.9%	-98.6%	-98.6%	-98.4%	-98.2%	-98.6%	-98.6%
2H2 Food and Beverages Industry	NMVOC	%	6.5%	6.1%	1.0%	36.8%	24.3%	34.9%	31.4%	46.7%	42.4%	48.1%	50.0%	54.4%	26.7%	21.9%

Chapter Five

Agriculture

5.1 Overview of the Agriculture (NFR 3) Sector

The Agriculture sector is the largest source of NH₃ in Ireland and at 115.53 kt accounted for 99.0 per cent of national total NH₃ emissions (116.70 kt) in 2016. The majority of the remainder of national total emissions is attributable to Road Transport (1A3b) and smaller fractions to combustion in the Manufacturing Industry and Construction (1A2), Commercial (1A4a) and Residential (1A4b), Agriculture/Forestry/Fishing: Off-road vehicles and other machinery (1A4cii) sectors, Other product Use (2G) and Biological treatment of waste – Composting (5B1).

Emissions of NH₃ from agriculture in Ireland's inventory are calculated using a Tier 2 approach developed by a member of the inventory team. The methodologies employed follow those utilised by Misselbrook et al. (2004, 2010, 2016) and the Inventory Guidebook (EMEP/EEA, 2016) and are described in the following sections. The methodology is based largely on the UK National Ammonia Reduction Strategy Evaluation System (NARSES) model for emissions from livestock and where required the Inventory Guidebook (EMEP/EEA, 2016) is also used (poultry and minor livestock species). For the calculation of NH₃ emissions from nitrogen fertilizer application the Tier 2 approach provided in the Inventory Guidebook (EMEP/EEA, 2016) is adopted.

The trend in emissions of NH₃ from agricultural sources is shown in Figure 5.1. Management of animal manures (3B) produced 47.7 per cent of NH₃ emissions from agriculture in 2016, with the application of inorganic fertiliser, sewage sludge and animal manures deposited and applied to soils (52.3 per cent) accounting for the remainder. The NH₃ emission trend is largely determined by the cattle population and shows a steady increase up to 122.20 kt in 1998 (an increase by 11.6 per cent from 1990). There has been some decline in the populations of cattle and sheep since 1999, as well as a decrease in fertiliser use, which contributed to a downturn in NH₃ emissions in the 1999–2011 period. The NH₃ emissions from the agriculture sector in 2016 were 5.5 per cent higher than the emission levels in 1990 (109.48 kt) and 5.5 per cent lower than the peak levels in 1998.

Emissions of NMVOC, TSP, PM₁₀ and PM_{2.5} from manure management (3B) were estimated for the first time in the 2015 submission. Emissions of NO_x are estimated for the first time in this submission.

NMVOCs from agriculture are estimated to be 45.90 kt in 2016, accounting for 42.4 per cent of the NMVOC inventory total (108.25 kt). Emissions from manure management (3B) make up 91.7 per cent of agriculture related NMVOC in 2016, with the remaining 8.3 per cent from cultivated crops (3De). For the calculation of NMVOC emissions, the Inventory Guidebook (EMEP/EEA, 2016) Tier 1 approach is adopted for cultivated crops (3De) and the Tier 2 approach is utilised for manure management (3B). The trend in NMVOC emissions from agriculture is shown in Figure 5.2.

NO_x emissions from agriculture are estimated to be 31.77 kt in 2016, accounting for 29.6 per cent of the national total. Emissions from manure management (3B) make up 2.6 per cent of the agriculture related NO_x with Inorganic N-fertilizers (3Da1), Animal manure applied to soils (3Da2a), Sewage sludge applied to soils (3Da2b), and Urine and dung deposited by grazing animals (3Da3) accounting for 41.1 per cent, 18.7 per cent, 0.0 per cent and 37.6 per cent of total agriculture emissions, respectively. Emissions in 2016 were 3.4 per cent below those in 1990 (32.90 kt). Estimates from manure management (3B) are derived from the Tier 2 mass flow approach to estimate NH₃ emissions, whilst emissions from agricultural soils (3D) utilise the Tier 1 emission factor presented in the Inventory Guidebook. The trend in NO_x emissions from agriculture is shown in Figure 5.3.

Estimates for agriculture emissions of TSP, PM₁₀ and PM_{2.5} contribute significantly to national totals for these pollutants, accounting for 24.3 per cent, 35.8 per cent and 12.1 per cent of national totals respectively. For these calculations, Tier 1 and 2 approaches provided in the Inventory Guidebook (EMEP/EEA, 2016) are adopted. For PM_{2.5} Dairy cattle (3B1a) and Non-dairy cattle (3B1b) are the main drivers of the trend accounting for 74.2 per cent of emissions in 2016. For PM₁₀ the main driver of the trend in emissions from agriculture is Inorganic N-fertilizers (3Da1) which accounted for 68.6 per cent of emissions in 2016. The second and third largest sources are Non-dairy cattle and Dairy cattle (3B1a) accounting for 10.9 per cent and 9.1 per cent, respectively of the agricultural total in 2016. A similar trend is evident for TSP with Inorganic N-fertilizers (3Da1), Non-dairy cattle and Dairy cattle (3B1a) accounting for 46.1 per cent, 16.3 per cent and 13.4 per cent, respectively of the agriculture sector total in 2016. The trend in PM_{2.5} is presented in Figure 5.4.

The use of some pesticides for arable farming can be a source of POP emissions, notably due to the trace content of HCB within some pesticides as a contaminant from the manufacturing process. Emissions of HCB from NFR Sector 3Df (Use of pesticides) are a key category in 2016, accounting for 68.6 per cent of national total HCB emissions.




Figure 5.1 Emission Trend for Ammonia from Agriculture 1990–2016




Figure 5.2 Emission Trend for NMVOC from Agriculture 1990–2016




Figure 5.4 Emission Trend for NO_x from Agriculture 1990–2016




Figure 5.3 Emission Trend for PM_{2.5} from Agriculture 1990–2016

5.2 Manure Management (NFR 3B)

The following sections outline the activity data, assumptions and calculations utilised in estimating NH₃, NMVOC, NO_x, TSP, PM₁₀ and PM_{2.5} emissions from agriculture in Ireland. For NH₃ and NO_x, the Tier 2 methodology uses a mass flow approach based on the concept of the flow of Total Ammoniacal Nitrogen (TAN) through the manure management system. Emissions are calculated for the same animal sub-categories as those utilised in Ireland's national greenhouse gas inventory (Table E.1 Annex E). The first step in the mass flow approach is the estimation of total annual nitrogen excretion by the animals. For dairy cows and other cattle, Ireland utilises the method described in IPCC (2006), chapter 10, further enhanced by country specific data on feeding practices and milk production (O'Mara, 2007, Duffy et al., 2018) to estimate N excretion. For all other categories of livestock, national values are utilised. Total nitrogen excretion is then apportioned to that which is deposited in buildings, collection yards (only applicable to dairy cows during lactation) and grazing. See Table E.3 Annex E for animal nitrogen excretion.

The method used to estimate NMVOC emissions is the tier 2 approach based on emission factors from Table 3.11 and Table 3.12 of the Inventory Guidebook (EMEP/EEA, 2016) for all livestock categories. The tier 2 approach considers NMVOC emissions from the following; silage stores, silage for feeding, housing, outdoor manure stores, manure application and from grazing animals as outlined in section 3.4 of the Inventory Guidebook (EMEP/EEA, 2016).

For TSP, PM₁₀ and PM_{2.5} emissions are estimated based upon animal numbers using the Tier 1 method and emission factors from the Inventory Guidebook (EMEP/EEA, 2016). See Tables E.6 to E.9 of Annex E for additional information of the EFs used for NMVOC, TSP, PM₁₀ and PM_{2.5} emission estimation.

5.2.1 Cattle (NFR 3B1)

A Farm Facilities Survey conducted in 2003 (Hyde et al., 2008) provides the basis for the calculation of the number of days housed and the number of days spent grazing by cattle on farms in Ireland. National averages are used for the purpose of inventory calculations. Data for the number of days housed are presented in Table E.2.1. The number of days spent grazing is then calculated by subtracting these values from 365 (i.e. days in a year).

Two housing types are distinguished for cattle production systems in Ireland – liquid (slurry-based) and solid-manure-based housing. As a result of differing management practices on farms, a proportion of each of the cattle subdivisions is not housed (out-wintered) and therefore graze pasture for the full year. The proportion of each sub-category of cattle that is managed in this manner is accounted for in Table E.2.1. For liquid manure based housing an emission factor of 27.7 per cent of the TAN available in liquid based housing is applied (Misselbrook et al., 2016). The TAN in cattle manure is assumed to be 60 per cent. For solid manure based housing, emission factors of 16.8 per cent and 4.2 per cent of the TAN available in solid manure based housing are applied to cattle housed on straw and calves housed on straw, respectively (Misselbrook et al., 2016).

In addition to animal housing, emissions are estimated for cow collecting yards used during milking utilising an emission factor of 22.5 per cent of the TAN available (Misselbrook et al., 2016).

The storage of both liquid and solid manure is considered. Liquid manure is stored either below the animals in slatted floor housing or removed from the house to outdoor storage. Emissions are calculated separately for indoor and outdoor storage. It is assumed that a crust will form in the indoor under slat storage. In addition, a fraction of the organic nitrogen in liquid manure is mineralised to TAN before emissions are calculated. A value of 0.1 (Dammgen et al., 2007) is applied as suggested in the Inventory Guidebook (EMEP/EEA, 2016). To fully account for all losses of N from liquid manure during storage estimates are made of N_2O , NO and N_2 losses during storage utilising the emission factors provided in the Inventory Guidebook (EMEP/EEA, 2016). An emission factor of 5 per cent of the TAN available in liquid manure stores is applied to estimate NH_3 emissions from liquid manure storage in covered stores and 10 per cent of the TAN available in liquid manure stored in uncovered stores (Misselbrook et al., 2016).

Solid manure is generally stored in the shed or outside in heaps. The contribution of the nitrogen content of straw used for bedding is accounted for based on national data on straw used for bedding and the length of the housing period (Hyde et al., 2008). Where manures are managed as solid, a fraction of the TAN is immobilised in organic matter. Immobilisation of nitrogen reduces the potential for NH_3 emissions from solid manures during storage and after landspreading. The value proposed in the Inventory Guidebook (EMEP/EEA, 2016) of 0.0067 kg kg⁻¹ (Kirchmann and Winter, 1989) is applied. To fully account for all losses of N from solid manure during storage estimates are made of N_2O , NO and N_2 losses during storage utilising the emission factors provided in the Inventory Guidebook (EMEP/EEA, 2016). An emission factor of 35 per cent of the TAN available in solid manure stores is utilised (Misselbrook et al., 2016).

Landspreading emissions are calculated by estimating the quantity of TAN available post storage of the manure (both liquid and solid) and accounting for the period of the year in which it is spread (i.e. spring, summer, autumn and winter) as outlined in Table E.5 Annex E. For liquid manure, a dry matter range of 6 per cent is assumed and two emission factors are applied, 48.4 per cent of the TAN available for the proportion applied in summer and 26.1 per cent for proportion applied in spring, autumn and winter (Misselbrook et al., 2016). For solid manure an emission factor of 68.3 per cent of the TAN available is applied regardless

of the period of the year in which it is spread (Misselbrook et al., 2016). It is assumed that all cattle slurry is applied using the splashplate method and that all solid manure is broadcast spread. The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO per kg of nitrogen is applied to calculate NO_x emissions from manure (solid and liquid) application to soil.

For cattle grazing an emission factor of 6 per cent of the TAN available at grazing is applied for all cattle categories (Misselbrook et al., 2016). The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO per kg of nitrogen is applied to calculate NO_x emissions from excreta deposited during grazing.

NM VOC emissions are estimated using Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2016). Estimates are made for housed cattle, silage feeding, silage store, manure storage, manure application and grazing. The emission factors for silage feeding and silage storage, housing and grazing are combined with the feed intake values (MJ feed intake) from Ireland's Tier 2 approach to the estimation of CH₄ emissions from enteric fermentation and manure management. The feed intake values for silage and grass are presented in Table E6.2, Annex E with the emission factors used presented in Table E6.1. Emissions of NM VOC from manure storage and manure application used the Inventory Guidebook (EMEP/EEA) approach whereby the NM VOC emissions from livestock buildings are coupled with the ratio of NH₃ emissions from storage and application with those in livestock buildings, respectively.

Emissions of TSP, PM₁₀ and PM_{2.5} are estimated using Tier 1 emission factors split by cattle type and housing category (slurry/solid housing). The emission factors used are presented in Tables E.7, E.8 and E.9, Annex E.

5.2.2 Sheep (3B2)

Sheep in Ireland are categorised into those on upland and those on lowland areas. Four subcategories exist within both upland and lowland areas, namely ewes, rams, lambs and other sheep more than 1 year old. The CSO publishes sheep population statistics on an annual basis and, to derive the number of head on both lowland and upland areas, several assumptions are made as follows based on expert opinion. Total ewe and ram numbers are taken as the mean of the June and December CSO censuses. On this basis, the number of ewes and rams are subdivided using the ratio 55:45 (lowland/upland) for years up to and including 1997. For 1998 to 2004 inclusive, a ratio of 70:30 is used, reflecting the destocking of upland areas. For 2005 onwards, a ratio of 80:20 is used. The total number of lambs slaughtered in any 1-year period is used as the activity data for lambs. Monthly lamb slaughtering figures are available from the CSO. From 2001 onwards, these numbers are adjusted for the number of lambs that originate in Northern Ireland and that are slaughtered in Ireland and for the number of lambs that are reared in the Republic but slaughtered in Northern Ireland. National totals are then subdivided similarly to the other categories of sheep. The numbers of other sheep over 1 year old are calculated from unpublished CSO data. Population statistics for each subcategory of sheep are presented in Table E.1, Annex E. Input data with respect to manure management practices are presented in Table E2.2 Annex E. Nitrogen excretion coefficients for all sub categories of sheep are provided in Table E.3 Annex E.

Similar to cattle, the proportion of TAN in the nitrogen excreted by sheep is assumed to be 60 per cent. The emission factor for sheep housing is derived by adjusting the housing emission factor for beef cattle on solid manure systems by the ratio of excretal outputs of sheep and beef cattle which is then back-calculated to derive a value of 21.6 per cent of the TAN available in sheep housing (Misselbrook et al. 2016). Information on the number of days that sheep are housed during the winter period is derived from the Farm Facilities

Survey (Hyde et al., 2008), which suggests sub-category specific housing period lengths (Table E.2.2 Annex E). More specifically, lowland and upland ewes are assumed to be housed for 61 and 85 days/year. No differentiation is made for upland and lowland rams, lambs and other sheep >1 year old- being housed for 85, 58 and 61 days/year respectively. In Ireland, sheep are generally housed in solid-manure-based housing systems.

The NH_3 emission factors used for the storage and landspreading of solid manure used for Cattle (3B1) are also considered appropriate for sheep manure (Misselbrook et al., 2016). For storage of solid manure and subsequent landspreading the emission factor for solid manure from cattle is used (68.3 per cent of TAN available at landspreading). Account is also taken of the nitrogen added from straw used for bedding and the immobilisation of TAN in organic matter when solid manures are managed following the approach adopted for solid manure from cattle. The emission factors presented in the Inventory Guidebook (EMEP/EEA, 2016) are used to estimate NO , N_2O and N_2 emissions from manure management. The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO per kg of nitrogen is applied to calculate NO_x emissions from excreta deposited during grazing and the application of solid manure to soil. For sheep grazing, the emission factor applied for cattle of 6 per cent of the TAN available at grazing is utilised to estimate NH_3 emissions (Misselbrook et al., 2016).

NM VOC emissions are estimated using Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2016) based on volatile solid (VS) excretion data as estimated in the national greenhouse gas inventory for agriculture. Estimates are made for housed sheep, manure storage, manure application and grazing sheep. The emission factors used are presented in Table E.6, Annex E.

Emissions of TSP, PM_{10} and $\text{PM}_{2.5}$ are estimated using Tier 1 emission factors in the Inventory Guidebook (EMEP/EEA, 2016) The emission factors used are presented in Tables E.7, E.8 and E.9, Annex E.

5.2.3 Swine (NFR 3B3)

Detailed population statistics are available for seven subcategories of pigs in Ireland using national statistics published by the CSO as follows: sows in pig, gilts in pig, other breeding sows, boars, gilts not yet served and two categories of fattening pigs (<20 kg and >20 kg live weight). The CSO undertakes and publishes two censuses per year, one in June and one in December. The average of the two census values is used in deriving the pig populations in the seven subcategories, thus providing an appropriate measure of the number of pigs on farms for the purposes of the annual NH_3 emission inventory.

For the NH_3 inventory, it is assumed that all pigs are housed and that the housing systems are liquid/slurry-based. Furthermore, it is assumed that the proportion of TAN in nitrogen excreted is 70 per cent. An emission factor of 33.2 per cent of the TAN in slurry produced by gilts in pig, gilts not yet served and pigs greater than 20 kgs is applied. For sows in pig, other sows for breeding and boars, an emission factor of 19.0 per cent of the TAN produced in the slurry of these pig sub-categories is used and an emission factor of 14.8 per cent of the TAN in slurry produced by pigs under 20 kg (Misselbrook et al., 2016).

For slurry storage, emissions of NH_3 are calculated separately for covered and uncovered stores based on the proportion of slurry stored in covered and uncovered storage (Hyde et al., 2008). Emission factors of 13.0 per cent of the TAN available in covered storage and 52.0 per cent of the TAN available in uncovered slurry stores are applied (Misselbrook et al., 2016). As is the case with cattle slurry account is taken of the losses of N_2O , NO and N_2 using the emission factors presented in the Inventory Guidebook (EMEP/EEA, 2016) and the

mineralisation of organic nitrogen in the liquid manure is also accounted for (Dammgen et al., 2007). It is assumed that all pig slurry is applied using the splashplate method. An emission factor of 19.0 per cent (Misselbrook et al., 2016) of the TAN available post storage is applied on the basis that pig slurry contains less than 4 per cent DM.

The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO per kg of nitrogen is applied to calculate NO emissions from the application of liquid manure to soil.

NMVOC emissions are estimated using Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2016) based on volatile solid (VS) excretion data as estimated in the national greenhouse gas inventory for agriculture. Estimates are made for housed pigs, manure storage and manure application. The emission factors used are presented in Table E.6, Annex E.

Emissions of TSP, PM₁₀ and PM_{2.5} are estimated using Tier 1 emission factors the Inventory Guidebook (EMEP/EEA, 2016) The emission factors used are presented in Tables E.7, E.8 and E.9, Annex E.

5.2.4 Poultry (NFR 3B4g)

Detailed population statistics are available for eight subcategories of poultry in Ireland using national statistics collated by the Department of Agriculture and CSO as follows: layers, broilers, layer breeders, broiler breeders, turkeys, turkey breeders, geese and ducks. The population statistics are provided in Table E.1, Annex E. The estimation of NH₃ emissions from poultry production utilises bird places as opposed to bird numbers so that production cycles are considered. The number of bird places is estimated from the annual bird population assuming that all bird places are full throughout the year after rest periods have been taken into account (rest periods are those periods after a production cycle in which the housing systems are emptied of all manure and bedding, thoroughly washed and prepared for the next batch of birds). In the case of broilers, there are 5.5 production cycles per year, with a 3- to 4-week rest period between production cycles. In the case of turkeys, there are 2.5 production cycles per year of approximately 120 days in length, with a 3- to 4-week rest period also applied. All other poultry subcategories have production cycles of over 1 year, and therefore no adjustments to population statistics are made.

It is assumed that all poultry are housed in some form of solid manure housing system and that the proportion of TAN in nitrogen excreted by poultry is 70 per cent. For a proportion of laying birds, free-range systems are in use, which consist of an area of grassland beside the bird house which the birds are allowed onto for a period of hours during the day. The percentage of laying birds that are housed in this type of system is based on statistics supplied by the Department of Agriculture, Food and Marine. An emission factor of 35 per cent of TAN is applied to the quantity of TAN which is deposited outdoors. Emission factors of 41.0 per cent, 28.0 per cent, and 35.0 per cent, 57 per cent and 24.0 per cent of TAN are applied to layer, broiler, turkey, geese and duck housing, respectively (Inventory guidebook, 2016). The storage of poultry manure is separated into two classes, litter and layer manure. An emission factor of 17.0 per cent of TAN is applied to broiler manure, 14.0 for layer manure, 24.0 per cent for turkey manure, 16.0 per cent for geese manure and 24.0 per cent for duck manure (Inventory Guidebook, 2016). To account for the loss of other nitrogen compounds, account is taken of the losses of N₂O, NO and N₂ using the emission factors presented in the Inventory Guidebook (EMEP/EEA, 2016). It is assumed that all poultry manure is broadcast spread and emission factors of 69.0 per cent, 66.0 per cent, 54.0 per cent, 45 per cent and 54.0 per cent of the TAN available post storage, are adopted for layers, broilers, turkeys, geese and ducks respectively (EMEP/EEA, 2016). The emission factor presented in Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO per kg of

nitrogen is applied to calculate NO emissions from excreta deposited during grazing and the application of manure to soil.

NMVOC, TSP, PM₁₀ and PM_{2.5} emissions are estimated using Tier 1 emission factors from the Inventory Guidebook (EMEP/EEA, 2016) The emission factors used are presented in Tables E.6 – E.9, Annex E.

5.2.5 Other livestock – Goats (NFR 3B4d), Horses (3B4e), Mules and Asses (NFR 3B4f) and Other Animals (3B4h)

The remaining livestock categories include goats, horses, mules and asses, deer, mink and foxes and estimates are made for emissions of NH₃, NO, NMVOC, TSP, PM₁₀ and PM_{2.5}.

For NH₃ emissions the emission factors used are the default Tier 2 factors presented in the Inventory guidebook (EMEP/EEA, 2016) for each step in the manure management chain. As part of the mass flow approach inventory guidebook emission factors for NO, N₂O and N₂ emissions from manure management are also estimated. The emission factor presented the Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO per kg of nitrogen is applied to calculate NO_x emissions from excreta deposited during grazing and the application of manure to soil.

NMVOC emissions are estimated for goats (3B4d), Horses (3B4e) and Mules and asses (3B4f) using the Tier 2 emission factors from the Inventory Guidebook (EMEP/EEA, 2016). In the case of other animals (3B4h) the Tier 1 emission factors presented in the inventory guidebook (EMEP/EEA, 2016) are used. The emission factors applied are presented in Table E.6, Annex E.

Tier 1 emission factors from the Inventory Guidebook (EMEP/EEA, 2016) are applied to estimate emissions of TSP, PM₁₀ and PM_{2.5} from the livestock categories described except for deer for which no emission factor is supplied in the Inventory Guidebook (EMEP/EEA, 2016).

5.2.6 Uncertainties

There is extensive and up-to-date statistical data on all aspects of the agriculture sector in Ireland. Most this data is compiled and published by the Central Statistics Office and is the official source of the basic data for inventory purposes. The exception is for statistics on synthetic fertiliser use and the poultry population which are obtained from the Department of Agriculture Food and the Marine (DAFM). The CSO and DAFM are key data providers whose annual statistical inputs to the inventory agency are covered by Memorandum of Understanding (MOU) in Ireland's national inventory system. As a result, the uncertainty associated with animal population statistics is low at 1 per cent. The emission factor uncertainty associated with NH₃ emission factors for dairy and other cattle is 50 per cent and for all other livestock categories 100 per cent. Much of NH₃ emission research is aimed at dairy cattle and other cattle therefore emissions from these categories are relatively well quantified in comparison to the other livestock categories. In comparison, the uncertainties associated with NMVOC and PM₁₀ emissions in agriculture are large due to the uncertainty (300 per cent) associated with the emission factors for both pollutants as discussed in the Inventory Guidebook (EMEP/EEA, 2016).

5.3 Agricultural Soils (NFR 3D)

5.3.1 Direct Soil Emissions - Inorganic N-fertilizers (NFR 3Da1)

The calculation of NH_3 emissions from nitrogen fertilizer application to agricultural soils utilises the Tier 2 approach outlined in the Inventory Guidebook (EMEP/EEA, 2016). Total fertilizer sales and emission estimates for each year of the time series 1990-2016 (Table E.4, Annex E) are apportioned into the categories, Ammonium sulphate, CAN, NK mixtures, NPK mixtures, NP mixtures, Other straight N compounds, Urea, and protected urea products according to the known sales of these compounds in each year as supplied to the inventory agency by the DAFM. Table E.10 of Annex E includes additional information on the methodological approach, quantities of individual fertilizer types, the emission factors used and resultant emissions. The emission factor presented in the Inventory Guidebook (EMEP/EEA, 2016) of 0.04 kg NO_x per kg of nitrogen is applied to calculate NO_x emissions from the application of inorganic N-fertilizers to soil. Ireland reports emissions of PM_{10} , $\text{PM}_{2.5}$ and TSP from farm level agricultural operations under 3Da1 to separate these emissions from those previously reported under 2B10b which are now reported under 3Dc following a recommendation in the NECD expert review of Ireland's 2017 submission. The default emission factors presented in the Inventory Guidebook (EMEP/EEA, 2016) are applied (Table E.10, Annex E) coupled with the total utilisable agricultural area (Table E.11, Annex E).

5.3.2 Direct Soil Emissions – Livestock manure applied to soils (NFR 3Da2a)

The calculation of NH_3 emissions from livestock manure applied to soil is discussed in sections 5.2.1 to 5.2.5 inclusive. Emissions for each of the livestock species are summed and reported under 3Da2a. The default emission factor of 0.04 kg NO_x per kg of nitrogen is applied to calculate NO_x emissions from the application of livestock manure to soils.

5.3.3 Direct Soil Emissions – Sewage sludge applied to soils (NFR 3Da2b)

Estimates of NH_3 were estimated from the application of sewage sludge (3Da2b) for the first time in Ireland's 2015 submission. The quantity of sewage sludge applied to land is estimated as part of the calculations for emissions of CH_4 and N_2O from wastewater in Ireland's Greenhouse Gas Inventory. The fraction of nitrogen volatilised is 0.13 kg $\text{NH}_3\text{-N}$ per kg N applied (EMEP/EEA, 2016). With respect to NO emissions the default Tier 1 emission factor of 0.002 kg NO_2 capita⁻¹ is utilised to estimate NO emissions.

5.3.4 Direct Soil Emissions – Other organic fertilizers applied to soils (NFR 3Da2c)

Emissions of NH_3 and NO from this source category are currently reported as NE. Information on the quantities of other organic fertilizers applied to agricultural soils does not exist in Ireland and is in comparison to the total quantity of nitrogen applied in manures or deposited during grazing negligible.

5.3.5 Direct Soil Emissions – Urine and dung deposited during grazing (NFR 3Da3)

The calculation of NH_3 emissions from urine and dung deposited on soil during grazing is discussed in sections 5.2.1 to 5.2.5 inclusive. Emissions for each of the livestock species are summed and reported under 3Da3. The default emission factor of 0.04 kg NO_x per kg of nitrogen is applied to calculate NO_x emissions from urine and dung deposited on soil during grazing.

5.3.6 Farm-level agricultural operations (NFR 3Dc)

Emissions of PM₁₀, PM_{2.5} and TSP are reported under NFR 3Dc. Previously the emission reported in this category were reported under 2B10b, however in response to the 2017 NECD expert review, these emissions have now been reallocated to 3Dc. Emissions are estimated using the total quantity of fertilizers (by weight) applied to soils in Ireland. The emission factors applied are those provided in CEPMEIP (2001) and are presented in Table E.10 Annex E. The total quantity of fertilizers (by weight) applied to agricultural soils is presented in Table E.11, Annex E.

5.3.7 Off-farm storage, handling and transport (NFR 3Dd)

In this category, fugitive PM emissions in the form of TSP, PM₁₀ and PM_{2.5} are estimated from the bulk handling of cereal grains. The general method for estimating fugitive PM emissions involves multiplying the amount of material, which in this case is cereal grain (barley, wheat and oats), by an emission factor. Given the importance of agriculture to Ireland's economy, production statistics are freely available (Table E.11, Annex E). Data in relation to the production of cereal grains are collated and provided by the Central Statistics Office (CSO). Emission factors of 100 g/t, 25 g/t and 4 g/t from CEPMEIP (2001) are utilised in the calculation of emissions of TSP, PM₁₀ and PM_{2.5}, respectively.

5.3.8 Cultivated crops (NFR 3De)

Emissions of NMVOCs are estimated using the default emission factor of 0.86 kg/ha and the total utilisable agricultural area presented in Table E.11, Annex E.

5.3.9 Use of Pesticides (NFR 3Df)

The main source of POPs from pesticides is HCB contamination of currently used pesticides. Where available, annual pesticides usage data have been used, although for some years only import data are available and the use of import data to inform the annual usage estimates discounts any consideration of stockpiling (or use of existing supplies) of pesticides. In addition, the emission factors for HCB content in pesticides are typically based on quantities of the "active" ingredient, not the total weight of the pesticide. Import data are available for the years 1998–2006 for chlorothalonil and quintozone. Further to the import data, information has been gathered on pesticide usage. The Pesticide Usage Survey (No. 1) (DAF, 2003) for grassland and fodder crops quotes the quantity of chlorothalonil (weight of active ingredient) used as 6,903 kg. It should be noted that chlorothalonil is a fungicide mainly used with silage and therefore the greater use of this product is outside the spectrum of the 2003 study. The Pesticide Usage Survey (No. 2) (DAF, 2004) for arable crops quotes the use of chlorothalonil (weight of active ingredient) used as 190,776 kg. This is in comparison with the import that year which was 241,285 kg (weight of active ingredient).

Most emissions from pesticide usage enter the atmosphere due to the spraying application and subsequent volatilisation of POPs from the surface of plants. A paper investigating the ultimate fate of HCB from pesticide application (Xu, 2008) calculates the split of emissions to air, soil and water to be 70.2 per cent, 28.8 per cent and 1 per cent, respectively.

Total HCB emissions have been estimated on this basis. In addition, European legal approval sets a limit for HCB in chlorothalonil as 0.01 g/kg, whilst research in Canada (Benezon, 1999) indicates a working contamination of 0.018–0.026 g/kg, based on data from 1988 to 1998. European legislation also sets a limit for quintozone at 1 g/kg pesticide. Estimates presented here use 0.026 g/kg as a worst-case assumption on a linear progression to 2001 (the year that quintozone was banned), and from 2001 the European standard value has been applied. Emission estimates are presented in Table 5.1.

Table 5.1. Emission Estimates for Hexachlorobenzene Emissions from Pesticide Use

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
HCb (kg)	0.186	0.327	0.476	1.234	1.151	1.151	1.151	1.151	1.151	1.151	1.151	1.151	1.151	1.151	1.151

5.4 Field burning of agricultural residues (NFR 3F)

Field burning of agricultural residues is the practice whereby crop residues are burnt as a means of clearing land to allow tillage operations to proceed. As stated in chapter 3F of the EMEP/EEA Guidebook 2016 the practice is largely outlawed. Prescribed burning of agricultural residues in Ireland may only be allowed in the case of disease or maintenance of soil organic matter as detailed in GAEC 6 page 52 of the cross compliance handbook (DAFM, 2015). In addition, the inventory agency has undertaken some analysis on this issue (Zimmerman, 2014) which suggests that fires on agricultural land are in general fires that have occurred due to fires in adjoining forestry or peatland for example and are not the anthropogenic burning of agricultural residues. The inventory agency continues to discuss with relevant agencies with respect to the identification of fires due to its importance in the estimation of non-CO₂ gases within the Land Use, Land Use Change and Forestry sector.

5.5 Uncertainties associated with Synthetic Fertilizer (3Da1), Organic fertilizers (3Da2c) and urine and dung deposited by grazing animals (3Da3).

Although losses of NH₃ from N-fertilisers applied to grass grazed by livestock are difficult to distinguish from subsequent NH₃ emissions from urine patches produced by grazing animals, those two emissions are calculated separately with emissions from grazing reported in 3Da3. The sources making the largest contributions to the overall uncertainty are ammonia losses from synthetic fertiliser use and animal manures deposited to pasture, range and paddock. The emission factors for these sources are currently assigned an uncertainty of ±200 per cent, and they contribute to 95.6 per cent of the overall uncertainty.

5.6 Recalculations in the Agriculture Sector

Recalculations in the agriculture sector are highlighted in tables 5.2 to 5.6. For NH₃ emissions from agriculture, there is on average 2.4 per cent increase in emission across the timeseries. However, within the sector there are several signification recalculations (Table 5.2) as follows.

Emissions from Dairy cattle (3B1a) are 8.5 per cent lower on average across the timeseries as a result of the application of a revised housing emission factor for slurry based housing from Misselbrook et al (2016). Similarly, emissions from Non-dairy cattle (3B1b) are also reduced, however the development of Tier 2 N excretion values for all cattle categories in this submission has negated any reduction in emissions from the category due to the application of a lower emission factor to livestock housing. Other important recalculations were the correction of a transcription error in the estimation of emissions from swine (3B3), revisions to the AWMS data for sheep (3B2), goats (3B4d), Horses (3B4e) and Mules and asses (3B4f) and the incorporation of the Inventory Guidebook (EMEP/EEA, 2016) emission factors for poultry (3B4gi, 3B4gii, 3B4giii and 3B4giv), other animals (3B4h), Inorganic N-fertilizers (3Da1) and sewage sludge applied to soils (3Da2b) into national emission estimates. The 15.2 per cent on average and 6.3 per cent on average increases in emissions evident for Animal manure applied to soils (3Da2a) and Urine and dung deposited by grazing animals (3Da3), respectively are a direct result of the implementation of Tier 2 N excretion values for Non—dairy cattle (3B1b) in this submission.

Revisions to AWMS data for sheep (3B2), goats (3B4d), Horses (3B4e) and mules and asses (3B4f) are responsible for most the recalculations evident in NMVOC emissions (Table 5.3). Emissions in this submission are on average 1.9 per cent higher across the timeseries when compared against the previous submission. Revisions to some poultry statistics for 2014 and 2015 are responsible for a 15.6 per cent increase in emissions from laying hens (3B4gi) in 2015 and 12.7 per cent and 13.5 per cent, respectively for broilers in 2014 and 2015. There were also minor revisions to the goat population (3B4d). Revised ratios of NH₃ emissions from housing and storage used in Tier 2 estimates are responsible for the minor recalculations to emissions from dairy cattle (3B1a) and non-dairy cattle (3B1b). Finally, emissions from the application of fertilizers to agricultural soils are no longer reported under 3Da1 and are now reported under Cultivated crops (3De) using the approach in the Inventory Guidebook (EMEP/EEA, 2016). This reallocation and revision of emission estimates was undertaken in response to the 2017 NECD expert review of Ireland's 2017 submission.

The inclusion of the emission factors presented in the latest version of the Inventory Guidebook (EMEP/EEA, 2016) are responsible for most the recalculations evident in Tables 5.4, 5.5 and 5.6 for emissions of TSP, PM₁₀ and PM_{2.5}. Revisions to poultry and goat population statistics for the years 2014 and 2015 are a minor source of the recalculations evident for those years. The reallocation of emissions from 2B10b to 3Dc as discussed in section 5.3.6 is also evident.

5.7 Quality Assurance/Quality Control

The general QA/QC procedures set down in Ireland's QA/QC plan have been undertaken for the Agriculture sector. The spreadsheets incorporate transparent linking between input data statistics and calculations, as well as internal checks on the calculations and the outputs. The inventory experts are actively involved in assessing the outcomes of NH₃ emission research in Ireland and continually re-examine the underlying assumptions in inventory estimates with sector-specific experts in the Department of Agriculture and other related bodies.

5.8 Planned Improvements

A large number of input variables determine emissions in the Agriculture sector and the final results are very sensitive to changes in many of these variables. Assumptions relating to some parameters have an important bearing on the outcome. Whilst methodologies for the agricultural emission sources that are relevant in Ireland are now very comprehensive, they remain generalised and necessarily simplified considering the complex systems and processes involved. The key to developing better estimates and reducing uncertainty is to take full account of national circumstances of climate, soil types, livestock- and crop-production practices, manure management systems and other influencing factors in a robust and justifiable manner when applying these methodologies. This requires detailed data from research programmes, and large amounts of statistical data. Nevertheless, the inventory agency is continually developing emission estimates so that they fully reflect national circumstances within the availability of reliable statistics and research studies.

The inventory agency will consider revising the ammonia emission factors, if necessary, based on any updates in the Inventory Guidebook and following the publication of any relevant country specific research for the next annual submission.

Table 5.2. NH₃ Recalculations for Agriculture 1990–2015

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017															
3B1a Manure management - Dairy cattle	NH3	kt	13.89	13.06	12.52	11.34	11.61	11.55	11.43	11.48	11.91	11.99	12.25	12.77	13.89
3B1b Manure management - Non-dairy cattle	NH3	kt	27.42	29.91	30.56	32.57	31.40	31.66	31.23	29.93	29.12	30.92	31.28	30.88	30.67
3B2 Manure management - Sheep	NH3	kt	1.73	1.76	1.76	1.52	1.28	1.19	1.11	1.10	1.12	1.17	1.16	1.15	1.13
3B3 Manure management - Swine	NH3	kt	5.86	7.28	7.92	7.66	7.11	6.93	6.78	7.06	7.12	7.03	6.88	6.98	6.89
3B4d Manure management - Goats	NH3	kt	0.04	0.04	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.04
3B4e Manure management - Horses	NH3	kt	0.53	0.58	0.60	0.68	0.76	0.82	0.84	0.91	0.91	0.95	0.87	0.81	0.80
3B4f Manure management - Mules and asses	NH3	kt	0.04	0.04	0.03	0.03	0.04	0.05	0.05	0.04	0.05	0.05	0.04	0.04	0.05
3B4gi Manure management - Laying hens	NH3	kt	0.52	0.37	0.35	0.40	0.37	0.37	0.44	0.44	0.42	0.53	0.58	0.61	0.67
3B4gii Manure management - Broilers	NH3	kt	0.40	0.55	0.62	0.64	0.47	0.47	0.58	0.58	0.56	0.56	0.52	0.52	0.52
3B4giii Manure management - Turkeys	NH3	kt	0.52	0.55	0.45	0.44	0.46	0.46	0.30	0.30	0.37	0.42	0.39	0.39	0.39
3B4giv Manure management - Other poultry	NH3	kt	0.08	0.08	0.08	0.12	0.11	0.09	0.08	0.06	0.06	0.06	0.06	0.06	0.06
3B4h Manure management - Other animals (please specify in IIR)	NH3	kt	0.57	0.33	0.36	0.35	0.35	0.34	0.43	0.41	0.38	0.40	0.40	0.41	0.40
3Da1 Inorganic N-fertilizers (includes also urea application)	NH3	kt	12.93	12.46	12.74	9.29	8.07	9.42	11.43	12.86	9.60	7.41	8.07	8.53	9.16
3Da2a Animal manure applied to soils	NH3	kt	27.81	28.95	28.96	29.27	28.44	28.53	28.23	26.54	26.46	27.60	27.84	27.98	28.65
3Da2b Sewage sludge applied to soils	NH3	kt	0.04	0.04	0.18	0.73	0.73	0.78	0.80	1.00	0.70	0.83	0.63	0.52	0.57
3Da3 Urine and dung deposited by grazing animals	NH3	kt	12.86	13.39	13.59	13.21	12.89	12.87	12.71	12.49	12.34	12.75	12.91	12.98	13.00
Total	NH3	kt	105.22	109.40	110.74	108.27	104.12	105.54	106.46	105.22	101.14	102.71	103.91	104.64	106.88
Submission 2018															
3B1a Manure management - Dairy cattle	NH3	kt	12.70	11.95	11.45	10.38	10.62	10.57	10.45	10.50	10.90	10.97	11.20	11.68	12.71
3B1b Manure management - Non-dairy cattle	NH3	kt	27.41	29.46	29.80	31.61	30.33	30.57	30.15	28.74	28.15	30.04	30.49	29.91	30.08
3B2 Manure management - Sheep	NH3	kt	1.72	1.75	1.75	1.51	1.28	1.18	1.11	1.09	1.11	1.16	1.16	1.14	1.13
3B3 Manure management - Swine	NH3	kt	3.85	4.79	5.20	5.03	4.68	4.57	4.47	4.65	4.70	4.64	4.53	4.60	4.54
3B4d Manure management - Goats	NH3	kt	0.03	0.03	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4e Manure management - Horses	NH3	kt	0.53	0.58	0.60	0.68	0.76	0.82	0.84	0.91	0.91	0.95	0.87	0.81	0.80
3B4f Manure management - Mules and asses	NH3	kt	0.05	0.04	0.03	0.03	0.04	0.05	0.05	0.04	0.05	0.06	0.05	0.05	0.05
3B4gi Manure management - Laying hens	NH3	kt	0.64	0.45	0.43	0.49	0.46	0.46	0.54	0.54	0.54	0.66	0.72	0.74	0.83
3B4gii Manure management - Broilers	NH3	kt	0.97	1.35	1.51	1.55	1.18	1.18	1.44	1.44	1.40	1.40	1.31	1.47	1.48
3B4giii Manure management - Turkeys	NH3	kt	1.00	1.07	0.87	0.84	0.88	0.88	0.58	0.58	0.71	0.81	0.74	0.79	0.81
3B4giv Manure management - Other poultry	NH3	kt	0.11	0.11	0.11	0.16	0.14	0.12	0.10	0.08	0.08	0.08	0.08	0.08	0.08
3B4h Manure management - Other animals (please specify in IIR)	NH3	kt	0.60	0.35	0.38	0.37	0.37	0.36	0.46	0.43	0.39	0.42	0.42	0.42	0.42
3Da1 Inorganic N-fertilizers (includes also urea application)	NH3	kt	13.69	13.01	13.17	9.76	8.63	9.75	11.58	13.16	9.96	7.84	8.69	9.03	9.65
3Da2a Animal manure applied to soils	NH3	kt	32.24	33.42	33.33	33.70	32.63	32.72	32.39	30.34	30.34	31.76	32.11	32.20	33.15
3Da2b Sewage sludge applied to soils	NH3	kt	0.03	0.03	0.12	0.47	0.48	0.50	0.52	0.65	0.46	0.54	0.41	0.34	0.37
3Da3 Urine and dung deposited by grazing animals	NH3	kt	13.92	14.34	14.44	13.96	13.58	13.56	13.41	13.17	12.99	13.43	13.63	13.69	13.77

Total	NH3	kt	109.48	112.72	113.19	110.57	106.06	107.30	108.12	106.36	102.70	104.78	106.41	106.96	109.89
% Change in Emissions															
3B1a Manure management - Dairy cattle	NH3	%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%	-8.5%
3B1b Manure management - Non-dairy cattle	NH3	%	-0.1%	-1.5%	-2.5%	-3.0%	-3.4%	-3.4%	-3.4%	-4.0%	-3.3%	-2.8%	-2.5%	-3.1%	-1.9%
3B2 Manure management - Sheep	NH3	%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%	-0.5%
3B3 Manure management - Swine	NH3	%	-34.4%	-34.2%	-34.3%	-34.3%	-34.2%	-34.1%	-34.0%	-34.1%	-34.1%	-34.0%	-34.1%	-34.1%	-34.0%
3B4d Manure management - Goats	NH3	%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-15.9%	-40.7%	-41.8%
3B4e Manure management - Horses	NH3	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	NH3	%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%	10.0%
3B4gi Manure management - Laying hens	NH3	%	23.8%	23.9%	23.5%	23.3%	23.3%	23.3%	23.3%	23.3%	28.4%	23.3%	23.3%	22.0%	23.3%
3B4gii Manure management - Broilers	NH3	%	144.6%	144.2%	143.1%	143.9%	150.0%	150.0%	150.0%	150.0%	150.0%	150.0%	150.0%	181.7%	183.9%
3B4giii Manure management - Turkeys	NH3	%	92.9%	92.9%	92.9%	92.9%	92.9%	92.9%	92.9%	92.9%	92.9%	92.9%	92.9%	103.9%	111.1%
3B4giv Manure management - Other poultry	NH3	%	33.3%	33.3%	33.3%	34.2%	34.2%	34.0%	33.7%	33.3%	33.3%	33.3%	33.7%	33.7%	33.7%
3B4h Manure management - Other animals	NH3	%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	3.1%	3.1%	3.0%	3.1%	3.0%
3Da1 Inorganic N-fertilizers (includes urea application)	NH3	%	5.9%	4.4%	3.3%	5.1%	6.9%	3.5%	1.3%	2.3%	3.7%	5.9%	7.7%	5.9%	5.3%
3Da2a Animal manure applied to soils	NH3	%	15.9%	15.4%	15.1%	15.1%	14.7%	14.7%	14.7%	14.3%	14.6%	15.1%	15.3%	15.1%	15.7%
3Da2b Sewage sludge applied to soils	NH3	%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%	-35.0%
3Da3 Urine and dung deposited by grazing animals	NH3	%	8.3%	7.1%	6.3%	5.6%	5.4%	5.3%	5.5%	5.5%	5.3%	5.3%	5.6%	5.5%	5.9%
Total	NH3	%	4.0%	3.0%	2.2%	2.1%	1.9%	1.7%	1.6%	1.1%	1.5%	2.0%	2.4%	2.2%	2.8%

Table 5.3. NMVOC Recalculations for Agriculture 1990–2015

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017															
3B1a Manure management - Dairy cattle	NMVOC	kt	9.16	8.39	7.82	6.96	7.16	7.13	7.08	7.11	7.37	7.43	7.64	7.99	8.74
3B1b Manure management - Non-dairy cattle	NMVOC	kt	22.32	24.54	25.11	26.38	26.02	25.81	25.86	24.77	24.20	25.32	26.13	25.60	25.37
3B2 Manure management - Sheep	NMVOC	kt	0.56	0.59	0.56	0.45	0.39	0.36	0.33	0.31	0.31	0.34	0.34	0.35	0.34
3B3 Manure management - Swine	NMVOC	kt	1.02	2.47	2.75	2.67	2.46	2.37	2.30	2.40	2.46	2.43	2.40	2.43	2.39
3B4d Manure management - Goats	NMVOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	NMVOC	kt	0.11	0.12	0.12	0.14	0.15	0.17	0.17	0.18	0.18	0.19	0.18	0.16	0.16
3B4f Manure management - Mules and asses	NMVOC	kt	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gi Manure management - Laying hens	NMVOC	kt	0.31	0.23	0.26	0.32	0.30	0.30	0.35	0.35	0.34	0.43	0.47	0.47	0.47
3B4gii Manure management - Broilers	NMVOC	kt	0.87	1.20	1.34	1.38	1.05	1.05	1.29	1.29	1.24	1.24	1.16	1.16	1.16
3B4giii Manure management - Turkeys	NMVOC	kt	0.74	0.79	0.65	0.62	0.65	0.65	0.43	0.43	0.53	0.60	0.55	0.55	0.55
3B4giv Manure management - Other poultry	NMVOC	kt	0.18	0.18	0.18	0.26	0.24	0.21	0.17	0.14	0.14	0.14	0.13	0.13	0.13
3B4h Manure management - Other animals	NMVOC	kt	0.41	0.25	0.29	0.29	0.29	0.29	0.37	0.36	0.36	0.38	0.38	0.38	0.38
3Da1 Inorganic N-fertilizers (includes urea application)	NMVOC	kt	3.36	3.45	3.39	3.29	3.29	3.55	3.53	3.55	3.53	3.48	3.44	3.42	3.38
3De Cultivated crops	NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Total	NMVOC	kt	39.04	42.21	42.47	42.79	42.01	41.88	41.89	40.90	40.69	42.00	42.83	42.65	43.08
Submission 2018															
3B1a Manure management - Dairy cattle	NMVOC	kt	9.23	8.45	7.87	7.01	7.22	7.18	7.14	7.16	7.43	7.48	7.69	8.05	8.80
3B1b Manure management - Non-dairy cattle	NMVOC	kt	22.56	24.81	25.39	26.67	26.31	26.09	26.15	25.04	24.46	25.59	26.41	25.87	25.64
3B2 Manure management - Sheep	NMVOC	kt	0.59	0.62	0.59	0.48	0.42	0.38	0.35	0.32	0.33	0.36	0.36	0.37	0.36
3B3 Manure management - Swine	NMVOC	kt	1.02	2.50	2.74	2.66	2.46	2.39	2.33	2.43	2.47	2.44	2.39	2.43	2.39
3B4d Manure management - Goats	NMVOC	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	NMVOC	kt	0.11	0.13	0.13	0.15	0.16	0.18	0.18	0.20	0.20	0.20	0.19	0.18	0.17
3B4f Manure management - Mules and asses	NMVOC	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gi Manure management - Laying hens	NMVOC	kt	0.31	0.23	0.26	0.32	0.30	0.30	0.35	0.35	0.35	0.43	0.47	0.48	0.54
3B4gii Manure management - Broilers	NMVOC	kt	0.87	1.20	1.34	1.38	1.05	1.05	1.29	1.29	1.24	1.24	1.16	1.31	1.32
3B4giii Manure management - Turkeys	NMVOC	kt	0.74	0.79	0.65	0.62	0.65	0.65	0.43	0.43	0.53	0.60	0.55	0.58	0.60
3B4giv Manure management - Other poultry	NMVOC	kt	0.18	0.18	0.18	0.26	0.24	0.21	0.17	0.14	0.14	0.14	0.13	0.13	0.13
3B4h Manure management - Other animals	NMVOC	kt	0.41	0.25	0.29	0.29	0.29	0.29	0.37	0.36	0.36	0.38	0.38	0.38	0.38
3Da1 Inorganic N-fertilizers (includes urea application)	NMVOC	kt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3De Cultivated crops	NMVOC	kt	3.82	3.77	3.82	3.70	3.68	3.98	3.95	3.93	3.92	3.90	3.85	3.84	3.81
Total	NMVOC	kt	39.85	42.93	43.27	43.56	42.77	42.70	42.71	41.65	41.44	42.79	43.60	43.63	44.17
% Change in Emissions															
3B1a Manure management - Dairy cattle	NMVOC	%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%
3B1b Manure management - Non-dairy cattle	NMVOC	%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%	1.1%
3B2 Manure management - Sheep	NMVOC	%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%	5.7%
3B3 Manure management - Swine	NMVOC	%	-0.1%	1.0%	-0.3%	-0.4%	0.0%	1.0%	1.3%	1.2%	0.3%	0.4%	-0.2%	-0.1%	0.1%
3B4d Manure management - Goats	NMVOC	%	15.4%	15.4%	15.4%	15.4%	15.4%	15.4%	15.4%	15.4%	15.4%	15.4%	15.4%	-18.7%	-20.1%
3B4e Manure management - Horses	NMVOC	%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%
3B4f Manure management - Mules and asses	NMVOC	%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%	6.2%
3B4gi Manure management - Laying hens	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	4.1%	0.0%	0.0%	3.2%	15.6%
3B4gii Manure management - Broilers	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	12.7%	13.5%
3B4giii Manure management - Turkeys	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	5.7%	9.4%
3B4giv Manure management - Other poultry	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	NMVOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	NMVOC	%	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total	NMVOC	%	2.1%	1.7%	1.9%	1.8%	1.8%	2.0%	2.0%	1.8%	1.8%	1.9%	1.8%	2.3%	2.5%

Table 5.4. TSP Recalculations for Agriculture 1990–2015

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017															
3B1a Manure management - Dairy cattle	TSP	kt	2.31	2.28	2.14	1.90	1.94	1.95	1.95	1.94	2.01	2.07	2.11	2.23	2.39
3B1b Manure management - Non-dairy cattle	TSP	kt	2.71	2.72	2.77	2.78	2.70	2.69	2.69	2.57	2.45	2.53	2.62	2.58	2.53
3B2 Manure management - Sheep	TSP	kt	1.11	1.16	1.11	0.89	0.79	0.71	0.66	0.60	0.62	0.67	0.68	0.70	0.68
3B3 Manure management - Swine	TSP	kt	0.82	1.03	1.13	1.10	1.01	0.98	0.96	1.00	1.02	1.00	0.98	1.00	0.98
3B4d Manure management - Goats	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	TSP	kt	0.03	0.03	0.03	0.04	0.04	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.04
3B4f Manure management - Mules and asses	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	TSP	kt	0.22	0.16	0.19	0.23	0.22	0.22	0.26	0.26	0.25	0.31	0.34	0.34	0.34
3B4gii Manure management - Broilers	TSP	kt	0.55	0.77	0.86	0.88	0.67	0.67	0.82	0.82	0.79	0.79	0.74	0.74	0.74
3B4giii Manure management - Turkeys	TSP	kt	0.78	0.84	0.69	0.66	0.69	0.69	0.45	0.45	0.56	0.64	0.58	0.58	0.58
3B4giv Manure management - Other poultry	TSP	kt	0.05	0.05	0.05	0.08	0.07	0.06	0.05	0.04	0.04	0.04	0.04	0.04	0.04
3B4h Manure management - Other animals	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	TSP	kt	6.10	6.27	6.14	5.98	5.96	6.44	6.40	6.45	6.41	6.32	6.25	6.20	6.12
3Dc Farm-level operations including storage, handling and transport of agricultural products	TSP	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
3Dd Off-farm storage, handling and transport of bulk agricultural products	TSP	kt	0.20	0.18	0.22	0.19	0.20	0.25	0.21	0.20	0.25	0.21	0.24	0.26	0.26
Submission 2018															
3B1a Manure management - Dairy cattle	TSP	kt	1.82	1.81	1.69	1.51	1.54	1.55	1.55	1.54	1.60	1.64	1.67	1.77	1.90
3B1b Manure management - Non-dairy cattle	TSP	kt	2.53	2.54	2.59	2.59	2.52	2.51	2.51	2.40	2.29	2.37	2.44	2.41	2.37
3B2 Manure management - Sheep	TSP	kt	1.12	1.17	1.11	0.90	0.79	0.71	0.66	0.61	0.62	0.68	0.69	0.70	0.68
3B3 Manure management - Swine	TSP	kt	0.98	1.24	1.36	1.32	1.22	1.20	1.17	1.22	1.24	1.23	1.20	1.22	1.20
3B4d Manure management - Goats	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	TSP	kt	0.03	0.03	0.03	0.04	0.04	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.04
3B4f Manure management - Mules and asses	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	TSP	kt	0.35	0.26	0.30	0.37	0.34	0.34	0.41	0.41	0.41	0.49	0.54	0.55	0.62
3B4gii Manure management - Broilers	TSP	kt	0.32	0.44	0.50	0.51	0.39	0.39	0.48	0.48	0.46	0.46	0.43	0.49	0.49
3B4giii Manure management - Turkeys	TSP	kt	0.17	0.18	0.15	0.14	0.15	0.15	0.10	0.10	0.12	0.13	0.12	0.13	0.14
3B4giv Manure management - Other poultry	TSP	kt	0.05	0.05	0.05	0.08	0.07	0.06	0.05	0.04	0.04	0.04	0.04	0.04	0.04
3B4h Manure management - Other animals	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	TSP	kt	6.93	6.85	6.93	6.71	6.67	7.22	7.17	7.13	7.11	7.07	6.99	6.97	6.91
3Dc Farm-level operations including storage, handling and transport of agricultural products	TSP	kt	0.18	0.19	0.17	0.15	0.13	0.12	0.12	0.14	0.12	0.12	0.15	0.14	0.14
3Dd Off-farm storage, handling and transport of bulk agricultural products	TSP	kt	0.20	0.18	0.22	0.19	0.20	0.25	0.21	0.20	0.25	0.21	0.24	0.26	0.26
% Change in Emissions															
3B1a Manure management - Dairy cattle	TSP	%	-21.4%	-20.7%	-20.7%	-20.6%	-20.7%	-20.7%	-20.7%	-20.5%	-20.5%	-20.5%	-20.5%	-20.4%	-20.4%
3B1b Manure management - Non-dairy cattle	TSP	%	-6.7%	-6.6%	-6.6%	-6.7%	-6.6%	-6.6%	-6.6%	-6.6%	-6.5%	-6.5%	-6.6%	-6.6%	-6.5%
3B2 Manure management - Sheep	TSP	%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%
3B3 Manure management - Swine	TSP	%	19.8%	20.9%	20.2%	20.5%	20.8%	22.2%	22.3%	22.1%	22.1%	23.0%	21.9%	22.0%	22.5%
3B4d Manure management - Goats	TSP	%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	0.7%	-29.0%	-30.3%
3B4e Manure management - Horses	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	TSP	%	59.7%	59.7%	59.7%	59.7%	59.7%	59.7%	59.7%	59.7%	66.3%	59.7%	59.7%	64.7%	84.5%
3B4gii Manure management - Broilers	TSP	%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-42.0%	-34.7%	-34.2%
3B4giii Manure management - Turkeys	TSP	%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-77.6%	-76.9%
3B4giv Manure management - Other poultry	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3Da1 Inorganic N-fertilizers (includes urea application)	TSP	%	13.6%	9.3%	12.9%	12.3%	11.9%	12.1%	12.0%	10.6%	10.9%	11.9%	11.8%	12.4%	12.8%
3Dd Off-farm storage, handling and transport of bulk agricultural products	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Table 5.5. PM₁₀ Recalculations for Agriculture 1990–2015

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017															
3B1a Manure management - Dairy cattle	PM10	kt	1.06	1.05	0.98	0.87	0.89	0.89	0.89	0.89	0.92	0.95	0.97	1.02	1.10
3B1b Manure management - Non-dairy cattle	PM10	kt	1.25	1.25	1.27	1.27	1.24	1.24	1.24	1.18	1.12	1.16	1.20	1.19	1.16
3B2 Manure management - Sheep	PM10	kt	0.45	0.47	0.44	0.36	0.31	0.28	0.26	0.24	0.25	0.27	0.27	0.28	0.27
3B3 Manure management - Swine	PM10	kt	0.37	0.46	0.50	0.49	0.45	0.44	0.43	0.44	0.45	0.44	0.44	0.44	0.44
3B4d Manure management - Goats	PM10	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM10	kt	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4f Manure management - Mules and asses	PM10	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM10	kt	0.22	0.16	0.19	0.23	0.22	0.22	0.26	0.26	0.25	0.31	0.34	0.34	0.34
3B4gii Manure management - Broilers	PM10	kt	0.55	0.77	0.86	0.88	0.67	0.67	0.82	0.82	0.79	0.79	0.74	0.74	0.74
3B4giii Manure management - Turkeys	PM10	kt	0.78	0.84	0.69	0.66	0.69	0.69	0.45	0.45	0.56	0.64	0.58	0.58	0.58
3B4giv Manure management - Other poultry	PM10	kt	0.05	0.05	0.05	0.08	0.07	0.06	0.05	0.04	0.04	0.04	0.04	0.04	0.04
3B4h Manure management - Other animals	PM10	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM10	kt	6.10	6.27	6.14	5.98	5.96	6.44	6.40	6.45	6.41	6.32	6.25	6.20	6.12
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM10	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM10	kt	0.05	0.04	0.05	0.05	0.05	0.06	0.05	0.05	0.06	0.05	0.06	0.06	0.07
Submission 2018															
3B1a Manure management - Dairy cattle	PM10	kt	0.83	0.82	0.77	0.69	0.70	0.70	0.70	0.70	0.73	0.75	0.76	0.81	0.87
3B1b Manure management - Non-dairy cattle	PM10	kt	1.14	1.14	1.16	1.17	1.13	1.13	1.13	1.08	1.03	1.07	1.10	1.09	1.07
3B2 Manure management - Sheep	PM10	kt	0.48	0.50	0.48	0.39	0.34	0.31	0.28	0.26	0.27	0.29	0.30	0.30	0.29
3B3 Manure management - Swine	PM10	kt	0.15	0.18	0.20	0.19	0.18	0.18	0.17	0.18	0.18	0.18	0.18	0.18	0.18
3B4d Manure management - Goats	PM10	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM10	kt	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4f Manure management - Mules and asses	PM10	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM10	kt	0.07	0.05	0.06	0.08	0.07	0.07	0.09	0.09	0.09	0.10	0.11	0.12	0.13
3B4gii Manure management - Broilers	PM10	kt	0.16	0.22	0.25	0.26	0.19	0.19	0.24	0.24	0.23	0.23	0.22	0.24	0.24
3B4giii Manure management - Turkeys	PM10	kt	0.17	0.18	0.15	0.14	0.15	0.15	0.10	0.10	0.12	0.13	0.12	0.13	0.14
3B4giv Manure management - Other poultry	PM10	kt	0.05	0.05	0.05	0.08	0.07	0.06	0.05	0.04	0.04	0.04	0.04	0.04	0.04
3B4h Manure management - Other animals	PM10	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM10	kt	6.93	6.85	6.93	6.71	6.67	7.22	7.17	7.13	7.11	7.07	6.99	6.97	6.91
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM10	kt	0.06	0.06	0.06	0.05	0.04	0.04	0.04	0.05	0.04	0.04	0.05	0.04	0.04
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM10	kt	0.05	0.04	0.05	0.05	0.05	0.06	0.05	0.05	0.06	0.05	0.06	0.06	0.07
% Change in Emissions															
3B1a Manure management - Dairy cattle	PM10	%	-21.7%	-21.2%	-21.2%	-21.1%	-21.2%	-21.2%	-21.2%	-21.1%	-21.1%	-21.0%	-21.0%	-21.0%	-21.0%
3B1b Manure management - Non-dairy cattle	PM10	%	-8.8%	-8.7%	-8.8%	-8.5%	-8.4%	-8.4%	-8.5%	-8.6%	-8.3%	-8.0%	-8.3%	-8.4%	-8.0%
3B2 Manure management - Sheep	PM10	%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%
3B3 Manure management - Swine	PM10	%	-60.1%	-59.9%	-60.0%	-60.0%	-59.9%	-59.7%	-59.7%	-59.7%	-59.7%	-59.6%	-59.8%	-59.7%	-59.7%
3B4d Manure management - Goats	PM10	%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	7.9%	-23.9%	-25.3%
3B4e Manure management - Horses	PM10	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	PM10	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	PM10	%	-66.4%	-66.4%	-66.4%	-66.4%	-66.4%	-66.4%	-66.4%	-66.4%	-65.0%	-66.4%	-66.4%	-65.3%	-61.2%
3B4gii Manure management - Broilers	PM10	%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-71.0%	-67.3%	-67.1%
3B4giii Manure management - Turkeys	PM10	%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-78.8%	-77.6%	-76.9%
3B4giv Manure management - Other poultry	PM10	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	PM10	%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%	-1.2%
3Da1 Inorganic N-fertilizers (includes urea application)	PM10	%	13.6%	9.3%	12.9%	12.3%	11.9%	12.1%	12.0%	10.6%	10.9%	11.9%	11.8%	12.4%	12.8%
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM10	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Table 5.6. PM_{2.5} Recalculations for Agriculture 1990–2015

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017															
3B1a Manure management - Dairy cattle	PM2.5	kt	0.69	0.68	0.64	0.57	0.58	0.58	0.58	0.58	0.60	0.62	0.63	0.67	0.72
3B1b Manure management - Non-dairy cattle	PM2.5	kt	0.82	0.82	0.84	0.84	0.81	0.81	0.81	0.78	0.74	0.76	0.79	0.78	0.76
3B2 Manure management - Sheep	PM2.5	kt	0.13	0.14	0.13	0.11	0.09	0.09	0.08	0.07	0.07	0.08	0.08	0.08	0.08
3B3 Manure management - Swine	PM2.5	kt	0.07	0.09	0.10	0.09	0.09	0.08	0.08	0.08	0.09	0.08	0.08	0.08	0.08
3B4d Manure management - Goats	PM2.5	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01
3B4f Manure management - Mules and asses	PM2.5	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM2.5	kt	0.04	0.03	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.06	0.07	0.07	0.07
3B4gii Manure management - Broilers	PM2.5	kt	0.07	0.10	0.11	0.12	0.09	0.09	0.11	0.11	0.10	0.10	0.10	0.10	0.10
3B4giii Manure management - Turkeys	PM2.5	kt	0.11	0.11	0.09	0.09	0.09	0.09	0.06	0.06	0.08	0.09	0.08	0.08	0.08
3B4giv Manure management - Other poultry	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4h Manure management - Other animals	PM2.5	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM2.5	kt	0.23	0.24	0.24	0.23	0.23	0.25	0.25	0.25	0.25	0.24	0.24	0.24	0.24
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM2.5	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Submission 2018															
3B1a Manure management - Dairy cattle	PM2.5	kt	0.54	0.54	0.50	0.45	0.46	0.46	0.46	0.46	0.48	0.49	0.50	0.53	0.57
3B1b Manure management - Non-dairy cattle	PM2.5	kt	0.77	0.77	0.78	0.78	0.76	0.76	0.76	0.73	0.69	0.72	0.74	0.73	0.72
3B2 Manure management - Sheep	PM2.5	kt	0.16	0.17	0.16	0.13	0.11	0.10	0.09	0.09	0.09	0.10	0.10	0.10	0.10
3B3 Manure management - Swine	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4d Manure management - Goats	PM2.5	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Manure management - Horses	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01
3B4f Manure management - Mules and asses	PM2.5	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4gi Manure management - Laying hens	PM2.5	kt	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4gii Manure management - Broilers	PM2.5	kt	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4giii Manure management - Turkeys	PM2.5	kt	0.03	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02
3B4giv Manure management - Other poultry	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
3B4h Manure management - Other animals	PM2.5	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic N-fertilizers (includes urea application)	PM2.5	kt	0.27	0.26	0.27	0.26	0.26	0.28	0.28	0.27	0.27	0.27	0.27	0.27	0.27
3Dc Farm-level agricultural operations including storage, handling and transport of agricultural products	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.01	0.00	0.00	0.01	0.01	0.01
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM2.5	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
% Change in Emissions															
3B1a Manure management - Dairy cattle	PM2.5	%	-21.7%	-21.0%	-21.0%	-20.9%	-21.0%	-21.0%	-21.0%	-20.8%	-20.8%	-20.7%	-20.8%	-20.6%	-20.7%
3B1b Manure management - Non-dairy cattle	PM2.5	%	-6.6%	-6.5%	-6.5%	-6.5%	-6.4%	-6.4%	-6.5%	-6.5%	-6.4%	-6.3%	-6.5%	-6.5%	-6.3%
3B2 Manure management - Sheep	PM2.5	%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%
3B3 Manure management - Swine	PM2.5	%	-90.6%	-90.6%	-90.7%	-90.7%	-90.7%	-90.6%	-90.6%	-90.6%	-90.6%	-90.6%	-90.7%	-90.7%	-90.6%
3B4d Manure management - Goats	PM2.5	%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	19.8%	-15.6%	-17.1%
3B4e Manure management - Horses	PM2.5	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4f Manure management - Mules and asses	PM2.5	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4gi Manure management - Laying hens	PM2.5	%	-87.0%	-87.0%	-87.0%	-87.0%	-87.0%	-87.0%	-87.0%	-87.0%	-86.4%	-87.0%	-87.0%	-86.5%	-84.9%
3B4gii Manure management - Broilers	PM2.5	%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-77.8%	-75.0%	-74.8%
3B4giii Manure management - Turkeys	PM2.5	%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-71.4%	-69.8%	-68.7%
3B4giv Manure management - Other poultry	PM2.5	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
3B4h Manure management - Other animals	PM2.5	%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%	-4.8%
3Da1 Inorganic N-fertilizers (includes urea application)	PM2.5	%	13.6%	9.3%	12.9%	12.3%	11.9%	12.1%	12.0%	10.6%	10.9%	11.9%	11.8%	12.4%	12.8%
3Dd Off-farm storage, handling and transport of bulk agricultural products	PM2.5	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Chapter Six

Waste

6.1 Overview of the Waste (NFR 5) Sector

Emissions from the Waste sector cover a number of different source categories and pollutants. These are detailed below in Table 6.1. All sources are considered in detail in this chapter.

Table 6.1. Pollutant Emissions by Waste Source Category

NFR Source Category	Pollutants
5A Biological treatment of waste - Solid waste disposal on land	NMVOC, Hg, PCDD/F, PCB, TSP, PM ₁₀ , PM _{2.5}
5B1 Biological treatment of waste - Composting	NH ₃ , CO
5B2 Biological treatment of waste - Anaerobic digestion at biogas facilities	NO, NA, NE
5C1a Municipal waste incineration	NO
5C1bi Industrial waste incineration	NO _x , SO ₂ , NMVOC, CO, TSP, PM _{2.5} , PM ₁₀ , BC, Pb, Cd, Hg, As, Cr, Cu, Ni, PCDD/F, B[a]P, B[b]F, B[k]F, HCB, PCB
5C1bii Hazardous waste incineration	IE (5C1bi)
5C1biii Clinical waste incineration	NO _x , SO ₂ , NMVOC, CO, TSP, PM _{2.5} , PM ₁₀ , BC, Pb, Cd, Hg, As, Cr, Cu, Ni, PCDD/F, B[a]P, B[b]F, B[k]F, HCB, PCB
5C1biv Sewage sludge incineration	NO, NA, IE
5C1bv Cremation	NO _x , SO ₂ , NMVOC, CO, TSP, PM _{2.5} , PM ₁₀ , BC, Pb, Cd, Hg, As, Cr, Cu, Ni, PCDD/F, B[a]P, B[b]F, B[k]F, HCB, PCB
5C1bvi Other waste incineration	NO, NA, NE
5C2 Open burning of waste	PCDD/F, B[a]P, B[b]F, B[k]F, PCB
5D Waste-water handling	NA, NE
5E Other waste	PM _{2.5} , PM ₁₀ , TSP, PCDD/F, B[a]P, B[b]F, B[k]F, I[123-cd]P, PCB

The Waste sector contains three key categories for five pollutants. Category Industrial waste incineration (5C1bi) is a key category for two pollutants: As and Cr, accounting for 51.3 per cent and 25.2 per cent of national total emissions, respectively. This category is the largest source of As and Cr in Ireland's 2016 data inventory. The second key category in the waste sector is Other waste (5E) for two pollutants: PCDD/F and PCBs. The category accounts for 20.6 per cent of national total PCB emissions in Ireland's 2016 data inventory. The category also accounts for 10.6 per cent of national total PCDD/F emissions. The third key category is Solid waste disposal on land (5A) for Hg, this sector accounts for 6.4 per cent of national total Mercury in 2016.

6.2 Biological treatment of waste - Solid waste disposal on land (NFR 5A)

6.2.1 Main Pollutants

Landfill gas generated at solid waste disposal sites is a source of NMVOC emissions. In Ireland sector 5A has been responsible on average for 0.5 per cent of national total

emissions across the time series 1990-2016, showing a decrease of 41.7 per cent from 0.83 kt in 1990 to 0.49 kt in 2016. Emission factors for NMVOC were sourced from the Inventory Guidebook (EMEP/EEA, 2016) and are listed in Table F1 of Annex F. The activity data used is the net fugitive methane emissions from SWDS (kt) as calculated in the national greenhouse gas inventory, which are converted to landfill gas data (Gm³) using STP molar conversion factors.

Emission of TSP, PM₁₀ and PM_{2.5} are also estimated. Emission factors from the Inventory Guidebook (EMEP/EEA, 2016) are applied to annual MSW data. Resulting emission estimates are included in Table 6.2 below.

Table 6.2. Emission Time Series for NMVOC TSP, PM₁₀ and PM_{2.5} from Solid Waste Disposal on Land

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NMVOC (kt)	0.83	1.01	0.80	0.64	0.66	0.39	0.29	0.18	0.18	0.24	0.19	0.29	0.41	0.47	0.49
TSP (t)	0.87	0.88	0.98	0.87	0.95	0.97	0.92	0.82	0.68	0.62	0.51	0.36	0.29	0.30	0.37
PM ₁₀ (t)	0.41	0.42	0.46	0.41	0.45	0.46	0.44	0.39	0.32	0.30	0.24	0.17	0.13	0.14	0.18
PM _{2.5} (t)	0.06	0.06	0.07	0.06	0.07	0.07	0.07	0.06	0.05	0.04	0.04	0.03	0.02	0.02	0.03

6.2.2 Heavy Metals

The relevant emissions in the Category 5A Solid Waste Disposal on Land sector in Ireland's air pollutant inventory include emissions of Hg from the disposal of batteries, electrical equipment, fluorescent lighting tubes, and measurement and control equipment in solid waste disposal sites (landfills). There is no direct estimate of the scale of disposal of the items mentioned at landfills in Ireland. However, Netcen/CTC (2006) provides a methodology to estimate emissions in Ireland using UK emission estimates, scaling by population for batteries and by household numbers for electrical equipment, fluorescent lighting and measurement and control equipment. Emission estimates for the above sources of Hg are presented in Table 6.3.

Table 6.3. Emission Time Series for Mercury from Solid Waste Disposal on Land

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Hg (kg)	28.96	18.27	18.68	20.08	21.50	21.86	22.01	21.94	21.71	21.49	21.21	21.05	20.91	20.63	20.94

No data are available on emissions of POPs to air from municipal solid waste (MSW) disposal, although the emissions are likely to be negligible. There is potential for POP emissions to air from landfill gas (LFG) as described in the following section.

6.2.3 Landfill Gas and Persistent Organic Pollutants

There is potential for releases of POPs in landfill gas (LFG) through the transfer of POPs present in solid waste to the LFG that is generated by the waste. A proportion of this LFG escapes to air, with the remainder captured for flaring or utilisation for energy recovery. The proportion of LFG that is utilised for electricity generation is not considered in this chapter as it is accounted for in NFR Category 1A1a Public Energy and Heat Production (Chapter Three), from which POP emissions are negligible. There is currently no information in Ireland on the release of POPs from LFG and therefore emission estimates are based on the UK inventory.

Activity data on the quantity of LFG flared and LFG that escapes to the atmosphere have been obtained from Ireland's GHG emission inventory for the 1990–2016 time series.

Emission factors have been taken from the UK NAEI (2006) for PCDD/F as 0.953 µg I-TEQ/t of escaping LFG and 0.614 µg I-TEQ/t of flared LFG (Table F.1 of Annex F). The emission factor for PCBs is 0.0008 kg/t of escaping gas (UK NAEI). Emission estimates and activity data for PCDD/F and PCBs are presented in Table 6.4.

Table 6.4. Time Series of Activity Data and Emissions of Dioxins and Furans and Polychlorinated Biphenyls from Landfill Gas

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
LFG emitted (kt)	105.4	127.4	101.5	80.6	83.9	49.3	37.1	22.8	22.3	30.5	24.2	36.9	51.8	59.4	61.4
LFG flared (kt)	NO	NO	3.9	28.6	29.0	40.4	46.6	52.1	49.9	44.2	45.1	39.0	25.5	15.6	13.0
PCDD/F (g I-TEQ)	0.100	0.121	0.101	0.112	0.116	0.097	0.093	0.086	0.083	0.083	0.078	0.083	0.081	0.076	0.075
PCBs (kg)	0.084	0.102	0.081	0.064	0.067	0.039	0.030	0.018	0.018	0.024	0.019	0.029	0.041	0.047	0.049

6.3 Biological treatment of waste - Composting (NFR 5B1)

Composting of organic waste, such as food waste, garden and park waste has taken place in Ireland since 2001. It consists of organic waste collected at kerbside and brought to civic amenity/temporary collections sites, as well as organic material composted at households. Activity data is sourced from National Waste Database Reports published by the EPA on a regular basis. Composting is a source of emissions of NH₃ and CO and the Tier 2 emission factors in the Inventory Guidebook (EMEP/EEA, 2016) of 0.66 kg/Mg waste and 0.56 kg/Mg waste are used, respectively. Emission estimates and activity data are presented in Table 6.5.

Table 6.5. Time Series of Activity Data and Emissions of NH₃ and CO from Composting

Year	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Waste composted (kt)	22.233	34	47.3	49.58	80.27	79.88	72.79	95.85	122.9	122.4	133.6	130.7	132.5	112.5	120.5	115.8
NH ₃ (kt)	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
CO (kt)	0.01	0.02	0.03	0.03	0.04	0.04	0.04	0.05	0.07	0.07	0.07	0.07	0.07	0.06	0.07	0.06

6.4 Waste Incineration (NFR 5C)

Ireland carries out relatively little waste incineration compared with some European countries and there is currently no incineration, without energy recovery, of MSW. The Waste Incineration (5C) category includes estimates of air pollutant emissions from the incineration of clinical waste (5C1biii), industrial waste (5C1bi), including hazardous waste (5C1bii, emissions from crematoria (5C1bv), and the open burning/combustion of waste materials such as farm plastics (5C2).

There is incineration of municipal wastes in Ireland following the commissioning of one incinerator in 2011 and a further in 2017. However, these are waste-to-energy facilities, and as such emissions are reported under electricity generation (1A1a).

Approximately 50 per cent of health-care waste was incinerated during the 1990s, with a total of 150 incinerators in operation. By 1999, only two of these remained in operation and both closed the following year.

Most of the industrial installations that incinerate hazardous industrial wastes are in the pharmaceutical sector.

The practice of cremation is also less common in Ireland than in other countries but has increased in recent years due to the decrease in available burial plots, particularly in larger cities and towns in Ireland.

All the above are sources of heavy metals, POPs and combustion pollutants (NO_x , SO_2 , NMVOC, CO, TSP, PM_{10} , $\text{PM}_{2.5}$ and BC) in Ireland and are discussed in detail in the following sections.

6.4.1 Clinical Waste (5C1biii)

The incineration of Clinical Waste is no longer carried out in Ireland. The bulk of hazardous clinical waste in Ireland is now treated using non-incineration technologies (namely sterilisation and shredding), with the remaining waste disposed of through landfilling, exported for incineration or used as a fuel in cement kilns. In the early 1990s, most hospitals operated on-site incinerator units where hazardous clinical waste was incinerated. A number of hospitals operated the practice of incinerating both hazardous and non-hazardous waste. Due to the implementation of stricter standards on incineration and the requirement for facilities to be licensed by the EPA, all incinerators were closed by the mid- to late-1990s. Prior to the closure of these facilities, a number of applications were made to the EPA in respect of IPPC licences. National reports and Government records contain some information on the quantity of health-care waste incinerated during the period of operation of the incinerators. From these sources, it was determined that an estimated 4,000 t of health-care waste was incinerated per annum. This value was used across the time series for the period 1990–1997, after which negligible quantities of health-care waste were incinerated up until the closure of the two remaining incinerators in 2000.

Emission estimates were derived for heavy metals using the quantity of health-care waste determined to be incinerated and Inventory Guidebook (EMEP/EEA, 2016) emission factors for As, Cd, Cr, Cu, Pb, Hg and Ni, assuming controlled air flow with no abatement. The emission factor for Zn was sourced from the UK NAEI. Emission factors are provided in Table F.1 of Annex F. Emission estimates for heavy metals are presented in Table 6.6.

Emissions of POPs from clinical wastes have been estimated using emission factors sourced from the UK NAEI. Dioxin and furan emission estimates are made utilising an emission factor of $372.1 \mu\text{g I-TEQ/t}$ health-care waste incinerated. This emission factor is used in the inventory for the period 1990–1997 until closure of all major plants. The PCB emission factor of 3.15 kg/Mt in 1990 reduces to 2.87 kg/Mt by 1995 and 2.36 kg/Mt by 1997, the last year of clinical waste incineration on the basis that environmental performance at the plants would have improved as in the UK. The emission factors for HCB from health-care waste incinerators have been estimated by taking the UK NAEI factor of 0.5 kg/Mt for 2006 and estimating the historical emission factors for 1990 and 1995 in proportion to those for PCBs in order to take account of the improvements in environmental performance that would have been introduced at some incinerators. Emission factors for intervening years are interpolated. Emission factors for 2006 are also available from the UK NAEI for benzo[a]pyrene, benzo[b]fluoranthene and benzo[k]fluoranthene, but there are no data for indeno[1,2,3-cd]pyrene. The emission factors for 2006 have been used to estimate emission factors for 1990 and 1995, scaling back in proportion to the emission factors for PCBs similar to that undertaken for HCB emission factors. Emission factors are given in Table F.1 of Annex F and the estimates for POPs are presented in Table 6.6.

Emission factors for: NO_x , CO, NMVOC, SO_x , TSP, PM_{10} and $\text{PM}_{2.5}$ were sourced from the Inventory guidebook (EMEP/EEA, 2016), using a Tier 1 approach, and are listed in Table F1 of Annex F. Resulting emission estimates 1990–1997 are included in Table 6.6.

Table 6.6. Time Series of Emissions from the Incineration of Clinical Waste

Year	1990	1991	1992	1993	1994	1995	1996	1997
NO _x (kt)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO (t)	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76
NM VOC (t)	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80
SO _x (t)	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16
TSP (kt)	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07
PM ₁₀ (kt)	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
PM _{2.5} (kt)	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
As (kg)	0.40	0.40	0.40	0.40	0.40	0.40	NO	NO
Cd (t)	0.01	0.01	0.01	0.01	0.01	0.01	NO	NO
Cr (kg)	1.60	1.60	1.60	1.60	1.60	1.60	NO	NO
Cu (kg)	2.40	2.40	2.40	2.40	2.40	2.40	NO	NO
Pb (t)	1.46	1.46	1.46	1.46	1.46	1.46	NO	NO
Hg (t)	0.22	0.22	0.22	0.22	0.22	0.22	NO	NO
Ni (kg)	1.20	1.20	1.20	1.20	1.20	1.20	NO	NO
Zn (t)	0.07	0.07	0.07	0.07	0.07	0.07	NO	NO
PCDD/F (g-I-TEQ)	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49
PCBs (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HCB (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
B[a]p (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
B[b]F (kg)	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.04
B[k]F (kg)	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.04

6.4.2 Industrial Waste (5C1bi), Hazardous Waste (5C1bii) and Sewage Sludge (5C1biv)

The category Hazardous Waste Incineration (5C1bii) is reported in Industrial Waste Incineration (5C1bi) and reported as IE under the latter category. EU Directives on waste management have set the basis for strict regulatory control on the environmental performance of hazardous industrial waste incinerators. The incineration of Industrial Waste (5C1bi) (including hazardous waste) is now highly regulated in Ireland. There are currently only a small number of facilities based in the pharmaceutical and chemical sectors that operate incinerators for the treatment of hazardous waste. The facilities that operate these units report emissions to the atmosphere to the EPA as part of IPPC licensing requirements. The disposal of CCA treated wood (CCA being a preservative containing copper, chromium and arsenic) by burning is also included as a source in this sector.

Estimates of the quantity of hazardous waste incinerated at the relevant facilities, determined from returns to the National Waste Database (Carey et al, 1996; Crowe et al, 2000; Meaney et al, 2003; Collins et al, 2004a; Collins et al, 2004b; Collins et al, 2005; Le Bolloch et al, 2006; Le Bolloch et al, 2007; Le Bolloch et al, 2009; McCoole et al, 2009; McCoole et al, 2011; McCoole et al, 2012; McCoole et al, 2013, <http://www.epa.ie/nationalwastestatistics/>), and information supplied by the facilities involved allows for the calculation of heavy metal emission estimates. Emission factors sourced from the Inventory Guidebook (EMEP/EEA 2016) for As, Cd, Pb, Hg, Ni and Zn and the UK NAEI for Cr and Cu are used to estimate emissions. Emission estimates are presented in Table 6.7, while the emission factors used are presented in Table F.1 of Annex F.

The hazardous waste incinerators currently in use in Ireland are relatively modern units designed to optimise the burning process, with wet scrubber abatement systems in place to reduce the emissions of POPs to air. Further to the use of incinerators, there are also a number of facilities that use thermal oxidisers, which are subject to emission limit values.

Annual Environmental Reports and IPPC Licence Applications provide adequate information in relation to the monitoring of PCDD/F emissions to air with limited information on the other relevant POPs. With respect to emissions of PAHs, emission factors sourced from the UK NAEI were applied to the tonnage of waste incinerated for each year. Pollutant-specific emission factors are presented in Table F.1 of Annex F. Emission estimates for POPs are presented in Table 6.7.

Emission factors for: NO_x, CO, NMVOC, SO_x, TSP, PM₁₀ and PM_{2.5} were sourced from the Inventory Guidebook (EMEP/EEA, 2016), using a Tier 1 approach, and are listed in Table F.1 of Annex F. Resulting emission estimates are included in Table 6.7 below.

Table 6.7. Time Series of Emissions from the Incineration of Industrial (incl. Hazardous & Sludge) Waste

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NO _x (kt)	0.02	0.02	0.02	0.03	0.03	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01
CO (t)	1.90	1.90	1.40	2.54	2.45	1.96	1.46	1.50	1.28	0.88	1.06	1.01	0.92	0.93	0.52
NMVOC (kt)	0.20	0.20	0.15	0.27	0.26	0.21	0.15	0.16	0.13	0.09	0.11	0.11	0.10	0.10	0.06
SO _x (t)	1.27	1.27	0.94	1.70	1.65	1.31	0.98	1.00	0.86	0.59	0.71	0.68	0.62	0.62	0.35
TSP (t)	0.27	0.27	0.20	0.36	0.35	0.28	0.21	0.21	0.18	0.13	0.15	0.14	0.13	0.13	0.07
PM ₁₀ (t)	0.19	0.19	0.14	0.25	0.25	0.20	0.15	0.15	0.13	0.09	0.11	0.10	0.09	0.09	0.05
PM _{2.5} (t)	0.11	0.11	0.08	0.14	0.14	0.11	0.08	0.09	0.07	0.05	0.06	0.06	0.05	0.05	0.03
As (t)	0.55	0.56	0.58	0.62	0.63	0.64	0.65	0.66	0.65	0.65	0.65	0.65	0.64	0.64	0.65
Cd (kg)	2.71	2.71	2.00	3.62	3.50	2.80	2.09	2.14	1.82	1.26	1.51	1.44	1.31	1.33	0.75
Cr (t)	0.53	0.54	0.56	0.60	0.61	0.62	0.63	0.63	0.63	0.63	0.63	0.62	0.62	0.61	0.63
Cu (t)	0.29	0.30	0.31	0.33	0.33	0.34	0.35	0.35	0.35	0.35	0.35	0.34	0.34	0.34	0.35
Pb (t)	0.04	0.04	0.03	0.05	0.05	0.04	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.01
Hg (kg)	1.52	1.52	1.12	2.03	1.96	1.57	1.17	1.20	1.02	0.71	0.85	0.81	0.73	0.74	0.42
Ni (kg)	3.79	0.38	0.28	0.51	0.49	0.39	0.29	0.30	0.26	0.18	0.21	0.20	0.18	0.19	0.10
Zn (t)	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
PCDD/F (g-I-TEQ)	0.03	0.03	0.08	0.02	0.01	0.60	1.11	0.42	0.01	0.01	0.01	0.01	0.00	0.01	0.01
PCBs (kg)	0.09	0.08	0.03	0.03	0.03	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HCB (kg)	0.05	0.04	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00
B[a]p (kg)	0.02	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B[b]F (kg)	0.07	0.05	0.02	0.03	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00
B[k]F (kg)	0.07	0.05	0.02	0.03	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00

6.4.3 Crematoria (5C1bv)

The practice of cremation is less popular in Ireland than in other countries. However, due to the decrease in the number of burial plots available, particularly in larger cities and towns, the number of cremations in Ireland has been steadily increasing. There are currently five crematoria operating in Ireland. Cremation has been in operation in Ireland for over a decade, with one of the crematoria open since the early 1990s. A pet crematorium is also currently operating in Ireland; however, emissions from this source are regarded as negligible. Data on the number of cremations in Ireland have been obtained via correspondence with crematoria operators or, in some cases, assumed capacity of the facility based on the equipment present and market share where no information on the number of cremations conducted was provided.

Heavy metals emissions are estimated using Inventory Guidebook (EMEP/EEA, 2016) emission factors for As, Cd, Pb, Cr, Hg, Ni, Cu Se and Zn and are presented in Table F.1 of Annex F. Emission estimates for each of the metals outlined are presented in Table 6.8.

Emissions of POPs from crematoria include PCDD/F, HCB and benzo[a]pyrene. Inventory Guidebook (EMEP/EEA, 2016) emission factors are used to derive emission estimates for the years 1990–2016 (Table F.1 of Annex F). Emission estimates are presented in Table 6.8.

Emission factors for: NO_x, CO, NMVOC, SO_x, TSP, PM₁₀ and PM_{2.5} were sourced from the Inventory guidebook (EMEP/EEA, 2016), using a Tier 1 approach, and are listed in Table F.1 of Annex F. Resulting emission estimates are included in Table 6.8.

Table 6.8. Time Series of Emissions from Crematoria

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NO _x (t)	1.24	1.24	1.82	2.03	2.06	3.14	3.14	3.14	2.54	2.75	3.07	3.37	3.73	4.02	4.17
CO (t)	0.21	0.21	0.31	0.34	0.35	0.53	0.53	0.53	0.43	0.47	0.52	0.57	0.63	0.68	0.71
NMVOC (t)	0.02	0.02	0.03	0.03	0.03	0.05	0.05	0.05	0.04	0.04	0.05	0.05	0.06	0.06	0.07
SO _x (t)	0.17	0.17	0.25	0.28	0.28	0.43	0.43	0.43	0.35	0.38	0.42	0.46	0.51	0.55	0.57
TSP (t)	0.06	0.06	0.08	0.09	0.10	0.15	0.15	0.15	0.12	0.13	0.14	0.16	0.17	0.19	0.19
PM ₁₀ (t)	0.04	0.04	0.06	0.07	0.07	0.10	0.10	0.10	0.08	0.09	0.10	0.11	0.12	0.13	0.14
PM _{2.5} (t)	0.02	0.02	0.03	0.04	0.04	0.06	0.06	0.06	0.05	0.05	0.06	0.06	0.07	0.08	0.08
As (g)	20.42	20.42	29.94	33.45	34.03	51.72	51.72	51.72	41.96	45.38	50.62	55.54	61.48	66.25	68.75
Cd (g)	7.55	7.55	11.07	12.36	12.58	19.11	19.11	19.11	15.51	16.77	18.71	20.53	22.72	24.48	25.41
Cr (g)	20.34	20.34	29.83	33.32	33.90	51.53	51.53	51.53	41.81	45.21	50.43	55.34	61.25	66.00	68.50
Cu (g)	18.65	18.65	27.35	30.55	31.08	47.23	47.23	47.23	38.32	41.44	46.23	50.73	56.15	60.50	62.79
Pb (g)	45.05	45.05	66.07	73.80	75.08	114.11	114.11	114.11	92.58	100.12	111.68	122.55	135.65	146.17	151.70
Hg (kg)	2.24	2.24	3.28	3.66	3.73	5.66	5.66	5.66	4.59	4.97	5.54	6.08	6.73	7.25	7.53
Ni (kg)	0.03	0.03	0.04	0.04	0.04	0.07	0.07	0.07	0.05	0.06	0.06	0.07	0.08	0.08	0.09
PCDD/F (g-I-TEQ)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
HCBs (g)	0.23	0.23	0.33	0.37	0.38	0.57	0.57	0.57	0.46	0.50	0.56	0.61	0.68	0.73	0.76
B[a]p (g)	0.02	0.02	0.03	0.03	0.03	0.05	0.05	0.05	0.04	0.04	0.05	0.05	0.06	0.06	0.07

6.4.4 Open Burning of Agricultural Wastes (5C2)

Open Burning of Agricultural Wastes, which includes the burning of crop residues, animal carcasses and poultry litter is a practice generally not undertaken in Ireland. Therefore, emission estimates from these sources are reported as “NO” (not occurring) for Heavy Metals.

Emissions from the open burning of farm plastics are the only source of emissions from agricultural wastes for which estimates are made. Information on the quantity of waste farm plastics that are burned in open fires is difficult to obtain. One of the largest sources of waste farm plastic, is waste plastic silage wrap and to a lesser extent synthetic fertiliser bags. The increased replacement of conventional silage with plastic wrapped silage bales, which use substantially more plastic, has seen an increase in the quantity of this waste stream. A number of different sources of information were utilised in the derivation of emission estimates. Information on the quantities of silage plastic on the market was obtained from the Irish Farm Film Producers Group (IFFPG), and national agricultural statistics were provided by the CSO and the National Farm Survey. Using the area of land utilised for silage for each year of the time series, an estimate of the plastic used for conventional (pit) silage and baled silage is made. Account is taken of plastic recovery under the silage plastics collection service operated by both the IFFPG and the Farm Relief Services. The plastic collected is recycled and used to make products such as park benches, plastic bags, garden furniture and plastic piping.

Dioxin and furan emissions from the open burning of farm plastics are determined using estimates of the quantities of material burned and the UNEP Toolkit (2013) emission factor of 300 µg I-TEQ/t burned for the open burning of municipal wastes. The UK NAEI provides an emission factor of 510 kg/Mt burned for the estimation of PCB emissions. There is minimal data available on emission factors for PAHs; however, the emission factors from the UK NAEI for small-scale waste burning are used as a best estimate. Emission factors of 89.5 kg/Mt for benzo[a]pyrene, 405 kg/Mt for benzo[b]fluoranthene and 405 kg/Mt for benzo[k]fluoranthene are applied. No data is available for indeno[1,2,3-cd]pyrene. Emission factors are compiled in Table F.2 of Annex F and the emission estimates are presented in Table 6.9.

Emissions of NO_x, SO₂, NMVOC, CO and particles from this small source are not estimated, and are therefore reported as “NE”.

Table 6.9. Time Series of Emissions from the Open Burning of Farm Plastics

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCDD/F (g-I-TEQ)	0.94	1.10	0.65	0.23	0.32	0.06	1.14	1.15	1.15	0.30	0.39	0.28	0.45	0.49	0.46
PCBs (kg)	1.59	1.86	1.11	0.39	0.54	0.10	1.94	1.95	1.96	0.50	0.67	0.48	0.77	0.84	0.78
B[a]p (kg)	0.28	0.33	0.20	0.07	0.09	0.02	0.34	0.34	0.34	0.09	0.12	0.08	0.13	0.15	0.14
B[b]F (kg)	1.26	1.48	0.88	0.31	0.43	0.08	1.54	1.55	1.55	0.40	0.53	0.38	0.61	0.67	0.62
B[k]F (kg)	1.26	1.48	0.88	0.31	0.43	0.08	1.54	1.55	1.55	0.40	0.53	0.38	0.61	0.67	0.62

6.5 Wastewater handling (NFR 5D)

Emissions of NMVOC from wastewater treatment plants may in some cases be significant in urban areas and may contribute at a national level. In Ireland's case, based on the use of the emission factor of 15mg/m³ of wastewater handled from EMEP/EEA 2016 guidebook and wastewater quantities handled in Ireland the emissions were estimated to be 0.006% of national NMVOC emissions. Therefore, Ireland considers emissions from this source as negligible and NE is reported for this source.

6.6 Other Waste (NFR 5E)

This NFR category includes emissions from accidental vehicle and building fires and other burning, which constitutes bonfires, domestic burning of MSW and burning of construction wastes. These are all sources of POPs, TSP, PM₁₀ and PM_{2.5}. Each of these combustion sources is described in the following sections.

6.6.1 Accidental Fires

Accidental fires are poorly controlled combustion events that can release large quantities of POPs and particulates into the environment. These include accidental fires of houses, other buildings and cars. A variety of materials can be burned in accidental fires, which can lead to some difficulty in obtaining detailed activity data and applying emission factors correctly. However, there are some data available in Ireland in relation to accidental building and vehicle fires from the Fire Services Department.

Vehicle fire statistics are only available since 2000, with the number of fires ranging from 1,600 to 7,700 per annum. With respect to earlier years, the URS Dames & Moore (2000) report suggests that the number of vehicle fires in 1998 was 4,130. It is assumed that, in the absence of any information, the number of vehicle fires per year in the period 1990–1997 is equal to that in 1998. Dioxin and furan emissions from vehicle fires are estimated using the UNEP Toolkit 2013 emission factor of 100 µg I-TEQ/vehicle fire. An emission factor of 25.5 mg/vehicle fire based on the NAEI emission factor of 510 kg/Mt burned for small scale

domestic waste burning (UK NAEI) for PCB emissions from the open burning of MSW is used for accidental vehicle fires, assuming that on average 50 kg of material are burnt per fire (Dyke, 1997), while those in relation to PAHs, also based on the mass of material burnt, sourced from the UK NAEI, suggest values for benzo[a]pyrene of 0.06 mg/vehicle fire, for benzo[b]fluoranthene of 0.10 mg/vehicle fire, for benzo[k]fluoranthene of 0.03 mg/vehicle fire, and for indeno[1,2,3-cd]pyrene of 0.07 mg/vehicle fire.

The Fire Services Department also provides information in relation to building fires, which is disaggregated into the type of building and the number of fires that are chimney fires. Information is only available for the years 2000–2007 at this level of disaggregation. For data prior to 2000, no differentiation was made between chimney fires and other types of building fires. The proportion of chimney fires to the total number of building fires post-2000 is therefore used to estimate the number of chimney fires annually prior to 2000. Limited information is available on the quantity of material burnt in accidental fires both in Ireland and internationally. The assumed quantity of material burnt in each building fire is 2.28 t per fire (Lorenz et al., 1996) and approximately 10 kg in each chimney fire. Dioxin and furan emissions are estimated using an emission factor of 400 µg I-TEQ/t of material burned (UNEP Toolkit, 2013). For PCB emissions, the emission factor of 510 kg/Mt burned (UK NAEI) for the open burning of MSW is applied, while, for PAH emissions (UK NAEI), the emission factors equate to 1.2 kg/Mt for benzo[a]pyrene, 1.9 kg/Mt for benzo[b]fluoranthene, 0.67 kg/Mt for benzo[k]fluoranthene and 1.3 kg/Mt for indeno[1,2,3-cd]pyrene. Accidental vehicle fires and building fires emission estimates are summed to provide an estimate of the total emissions from accidental fires. Emission estimates for the 1990–2016 time series are presented in Table 6.10. Emission factors are compiled in Table F.2 of Annex F. Ireland has included estimates of TSP, PM₁₀ and PM_{2.5} emissions in this using the emission factors from the Inventory Guidebook (EMEP/EEA, 2016).

Table 6.10. Time Series of Emissions from Accidental Fires from Vehicles and Buildings

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCDD/F (g-I-TEQ)	5.04	5.04	3.87	3.39	3.47	4.45	4.40	4.45	4.28	3.71	3.29	2.00	1.93	3.46	1.59
PCBs (kg)	6.00	6.00	4.14	3.82	3.88	5.05	4.93	5.06	5.00	4.37	3.89	2.39	2.30	4.13	1.86
B[a]p (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.004
B[b]F (kg)	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.02	0.01
B[k]F (kg)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.003	0.00	0.01	0.002
I(123-cd)P (kg)	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.005
TSP (kt)	0.36	0.36	0.37	0.34	0.35	0.43	0.43	0.43	0.43	0.38	0.34	0.21	0.20	0.36	0.17
PM10 (kt)	0.36	0.36	0.37	0.34	0.35	0.43	0.43	0.43	0.43	0.38	0.34	0.21	0.20	0.36	0.17
PM2.5 (kt)	0.36	0.36	0.37	0.34	0.35	0.43	0.43	0.43	0.43	0.38	0.34	0.21	0.20	0.36	0.17

6.6.2 Other Burning

This section includes the emission of POPs from domestic bonfires, the burning of domestic waste both indoors and outdoors and the open burning of construction waste. Domestic bonfires normally include a variety of garden wastes (e.g. wood, leaves, etc.), and their importance with respect to POP emissions is greatly increased in cases where other wastes are added to the bonfires (e.g. plastics). Some households are not covered, or opt not to be covered, by waste collection systems and may burn household waste. Combustion of treated wood that has been used for construction, fencing and furniture can be a particularly significant source of POP emissions. For example, where wood is pre-treated with chlorinated fungicides, such as lindane or pentachlorophenol, its combustion can be a potentially significant source of PCDD/F emissions to air, whilst wood pre-treatment with

creosote is a potential source for PAH emissions. However, the use of these chemicals has been significantly reduced in Ireland since the early 1990s.

For domestic bonfires, activity data are determined on a per-capita basis using the UK inventory as the reference, as no information is available in Ireland. For the burning of household waste, estimates for uncollected household waste were obtained for each of the years 2001 through to 2013, as well as for 1998 and 1995 from National Waste Reports (Carey et al., 1996; Crowe et al., 2000; Collins et al., 2004a, 2004b; Le Bolloch et al., 2006, 2007, 2009, McCooile et al, 2009; McCooile et al, 2011; McCooile et al, 2012; McCooile et al, 2013<http://www.epa.ie/nationalwastestatistics>), with annual data interpolated for other years. “Uncollected waste” refers to the waste produced by the portion of the population not provided with, or not availing of, a waste collection service, corrected to take account of local conditions. This is calculated according to a standard methodology at the local authority level, based on total numbers of households, numbers of households served with waste collection, and quantities of waste collected per household in each local authority area. In addition, a proportion of households share waste collections services. Only the fraction of household waste that is combustible is burned. Compositional statistics at a national level are applied to estimate the quantities of combustible materials burnt.

Information on construction and demolition waste is available from National Waste Reports. The proportion of wood within this waste stream is estimated using data collected but not published in the National Waste Reports for the years 2004 and 2006, based on estimates of both authorised and unauthorised construction and demolition waste disposal. These values have been used for all other years in the absence of any other information. The URS Dames & Moore study (2000) suggests that 5 per cent of construction and demolition waste wood arising is burned on construction sites, whereas the UK NAEI suggests a value of 0.1 per cent. The value of 5 per cent is applied for the years 1990–1998, linearly decreasing for the years 1999–2003, with the value of 0.1 per cent applied for the period 2004–2016, based on correspondence with representatives from the National Construction and Demolition Waste Council who indicate that they would expect virtually no uncontrolled burning in urban areas.

The UNEP toolkit for open burning of construction and demolition waste wood (60 µg I-TEQ/t burned) is applied to estimate PCDD/F emissions from bonfires on the basis that bonfires contain mainly wood and garden waste. Domestic burning of MSW contains material that varies and that often includes plastics and sometimes specific chemicals that potentially affect PCDD/F emissions. The UK NAEI suggests an emission factor of 173 µg I-TEQ/t burned. This emission factor not only takes into account the wide range of materials in household waste but also other materials such as treated and untreated wood. In relation to PCDD/F emissions from wood burning, an emission factor of 60 µg I-TEQ/t is applied.

The estimated emission factor of 1.14 kg/Mt burned for PCBs from bonfires has been taken as the average of the UK NAEI emission factors for domestic wood combustion (e.g. fireplaces) (1.99 kg/Mt burned) and open burning of crop residues (0.29 kg/Mt). For the open burning of domestic wastes, the UK NAEI emission factor of 510 kg/Mt burned has been adopted for PCBs. There are no specific data on PCB emissions from the open burning of construction wood, but emission factors from the NAEI for industrial combustion of wood indicate no difference for treated and untreated wood. Emission factors for domestic wood combustion from the NAEI and the Inventory Guidebook range from 1.99 to 6 kg/Mt burned. The emission factor of 1.99 kg/Mt has been adopted for open burning of construction waste wood in Ireland.

Emission factors with respect to PAH emission estimates are also sourced from the UK NAEI. For bonfires, the emission factors are 1,300 kg/Mt for benzo[a]pyrene, 1,500 kg/Mt for benzo[b]fluoranthene, 500 kg/Mt for benzo[k]fluoranthene and 90 kg/Mt for indeno[1,2,3-cd]pyrene. For the open burning of domestic wastes, emission factors for small-scale waste

burning are applied as follows: 89.5 kg/Mt for benzo[a]pyrene, 405 kg/Mt for both benzo[b]fluoranthene and benzo[k]fluoranthene. No data are available for indeno[1,2,3-cd]pyrene. These emission factors are also used to estimate emissions from the open burning of wood at construction sites.

The three sources of emissions described in previous paragraphs are summed to provide total emission estimates for Category 5.E Other Waste. Emission factors are compiled in Table F.2 of Annex F. Emission estimates for the 1990–2016 time series are presented in Table 6.11.

Table 6.11. Time Series of Emissions from Other Waste Burning

Year	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
PCDD/F (g-I-TEQ)	1.15	1.29	1.72	2.24	2.30	0.66	0.66	0.67	0.67	0.66	0.66	0.66	0.66	0.65	0.66
PCBs (kg)	7.73	9.46	14.75	21.21	21.77	0.73	0.65	0.69	0.69	0.69	0.69	0.69	0.69	0.69	0.69
B[a]p (t)	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
B[b]F (t)	0.02	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
B[k]F (t)	0.01	0.01	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
I(123-cd)P (kg)	0.82	0.83	0.85	0.87	0.88	0.91	0.92	0.92	0.92	0.92	0.91	0.91	0.91	0.90	0.92

6.7 Recalculations in the Waste Sector

Recalculations in the waste sector (Table 6.12) in this submission are limited to estimates from:

Other waste (5E)

Recalculations to emission estimates of dioxins and PCBs are the result change in the emission factor for Household waste burned (in the open or indoors in domestic fires) for Dioxins from 173 ug-I-TEQ/Mt burned (NAEI) to 40 ug-I-TEQ/Mt burned (UNEP, 2012). In addition the methodology for estimating emissions from Accidental vehicle and building fires was updated which resulted in a decrease in the activity data across the time series 1990–2016 and a resultant decrease in emissions.

6.8 Quality Assurance/Quality Control

Previous work has ensured that the estimates in the Waste sector are now fully consistent with other sectors in the inventory. This has allowed the detailed QA/QC procedures in the national inventory system to be implemented on data in the Waste sector in a manner consistent with other sectors.

6.9 Planned Improvements

The inventory team will continue to review emission estimates for this sector in light of any new information that may become available for future submissions. The inventory team also plans to continue to outsource contracts on a periodic basis to re-examine and extend the inventory time series with respect to heavy metals and persistent organic pollutants.

Table 6.12. Recalculations for Waste 1990–2015

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
5A Solid waste disposal on land	NM VOC	kt	0.83	1.01	0.80	0.64	0.66	0.39	0.29	0.18	0.18	0.24	0.19	0.29	0.41	0.47
5A Solid waste disposal on land	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM ₁₀	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM _{2.5}	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	Hg	t	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5A Solid waste disposal on land	PCDD/F	g-I-TEQ	0.10	0.12	0.10	0.11	0.12	0.10	0.09	0.09	0.08	0.08	0.08	0.08	0.08	0.08
5A Solid waste disposal on land	PCBs	kg	0.08	0.10	0.08	0.06	0.07	0.04	0.03	0.02	0.02	0.02	0.02	0.03	0.04	0.05
Submission 2018																
5A Solid waste disposal on land	NM VOC	kt	0.83	1.01	0.80	0.64	0.66	0.39	0.29	0.18	0.18	0.24	0.19	0.29	0.41	0.47
5A Solid waste disposal on land	TSP	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM ₁₀	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	PM _{2.5}	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5A Solid waste disposal on land	Hg	t	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5A Solid waste disposal on land	PCDD/F	g-I-TEQ	0.10	0.12	0.10	0.11	0.12	0.10	0.09	0.09	0.08	0.08	0.08	0.08	0.08	0.08
5A Solid waste disposal on land	PCBs	kg	0.08	0.10	0.08	0.06	0.07	0.04	0.03	0.02	0.02	0.02	0.02	0.03	0.04	0.05
5A Solid waste disposal on land	NM VOC	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
5A Solid waste disposal on land	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	PM ₁₀	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	PM _{2.5}	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5A Solid waste disposal on land	PCDD/F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%
5A Solid waste disposal on land	PCBs	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
NFR Category			1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
5B Biological treatment of waste - Composting	NH ₃	kt	NO	NO	NO	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
5B Biological treatment of waste - Composting	CO	kt	NO	NO	NO	0.04	0.04	0.04	0.05	0.07	0.07	0.07	0.07	0.07	0.06	0.07
Submission 2018																
5B Biological treatment of waste - Composting	NH ₃	kt	NO	NO	NO	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
5B Biological treatment of waste - Composting	CO	kt	NO	NO	NO	0.04	0.04	0.04	0.05	0.07	0.07	0.07	0.07	0.07	0.06	0.07
% Change in Emission																
5B Biological treatment of waste - Composting	NH ₃	%				0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5B Biological treatment of waste - Composting	CO	%				0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

*Prior to 1998, emissions were only reported for the sub-category 5Cbi (Clinical waste incineration). For the years 1998-2015 this sub-category is NO. Other sub-categories of 5C are reported as NO, NA, IE, and NE.

Table 6.12. Recalculations for Waste 1990–2015 (continued)

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Submission 2017																
5C Waste Incineration	NO _x	kt	0.03	0.03	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.02
5C Waste Incineration	NMVOG	kt	0.20	0.20	0.15	0.27	0.26	0.21	0.15	0.16	0.13	0.09	0.11	0.11	0.10	0.10
5C Waste Incineration	SO ₂	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	TSP	kt	0.07	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM ₁₀	kt	0.05	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM _{2.5}	kt	0.03	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Cd	t	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Pb	t	1.49	1.49	0.03	0.05	0.05	0.04	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02
5C Waste Incineration	Hg	t	0.22	0.22	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste Incineration	PCDD/F	g-I-TEQ	2.46	2.62	0.73	0.25	0.33	0.66	2.26	1.56	1.16	0.31	0.40	0.29	0.46	0.50
5C Waste Incineration	PCBs	kg	1.69	1.95	1.14	0.42	0.57	0.12	1.96	1.97	1.97	0.51	0.68	0.49	0.78	0.85
5C Waste Incineration	HCB	t	0.06	0.05	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste Incineration	B[a]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[b]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[k]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Submission 2018																
5C Waste Incineration	NO _x	kt	0.03	0.03	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.02
5C Waste Incineration	NMVOG	kt	0.20	0.20	0.15	0.27	0.26	0.21	0.15	0.16	0.13	0.09	0.11	0.11	0.10	0.10
5C Waste Incineration	SO ₂	kt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	TSP	kt	0.07	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM ₁₀	kt	0.05	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	PM _{2.5}	kt	0.03	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Cd	t	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	Pb	t	1.49	1.49	0.03	0.05	0.05	0.04	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02
5C Waste Incineration	Hg	t	0.22	0.22	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste Incineration	PCDD/F	g-I-TEQ	2.46	2.62	0.73	0.25	0.33	0.66	2.26	1.56	1.16	0.31	0.40	0.29	0.46	0.51
5C Waste Incineration	PCBs	kg	1.69	1.95	1.14	0.42	0.57	0.12	1.96	1.97	1.97	0.51	0.68	0.49	0.78	0.85
5C Waste Incineration	HCB	t	0.06	0.05	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste Incineration	B[a]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[b]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	B[k]F	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Waste Incineration	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
% Change in Emission																
5C Waste Incineration	NO _x	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	NMVOG	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	SO ₂	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	TSP	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

NFR Category	Pollutant	Unit	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
5C Waste Incineration	PM ₁₀	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	PM _{2.5}	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	Cd	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	Pb	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	Hg	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	PCDD/F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.8%
5C Waste Incineration	PCBs	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	HCB	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	B[a]P	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	B[b]F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	B[k]F	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
5C Waste Incineration	I[123-cd]P	%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Submission 2017																
5E Other Waste	TSP		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
5E Other Waste	PM ₁₀		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
5E Other Waste	PM _{2.5}		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
5E Other Waste	PCDD/F	g-I-TEQ	10.17	10.76	11.69	13.39	13.55	6.91	6.95	6.91	6.76	6.66	6.62	6.50	6.48	6.48
5E Other Waste	PCBs	kg	16.27	18.00	21.83	27.91	28.39	7.89	7.82	7.85	7.81	7.78	7.77	7.73	7.73	7.73
5E Other Waste	B[a]P	t	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	B[b]F	t	0.02	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5E Other Waste	B[k]F	t	0.01	0.01	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Submission 2018																
5E Other Waste	TSP		0.36	0.36	0.37	0.34	0.35	0.43	0.43	0.43	0.43	0.38	0.34	0.21	0.20	0.36
5E Other Waste	PM ₁₀		0.36	0.36	0.37	0.34	0.35	0.43	0.43	0.43	0.43	0.38	0.34	0.21	0.20	0.36
5E Other Waste	PM _{2.5}		0.36	0.36	0.37	0.34	0.35	0.43	0.43	0.43	0.43	0.38	0.34	0.21	0.20	0.36
5E Other Waste	PCDD/F	g-I-TEQ	6.19	6.33	5.59	5.63	5.76	5.11	5.06	5.12	4.95	4.37	3.95	2.66	2.59	4.11
5E Other Waste	PCBs	kg	13.73	15.46	18.89	25.04	25.66	5.78	5.59	5.75	5.69	5.06	4.59	3.08	2.99	4.83
5E Other Waste	B[a]P	t	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	B[b]F	t	0.02	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5E Other Waste	B[k]F	t	0.01	0.01	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5E Other Waste	I[123-cd]P	t	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
% Change in Emission																
5E Other Waste	PCDD/F	%	-39.1%	-41.2%	-52.2%	-57.9%	-57.5%	-26.1%	-27.1%	-25.9%	-26.8%	-34.4%	-40.3%	-59.0%	-60.0%	-36.6%
5E Other Waste	PCBs	%	-15.6%	-14.1%	-13.5%	-10.3%	-9.6%	-26.7%	-28.6%	-26.7%	-27.1%	-35.0%	-40.9%	-60.1%	-61.3%	-37.6%
5E Other Waste	B[a]P	%	0.0%	0.0%	0.0%	-0.3%	-0.3%	-0.6%	-0.7%	-0.8%	-0.8%	-0.2%	-2.6%	-3.2%	-0.1%	-0.9%
5E Other Waste	B[b]F	%	0.0%	0.0%	0.0%	-0.2%	-0.2%	-0.6%	-0.7%	-0.8%	-0.8%	-0.2%	-2.6%	-3.2%	-0.1%	-0.9%
5E Other Waste	B[k]F	%	0.0%	0.0%	0.0%	-0.1%	-0.1%	-0.5%	-0.7%	-0.7%	-0.8%	-0.2%	-2.4%	-3.0%	-0.1%	-0.8%
5E Other Waste	I[123-cd]P	%	-0.8%	-0.8%	-0.9%	-1.2%	-1.2%	-1.1%	-1.3%	-1.3%	-1.3%	-0.9%	-3.4%	-4.4%	-1.3%	-1.6%

*Prior to 1998, emissions were only reported for the sub-category 5Cbiii (Clinical waste incineration). For the years 1998-2014 this sub-category is NO. Other sub-categories of 5C are reported as NO, NA, IE, and NE.

Chapter Seven

Gridded and LPS data

7.1 Overview of Gridded and LPS data reporting

It is mandatory to report gridded emissions and emissions from large point sources every four years both under the Convention on Long-Range Transboundary Air Pollution and under Directive (EU) 2016/2284 on the reduction of national emissions of certain atmospheric pollutants (the revised National Emission Ceilings Directive).

The Guidelines for Reporting Emissions and Projections Data under the Convention on Long-Range Transboundary Air Pollution (ECE/EB.AIR/125), adopted in 2014, specify the scope, methodologies, formats and deadlines for annual inventory submissions by Parties to the Convention. The guidelines specify that at four yearly intervals, starting in 2017, Parties shall report updated aggregated sectoral (GNFR) gridded emissions and LPS emissions. The EMEP grid is defined in paragraph 14 of the reporting guidelines and refers to a 0.1°x0.1° latitude-longitude projection in the geographic coordinate World Geodetic System (WGS) latest revision, WGS 84. The EMEP domain covers the geographic domain between 30°N-82°N latitude and 30°W-90°E longitude.

The reporting guidelines under the Convention refers to the EMEP/EEA Guidebook for technical guidance on the spatial distribution of emissions. Directive (EU) 2016/2284 refers to the reporting guidelines under the Convention and as such the requirements under the two reportings are identical.

The development of a high-resolution model for distribution of emissions is part of an on-going research project funded by the EPA ("National mapping of GHG and non-GHG emissions sources". Ref: 2015-CCRP-MS.26). The project has developed a model for distributing emissions at a resolution of 1 km x 1 km covering all sectors and pollutants included in the official Irish emission inventory. The generated spatial emissions data (GNFR) is fully consistent with the reported emission inventories (NFR) under the LRTAP Convention. A list of GNFR categories is presented in Table 7.1 and the relationship between NFR and GNFR is presented in Table 7.2.

Table 7.1. List of GNFR categories

GNFR
A_PublicPower
B_Industry
C_OtherStationaryComb
D_Fugitive
E_Solvents
F_RoadTransport
G_Shipping
H_Aviation
I_Offroad
J_Waste
K_AgriLivestock
L_AgriOther
M_Other
'MEMO' ITEMS - NOT TO BE INCLUDED IN NATIONAL TOTALS
O_AviCruise
P_IntShipping
z_Memo
N_Natural

Table 7.2. Correspondence list for GNFR and NFR categories

NFR	NFR name	GNFR
1A1a	Public electricity and heat production	A_PublicPower
1A1b	Petroleum refining	B_Industry
1A1c	Manufacture of solid fuels and other energy industries	B_Industry
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	B_Industry
1A2b	Stationary combustion in manufacturing industries and construction: Non-ferrous metals	B_Industry
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	B_Industry
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	B_Industry
1A2e	Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	B_Industry
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	B_Industry
1A2gvii	Mobile Combustion in manufacturing industries and construction	I_Offroad
1A2gviii	Stationary combustion in manufacturing industries and construction: Other	B_Industry
1A3ai(i)	International aviation LTO (civil)	H_Aviation
1A3aii(i)	Domestic aviation LTO (civil)	H_Aviation
1A3bi	Road transport: Passenger cars	F_RoadTransport
1A3bii	Road transport: Light duty vehicles	F_RoadTransport
1A3biii	Road transport: Heavy duty vehicles and buses	F_RoadTransport
1A3biv	Road transport: Mopeds & motorcycles	F_RoadTransport
1A3bv	Road transport: Gasoline evaporation	F_RoadTransport
1A3bvi	Road transport: Automobile tyre and brake wear	F_RoadTransport
1A3bvii	Road transport: Automobile road abrasion	F_RoadTransport
1A3c	Railways	I_Offroad
1A3di(ii)	International inland waterways	G_Shipping
1A3dii	National navigation (shipping)	G_Shipping

NFR	NFR name	GNFR
1A3ei	Pipeline transport	I_Offroad
1A3eii	Other (please specify in the IIR)	I_Offroad
1A4ai	Commercial/institutional: Stationary	C_OtherStationaryComb
1A4aii	Commercial/institutional: Mobile	I_Offroad
1A4bi	Residential: Stationary	C_OtherStationaryComb
1A4bii	Residential: Household and gardening (mobile)	I_Offroad
1A4ci	Agriculture/Forestry/Fishing: Stationary	C_OtherStationaryComb
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	I_Offroad
1A4ciii	Agriculture/Forestry/Fishing: National fishing	I_Offroad
1A5a	Other stationary (including military)	C_OtherStationaryComb
1A5b	Other, Mobile (including military, land based and recreational boats)	I_Offroad
1B1a	Fugitive emission from solid fuels: Coal mining and handling	D_Fugitive
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	D_Fugitive
1B1c	Other fugitive emissions from solid fuels	D_Fugitive
1B2ai	Fugitive emissions oil: Exploration, production, transport	D_Fugitive
1B2aiv	Fugitive emissions oil: Refining / storage	D_Fugitive
1B2av	Distribution of oil products	D_Fugitive
1B2b	Fugitive emissions from natural gas (exploration, production, processing, transmission, storage, distribution and other)	D_Fugitive
1B2c	Venting and flaring (oil, gas, combined oil and gas)	D_Fugitive
1B2d	Other fugitive emissions from energy production	D_Fugitive
2A1	Cement production	B_Industry
2A2	Lime production	B_Industry
2A3	Glass production	B_Industry
2A5a	Quarrying and mining of minerals other than coal	B_Industry
2A5b	Construction and demolition	B_Industry
2A5c	Storage, handling and transport of mineral products	B_Industry
2A6	Other mineral products	B_Industry
2B1	Ammonia production	B_Industry
2B10a	Chemical industry: Other (please specify in the IIR)	B_Industry
2B10b	Storage, handling and transport of chemical products (please specify in the IIR)	B_Industry
2B2	Nitric acid production	B_Industry
2B3	Adipic acid production	B_Industry
2B5	Carbide production	B_Industry
2B6	Titanium dioxide production	B_Industry
2B7	Soda ash production	B_Industry
2C1	Iron and steel production	B_Industry
2C2	Ferroalloys production	B_Industry
2C3	Aluminium production	B_Industry
2C4	Magnesium production	B_Industry
2C5	Lead production	B_Industry
2C6	Zinc production	B_Industry
2C7a	Copper production	B_Industry
2C7b	Nickel production	B_Industry
2C7c	Other metal production	B_Industry
2C7d	Storage, handling and transport of metal products	B_Industry
2D3a	Domestic solvent use including fungicides	E_Solvents
2D3b	Road paving with asphalt	E_Solvents
2D3c	Asphalt roofing	B_Industry
2D3d	Coating applications	B_Industry
2D3e	Degreasing	E_Solvents
2D3f	Dry cleaning	E_Solvents
2D3g	Chemical products	E_Solvents
2D3h	Printing	E_Solvents
2D3i	Other solvent use	E_Solvents
2G	Other product use	E_Solvents
2H1	Pulp and paper industry	B_Industry
2H2	Food and beverages industry	B_Industry
2H3	Other industrial processes	B_Industry

NFR	NFR name	GNFR
2I	Wood processing	B_Industry
2J	Production of POPs	B_Industry
2K	Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)	B_Industry
2L	Other production, consumption, storage, transportation or handling of bulk products	B_Industry
3B1a	Manure management - Dairy cattle	K_AgriLivestock
3B1b	Manure management - Non-dairy cattle	K_AgriLivestock
3B2	Manure management - Sheep	K_AgriLivestock
3B3	Manure management - Swine	K_AgriLivestock
3B4a	Manure management - Buffalo	K_AgriLivestock
3B4d	Manure management - Goats	K_AgriLivestock
3B4e	Manure management - Horses	K_AgriLivestock
3B4f	Manure management - Mules and asses	K_AgriLivestock
3B4gi	Manure management - Laying hens	K_AgriLivestock
3B4gii	Manure management - Broilers	K_AgriLivestock
3B4giii	Manure management - Turkeys	K_AgriLivestock
3B4giv	Manure management - Other poultry	K_AgriLivestock
3B4h	Manure management - Other animals	K_AgriLivestock
3Da1	Inorganic N-fertilizers (includes also urea application)	L_AgriOther
3Da2a	Animal manure applied to soils	L_AgriOther
3Da2b	Sewage sludge applied to soils	L_AgriOther
3Da2c	Other organic fertilisers applied to soils (including compost)	L_AgriOther
3Da3	Urine and dung deposited by grazing animals	L_AgriOther
3Da4	Crop residues applied to soils	L_AgriOther
3Db	Indirect emissions from managed soils	L_AgriOther
3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products	L_AgriOther
3Dd	Off-farm storage, handling and transport of bulk agricultural products	L_AgriOther
3De	Cultivated crops	L_AgriOther
3Df	Use of pesticides	L_AgriOther
3F	Field burning of agricultural residues	L_AgriOther
3I	Agriculture other	L_AgriOther
5A	Biological treatment of waste - Solid waste disposal on land	J_Waste
5B1	Biological treatment of waste - Composting	J_Waste
5B2	Biological treatment of waste - Anaerobic digestion at biogas facilities	J_Waste
5C1a	Municipal waste incineration	J_Waste
5C1bi	Industrial waste incineration	J_Waste
5C1bii	Hazardous waste incineration	J_Waste
5C1biii	Clinical waste incineration	J_Waste
5C1biv	Sewage sludge incineration	J_Waste
5C1bv	Cremation	J_Waste
5C1bvi	Other waste incineration	J_Waste
5C2	Open burning of waste	J_Waste
5D1	Domestic wastewater handling	J_Waste
5D2	Industrial wastewater handling	J_Waste
5D3	Other wastewater handling	J_Waste
5E	Other waste	J_Waste
6A	Other (included in national total for entire territory)	M_Other
'MEMO' ITEMS - NOT TO BE INCLUDED IN NATIONAL TOTALS		
1A3ai(ii)	International aviation cruise (civil)	O_AviCruise
1A3aii(ii)	Domestic aviation cruise (civil)	O_AviCruise
1A3di(i)	International maritime navigation	P_IntShipping
1A5c	Multilateral operations	z_Memo
1A3	Transport (fuel used)	z_Memo
6B	Other not included in national total of the entire territory	z_Memo
11A	Volcanoes	N_Natural
11B	Forest fires	N_Natural
11C	Other natural emissions	N_Natural

7.2 Mapping methodology

The methodology used in the emissions mapping follow the guidelines in the 2016 EMEP/EEA emission inventory guidebook. The overall approach aims to allocate the national total emissions to the geographical location where they occur as accurately as possible. A distinction is made between point sources and area sources. Point sources are sources that can be treated individually and have an exact location, e.g. industrial plants. Area sources cover a group of minor emission sources with similar characteristics that cannot be treated individually because of the number of sources, e.g. residential plants. Some sectors are covered only by point sources or only by area sources, but many sectors cover both point and area sources. In the latter case point and area sources are treated separately in the data processing and following they are combined on sectoral level in the spatial emission mapping.

Emissions from point sources can be allocated to an exact location, e.g. the location of a power plant or an industrial plant. Activity data and/or emissions are available for a number of large plants e.g. from PRTR/E-PRTR reporting. These data are used either directly (emissions) or indirectly (activity data) to allocate point source emissions. Both locations and emissions are generally very accurate for point sources.

The individual source contribution cannot be determined for area sources, and emissions allocations are based on a number of spatial data sets. For each area source related available spatial data are evaluated and the closest related are used for emission mapping, taking into account completeness (must cover the entire national area), spatial resolution, accuracy, update frequency etc.

The 2016 EMEP/EEA Guidebook describe a tiered approach for spatial distribution of emissions, depending on the data availability and level of detail for the individual emission sources/sectors. Furthermore, different methodological tiers can be used for different pollutants from a source, e.g. point source emission data are most often available only for some pollutants, while emission mapping for remaining pollutants follow a lower tier method.

The concept of tiered mapping is summarised as follows:

- **Tier 3 methods** are based on closely related spatial emission or activity data, e.g. data for regulated processes and industries, and road traffic flows by vehicle type derived from surveys.
- **Tier 2 methods** are based on the use of surrogate statistics relate to the sector, e.g. heat demand for the residential sector, agricultural animal statistics, and land parcel identification system data
- **Tier 1 methods** are based on loosely related surrogate statistics, e.g. building use, population density, and land use.

The tiered methodology is outlined in the decision tree in Figure 7.1




Figure 7.1 General decision tree for emissions mapping (EMEP/EEA 2016)

Mapping of Ireland's emissions is done on a highly disaggregated level both regarding sectoral and spatial resolution. Spatial distribution keys (GeoKeys) are set up for each NFR category with a spatial resolution of 1 km x 1 km. GeoKeys are normalised tables including the share of an emission source that should be allocated to each grid cell. Some GeoKeys are used for all pollutants from a sector, while others are pollutant specific. Further, some sectors have been disaggregated into different sources and GeoKeys have been set up on source level and afterwards combined to create one overall GeoKey for the NFR sector. In this way, a high level of accuracy is ensured in the emission mapping as the highest tier level methodology is applied for all sources.

GeoKeys for the individual sectors/sources are built from a number of different spatial data. Some of the spatial data sets describe the emission allocation very accurate, e.g. E-PRTR reporting, while others are proxies for activity level or other related parameters; some being good proxies, e.g. mileage data for emissions from road transport, others being less good proxies, e.g. population density for domestic solvent use.

The common methodological approach is to make an overlay analysis of one or more spatial layers and the 1 km x 1 km grid in a Geographical Information System (GIS).

Preparation of the GeoKey for railway transport is described here as an example, and related maps are shown in Figure 7.2. Railway transport is an area source, and the emissions are spatially allocated to the railway network. The railway network is available as a digital map including the network as lines (Figure 7.2 a), which allow for the emissions to be distributed evenly to the railway network. Activity data based on railway statistics are added to the map's attribute data, and are used to improve the spatial distribution to allocate emissions according to the activity levels. An overlay of the railway network including activity data and the 1 km x 1 km grid is made in GIS (Figure 7.2 b), and the layers are intersected to cut the railway lines by the grid (Figure 7.2 c). The length of each line segment is calculated using standard GIS tools, and the share of each railway line in each grid cell is calculated. As more line segments can occur in the same grid cell, e.g. when two railway lines meet, the shares are summarised by grid cell to generate the GeoKey, which holds the share of the national emission by grid cell (Figure 7.2 d).



Figure 7.2 Example of GeoKey preparation

The emission mapping is made using an orthogonal grid with a resolution of 1 km x 1 km in the Irish projection TM65. The spatial emissions are redistributed into the reporting grid, the EMEP/EEA grid with a spatial resolution of 0.1 degree x 0.1 degree, using the share of each 1 km x 1 km grid cell that intersects the individual 0.1 degree x 0.1 degree grid cells.

7.3 Mapping methods for each GNFR

A summary of the data and mapping approaches used in compilation of the spatial inventory for Ireland are outlined in the tables below by GNFR sector. An indication of the tier 1-3 categorisation has also been provided as a simple measure of uncertainty in the approach applied. A more detailed description of the methodologies applied is available in Plejdrup et al. (2017)

7.3.1 Public Power (GNFR A_PublicPower)

Detailed location information for this sector were available on the individual large point sources for the NFR sector 1A1a Public Power. The emissions for some pollutants were available for the individual point sources, and in these cases, the data were used directly. For the pollutants where plant specific data were not available, the distribution is based on the activity data. The use of plant specific data and exact location of the emissions corresponds to a tier 3 method.

7.3.2 Industry (GNFR B_Industry)

As shown in Table 7.1, this GNFR covers many different source categories and hence the available spatial data vary across sectors. The categories include both combustion related categories and categories where the emissions are related to the process.

Where detailed emissions and location information were available, e.g. from the E-PRTR or the EU ETS on the individual point source emissions for the NFR sectors in GNFR B Industrial Combustion sector, these were used to map emissions to the known location. This is the case for e.g. emissions from refining and other energy industries (NFR categories 1A1b and 1A1c respectively).

For other source categories, some data are available at point source level, but the coverage does not match the national total. In these cases the emissions covered by point sources are allocated to the relevant point sources and the residual emission is distributed according to a more general spatial distribution key, e.g. industrial heat demand. These two distributions are then combined to one GeoKey covering the total sectoral emission.

This approach is considered a tier 2 or tier 3 method.

7.3.3 Other stationary combustion (GNFR C_OtherStationaryComb)

This GNFR category covers combustion in three subsectors, i.e. commercial/institutional, residential and agriculture. The most important sector in terms of emission contribution is residential combustion.

For commercial/institutional plants, the distribution is based on heat demand for commercial and public buildings as calculated by the Irish Heat Map. The Heat Map is based on a study from 2015 commissioned by SEAI to fulfil Ireland's requirements under article 14 of the Energy Efficiency Directive (2012/27/EU). As part of this study a spatial representation of Ireland's heat demand was developed.

For residential plants, the distribution is based on information from the 2011 census on primary fuel types in households combined with an estimated unit consumption calibrated with the estimated national residential fuel consumption and the emission factors used in the emission inventory.

For the agricultural sector, the spatial data on farmyards and buildings from the Land Parcel Information System (LPIS) were used.

This is considered tier 2/3 methodologies.

7.3.4 Fugitive Emissions (GNFR D_Fugitive)

This sector covers both categories estimated as point sources (e.g. coal mining/handling, service stations and flaring) and area sources (e.g. natural gas distribution).

The point source data have been used to allocate emissions and, where available, activity data have been incorporated to further improve the distribution of emissions. Information on coal mining areas as well as coal consumers were provided by the EPA, while a list of service stations was provided by the CSO. For natural gas distribution, the spatial information included for gas use in the Heat Map was utilised to distribute emissions.

This is considered tier 2/3 methodologies.

7.3.5 Solvents (GNFR E_Solvents)

The national emissions from domestic solvent use were mapped across the country using population density as spatial proxy. This approach is a tier 2 method.

For the remaining solvent use categories, there was some spatial information available, e.g. location of dry cleaners. However, both population density and industrial heat demand were used as spatial proxies to map emissions from coating applications, chemical products and printing. This is considered tier 1/2/3 methodologies.

7.3.6 Road transport (GNFR F_RoadTransport)

Spatial mileage data for national roads (NR) provided by TII for total mileage and % heavy vehicles are used to allocate emissions from road transport on NR. Road transport on other roads is estimated as the residual of the national total mileage used in the inventory, and emissions are allocated to roads other than national roads. As mileage data is not available for other roads than NR a polygon map of the road network is applied for mapping, thereby using road area as a proxy for the activity level. Separate GeoKeys are prepared for passenger cars including vans and 2-wheelers (PC), heavy vehicles including busses (HV), and all vehicles (PC+HV).

The approaches used are tier 3 for national roads and tier 2 for remaining roads.

7.3.7 National navigation (GNFR G_Shipping)

The estimates of the emissions from national navigation were mapped using a buffer zone of six nautical miles around the coast of Ireland. The buffer zone was adjusted to take into account the shortest path between headlands in Ireland. This approach is a tier 2 method.

7.3.8 Aviation (LTO) (GNFR H_Aviation)

National total emissions from aircraft operating on the ground and in the air over Ireland, up to an altitude of 1000 m (equating to the take offs and landing – LTO) were mapped at the locations of the airports including a five-kilometre buffer zone. The number of LTOs at each airport was used to further improve the distribution of emissions. This approach is a tier 3 method.

7.3.9 Off road mobile sources (GNFR I_Offroad)

This GNFR category comprises several different activities such as railways, fishing and agricultural machinery.

For railways, the railway network and data for annual passages were provided by Irish Rail and this information has been used to develop a GeoKey for this sector.

For fishing, the emissions have been distributed based on data for fishing areas within the Irish exclusive economic zone and fishing statistics.

For agricultural machinery, data on the number of different types of machinery at county level were obtained from the CSO and this information was combined with the land information from LPIS on cropland and improved grassland.

This is considered tier 2/3 methodologies.

7.3.10 Waste handling and treatment (GNFR J_Waste)

The estimates of the emissions from solid waste disposal on land were mapped at the locations of landfill sites. For composting 75 % of the emission were allocated to the licensed facilities while the remaining 25 % were allocated to non-urban residential buildings.

Emissions from clinical waste incineration, industrial waste incineration and cremation were mapped at the locations of the known facilities. Activity and location data for industrial waste incinerators and crematoria were available and used to weight emissions to areas of known activity proportionally.

Estimates of the national emissions from other waste handling (e.g. accidental fires) were mapped according to population density.

This is considered tier 1/2/3 methodologies

7.3.11 Agricultural livestock (GNFR K_AgriLivestock)

National emissions from pigs and poultry were distributed based on detailed data on farms and animal numbers from the 2010 agricultural census provided by UCD.

National emissions from mink were distributed based on farm locations and animal numbers provided by EPA.

National emissions from cattle, sheep and horses were distributed based on data from the 2010 agricultural census provided by CSO on the number of animals per electoral district combined with the Land Parcel Identification System (LPIS) data on location of farmyards and buildings.

National emissions from goats, mules and asses, and deer were distributed based on data from the 2010 agricultural census provided by CSO on the number of animals per county combined with the Land Parcel Identification System (LPIS) data on location of farmyards and buildings.

This approach is a tier 2/3 method.

7.3.12 Agricultural soils (Other emissions) (GNFR L_AgriOther)

National emissions from sources related to agricultural soils, e.g. application of fertiliser and manure as well as grazing animals were distributed on cropland and/or grassland from LPIS, taking into account the animal density when distributing emissions from animal manure. This approach is a tier 1/2 method.

7.3.13 Aviation (Cruise) (GNFR O_AviCruise)

This category includes cruise emissions from both national and international aviation. For national cruise emissions, the distribution is based on information on the number of flights between Irish airports and emissions are allocated to great circle lines between these airports.


For international cruise emissions, the majority of emissions will occur outside the Irish territory, but for the purposes of the submission, emissions are allocated evenly across the entire Irish area outlined by the Irish Exclusive Economic Zone (EEZ).


7.3.14 International navigation (GNFR P_IntShipping)


Emissions from international navigation will largely occur outside the Irish territory, but for the purposes of the submission, emissions are allocated evenly across the sea area outlined by the Irish Exclusive Economic Zone (EEZ).


7.4 National total emission mapped by the EMEP 0.1 degree x 0.1 degree grid

Figure 7.3 a-e present the mapped national total emissions for NO_x, SO₂, NMVOC, NH₃, and PM_{2.5} in Ireland by EMEP 0.1 x 0.1 degree grid.












Figure 7.3 National Total Emissions in 2015 for a) NO_x, b) SO₂, c) NMVOC, d) NH₃, and e) PM_{2.5}

Chapter Eight

Projections

8.1 Overview of Emissions Projections

Ireland is currently finalising updated projections and will submit the updated projections using the template contained within annex IV of the reporting guidelines. Ireland will update this chapter accordingly and resubmit it's IIR for 2018. The data presented in this chapter thus relates to projections data submitted in 2017.

This chapter presents the emission projections that have been compiled for reporting under the National Emissions Ceiling Directive and the Convention on Long Range Transboundary Air Pollution (CLRTAP).

Section 8.2 describes emission reduction targets for 2020 and 2030. Section 8.3 provides a short explanation of how energy forecasts are generated for both “*With Existing Measures*” and “*With Additional Measures*” scenarios. Information on key assumptions and underlying data are also provided.

Section 8.4 presents the emission projections for each of the pollutants covered, and considers the key trends across the time series which include the impact of national policies and measures (in response to European Directives and legislation) aimed at reducing greenhouse gas emissions. Sections 8.5 to 8.13 consider each of the main NFR source sectors, and provide a more detailed explanation of the emission projections compilation process for each sector.

The NEC Directive requires biennial reporting from 2017 of projected emissions for SO₂, NO_x, NH₃, NMVOC, PM_{2.5} and, if available, Black Carbon covering projection years 2020, 2025, 2030 and, where available, 2040 and 2050.

The CLRTAP guidelines for reporting emissions and projections data state that parties to the Gothenburg Protocol within the geographical scope of the EMEP shall regularly update their projections and report every four years from 2015 onwards their updated projections, for the years 2020, 2025 and 2030 and, where available, also for 2040 and 2050. Parties to the other protocols are encouraged to regularly update their projections and report every four years from 2015. In addition, parties should provide a “*With Existing Measures*” and where relevant a “*With Additional Measures*” projection estimate.

Projected emission estimates and supporting quantitative information were reported for SO₂, NO_x, NH₃, NMVOC, PM_{2.5} and, Black Carbon under the NEC Directive and CLRTAP utilising the reporting template contained within annex IV of the Guidelines for Reporting Emissions and Projections Data under the CLRTAP⁵.

⁵ http://www.ceip.at/reporting_instructions/annexes_to_guidelines/

This chapter details emission projections under both the *With Existing Measures* scenario and *With Additional Measures* scenario for the following pollutants for the period 2016-2030: NO_x, SO₂, NMVOC, NH₃ and PM_{2.5}.

8.2 Emission Reduction Targets for 2020 and 2030

The Gothenburg Protocol includes national emission reduction commitments to be achieved in 2020 and beyond for NO_x, SO_x, NMVOC and NH₃ and PM_{2.5}.

The National Emission Ceilings Directive (NECD, 2001/81/EC) was reviewed as part of the Clean Air Policy Package and a new Directive came into effect in December 2016⁶. Emission reduction commitments have been set for Ireland for 2020 and 2030 for NO_x, SO_x, NMVOC, NH₃, and PM_{2.5}. Table 8.1 details the emission reduction targets in place for each pollutant for 2020 under the Gothenburg Protocol and 2020 and 2030 under the new National Emissions Ceilings Directive.

Table 8.1. Emission Reduction Commitments for 2020 and 2030 (expressed as a percentage reduction of 2005 levels)

Pollutant	SO ₂	NO _x	NH ₃	NMVOC	PM _{2.5}
2020	65%	49%	1%	25%	18%
2030	85%	69%	5%	32%	41%

8.3 With Existing Measures and With Additional Measures Scenarios

Sustainable Energy Authority of Ireland (SEAI) compile national energy forecasts presenting energy trends into the future. The most recent forecast provides energy demand trends to 2035. These energy forecasts form the basis for almost all energy-related emission projections presented in this chapter. The latest energy forecasts include two scenarios which are used in national emission projections: *Baseline* and *NEEAP/NREAP* (adjusted for 2017 projections to reflect current progress and the trajectory towards achieving 2020 energy efficiency and renewable energy targets).

The *Baseline* energy forecast projects forward Ireland's energy demand, incorporating the expected impacts of policies and measures that were in place (e.g. legislatively provided for) by the end of 2015. It represents a hypothetical future scenario in which no further policy actions or measures have been taken. It excludes policies that are committed to but which do not yet have measures in place to deliver them. The *Baseline* energy forecast thus underpins the energy related *With Existing Measures* (WEM) air pollutant emission projections.

The *NEEAP/NREAP* energy forecast (adjusted for 2017 projections) presents an alternative view of future energy demand that accounts for further implementation of the National Renewable Energy Action Plan⁷ (NREAP) and the 3rd National Energy Efficiency Action Plan⁸ (NEEAP) based on current progress. Therefore, this forecast includes existing *and* further implementation of planned policies and measures based on current progress. For 2017 projections, the latest *NEEAP/NREAP* energy forecast has been adjusted to reflect

⁶ DIRECTIVE (EU) 2016/2284 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC

⁷ <http://www.dccae.gov.ie/energy/en-ie/Renewable-Energy/Pages/Action-Plan.aspx>

⁸ <http://www.dccae.gov.ie/energy/SiteCollectionDocuments/Energy-Efficiency/NEEAP%203.pdf>

current progress and the trajectory towards achieving 2020 targets. This includes an expected shortfall in achieving full energy efficiency targets and renewable energy targets for electricity, transport and heat. The *NEEAP/NREAP* energy forecast (adjusted) therefore underpins the energy related *With Additional Measures* (WAM) emission projections.

For the *Baseline* energy forecast, the Economic and Social Research Institute (ESRI) use macro-economic projections which are produced using the COSMO model⁹. The baseline projections and underlying assumptions are described here in Chapter 1 of “Ireland’s Economic Outlook: Perspectives and Policy Challenges”, which was published on 5 December 2016¹⁰. Projections on the global economic environment, including oil prices, as based in simulations using the NiGEM model (National Institute Global Econometric Model¹¹) maintained by the National Institute of Economic and Social Research¹². Projections from the COSMO model were used to produce projections of the energy demand equation time series variables (i.e. demand equations by fuel and sector). The integration of energy demand into the COSMO model is work that is due to be undertaken in 2017.

Annual electricity demand, which is an output of the electricity demand equations/COSMO was transferred, as well as fuel prices, as an input into an electricity dispatch model to determine fuels used at an hourly level to service aggregate electricity demand. This process provides a high level of accuracy on the fuels used in the electricity sector. The software used for the energy forecasts to model the Irish Electricity Market is PLEXOS 7.4 R01. PLEXOS is a power systems modelling tool used for electricity market modelling and planning.

The energy forecast includes sectoral output figures and other relevant key variables such as price, economic growth, population and housing stock. To produce the finalised *Baseline* energy forecast, SEAI amends the output of the energy demand produced by ESRI (described above) to take account of the expected impact of energy efficiency measures put in place before the end of 2015 but which are considered too recent to be detectable in any time-series analysis. The *NEEAP/NREAP* energy forecast (adjusted) builds on the *Baseline* forecast with adjustments made to account for further implementation of additional policies and measures outlined in the NEEAP⁵ and NREAP⁴. For 2017 projections, the *NEEAP/NREAP* energy forecast has been adjusted to reflect current progress and the trajectory towards achieving 2020 targets. This includes an expected shortfall in achieving energy efficiency and renewable energy targets.

The energy forecasts that underpin the energy-related emissions projections are based on macroeconomic projections as described above. Table 8.2 shows the key parameters underlying the macroeconomic outlook and therefore the *With Existing Measures* and *With Additional Measures* emission projections scenarios. The forecasts are based on international fuel import oil prices. Coal and gas prices were published by the United Kingdom’s Department of Energy and Climate Change. The carbon prices are those circulated by the European Commission in June 2016. Carbon dioxide price assumptions in the non-ETS sectors are based in the medium term on the Finance Bill 2010¹³ which saw the introduction of a carbon tax of €15 per tonne CO₂. In the longer term the carbon tax is assumed to follow the EU ETS carbon price.

⁹ <https://www.esri.ie/projects/modelling-the-irish-economy/>

¹⁰ <http://www.esri.ie/pubs/EO1.pdf>

¹¹ <https://nimodel.niesr.ac.uk/>

¹² <http://www.niesr.ac.uk/>

¹³ Finance Bill 2010. <http://www.finance.gov.ie/ga/news-centre/press-releases/finance-bill-2010>

Activity data forecasts and the methodological approach to emission projections from the agriculture and waste sectors are discussed in their respective sections.

Table 8.2. Key assumptions underpinning the energy forecasts

	2016 – 2020	2021-2025	2026-2030
Average Annual % Growth Rate			
GDP	+3.74%	+3.24%	+2.59%
GNP	+3.42%	+3.32%	+1.97%
Personal Consumption	+2.97%	+2.57%	+1.11%
	2020	2025	2030
Housing Stock ('000)	2,018	2,112	2,206
Population ('000)	4,834	5,027	5,209
EUETS: Carbon €₂₀₁₃/tCO₂	15	22.5	33.5
Carbon tax €₂₀₁₃/tCO₂	15	22.5	33.5
Coal \$₂₀₁₃/boe	9.9	11.6	10.6
Oil \$₂₀₁₃/boe	56.8	62.8	69.4
Gas \$₂₀₁₃/boe	20.4	24.6	27.3
Peat €/MWh	25	25	25

8.4 Key Trends

Air pollutant emission projections have been generated for a WEM scenario and a WAM scenario. In the following sections, both scenarios are presented in the context of emission reduction targets set for 2020 and 2030. The WAM scenario is then considered in more detail.

Figures in the following sections include historic air pollutant emission estimates for 2005 to 2015 as contextual information. Tables in the following sections present projected emission estimates up to and including 2030 under both the WEM and WAM scenarios.

8.4.1 Sulphur Dioxide (SO₂)

Emission projections for SO₂ for the WEM and WAM scenarios are presented in Figure 8.1. The emission reduction targets for 2020 and 2030 are also presented.

Total SO₂ emissions under the WAM scenario are projected to be 11.8 kt in 2020. The emission projections predict compliance with the 2020 emission reduction target by 14.1 kt. Total SO₂ emissions are projected to be 8.3 kt in 2030 under the WAM scenario which is below the emission reduction target for that year.

The difference between the WEM and WAM scenarios is attributed to the effect of further penetration of renewables in electricity generation in addition to renewables and energy efficiency measures in buildings in the residential, commercial and manufacturing industries sectors.




Figure 8.1 SO₂ Emission Projections for the With Existing Measures and With Additional Measures Scenarios

Sulphur dioxide emissions under the WAM scenario by source sector are presented in Figure 8.2. Emissions from Public Electricity and Heat Production (1A1a) currently contribute approximately one-third of national total emissions of SO₂. From 2025 onwards, emissions are projected to significantly reduce further which is attributed to an assumption that coal will be replaced with gas that is used for electricity generation.




Figure 8.2 SO₂ Emission Projections for the With Additional Measures Scenario by Source Sector

Table 8.3. Projected Emissions of SO₂ under the With Existing Measures and With Additional Measures Scenarios (kt)

With Existing Measures scenario					
	2015	2016	2020	2025	2030
Public Electricity and Heat Production	5.2	7.3	2.0	1.6	1.9
Residential & Commercial/Institutional	7.0	6.9	5.2	4.1	3.5
Manufacturing Industries and Construction	2.3	2.1	2.0	1.9	1.9
Agriculture/Forestry/Fishing	0.0	0.1	0.0	0.1	0.1
Transport	0.3	0.3	0.3	0.3	0.3
Other NFR sectors	2.8	3.1	3.1	3.3	3.3
TOTAL	17.6	19.7	12.6	11.2	10.9
With Additional Measures scenario					
Public Electricity and Heat Production	5.2	7.3	1.5	0.2	0.2
Residential & Commercial/Institutional	7.0	6.9	5.0	3.7	2.8
Manufacturing Industries and Construction	2.3	2.0	1.8	1.8	1.7
Agriculture/Forestry/Fishing	0.0	0.1	0.0	0.1	0.1
Transport	0.3	0.3	0.3	0.3	0.3
Other NFR sectors	2.8	3.1	3.1	3.3	3.3
TOTAL	17.6	19.5	11.8	9.2	8.3

8.4.2 Nitrogen Oxides (NO_x)

Emission projections for NO_x for the WEM and WAM scenarios are presented in Figure 8.3. The emission reduction targets for 2020 and 2030 are also presented.

Total NO_x emissions under the WAM scenario are projected to be 57.3 kt in 2020. The emission projections predict compliance with the 2020 emission reduction target. Total NO_x emissions in 2030 are projected to be 47.0 kt, which is an estimated 4.7 kt non-compliance with the emission reduction target.

Projected emissions across the time series show a steady decline for both the WEM and WAM scenarios. The forecasted replacement of coal with natural gas in Public Electricity and Heat Production (1A1a) under the WAM scenario reduces emissions of NO_x from the sector in 2025. The projected emissions trend is very much dominated by the transport (1A3) sector (Figure 8.4).

Similar to SO₂ the policies and measures aimed at the use of renewable fuels and improving energy efficiency in dwellings, commercial and public buildings and industry are responsible for the difference between the WEM and WAM scenarios.




Figure 8.3 NO_x Emission Projections for the With Existing Measures and With Additional Measures Scenarios

Figure 8.4 presents the NO_x emissions under the WAM scenario by source sector. Emissions from transport (1A3) are the largest contributor to the total accounting for approximately 50% of national total emissions for much of the time series. Emissions from the transport sector (1A3) show a decrease across the time series.




Figure 8.4 NO_x Emission Projections for the With Additional Measures Scenario by Source Sector

Table 8.4. Projected Emissions of NO_x under the With Existing Measures and With Additional Measures Scenarios (kt)

With Existing Measures Scenario					
	2015	2016	2020	2025	2030
Public Electricity and Heat Production	9.5	6.6	5.1	5.4	6.1
Residential & Commercial/Institutional	7.4	7.6	7.5	7.8	7.9
Manufacturing Industries and Construction	11.0	9.5	9.7	9.7	9.6
Agriculture/Forestry/Fishing	4.4	4.6	3.7	3.0	2.5
Transport	45.9	42.5	31.2	25.5	22.1
Other NFR sectors	1.4	1.5	1.5	1.6	1.6
Total	79.5	72.3	58.8	53.0	49.9
With Additional Measures Scenario					
Public Electricity and Heat Production	9.5	6.6	4.5	4.0	4.4
Residential & Commercial/Institutional	7.4	7.5	7.3	7.5	7.5
Manufacturing Industries and Construction	11.0	9.1	9.1	9.0	9.0
Agriculture/Forestry/Fishing	4.4	4.6	3.7	3.0	2.5
Transport	45.9	42.4	31.2	25.5	22.1
Other NFR sectors	1.4	1.5	1.5	1.6	1.6
Total	79.5	71.8	57.3	50.6	47.0

8.4.3 Ammonia (NH₃)

Figure 8.5 presents the emission projections for NH₃ for the WEM and WAM scenarios. The emission reduction targets for 2020 and 2030 are also presented.

Total NH₃ emissions under the WAM scenario are projected to be 116.5 kt in 2020. As a result an exceedance of the 2020 emission reduction target by 6.6 kt is projected. Post 2020, emissions decrease to 116.2 kt in 2030. Ireland's 2030 emission target of a 5 per cent reduction on 2005 levels results in a distance to the emission reduction target in 2030 of 10.7 kt under the WAM scenario. Compliance with the 2030 emission reduction target will present a significant challenge as a result.




Figure 8.5 NH₃ Emission Projections for the With Existing Measures and With Additional Measures Scenarios

The WAM scenario includes a reduction in fertilizer nitrogen use due to the introduction of urease and nitrification inhibitors into the market. It is estimated that approximately 0.7 kt NH₃ may be saved on an annual basis as a result of this measure and is responsible for majority of the difference identified in Figure 8.5 between the WEM and WAM scenarios.

Ammonia emissions by source sector under the WAM scenario are presented in Figure 8.6. Emissions from dairy cattle show considerable growth (20 per cent) between 2015 and 2020, driven by the target of increasing milk production as set out in the Food Wise 2025 strategy (see section 8.12). This strategy is discussed in more detail in section 8.12. Post 2020 emissions from the agriculture sector continue to grow up until 2024 after which time emissions fluctuate which is attributed to a reduction in the quantity of emissions from the other cattle sector as the beef herd contracts.

It is forecasted that nitrogen (N) fertilizer application will increase by 21 per cent between 2015 and 2020 and remain relatively constant post 2020.




Figure 8.6 NH₃ Emission Projections for the With Additional Measures Scenario by Source Sector

Table 8.5. Projected Emissions of NH₃ under the With Existing Measures and With Additional Measures Scenarios (kt)

With Existing Measures Scenario					
	2015	2016	2020	2025	2030
Dairy Cattle	13.9	13.8	16.7	18.6	19.7
Other Cattle	30.7	31.1	30.9	29.5	27.3
Other Livestock	10.9	11.0	12.2	12.6	12.6
Synthetic Fertiliser use	9.2	10.7	11.1	11.0	11.0
N-excreted on Pasture	13.0	13.0	13.7	13.7	13.3
Manure and sludge applied to soil	29.2	29.3	31.4	32.1	31.8
Road Transport	0.9	0.8	0.7	0.7	0.7
Other	0.4	0.4	0.4	0.4	0.4
Total	108.1	110.2	117.2	118.6	116.9
With Additional Measures Scenario					
Dairy Cattle	13.9	13.8	16.7	18.6	19.7
Other Cattle	30.7	31.1	30.9	29.5	27.3
Other Livestock	10.9	11.0	12.2	12.6	12.6
Synthetic Fertiliser use	9.2	10.7	10.3	10.2	10.2
N-excreted on Pasture	13.0	13.0	13.7	13.7	13.3
Manure and sludge applied to soil	29.2	29.3	31.4	32.1	31.8
Road Transport	0.9	0.8	0.7	0.7	0.7
Other	0.4	0.4	0.5	0.5	0.6
Total	108.1	110.2	116.5	117.9	116.2

8.4.4 Non-Methane Volatile Organic Compounds (NMVOCs)

Figure 8.7 presents the emissions projections for NMVOC for the WEM and WAM scenarios. The emission reduction targets for 2020 and 2030 are also presented.

Article 4 (3) of the National Emission Ceiling Directive provides that emissions of non-methane volatile organic compounds from categories 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with 2020 and 2030 targets. The *with existing measures* emissions and *with additional emissions* in addition to 2020 and 2030 targets displayed in Figure 8.7 exclude emissions from these categories (3B and 3D).

Total NMVOC emissions under the WAM scenario (excluding emissions from agricultural categories 3B and 3D) are projected to be 57.4 kt in 2020. The emission projections predict a distance to target of 4.8 ktg. Emissions in 2030 are projected to be 57.4 kt which is 8.7 kt above the emission reduction target.

Projected emissions across the time series show a slight increase in both the WEM and WAM scenarios up to 2020 followed by a gradual reduction out to 2030. The trend of reduced emissions in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors are counteracted by the increase in emissions from solvents and fugitive emissions.




Figure 8.7 NMVOC Emission Projections for the With Existing Measures and With Additional Measures Scenarios¹⁴

Projected NMVOC emissions by source sector under the WAM scenario are presented in Figure 8.8. For the purposes of this graph emissions from Agriculture categories 3B and 3D are included in the projections from all years out to 2030. Projected emissions from Agriculture and Solvents and other product use are the drivers of the trend, cumulatively accounting for approximately 87 per cent on average of total emissions in 2030 under the WAM scenario.




Figure 8.8 NMVOC Emission Projections for the With Additional Measures Scenario by Source Sector¹⁵

¹⁴ Article 4 (3) of the National Emission Ceiling Directive provides that emissions of non-methane volatile organic compounds from categories 3B (manure management) and 3D (agricultural soils) are not accounted for the purpose of complying with 2020 and 2030 targets. The *with existing measures* emissions and *with additional emissions* in addition to 2020 and 2030 targets displayed in this graph exclude emissions from these categories (3B and 3D)

Table 8.6. Projected Emissions of NMVOC under the With Existing Measures and With Additional Measures Scenarios (kt)

With Existing Measures Scenario					
	2015	2016	2020	2025	2030
Public Electricity and Heat Production	0.3	0.3	0.3	0.4	0.4
Residential & Commercial/Institutional	9.2	9.1	6.9	5.5	4.8
Manufacturing Industries and Construction	2.9	3.9	4.2	4.2	4.2
Agriculture	43.2	43.3	44.9	44.5	42.9
Transport	5.9	5.4	4.4	4.2	4.0
Fugitive and Solvents	39.2	39.8	41.0	42.8	44.3
Other NFR Sectors	0.9	0.9	0.7	0.5	0.5
Total	101.4	102.6	102.4	102.2	101.2
With Additional Measures Scenario					
Public Electricity and Heat Production	0.3	0.3	0.3	0.4	0.4
Residential & Commercial/Institutional	9.2	9.0	6.7	5.0	3.9
Manufacturing Industries and Construction	2.9	3.8	4.3	4.4	4.4
Agriculture	43.2	43.3	44.9	44.5	42.9
Transport	5.9	5.4	4.5	4.2	4.0
Fugitive and Solvents	39.2	39.8	41.0	42.8	44.2
Other NFR Sectors	0.9	0.9	0.7	0.5	0.5
Total	101.4	102.4	102.2	101.8	100.3

8.4.5 Particulate Matter < 2.5 µm in diameter (PM_{2.5})

Emissions projections for PM_{2.5} for the WEM and WAM scenarios are presented in Figure 8.9. The emission reduction targets for 2020 and 2030 are also presented.

Total PM_{2.5} emissions under the WAM scenario are projected to be 11 kt in 2020. The emission projections under the WAM scenario predict compliance with both the 2020 and 2030 emission reduction targets by a margin of 4.8 kt and 2.9 kt, respectively.

Projected emissions across the time series show a steady decline for both the WEM and WAM scenarios up to 2030 (Figure 8.9). This trend is largely the result of forecasted reductions in emissions in the Residential (1A4b) and Commercial/Institutional (1A4a) sectors.

¹⁵ Note that for the purposes of this graph emissions from Categories 3B and 3D are included in the projections from all years out to 2030




Figure 8.9 PM_{2.5} Emission Projections for the With Existing Measures and With Additional Measures Scenarios

Figure 8.10 presents the projected PM_{2.5} emissions under the WAM scenario by source sector. Emissions from Residential (1A4b) and Commercial/Institutional (1A4a) sectors cumulatively are the largest contribution to the projected emissions total by 2030, and also dominate the trend with time. Emissions from the Transport sector (1A3) decrease across the time series, due to the introduction of vehicles with improved emission control technologies.




Figure 8.10 PM_{2.5} Emission Projections for the With Additional Measures Scenario by Source Sector

Table 8.7. Projected Emissions of PM_{2.5} under the With Existing Measures and With Additional Measures Scenarios (kt)

With Existing Measures Scenario					
	2015	2016	2020	2025	2030
Public Electricity and Heat Production	0.4	0.4	0.3	0.3	0.3
Residential & Commercial/Institutional	7.5	7.4	5.6	4.4	3.7
Manufacturing Industries and Construction	1.5	1.7	1.7	1.7	1.6
Agriculture	2.2	2.1	2.2	2.3	2.2
Transport	2.1	1.6	1.2	1.1	1.0
Other NFR Sectors	0.2	0.2	0.1	0.1	0.1
Total	13.9	13.5	11.1	9.8	9.1
With Additional Measures Scenario					
Public Electricity and Heat Production	0.4	0.4	0.2	0.2	0.2
Residential & Commercial/Institutional	7.5	7.4	5.5	4.1	3.2
Manufacturing Industries and Construction	1.5	1.7	1.7	1.7	1.7
Agriculture	2.2	2.1	2.2	2.3	2.2
Transport	2.1	1.6	1.2	1.1	1.0
Other NFR Sectors	0.2	0.2	0.1	0.1	0.1
Total	13.9	13.4	11.0	9.5	8.4

8.5 Energy Industries (NFR 1A1)

Public Electricity and Heat Production covers all electricity generation including electricity generated from renewable sources. The Plexos_Ireland model was used to model electricity generation. As an electrical systems model, the core input data comprises technical details of generators, transmission lines and loads as well as fuel costs, operational costs and emission reduction rates and costs.

In the *Baseline* energy forecast the renewable energy generated shows Ireland reaching 22.7 per cent of electricity consumption from renewable energy by 2020. Renewable electricity generation capacity is dominated by wind but also includes, for example, the operation of a second waste to energy incinerator and the continued development of landfill gas electricity generation. It is also assumed that electricity trading occurs through the 500 MW East-West interconnector. In 2030 it is estimated that renewable energy generation increases to 25 per cent of electricity consumption.

In the *NEEAP/NREAP* energy forecast (adjusted) (and therefore the *With Additional Measures* emissions scenario) it is assumed that for 2020 there is a 37.3 per cent share of renewable energy in electricity generation as a result of additional expansion in wind energy, biomass electricity generating capacity in addition to solar photo voltaics and the continued development of landfill gas electricity generation. The largest contribution is from wind which at 896 ktoe in 2020 is 62 per cent above that included in the *Baseline* and therefore the *With Existing Measures Scenario*. This falls short of the full target of 40 per cent share of renewable energy in electricity generation in 2020. In 2030 it is estimated that renewable energy generation reduces to 29 per cent of electricity consumption.

The *With Existing Measures* scenario takes into account the impact of current policies and measures in the energy sector including:

- Increased efficiency in power generation
- Reduced transmission and distribution losses
- 22.7% renewables by 2020
- Reduced electricity demand from energy efficiency measures in the various sectors including residential, industry, commercial/institutional

The *With Additional Measures* scenario takes into account the impact of additional policies and measures in the energy sector including:

- Reduced electricity demand from additional energy efficiency measures in industry, services and residential
- Increased electricity demand from electric vehicles roll-out
- Replacement of coal fired generation with natural gas
- 37.3% renewable by 2020

Emission factors for NO_x are based on those in the national inventory. Emission abatement (e.g. Selective Catalytic Reduction used in coal fired electricity generation) and compliance with the Industrial Emissions Directive (2010/75/EC) are taken into account.

In Ireland, biomass used for electricity generation is co-fired, generally with peat. It is currently not possible to separate the biomass component from reported emission levels in individual plant's IPC / Industrial Emissions Licence Annual Environmental Reports and therefore emissions are split on an energy basis with implied emission factors thus calculated for peat and biomass. This approach is applied for all future years to account for the combustion of biomass for electricity generation.

Emission factors for SO₂ are based on those in the national inventory. Emission abatement (e.g. flue gas desulphurisation used in coal fired electricity generation) and compliance with the Industrial Emissions Directive (2010/75/EC) are taken into account.

There has been significant reduction in the use of oil in electricity generation due to the closure and decommissioning of oil fuelled generation plants. Oil is also used as a start-up fuel in coal and peat fired generation stations. The 2015 SO₂ inventory emission factor for oil is assumed for all future years. Utilising the emission factor applied in the national inventory for 2015 allows for a decrease compared with historic years which is assumed to implicitly cover the impact of Directive 2009/30/EC (concerning a reduction of sulphur content of certain liquid fuels).

The SO₂, NMVOC and PM_{2.5} emission factors for the other relevant fuel types (i.e. coal, peat biomass and non-renewable wastes) are assumed to remain constant at the value utilised in the national inventory for 2015. The same approach as that taken in NO_x emission estimates with respect to co-firing of biomass is applied to projected emission estimates for SO₂, NMVOC and PM_{2.5}.

8.5.1 Oil Refining and Solid Fuel Manufacturing (NFR 1A1b and 1A1c)

Projected NO_x, SO₂, NMVOC and PM_{2.5} emissions from oil refining (one plant) and solid fuel manufacture are estimated. Energy forecasts are not available for these sectors. Projections are based on the growth rate of projected greenhouse gas emissions, which are provided to the EPA by the relevant installation operators.

8.6 Manufacturing Industries and Construction (NFR 1A2)

Projected NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emissions are based on SEAI's *Baseline* and *NEEAP/NREAP* energy forecasts (2017 NEEAP and NREAP forecasts were adjusted) and are therefore estimated for both the WEM and WAM scenarios.

The *With Existing Measures* scenario takes into account the impact of current policies and measures in the industrial sector including:

- SEAI Large Industry Programme
- CHP deployment
- Accelerated Capital Allowance (ACA)
- Renewable Heat
- Carbon tax
- Better Energy Workplaces

The *With Additional Measures* scenario takes into account the impact of additional policies and measures in the industrial sector including:

- Increased share of renewable sources used for thermal energy
- Increased energy efficiency in buildings

The projected emission factors for NO_x and SO₂ from the combustion of coal, natural gas and petroleum coke are based on the weighted average emission factor for coal, natural gas and petroleum coke across the sub sectors 1A2a – 1A2g in 2015. Energy forecasts are only provided to the EPA at an aggregated industry level (i.e. 1A2). Emission factors for NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} for the combustion of other fuels are assumed to remain constant at 2015 values for all future years.

8.7 Transport (NFR 1A3)

Transport emissions cover Aviation (1A3a), Road Transportation (1A3b), Rail (1A3c), Navigation (1A3d) and Other transportation (Natural gas pipeline compressors, 1A3e). Projected emissions of NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} from road transport are based on SEAI's *Baseline* and *NEEAP/NREAP* energy forecasts (adjusted for 2017 forecasts) and are therefore estimated for both the WEM and WAM scenarios.

It is assumed that fuel use in rail will remain constant at 2015 levels for each year out to 2030.

Projected fuel combustion from navigation is assumed to be equal that combusted in the sector in 2015 for each projected year to 2030. Navigation (1A3d) emissions for each of the pollutants are assumed to remain constant at 2015 levels for each future year.

Other Transportation (1A3e) refers to the use of natural gas for combustion in natural gas pipeline compressor stations. Emissions from this sector are inferred from forecast gas demand in the residential, industrial and commercial and institutional services sectors from the energy forecasts provided by SEAI and are calculated for both the WEM and WAM scenarios.

The *With Existing Measures* scenario takes into account the impact of current policies and measures in the transport sector including:

- VRT and Motor Tax changes

- Improved fuel economy of private cars
- Public transport efficiency improvements
- Aviation efficiency
- Carbon tax
- Renewables

The *With Additional Measures* scenario takes into account the impact of additional policies and measures in the transport sector including:

- Electric vehicle deployment
- Natural gas transport savings between scenarios
- Increased share of renewable sources (e.g. biofuels) used for transport

8.7.1 Domestic aviation (NFR 1A3a)

NO_x, SO₂, NMVOC and PM_{2.5} emission projections from aviation are estimated using the 2015 inventory and also forecasted data, where available, related to aircraft movements as provided to the EPA by the management authorities of Ireland's main airports (Dublin, Cork and Shannon).

Emissions associated with all LTO (landing and take-off) cycles are calculated. It is assumed that NO_x, SO₂, NMVOC and PM_{2.5} emission factors remain constant at the values used in the 2015 national inventory.

8.7.2 Road transportation (NFR 1A3b)

Energy forecasts provide future demand for petrol, diesel, renewables and electricity use in the road transport sector for both the WEM and WAM scenarios.

In the energy forecast underpinning the WAM emission projection for road transport, it is assumed that renewables will account for 8 per cent of road transport fuel by 2020.

NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emissions from road traffic are estimated using the COPERT model (COPERT 4 Version 11.3) developed within the CORINAIR programme (Gkatzoflias et al., 2012).

8.7.3 Rail (NFR 1A3c)

NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emission projections from rail transport are estimated. It is assumed that fuel use in the sector will remain constant at 2015 levels for each year out to 2030. It is also assumed that NO_x, NMVOC, NH₃ PM_{2.5} emission factors remain the same as in the 2015 national inventory.

8.7.4 Navigation (NFR 1A3d)

NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emission projections from navigation are estimated. Gasoil/diesel consumption in in-land navigation is assumed to remain constant at the 2015 level out to 2030. NO_x, NMVOC, NH₃ and PM_{2.5} emission factors are assumed to remain constant at the 2015 level.

8.7.5 Gas Transmission (NFR 1A3e)

Emissions projections for NO_x, SO₂, NMVOC, and PM_{2.5} from natural gas transmission in Ireland's natural gas pipeline network are estimated. Future gas demand for "own use and transformation" is inferred based on forecast gas demand in the residential, commercial and institutional services and industrial sectors. Subtracting the amount of gas estimated to be lost from the distribution network allows "own use" gas demand and associated emissions to be estimated. It is assumed that NO_x, SO₂, NMVOC and PM_{2.5} emission factors remain the same as in the 2015 national inventory.

8.8 Residential and Commercial/Institutional (NFR 1A4)

8.8.1 Commercial/Institutional (NFR 1A4a)

Projected NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emissions are based on SEAI's *Baseline* and *NEEAP/NREAP* energy forecasts (2017 NEEAP and NREAP forecasts were adjusted) and are therefore estimated for both the WEM and WAM scenarios.

Oil and gas account for the majority of non-electricity energy demand in this sector. These fuels are used predominantly for space-heating purposes. Projected emissions of NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} are estimated.

It is assumed that NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emission factors remain the same as in the 2015 national inventory.

The *With Existing Measures* scenario takes into account the impact of current policies and measures in the commercial/institutional sector including:

- Public Sector Programme
- 2005 Building Regulations
- SEAI Small Business Support
- Supports for Exemplar Energy Efficient Projects (SEEEP) and Energy Efficiency Retrofit Fund (EERF)
- Accelerated Capital Allowance (ACA)
- Public Sector Building Demonstration Programme
- CHP deployment
- Renewable Heat
- Carbon tax
- Better Energy Workplaces
- Energy Supplier Obligation Scheme (non-residential)
- Better Energy Communities

The *With Additional Measures* scenario takes into account the impact of additional policies and measures in the commercial/institutional sector including:

- Increased share of renewable sources used for thermal energy
- Increased energy efficiency in buildings

8.8.2 Residential (1A4b)

Projected NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emissions are based on SEAI's *Baseline* and *NEEAP/NREAP* energy forecasts (2017 NEEAP and NREAP forecasts were adjusted) and are therefore estimated for both the WEM and WAM scenarios.

Projected emissions of NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} are estimated. It is assumed that NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} emission factors remain the same as in the 2015 national inventory.

The *With Existing Measures* scenario takes into account the impact of current policies and measures in the commercial/institutional sector including:

- 2002 Building Regulations
- 2008 Building Regulations
- 2011 Building Regulations
- Efficient Boiler Standard
- Greener Homes Scheme
- Warmer Homes Scheme
- Better Energy Homes
- Energy Supplier Obligation Scheme (residential)
- Better Energy Communities
- Carbon Tax

The *With Additional Measures* scenario takes into account the impact of additional policies and measures in the commercial/institutional sector including:

- Increased share of renewable sources used for thermal energy
- Increased energy efficiency in buildings

8.9 Combustion in Agriculture and Fishing (NFR 1A4c)

Projected emissions of NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} are estimated for the combustion of diesel in the Agriculture sector and the combustion of diesel and fuel oil in the Fishing sector. Forecast fuel use in the agriculture sector is included in the energy forecasts.

Forecasted NO_x, NMVOC, NH₃ and PM_{2.5} emission factors for mobile combustion in the agriculture sector are estimated following the Tier 2 approach outlined in the EMEP/EEA Inventory Guidebook. Emissions from the fishing sector are assumed to remain constant at the 2015 level for each projected year to 2030.

8.10 Mineral Products (NFR 2A)

Emission of air pollutants from Industrial Processes cannot usually be separated from the emissions from fuel combustion in industry. Emissions from industrial processes are therefore assumed to be included in projected estimates for the Manufacturing Industries and Construction (1.A.2) sector.

8.11 Non-Energy Products from Fuels and Non-Energy Products from Fuels and Solvent Use (NFR 2D)

Emissions projections of NMVOCs from solvent use and other products are estimated. GDP growth is used to project emissions out to 2030 with the 2015 inventory used as a starting point.

8.12 Agriculture (NFR 3)

The Agriculture sector is the largest source of NH₃ emissions in Ireland. Projected estimates of NH₃ from the Agriculture sector are undertaken using the same methodological approach as the current national inventory (Chapter 5). The methodology uses a mass flow approach based on the concept of the flow of Total Ammoniacal Nitrogen (TAN) through the manure management system.

Projected activity data (animal numbers, crop areas and fertiliser use) are provided by Teagasc (The Irish Agriculture and Food Development Authority) to the EPA in order to prepare agricultural emission projections. The activity data assumes that there is an expansion in the value of Irish agriculture over the period to 2025 to meet the targets set out in “Food Wise 2025”¹⁶ published by the Department of Agriculture, Food and the Marine in 2015.

The FAPRI-Ireland model was used for preparing agricultural forecast data. This model is linked to the FAPRI world modelling system and so takes account of and contributes to, the projections for prices obtained and quantities traded on the world markets.

The majority of the data supplied to the EPA is disaggregated at the level of that used in inventory estimates. As is currently the case in the national agricultural NH₃ inventory the use of data from a Farm Facilities Survey (Hyde et al., 2008) forms the basis of the underlying assumptions with respect to manure management practices and is used for all projected years (Chapter 5 provides further details of the manure management practices assumed). The WAM scenario includes an estimate of the savings associated with the introduction of nitrification and urease inhibitors/stabilizers in synthetic nitrogen fertilizer to meet nutrient efficiency gains in the Ireland’s Rural Development Programme 2014-2020. It is envisaged that under this measure that there will be a reduction in the requirement (as currently forecasted) for nitrogen fertilizer of 10,000 tons nitrogen in 2018, increasing linearly to 30,000 tonnes in 2020 and is maintained at that level thereafter to 2030.

Projected emissions of NMVOC and PM_{2.5} from manure management are estimated using the same approaches and methodologies as discussed in Chapter 5. The projected activity data that is utilized for NH₃ emission projections is also used to estimate projected emissions of NMVOC and PM_{2.5} for the Agriculture sector.

8.13 Waste (NFR 5)

Air pollutant emission projections in the form of NO_x, SO₂, NMVOC, NH₃ and PM_{2.5} are estimated for the waste sector. Non-methane volatile organic compounds are estimated from landfill gas production, whilst NO_x, SO₂, NH₃ and PM_{2.5} emissions are estimated from the incineration of industrial waste and from cremation.

Solid waste disposal to landfill (5A) produces significant quantities of landfill gas. Projected landfill gas production is based on greenhouse gas emission estimates for the sector undertaken by the EPA and submitted to the European Commission under Regulation 525/2013. The emission factor utilized in the national inventory of 5.65 gm⁻³ NMVOC/m³

¹⁶ Food Wise 2025. A 10-year vision for Irish agri-industry. Department of Agriculture, Food and the Marine, 2015.
<https://www.agriculture.gov.ie/foodwise2025/>

landfill gas is used in projected emission estimates. Ireland has met all Landfill Directive¹⁷ targets for diversion of biodegradable municipal waste from landfill to date.

The incineration of Industrial waste (5Cb) is now highly regulated in Ireland. There are currently only a small number of facilities based in the pharmaceutical and chemical sectors that operate incinerators for the treatment of industrial waste. It is assumed that the quantity of industrial waste incinerated and the emissions of NO_x, SO₂, NMVOC and PM_{2.5} at these facilities will remain constant at the 2015 level for each projected year to 2030.

The practice of Cremation (5C1bv) is less popular in Ireland than in other countries. However, due to the decrease in the number of burial plots available, particularly in larger cities and towns, the number of cremations in Ireland has increased. There are currently five crematoria operating in Ireland. It is assumed that the number of cremations will increase at the average of the 2013-2015 level of increase for each projected year to 2030.

¹⁷ Council Directive 1999/31/EC on the landfill of waste

Chapter Nine

Adjusted annual national emission inventories

9.1 Introduction

Ireland is a signatory to the Gothenburg Protocol but has not yet ratified it and so is not formally covered by its emission ceilings or the associated adjustment mechanism. However, Ireland had national emissions ceilings for sulphur dioxide (SO₂), nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOCs) and ammonia (NH₃) under the Directive 2001/81/EC (National Emissions Ceilings Directive) for 2010. Furthermore, article 4 of the new National Emissions Ceilings Directive (2016/2284/EU) ensures that the commitments of Directive 2001/81/EC continue to apply until 31st December 2019.

Article 21(2) of Directive (EU) 2016/2284 indicates that Member States may apply Article 5(1) of the Directive in relation to the ceilings in Annex I to Directive 2001/81/EC. Article 5(1) allows Member States to establish adjusted annual national emission inventories where non-compliance with emission ceilings or reduction commitments occur due to applying improved emission inventory methods in accordance with best science. The information provided in this chapter follows the reporting requirements of the adjustment process presented in Article 5 and Part 4 of Annex IV of the Directive (EU) 2016/2284.

Ireland's latest inventory submission estimates emissions of NO_x and NMVOC as presented in Table 9.1. For both NO_x and NMVOC national emission ceilings are exceeded for all years since 2010 and the emission ceiling for NH₃ is exceeded in 2016. Ireland considers that this has arisen due to several factors including;

- New sources that have been added to the inventory, which were either not known when the NECD (2001/81/EC) ceilings were set, or for which there was no available methodology at the time
- Emission factors and methodologies which have been improved in accordance with best practice, resulting in an increase to the estimated emissions. These revisions were not foreseen when the NECD (2001/81/EC) ceilings were set.

Table 9.1. Summary of National Emissions and the NECD Emission Ceilings

	2010	2011	2012	2013	2014	2015	2016
NO _x National Total (kt NO ₂)	113.32	101.35	104.72	105.43	104.14	104.39	107.30
NO _x NECD 2010 Ceiling (kt of NO ₂)	65.00	65.00	65.00	65.00	65.00	65.00	65.00
NMVOC (kt)	109.24	106.51	107.90	110.34	106.11	106.40	108.25
NMVOC NECD 2010 Ceiling (kt)	55.00	55.00	55.00	55.00	55.00	55.00	55.00

Ireland has undertaken a detailed assessment of the new and revised sources within the national inventory in the context of the requirements outlined in Part 4 of Annex IV in Directive (EU) 2016/2284, and is of the view that several changes to the national inventory

should be included in the adjusted annual national emission inventory. The proposed adjustments for NO_x and NMVOC are summarised in the Tables 9.2a and 9.2b below.

The resulting adjusted annual national emission inventories show Ireland to be in compliance with the NMVOC ceiling and the NO_x emission ceiling from 2011 onwards.

The justification and quantification of each adjustment application is provided in detail in the following sections and is also summarised in the excel spreadsheet (Annex VII Adjustment summary) that was submitted with the national emission inventory. The established adjusted annual emission inventory outlined here will be subject to a review by the European Commission as outlined in Article 5(6) of Directive (EU) 2016/2284, to assess whether Member States have fulfilled the relevant conditions set out in Article 5(1) and Part 4 of Annex IV. The portion of the NO_x adjustment which was agreed in 2017 and the portion to be reviewed is shown in table 9.2.a.

Without the adjustments indicated above, it is estimated that under the With Additional Measures scenario (WAM), Ireland will come into compliance with the NO_x emission ceiling in 2018, but would not reach compliance with the NMVOC ceiling before compliance requirements change to emission reduction commitments post 2020.

Table 9.2a. Summary of NO_x Adjustments

NFR	Source Sector Name	2010	2011	2012	2013	2014	2015	2016	Comments
	NO _x Adjustments (kt NO ₂)	-45.63	-42.37	-43.77	-47.31	-46.89	-46.05	-47.83	
1A3bi	Road transport - Passenger Cars	-4.80	-5.12	-6.05	-6.85	-7.27	-7.37	-8.53	<i>Current (COPERT5) - Original (COPERT2 EFs x Current Fleet)</i>
1A3bii	Road transport - Light Duty Vehicles	-1.01	-0.93	-0.82	-0.81	-0.73	-0.75	-0.97	
1A3biii	Road transport - Heavy Duty Vehicles	-2.04	-2.06	-1.80	-1.98	-1.70	-1.25	-0.82	
1A3biii	Road transport - Buses	-0.36	-0.27	-0.23	-0.25	-0.34	0.04	0.58	
1A3biv	Road transport - Mopeds & Motorcycles	-0.01	0.00	0.00	0.00	0.00	0.00	0.00	
1A3c	Rail	-0.67	-0.67	-0.64	-0.64	-0.59	-0.60	-0.61	<i>"Original EFs" confirmed as being used in 1999 submission</i>
1A3dii	National Navigation	-2.25	-1.95	-2.06	-2.02	-2.53	-2.49	-3.00	
1A4ai	Commercial Combustion (Stationary)	-1.27	-1.19	-1.17	-1.07	-0.96	-0.95	-1.00	
1A4bi	Residential	-0.46	-0.44	-0.44	-0.50	-0.39	-0.36	-0.30	
1A4ciii	National Fishing	-1.81	-1.50	-1.66	-1.85	-1.76	-1.55	-1.42	
3B1a	Manure management - Dairy cattle	-0.04	-0.04	-0.04	-0.04	-0.05	-0.05	-0.05	<i>New EF's in 2016 guidebook-these adjustments are yet to be reviewed and agreed by the EC and hence are not subtracted from the national total in Annex 1 as submitted.</i>
3B1b	Manure management - Non-dairy cattle	-0.51	-0.50	-0.54	-0.54	-0.53	-0.54	-0.56	
3B2	Manure management - Sheep	-0.05	-0.05	-0.05	-0.05	-0.05	-0.05	-0.05	
3B3	Manure management - Swine	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
3B4d	Manure management - Goats	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
3B4e	Manure management - Horses	-0.04	-0.04	-0.04	-0.04	-0.03	-0.03	-0.03	
3B4f	Manure management - Mules and asses	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
3B4gi	Manure management - Laying hens	-0.02	-0.02	-0.02	-0.02	-0.02	-0.03	-0.03	
3B4gii	Manure management - Broilers	-0.07	-0.07	-0.07	-0.06	-0.07	-0.07	-0.07	
3B4giii	Manure management - Turkeys	-0.02	-0.02	-0.03	-0.03	-0.03	-0.03	-0.03	
3B4giv	Manure management - Other poultry	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
3B4h	Manure management - Other animals (please specify in IIR)-solid	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	
3B4h	Manure management - Other animals (please specify in IIR)-slurry	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
3Da1	Inorganic N-fertilizers (includes also urea application)	-13.94	-11.38	-11.41	-13.58	-12.77	-12.69	-13.05	
3Da2a	Animal manure applied to soils	-5.23	-5.23	-5.47	-5.53	-5.54	-5.70	-5.94	
3Da2b	Sewage sludge applied to soils	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	
3Da3	Urine and dung deposited by grazing animals	-11.01	-10.85	-11.18	-11.41	-11.49	-11.55	-11.93	

NFR	Source Sector Name	2010	2011	2012	2013	2014	2015	2016	Comments
	Portion of adjustment for review (kt NO ₂)	-30.96	-28.23	-28.88	-31.34	-30.61	-30.77	-31.77	Not yet agreed
	Agreed portion of adjustment (kt NO ₂)	-14.67	-14.14	-14.89	-15.97	-16.28	-15.28	-16.07	Subtracted from Annex 1 total
	NOx National Total (kt NO ₂)	113.32	101.35	104.72	105.43	104.14	104.39	107.30	All adjustments (agreed and for review)
	NOx Adjusted Total (kt NO ₂)	67.69	58.98	60.95	58.12	57.26	58.34	59.47	
	NOx NECD Ceiling (kt NO ₂)	65.00	65.00	65.00	65.00	65.00	65.00	65.00	

Table 9.2b. Summary of NMVOC Adjustments

NFR	Source Sector Name	2010	2011	2012	2013	2014	2015	2016	Comments
	NMVOC Adjustments (kt)	-53.94	-54.74	-57.47	-59.88	-57.18	-58.54	-61.41	Agreed subtracted from Annex 1
2H2	Food and beverages industry	-12.29	-13.31	-14.69	-16.29	-13.56	-14.38	-15.52	"New" source
3B	Manure Management	-37.72	-37.52	-38.89	-39.75	-39.79	-40.36	-42.07	
3Da1	Inorganic N Fertilisers	-3.93	-3.92	-3.90	-3.85	-3.84	-3.81	-3.82	
	NMVOC National Total (kt)	109.24	106.51	107.90	110.34	106.11	106.40	108.25	
	NMVOC Adjusted Total (kt)	55.30	51.76	50.42	50.45	48.93	47.86	46.84	
	NMVOC NECD Ceiling (kt)	55.00	55.00	55.00	55.00	55.00	55.00	55.00	

9.2 Meeting the Requirements for an Adjustment

9.2.1 New Emission Sources

Part 4.1.d.i of Annex IV of Directive (EU) 2016/2284 indicates that for new emission source categories, evidence must be provided that:

1. The new source category is acknowledged in scientific literature (and/or the EMEP/EEA Guidebook).
2. The source category was not included in the relevant historic national emissions inventory when the ceilings were set.
3. The emissions from a new sources category contribute to the MS not complying with their 2010 emissions ceilings under Directive 2001/81/EC.

All of the new sources identified in this adjustment application are included in the current version of the EMEP/EEA Guidebook, and were not included in the national emissions inventory in 1999 (when the targets contained within Directive 2001/81/EC were set). Tables 9.2a and 9.2b demonstrate that the identified new sources contribute to the exceedance of the NECD emissions ceilings.

As such, it is considered that all of the criteria are met for the new sources that have been identified.

Quantification of the adjustment for each of the new sources is presented in detail in sections 9.3 and 9.4. But the following general approach has been used:

The adjustment for new source has been obtained by multiplying the emission by -1.

9.2.2 Significantly Different Emission Factors

Part 4.1.d.ii of Annex IV of Directive (EU) 2016/2284 indicates that where significantly different emission factors (EFs) are used, the following evidence is required:

1. The original EF, and information on its origin or derivation.
2. Evidence that the original EF was used in determining the emission ceilings when they were set.
3. The updated EF, and information on its origin or derivation.
4. A comparison of the original and updated EFs, demonstrating that the change contributes to a MS being in exceedance.
5. A rationale for deciding whether the changes in EF are significant.

Section 9.3 and 9.4 below present the original and updated EFs and the quantified impact on the emissions estimates of the change. The sources of the EFs are also presented, and the original EFs were all used in the emission inventory in 1999, when the emission ceilings were set.

All of the changes to emission factors that are presented contribute to moving Ireland's national total emissions into compliance. All of changes are therefore considered to be "significant" in the context of attaining compliance with emission ceilings.

In calculating adjustments for revised EFs, it is not necessary to present the current activity data (since this can be derived from the current emission and the current emission factor). In presenting information to quantify the adjustment, the following approach has been used:

$$\text{Adjustment} = (\text{EF}_{\text{Original}} \times \text{AD}_{\text{Current}}) - (\text{E}_{\text{Current}})$$

$$\text{Given that: } \text{AD}_{\text{Current}} = \text{E}_{\text{Current}} / \text{EF}_{\text{Current}}$$

The adjustment can be written:

$$\text{Adjustment} = (\text{EF}_{\text{Original}} \times \text{E}_{\text{Current}} / \text{EF}_{\text{Current}}) - \text{E}_{\text{Current}}$$

The information provided in the tables for each of the adjustments in the following sections should therefore be sufficient to allow a review of the adjustment quantification.

The final adjustment value has been determined and provided at the individual NFR category level. Where there have been revisions to emission factors for sources within an NFR category, all revisions have been included i.e. both increases and decreases to emission factors have been included. This avoids selectively including only EF revisions which would result in a favourable revision of the national emissions inventory total – a process which is not considered to be appropriate. As a result, the adjustment values that are shown for each NFR category can be considered “net” adjustment values.

9.3 NO_x Adjustment Applications

9.3.1 1A3b Road Transport (NO_x)

Justification – Significantly Different EFs

Emission factors for NO_x from road transport were included in the EMEP/CORINAIR Emissions Inventory Guidebook (version 2) in 1999 which were applicable when the emission ceilings were set. However, the EFs used in the current national emissions inventory are higher than these original EFs. Ireland considers that the current NO_x EFs for this source are significantly different, as defined by the Directive (EU) 2016/2284 and are eligible for an adjustment

Overall emissions from the different vehicle categories within road transport for the period 2010 to 2016 are;

- passenger cars (1A3bi) are now on average 90.5 per cent higher
- light duty vehicles (1A3bii) are now on average 13.3 per cent higher
- heavy duty vehicles (1A3biii) are now on average 52.6 per cent higher
- buses and coaches (1A3biii) are now on average 1.7 per cent higher
- mopeds and motorcycles (1A3biv) are now on average 12.0 per cent higher

Quantification

The adjustment quantification is explained for each mode of road transport in Tables 9.3 to 9.8.

The general approach for quantification is presented in section 9.2.2. In this case the original emission factors are those that were used in the national inventory submission in 1999 and were derived from the COPERT II model using a Tier 3 methodology. The current EFs were derived from the COPERT 5 model using a Tier 3 methodology. The 2018 adjustment application for NO_x submitted under the NECD is based on the same methodology as used for the original approved adjustment application with slight changes to IEFs due to the

revisions included in COPERT 5 in comparison to COPERT 4. The adjustment has been reviewed and approved by the EC.

Table 9.3. Adjustment Quantification for Road Transport-1A3bi Passenger Cars (NO_x)

Activity (NFR)	Activity technology	Emission Factors		Current NO _x emission (tonnes)						
		Current EF (g/km)	"Original" EF (g/km)	2010	2011	2012	2013	2014	2015	2016
1A3bi Passenger Cars	Conventional	0.543	0.571	88.44	68.72	63.00	29.22	25.93	17.83	13.25
	PC Euro 1 - 91/441/EEC	0.434	0.567	590.69	366.18	273.31	87.72	67.17	53.56	35.90
	PC Euro 2 - 94/12/EEC	0.307	0.280	3176.17	2565.15	2114.27	1849.36	1366.38	1078.99	770.17
	PC Euro 3 - 98/69/EC Stage2000	0.291	0.252	4819.44	4372.37	4152.94	3953.77	3259.35	2751.91	2477.84
	PC Euro 4 - 98/69/EC Stage2005	0.312	0.105	5156.65	4788.55	4962.35	5018.62	5256.19	4805.94	4896.06
	PC Euro 5 - EC 715/2007	0.458	0.121	0.56	1250.79	2343.45	3367.52	4258.59	4159.54	4589.82
	PC Euro 6 - EC 715/2007	0.365	0.119	0.19	0.22	0.42	0.57	0.82	997.43	2202.79
				13832.15	13411.96	13909.74	14306.79	14234.43	13865.21	14985.84
				Adjustment (tonnes)						
	Conventional			1.83	1.42	1.28	0.60	0.56	0.38	0.30
	PC Euro 1 - 91/441/EEC			136.01	88.36	66.02	25.02	20.16	16.40	12.81
	PC Euro 2 - 94/12/EEC			-427.44	-310.35	-258.40	-222.09	-102.93	-94.86	-52.58
	PC Euro 3 - 98/69/EC Stage2000			-1136.09	-872.22	-876.54	-841.81	-548.60	-370.05	-369.40
	PC Euro 4 - 98/69/EC Stage2005			-3368.78	-3100.52	-3248.45	-3316.86	-3490.25	-3184.60	-3242.48
	PC Euro 5 - EC 715/2007			-0.56	-923.31	-1733.36	-2493.63	-3148.07	-3062.03	-3388.94
	PC Euro 6 - EC 715/2007			-0.19	-0.22	-0.42	-0.57	-0.82	-673.22	-1486.66
				-4795.24	-5116.84	-6049.85	-6849.35	-7269.93	-7367.97	-8526.94

Table 9.4. Adjustment Quantification for Road Transport-1A3bii Light Duty Vehicles (NO_x)

Activity (NFR)	Activity technology	Emission Factors		Current NOx emission (tonnes)						
		Current EF (g/km)	"Original" EF (g/km)	2010	2011	2012	2013	2014	2015	2016
1A3bii Light Duty Vehicles	Conventional	1.315	1.192	80.47	57.64	32.39	22.43	18.32	10.91	8.75
	LD Euro 1 - 93/59/EEC	1.145	0.497	371.34	295.83	251.17	206.28	143.50	86.71	90.40
	LD Euro 2 - 96/69/EEC	1.145	0.298	1926.36	1734.38	1375.02	1271.05	600.41	429.77	414.35
	LD Euro 3 - 98/69/EC Stage2000	0.951	1.066	2603.56	2591.15	2263.46	2147.21	1782.66	1425.24	1253.50
	LD Euro 4 - 98/69/EC Stage2005	0.770	0.875	2344.51	2384.47	2164.30	2354.39	2501.07	2141.66	2009.40
	LD Euro 5 - 2008 Standards	1.283	0.876	0.00	355.21	719.98	1060.29	2395.87	2284.29	2392.30
	LD Euro 6	1.036	0.876	0.00	0.00	0.00	0.00	0.00	770.57	1804.28
				7326.25	7418.68	6806.33	7061.65	7441.85	7149.15	7972.98
				Adjustment (tonnes)						
	Conventional			-7.48	-5.35	-2.99	-2.08	-1.70	-1.02	-0.81
	LD Euro 1 - 93/59/EEC			-210.23	-167.49	-142.22	-116.81	-81.24	-49.07	-51.17
	LD Euro 2 - 96/69/EEC			-1424.81	-1282.87	-1017.11	-940.21	-444.14	-317.84	-306.45
	LD Euro 3 - 98/69/EC Stage2000			312.74	311.28	271.94	257.98	214.15	171.74	151.05
	LD Euro 4 - 98/69/EC Stage2005			321.70	327.21	297.02	323.12	343.17	294.70	276.49
	LD Euro 5 - 2008 Standards			0.00	-112.78	-228.61	-336.66	-760.70	-724.82	-759.10
	LD Euro 6			0.00	0.00	0.00	0.00	0.00	-119.44	-279.53
				-1008.07	-930.00	-821.96	-814.65	-730.46	-745.75	-969.52

Table 9.5. Adjustment Quantification for Road Transport-1A3biii Heavy Duty Vehicles (NO_x)

Activity (NFR)	Activity technology	Emission Factors		Current NOx emission (tonnes)						
		Current EF (g/km)	"Original" EF (g/km)	2010	2011	2012	2013	2014	2015	2016
1A3biii Heavy Duty Vehicles	Conventional	6.575	5.839	136.44	115.23	84.28	71.33	41.52	44.26	18.78
	HD Euro I - 91/542/EEC Stage I	4.222	3.787	221.98	187.44	159.00	148.95	63.69	46.08	26.45
	HD Euro II - 91/542/EEC Stage II	4.449	2.779	1500.06	1404.14	1128.29	1164.39	455.57	364.39	315.95
	HD Euro III - 2000 Standards	3.602	2.061	1861.93	1930.40	1724.23	1814.88	1625.35	1374.69	1334.16
	HD Euro IV - 2005 Standards	2.541	1.441	1473.86	1558.74	1444.10	1744.39	1965.21	1906.27	1950.36
	HD Euro V - 2008 Standards	1.436	1.468	0.00	73.11	160.36	257.50	652.03	739.32	814.74
	HD Euro VI	0.116	1.468	0.00	0.00	0.00	0.00	0.00	24.95	60.04
				5194.27	5269.04	4700.26	5201.44	4803.37	4499.95	4520.49
				Adjustment (tonnes)						
	Conventional			-29.21	-22.96	-14.00	-12.47	-5.16	-4.95	-0.08
	HD Euro I - 91/542/EEC Stage I			-22.41	-18.89	-16.10	-14.97	-6.44	-4.75	-2.63
	HD Euro II - 91/542/EEC Stage II			-556.77	-520.76	-418.60	-430.94	-169.82	-136.80	-116.55
	HD Euro III - 2000 Standards			-794.68	-824.62	-737.32	-775.66	-694.43	-587.99	-569.50
	HD Euro IV - 2005 Standards			-636.78	-673.93	-624.75	-754.12	-850.33	-825.23	-841.18
	HD Euro V - 2008 Standards			0.00	3.13	6.22	7.61	21.20	16.31	14.76
	HD Euro VI			0.00	0.00	0.00	0.00	0.00	291.04	693.98
				-2039.86	-2058.04	1804.54	1980.56	1704.98	1252.36	-821.21

Table 9.6. Adjustment Quantification for Road Transport-1A3biii Heavy Duty Vehicles, Buses (NO_x)

Activity (NFR)	Activity technology	Emission Factors		Current NOx emission (tonnes)						
		Current EF (g/km)	"Original" EF (g/km)	2010	2011	2012	2013	2014	2015	2016
1A3biii Heavy Duty Vehicles (Buses)	Conventional	10.264	9.719	357.90	283.10	229.82	240.05	363.44	65.98	20.43
	HD Euro I - 91/542/EEC Stage I	7.049	5.597	718.82	627.14	498.78	416.61	458.94	213.88	68.87
	HD Euro II - 91/542/EEC Stage II	7.742	11.593	3139.26	2777.19	2619.54	2339.33	2534.05	2418.14	2211.69
	HD Euro III - 2000 Standards	6.329	3.330	2829.71	2403.91	2250.06	2043.37	2259.48	2206.59	2240.81
	HD Euro IV - 2005 Standards	4.372	3.292	1312.09	1189.41	1169.41	1174.00	1477.30	1577.90	1663.52
	HD Euro V - 2008 Standards	3.816	3.718	84.27	132.63	261.29	352.61	401.88	428.45	546.06
	HD Euro VI	0.302	4.290	0.00	0.00	0.00	0.00	0.00	25.37	87.29
				8442.04	7413.39	7028.89	6565.96	7495.09	6936.31	6838.66
				Adjustment (tonnes)						
	Conventional			-8.01	-7.54	-6.45	-5.73	-5.44	-3.50	-5.11
	HD Euro I - 91/542/EEC Stage I			-148.61	-129.44	-102.73	-85.18	-93.42	-44.06	-12.75
	HD Euro II - 91/542/EEC Stage II			1380.80	1238.88	1195.85	1075.79	1176.56	1202.71	1102.54
					-	-		-	-	-
	HD Euro III - 2000 Standards			-1296.50	1103.24	1038.66	-942.90	1045.93	1045.66	1050.59
	HD Euro IV - 2005 Standards			-289.82	-266.34	-267.44	-271.40	-349.77	-390.03	-402.89
	HD Euro V - 2008 Standards			-0.80	-3.56	-15.12	-20.75	-19.63	-10.93	-22.27
	HD Euro VI			0.00	0.00	0.00	0.00	0.00	334.40	970.83
				-362.95	-271.24	-234.55	-250.16	-337.62	42.93	579.76

Table 9.7. Adjustment Quantification for Road Transport-1A3biv Mopeds and Motorcycles (NO_x)

Activity (NFR)	Activity technology	Emission Factors		Current NOx emission (tonnes)						
		Current EF (g/km)	"Original" EF (g/km)	2010	2011	2012	2013	2014	2015	2016
1A3biv Mopeds and Motorcycles	Conventional	0.052	0.041	0.10	0.08	0.06	0.05	0.05	0.05	0.05
	Mop - Euro I	0.184	0.116	0.49	0.40	0.34	0.30	0.32	0.35	0.38
	Mop - Euro II	0.157	0.083	0.26	0.22	0.19	0.17	0.17	0.19	0.22
	Mop - Euro III	0.164	0.087	0.32	0.30	0.29	0.27	0.33	0.40	0.53
	Conventional	0.232	0.124	10.24	8.74	7.15	6.77	7.05	7.28	7.84
	Mot - Euro I	0.356	0.221	13.15	11.80	10.50	10.12	11.61	12.96	14.75
	Mot - Euro II	0.159	0.221	3.83	3.52	3.11	3.05	3.29	3.76	4.55
	Mot - Euro III	0.097	0.221	2.81	2.92	2.79	2.92	3.66	4.57	6.17
				31.19	27.98	24.44	23.65	26.49	29.57	34.50
				Adjustment (tonnes)						
	Conventional			-0.02	-0.02	-0.01	-0.01	-0.01	-0.01	-0.01
	Mop - Euro I			-0.18	-0.15	-0.13	-0.11	-0.12	-0.13	-0.14
	Mop - Euro II			-0.12	-0.10	-0.09	-0.08	-0.08	-0.09	-0.11
	Mop - Euro III			-0.15	-0.14	-0.13	-0.13	-0.16	-0.19	-0.25
	Conventional			-4.87	-4.14	-3.36	-3.19	-3.30	-3.38	-3.58
	Mot - Euro I			-5.02	-4.50	-4.00	-3.84	-4.40	-4.91	-5.57
	Mot - Euro II			1.28	1.21	1.10	1.13	1.25	1.47	1.81
	Mot - Euro III			3.15	3.37	3.29	3.59	4.62	5.89	8.05
				-5.94	-4.47	-3.34	-2.64	-2.20	-1.36	0.19

Table 9.8. Adjustment Quantification for Road Transport-1A3b Road Transport (NO_x)

Activity (NFR)	Activity technology	Emission Factors		Current NOx emission (tonnes)						
		Current EF (g/km)	"Original" EF (g/km)	2010	2011	2012	2013	2014	2015	2016
1A3b Road Transport	1A3bi Passenger Cars			13832.15	13411.96	13909.74	14306.79	14234.43	13865.21	14985.84
	1A3bii Light Duty Vehicles			7326.25	7418.68	6806.33	7061.65	7441.85	7149.15	7972.98
	1A3biii Heavy Duty Vehicles and buses			13636.31	12682.42	11729.15	11767.40	12298.46	11436.26	11359.15
	1A3biv Mopeds and Motorcycles			31.19	27.98	24.44	23.65	26.49	29.57	34.50
				34825.90	33541.04	32469.66	33159.49	34001.23	32480.19	34352.47
				Adjustment (tonnes)						
	1A3bi Passenger Cars			-4795.24	-5116.84	-6049.85	-6849.35	-7269.93	-7367.97	-8526.94
	1A3bii Light Duty Vehicles			-1008.07	-930.00	-821.96	-814.65	-730.46	-745.75	-969.52
	1A3biii Heavy Duty Vehicles and buses			-2402.81	-2329.27	-2039.10	-2230.72	-2042.60	-1209.44	-241.45
	1A3biv Mopeds and Motorcycles			-5.94	-4.47	-3.34	-2.64	-2.20	-1.36	0.19
				-8212.05	-8380.57	-8914.25	-9897.36	-10045.20	-9324.51	-9737.72

9.3.2 1A3c Railways (NO_x)

Justification – Significantly Different EFs

Emission factors for NO_x from Railways (1A3c) were included in the EMEP/CORINAIR Emissions Inventory Guidebook (version 2) in 1999 which were applicable when the emission ceilings were set. Ireland used an EF from the “Handbook of Emission Factors, Non-Industrial Sources”, (Ministry of Health and Environmental Protection, The Netherlands, 1980) in the emission inventory when the emissions ceilings were set. However, the EF used in the current national emissions inventory is higher than this original EF. Ireland considers that the current NO_x EF for this source is significantly different, as defined by the Directive (EU) 2016/2284 and is eligible for an adjustment.

The current EF for NO_x is 49% higher than the original EF.

Quantification

The adjustment quantification is explained in Table 9.9.

The general approach for quantification is presented in section 9.2.2. The original EF for railways (1A3c) is from the “Handbook of Emission Factors, Non-Industrial Sources”, (Ministry of Health and Environmental Protection, The Netherlands, 1980). Tables 53, 55 and 57 on pages 80-81 refer to diesel locomotives for freight and passenger trains and for shunting locomotives and show an EF for NO_x of 35g/kg of fuel or 808.2 kg/TJ (NCV of Gasoil, 43.31 MJ/kg). This EF is rounded up in the original inventory to 810 kg/TJ.

Ireland now uses the EF from the Inventory Guidebook (EMEP\EEA, 2016) to replace the above referenced EF which was published in 1980. The 2018 adjustment application for NO_x submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.9. Adjustment Quantification for Railways (NO_x)

Activity (Fuel)	Emission Factors		Current NO _x emissions (tonnes)						
	Current EF (kg/TJ)	"Original" EF (kg/TJ)	2010	2011	2012	2013	2014	2015	2016
Gasoil/ Diesel	1,209.93	810	2,013.05	2,034.85	1,948.41	1,940.35	1,780.01	1,814.06	1,847.51
Gasoil/ Diesel			Adjustments (tonnes)						
			-665.39	-672.60	-644.03	-641.36	-588.36	-599.62	-610.68
Railways			Total Adjustment (ktonnes)						
			-0.67	-0.67	-0.64	-0.64	-0.59	-0.60	-0.61

9.3.3 1A3dii National Navigation (NO_x)

Justification – Significantly Different EFs

Emission factors for NO_x from National Navigation (1A3dii) were included in the EMEP/CORINAIR Emissions Inventory Guidebook (version 2) in 1999 when the emission ceilings were set. Ireland used an EF for shipping from the “Handbook of Emission Factors, Non-Industrial Sources”, (Ministry of Health and Environmental Protection, The Netherlands, 1980) in the emission inventory when the emissions ceilings were set. However, the EF used in the current national emission inventory is higher than this original EF. Ireland considers

that the current NO_x EF for this source is significantly different, as defined by the Directive (EU) 2016/2284 and is eligible for an adjustment.

The current EF for NO_x is 85% higher than the original EF.

Quantification

The adjustment quantification is explained in the Table 9.10.

The general approach for quantification is presented in section 9.2.2. The original EF for shipping/navigation (1A3dii) is from the “Handbook of Emission Factors, Non-Industrial Sources”, (Ministry of Health and Environmental Protection, The Netherlands, 1980). Table 66 on page 91 refer to sea-going vessels and show an EF for NO_x ranging from 35g/kg of fuel to 44 g/kg fuel depending on the percentage power in use. This would equate to an EF for NO_x between 808.2 kg/TJ to 1015.9 kg/TJ (NCV of Gasoil, 43.31 MJ/kg).

Ireland now uses the EF from the Inventory Guidebook (EMEP\EEA, 2016) to replace the above referenced EF which was published in 1980. The 2018 adjustment application for NO_x submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.10. Adjustment Quantification for National Navigation (NO_x)

Activity (Fuel)	Emission Factors		Current NO _x emissions (tonnes)						
	Current EF (kg/TJ)	"Original" EF (kg/TJ)	2010	2011	2012	2013	2014	2015	2016
Gasoil / Diesel	1,812.59	980	4,897.08	4,251.32	4,492.80	4,394.62	5,501.37	5,426.06	6,520.50
			Adjustments (tonnes)						
Gasoil/Diesel			-2249.40	-1952.79	-2063.70	-2018.61	-2526.98	-2492.39	-2995.10
			Total Adjustment (ktonnes)						
National Navigation			-2.25	-1.95	-2.06	-2.02	-2.53	-2.49	-3.00

9.3.4 1A4ai Commercial/Institutional Stationary Combustion (NO_x)

Justification – Significantly Different EFs

Emission factors for NO_x from Commercial/Institutional stationary combustion (1A4ai) are included in the EMEP/CORINAIR Emissions Inventory Guidebook (version 2) in 1999 when the emission ceilings were set. Ireland used EFs for combustion in commercial from Table 3.4 page 16 of “Corinair 1990 Emission Inventory for Ireland” report published by the Environmental Research Unit in July 1993. These EFs can also be referenced as Table 2.2 of “Corinair Technical annexes Volume 2, Default emission factors handbook” published by the European Commission in 1994. However, the EFs used in the current national emission inventory are higher than these original EFs. Ireland considers that the current NO_x EFs for these sources are significantly different, as defined by the Directive (EU) 2016/2284 and are eligible for an adjustment.

The current EFs for each fuel for this source;

- Fuel oil is 17% lower
- LPG is 48% higher
- Gasoil is 100% higher
- Natural Gas is 48% higher

- Biomass is 82% higher
- Biogas is 48% higher

than the original EFs.

Quantification

The adjustment quantification is explained in Table 9.11.

The general approach for quantification is presented in section 9.2.2. The original EFs for combustion in commercial (1A4ai) are from Table 3.4 page 16 of “Corinair 1990 Emission Inventory for Ireland” report published by the Environmental Research Unit in July 1993. These EFs can also be referenced in Table 2.2 of “Corinair Technical annexes Volume 2, Default emission factors handbook” published by the European Commission in 1994. There are additional older references also available if needed.

Ireland now uses the EFs from the Inventory Guidebook (EMEP/EEA, 2016) to replace the above referenced EFs which were published in 1987, 1993 and 1994. The 2018 adjustment application for NO_x submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.11. Adjustment Quantification for Commercial/Institutional Stationary Combustion (NO_x)

Activity (Fuel)	Emission Factors		Current NO _x emissions (tonnes)						
	Current EF (kg/TJ)	"Original" EF (kg/TJ)	2010	2011	2012	2013	2014	2015	2016
Fuel Oil	100	120	41.24	41.24	41.24	41.24	41.24	41.24	41.24
LPG	74	50	24.05	22.93	22.12	26.79	24.07	24.43	26.17
Gasoil / Diesel	100	50	1,575.75	1,547.39	1,436.99	1,195.48	978.94	944.54	957.98
Natural Gas	74	50	1,420.62	1,178.29	1,295.54	1,306.92	1,286.16	1,382.03	1,450.59
Biomass	91	50	45.74	60.04	72.82	95.58	106.01	63.59	98.79
Biogas	74	50	11.87	13.08	12.12	13.78	16.03	16.92	21.48
			Adjustments (tonnes)						
Fuel Oil			8.25	8.25	8.25	8.25	8.25	8.25	8.25
LPG			-7.80	-7.44	-7.17	-8.69	-7.81	-7.92	-8.49
Gasoil / Diesel			-787.88	-773.70	-718.50	-597.74	-489.47	-472.27	-478.99
Natural Gas			-460.74	-382.15	-420.17	-423.86	-417.13	-448.23	-470.46
Biomass			-20.61	-27.05	-32.81	-43.06	-47.76	-28.65	-44.51
Biogas			-3.85	-4.24	-3.93	-4.47	-5.20	-5.49	-6.97
			Total Adjustment (ktonnes)						
Comm/Instit Stationary Combustion			-1.27	-1.19	-1.17	-1.07	-0.96	-0.95	-1.00

9.3.5 1A4bi Residential Stationary Combustion (NO_x)

Justification – Significantly Different EFs

Emission factors for NO_x from Residential stationary combustion (1A4bi) were included in the EMEP/CORINAIR Emissions Inventory Guidebook (version 2) in 1999 when the emission ceilings were set. Ireland used EFs for combustion in residential from Table 3.4 page 16 of “Corinair 1990 Emission Inventory for Ireland” report published by the Environmental Research Unit in July 1993. These EFs can also be referenced as Table 2.2 of “Corinair Technical annexes Volume 2, Default emission factors handbook” published by

the European Commission in 1994. However, the EFs used in the current national emission inventory are generally higher than these original EFs. Ireland considers that the current NO_x EFs for these sources are significantly different, as defined by the Directive (EU) 2016/2284 and are eligible for an adjustment.

The current EFs for each fuel for this source;

- Bituminous coal and Anthracite are 120% higher
- Lignite, sod peat and peat briquettes are 10% higher
- Kerosene and Gasoil are 2% higher
- Petroleum coke is Natural Gas is 49% lower
- Natural Gas and LPG are 16% lower
- Biomass is 20% lower

than the original EFs.

Quantification

The adjustment quantification is explained in Table 9.12.

The general approach for quantification is presented in section 9.2.2. The original EFs for stationary combustion (1A4bi) in the residential sector are from Table 3.4 page 16 of “Corinair 1990 Emission Inventory for Ireland” report published by the Environmental Research Unit in July 1993. These EFs can also be referenced in Table 2.2 of “Corinair Technical annexes Volume 2, Default emission factors handbook” published by the European Commission in 1994. There are additional older references also available if needed.

Ireland now uses the EFs from the Inventory Guidebook (EMEP/EEA, 2016) to replace the above referenced EFs which were published in 1987, 1993 and 1994, respectively. The 2018 adjustment application for NO_x submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.12. Adjustment Quantification for Residential Stationary Combustion (NO_x)

Activity (Fuel)	Emission Factors		Current NO _x emissions (tonnes)						
	Current EF (kg/TJ)	"Original" EF (kg/TJ)	2010	2011	2012	2013	2014	2015	2016
Bituminous Coal	110.00	50.00	816.80	698.46	759.37	796.71	612.03	578.96	463.91
Anthracite ¹	110.00	50.00	306.56	310.39	300.19	381.04	338.19	312.50	315.74
Lignite	110.00	100.00	47.81	48.38	54.87	79.68	58.36	58.22	43.73
Sod Peat	110.00	100.00	761.99	749.14	588.14	588.14	588.14	588.14	588.14
Briquettes	110.00	100.00	405.68	362.01	399.88	415.41	332.81	335.73	318.62
Kerosene	51.00	50.00	2,157.51	1,705.30	1,457.41	1,507.66	1,428.64	1,653.96	1,740.79
LPG	42.00	50.00	65.67	60.27	58.13	70.41	63.25	64.20	68.78
Gasoil / Diesel	51.00	50.00	431.93	413.83	394.34	341.96	306.40	295.18	309.95
Petroleum Coke	51.00	100.00	28.36	18.49	21.14	23.36	18.07	13.89	11.75
Natural Gas	42.00	50.00	1,248.03	1,001.04	1,055.94	1,066.00	941.96	976.14	989.99
Biomass ²	50.00	100.00	29.23	27.78	29.81	32.08	27.32	26.42	23.72
			6,299.56	5,395.10	5,119.22	5,302.44	4,715.17	4,903.34	4,875.11
			Adjustments (tonnes)						
Bituminous Coal			-445.53	-380.98	-414.20	-434.57	-333.83	-315.80	-253.04
Anthracite ¹			-167.22	-169.30	-163.74	-207.84	-184.47	-170.46	-172.22
Lignite			-4.35	-4.40	-4.99	-7.24	-5.31	-5.29	-3.98
Sod Peat			-69.27	-68.10	-53.47	-53.47	-53.47	-53.47	-53.47
Briquettes			-36.88	-32.91	-36.35	-37.76	-30.26	-30.52	-28.97
Kerosene			-42.30	-33.44	-28.58	-29.56	-28.01	-32.43	-34.13
LPG			12.51	11.48	11.07	13.41	12.05	12.23	13.10
Gasoil / Diesel			-8.47	-8.11	-7.73	-6.71	-6.01	-5.79	-6.08
Petroleum Coke			27.25	17.77	20.31	22.44	17.36	13.34	11.29
Natural Gas			237.72	190.68	201.13	203.05	179.42	185.93	188.57
Biomass ²			39.92	35.50	41.00	43.03	37.87	42.95	40.94
			-456.614	-441.819	-435.548	-495.219	-394.654	-359.301	-297.989
			Total Adjustment (ktonnes)						
Residential Stationary Combustion			-0.46	-0.44	-0.44	-0.50	-0.39	-0.36	-0.30

¹ Includes Manufactured Ovoids, ² Biomass has a T2 method applied except to residential stoves and fireplaces for nontraded wood with EF 80kg/TJ

9.3.6 1A4ciii National Fishing (NO_x)

Justification – A New Source/Significantly Different EFs

Activity data to estimate emission of NO_x from National Fishing (1A4ciii) did not exist in Ireland's national energy balance until 2012, when they were first introduced for the 1990-2010 energy balance. Marine diesel used in National Fishing is now included in the energy balance for all years from 1990 to 2016.

For a new source, Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria are met, and it is therefore concluded that NO_x emissions from national fishing (1A4ciii) are a new source and are eligible for an adjustment.

The fuel allocated to National Fishing (1A4ciii) since 2012 did not increase the overall gasoil/diesel use in the energy balance estimates, as the fuel was re-allocated from the category Commercial/Institutional (1A4ai) stationary combustion. So while National Fishing is in effect a new source introduced in 2012, to avoid double counting it is more appropriate that an adjustment is considered on the basis of significantly different EFs.

Quantification

The adjustment quantification is explained in the table 9.13 below.

The general approach for quantification is presented in section 9.2.2. Ireland used EFs for stationary combustion in the commercial/Institutional category (1A4ai) from Table 3.4 page 16 of "Corinair 1990 Emission Inventory for Ireland" report published by the Environmental Research Unit in July 1993. These EFs can also be referenced in Table 2.2 of "Corinair Technical annexes Volume 2, Default emission factors handbook" published by the European Commission in 1994.

Ireland now uses an EF for NO_x for National Fishing from the 2016 version of the EMEP/EEA Emissions Inventory Guidebook and is used to estimate emissions of NO_x for all years from 1990 to 2016 in the current inventory submission.

The current EF for gasoil for this source is 3525% higher than the original EF. The 2018 adjustment application for NO_x submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.13. Adjustment Quantification for National Fishing (NO_x)

Activity (Fuel)	Emission Factors		Current NO _x emissions (tonnes)						
	Current EF (kg/TJ)	"Original" EF (kg/TJ)	2010	2011	2012	2013	2014	2015	2016
Gasoil/Diesel	1,812.59	50.00	1864.20	1544.13	1709.37	1902.07	1812.30	1593.33	1463.08
Adjustments (tonnes)									
Gasoil/Diesel			-	-	-	-	-	-	-
			1812.78	1501.54	1662.22	1849.60	1762.30	1549.38	1422.72
Total Adjustment (ktonnes)									
National Fishing			-1.86	-1.54	-1.71	-1.90	-1.81	-1.59	-1.46

9.3.7 3B1a-3B4h Manure Management (NO_x)

Justification – New emission source

Emissions of NO_x from Manure Management were not included in Ireland's national emissions inventory in 1999, when the ceilings were set. This sector is to be considered as a new source compared to when the emission ceilings were set, since default methodologies and emission factors were not available in the EMEP/EEA Guidebook until the 2009 edition. A methodology is presented in the 2016 EMEP/EEA Guidebook, and this is currently used to estimate emissions that are included in the Irish national emissions inventory.

For a new source, the Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria are met, and it is therefore concluded that NO_x emissions from manure management are a new source and are eligible for an adjustment.

Quantification

Emissions have been calculated by using the Tier 2 mass-flow methodology for NH₃ and the default emission factors presented in the Inventory Guidebook (EMEP/EEA, 2016), Chapter 3. B Manure Management Table 3-10. As a new source, quantification of the adjustment is achieved by subtracting the emissions from the national inventory total as shown in Table 9.14.

Table 9.14. Adjustments for Manure Management (NO_x)

Source Sector	Adjustments (ktonnes)						
	2010	2011	2012	2013	2014	2015	2016
Manure Management – 3B1a Dairy cattle	-0.041	-0.042	-0.043	-0.044	-0.045	-0.049	-0.052
Manure Management – 3B1b Non-dairy cattle	-0.512	-0.505	-0.541	-0.543	-0.530	-0.540	-0.558
Manure Management – 3B2 Sheep	-0.047	-0.048	-0.050	-0.050	-0.049	-0.049	-0.049
Manure Management – 3B3 Swine	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002
Manure Management – 3B4d Goats	-0.002	-0.002	-0.001	-0.001	-0.001	-0.002	-0.001
Manure Management – 3B4e Horses	-0.039	-0.039	-0.041	-0.037	-0.035	-0.034	-0.034
Manure Management – 3B4f Mules and asses	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002
Manure Management - 3B4gi Laying hens	-0.018	-0.018	-0.021	-0.023	-0.024	-0.027	-0.027
Manure Management - 3B4gii Broilers	-0.070	-0.068	-0.068	-0.063	-0.071	-0.072	-0.072
Manure Management - 3B4giii Turkeys	-0.020	-0.025	-0.028	-0.026	-0.027	-0.028	-0.029
Manure Management - 3B4giv Other poultry	-0.004	-0.004	-0.004	-0.004	-0.004	-0.004	-0.004
Manure Management – 3B4h Other animals (solid)	-0.012	-0.006	-0.006	-0.006	-0.006	-0.006	-0.006
Manure Management – 3B4h Other animals (slurry)	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total Adjustment – 3B Manure Management	-0.768	-0.760	-0.808	-0.802	-0.798	-0.815	-0.837

9.3.8 3D Fertilisers applied to soils (NO_x)

Justification – New Emission source

Emissions from fertiliser applied to soils were not included in Ireland's national emissions inventory in 1999, when the ceilings were set. This sector is to be considered as a new source compared to when the emission ceilings were set. A methodology is presented in the 2016 EMEP/EEA Guidebook, and this is currently used to estimate emissions that are included in the Irish national emissions inventory.

For a new source, the Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was

not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria are met, and it is therefore concluded that NO_x emissions from fertiliser applied to soils are a new source and are eligible for an adjustment.

Quantification

Emissions for 3Da1 (Inorganic N-fertilizers including urea application), 3D2a (Animal manure applied to soils), 3Da2b (Sewage sludge applied to soils) and 3Da3 (Urine and Dung deposited by grazing animals) have been calculated by using the Tier 1 methodology and emission factors presented in the Inventory Guidebook (EMEP/EEA, 2016), Chapter 3. D Crop production and agricultural soils Table 3-1. as shown in table 9.15.

Table 9.15. Adjustments for Fertilizer applied to soils (NO_x)

Source Sector	Adjustments (ktonnes)						
	2010	2011	2012	2013	2014	2015	2016
Inorganic N-fertilizers (includes also urea application)-3Da1	-13.943	-11.381	-11.409	-13.583	-12.765	-12.695	-13.047
Animal manure applied to soils 3Da2a	-5.232	-5.234	-5.472	-5.531	-5.542	-5.695	-5.943
Sewage sludge applied to soils 3Da2b	-0.009	-0.009	-0.009	-0.009	-0.009	-0.009	-0.010
Urine and dung deposited by grazing animals -3Da3	-11.009	-10.850	-11.183	-11.414	-11.494	-11.554	-11.929
Total Adjustment – 3D Fertiliser application	-30.193	-27.474	-28.073	-30.538	-29.811	-29.954	-30.929

9.4 NMVOC Adjustment Applications

9.4.1 Food and Beverage Industry, 2H2: Spirits (NMVOC)

Justification - A New Source

NMVOC emissions from the food and beverage industry were not included in Ireland's national emissions inventory in 1999, when the ceilings were set. A methodology is presented in the 2016 EMEP/EEA Guidebook, and this is currently used to estimate emissions that are included in the Irish national emissions inventory.

For a new source, the Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria were not met, however paragraph 2 bis (a) (i) of Decision 2012/12 (as amended by ECE.EB.AIR/127/Add.1 (2014/1)), indicates that a new source is eligible for an adjustment for a new source if a methodology existed in the relevant version of the EMEP/EEA Guidebook, but the "Party can demonstrate that it was unable to apply this methodology due to a lack of relevant national statistical data...". The national data for spirit manufacture was confidential and was therefore not available when the ceilings were set. It was therefore concluded that NMVOC emissions from the food and beverage industry are a new source and are eligible for an adjustment.

Quantification

Emissions have been calculated by using the Tier 2 methodologies presented in the Inventory Guidebook (EMEP/EEA, 2016), with EFs taken from Chapter 2H2, Table 3-28

As a new source, quantification of the adjustment is achieved by subtracting the emissions from the national inventory total as shown in Table 9.16. The 2018 adjustment application for

NMVOC submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.16. Adjustments for emissions from the Food and Beverage Industry (NMVOC)

Source Sector	Adjustments (ktonnes)						
	2010	2011	2012	2013	2014	2015	2016
Food and Beverages industry: Spirits	-12.29	-13.31	-14.69	-16.29	-13.56	-14.38	-15.52

9.4.2 Manure Management, 3B (NMVOC)

Justification - A New Source

NMVOC emissions from manure management were not included in Ireland's national emissions inventory in 1999, when the ceilings were set. Emission factors and methodological approaches were included for the first time in the 2013 edition of the Inventory Guidebook. A methodology is presented in the 2016 EMEP/EEA Guidebook, and this is currently used to estimate emissions that are included in the Irish national emissions inventory.

For a new source, the Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria are met, and it is therefore concluded that NMVOC emissions from manure management are a new source and are eligible for an adjustment.

Quantification

Emissions have been calculated by using the methodology presented in the Inventory Guidebook (EMEP/EEA, 2016), Chapter 3B, Section 3.4 using a Tier 2 approach. Estimates are undertaken for each livestock type. Animal population data are combined with data on time spent in housing or at pasture combined with EFs from the Inventory Guidebook (EMEP/EEA, 2016) to give NMVOC emission estimates from housing, storage and application for each livestock type.

As a new source, quantification of the adjustment is achieved by subtracting the emissions from the national inventory total as shown in Table 9.17. The 2018 adjustment application for NMVOC submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.17. Adjustments from Manure Management (NMVOC)

Source Sector	Adjustments (ktonnes)						
	2010	2011	2012	2013	2014	2015	2016
Manure Management – 3B1a Dairy cattle	-7.16	-7.43	-7.48	-7.69	-8.05	-8.80	-9.30
Manure Management – 3B1b Non-dairy cattle	-25.04	-24.46	-25.59	-26.41	-25.87	-25.64	-26.74
Manure Management – 3B2 Sheep	-0.32	-0.33	-0.36	-0.36	-0.37	-0.36	-0.36
Manure Management – 3B3 Swine	-2.43	-2.47	-2.44	-2.39	-2.43	-2.39	-2.48
Manure Management – 3B4d Goats	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Manure Management – 3B4e Horses	-0.20	-0.20	-0.20	-0.19	-0.18	-0.17	-0.17
Manure Management – 3B4f Mules and asses	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01
Manure Management - 3B4gi Laying hens	-0.35	-0.35	-0.43	-0.47	-0.48	-0.54	-0.55
Manure Management - 3B4gii Broilers	-1.29	-1.24	-1.24	-1.16	-1.31	-1.32	-1.33
Manure Management - 3B4giii Turkeys	-0.43	-0.53	-0.60	-0.55	-0.58	-0.60	-0.62
Manure Management - 3B4giv Other poultry	-0.14	-0.14	-0.14	-0.13	-0.13	-0.13	-0.13
Manure Management – 3B4h Other animals	-0.36	-0.36	-0.38	-0.38	-0.38	-0.38	-0.38

Total Adjustment – 3B Manure Management	-37.72	-37.52	-38.89	-39.75	-39.79	-40.36	-42.07
--	---------------	---------------	---------------	---------------	---------------	---------------	---------------

9.4.3 Inorganic N Fertilisers, 3D1a (NMVOC)

Justification - A New Source

NMVOC emissions from inorganic N fertilisers were not included in Ireland's national emissions inventory in 1999, when the ceilings were set. A methodology is presented in the 2016 EMEP/EEA Guidebook, and this is currently used to estimate emissions that are included in the Irish national emissions inventory.

For a new source, the Directive (EU) 2016/2284 requires that the source is currently included in scientific literature (such as the EMEP/EEA Guidebook), and that the source was not included in the historic national emissions inventory when emission ceilings were set. Both of these criteria are met, and it is therefore concluded that NMVOC emissions from manure management are a new source and are eligible for an adjustment.

Quantification

Emissions have been calculated by using the Tier 1 methodology presented in the Inventory Guidebook (EMEP/EEA, 2016), Chapter 3D, Section 3.2.

As a new source, quantification of the adjustment is achieved by subtracting the emissions from the national inventory total as shown in Table 9.18. The 2018 adjustment application for NMVOC submitted under the NECD is based on the same methodology as used for the original approved adjustment application which has been reviewed and approved by the EC.

Table 9.18. Adjustment from Inorganic N Fertilisers (NMVOC)

Source Sector	Adjustments (ktonnes)						
	2010	2011	2012	2013	2014	2015	2016
Inorganic N Fertilisers	-3.93	-3.92	-3.90	-3.85	-3.84	-3.81	-3.82

9.5 Impact of the flexibility on compliance

The established adjusted annual national emission inventories as allowed under the flexibility in Article 5(1) of Directive (EU) 2016/2284 show Ireland to be in compliance with the NMVOC ceiling and the NO_x emission ceiling from 2011 onwards.

The adjusted national inventories for NO_x and NMVOC are presented in Figures 9.1 and 9.2




Figure 9.1 Emission Trend for NO_x 1990–2016 (Adjusted Article 5(1))




Figure 9.2 Emission Trend for NMVOC 1990–2016 (Adjusted Article 5(1))

References

- ABFI, 2013 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2013, [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Home/\\$File/Beer+Market+Report+2013.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Home/$File/Beer+Market+Report+2013.pdf)
- ABFI, 2014 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2014, [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesABFI/Media~Newsroom~the-grape-depression-tax-take-on-a-standard-bottle-of-wine-is-now-over-50/\\$File/Irish+Beer+Market-2014.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesABFI/Media~Newsroom~the-grape-depression-tax-take-on-a-standard-bottle-of-wine-is-now-over-50/$File/Irish+Beer+Market-2014.pdf)
- ABFI, 2015 Alcohol Beverage Federation of Ireland. Irish Beer Market Report 2015, , [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Publications~2015-beer-market-report/\\$File/IBA+Report-2015+web+version.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesBeer/Publications~2015-beer-market-report/$File/IBA+Report-2015+web+version.pdf)
- AEA, 2012. Provision of Technical Assistance to support the National Inventory Agency of the Office of Climate, Licensing, Research and Resource Use in the EPA. Mapping Methods for Gridded Data (AEA/R3274/ED57379).
- AEA/CTC, 2008. Inventories of Persistent Organic Pollutants in Ireland 1990 and 1995-2006. A report produced for Ireland's Environmental Protection Agency. 152 pp.
- Barry S. and O'Regan, B. (2014). FS-2 Activity data for emissions of Non-methane VOC Fellowship 2006-2012. State of Knowledge Report. Submitted to the Environmental Protection Agency, Johnstown Castle, Co. Wexford.
- Basu, M.K., Wilson, H.J. and Krishnan, G., 1991. Mercury risk from teeth. Nature 349, 109.
- Benezon, N., 1999. HCB emissions for Ontario 1988, 1998 and 2000. Draft report for Environment Canada, Canada.
- BiPRO (Beratungsgesellschaft für integrierte Problemlösungen), 2005. Study to facilitate the implementation of certain waste related provisions of the Regulation on Persistent Organic Pollutants (POPs). BiPRO GmbH, Munich, Germany. European Commission, Brussels. http://ec.europa.eu/environment/waste/studies/pdf/pops_waste_full_report.pdf
- Carey, P. et al., 1996. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- CEPMEIP, 2001. CEPMEIP emission factors for particulate matter. Berdowski, J., Visschedijk, Creemers, E. and Pulles, T.
- CEU (Council of the European Union), 1999. Council Directive 1999/13/EC of 11 March on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations. OJ L85, 29 March 1999.
- Collins, C., Meaney, B., Nolan, K., Maher, H. and Murphy, D. 2004a. National Waste Database Interim Report 2002. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Collins, C., Meaney, B., Nolan, K., Maher, H. and Corish, C. 2004b. National Waste Database Interim Report 2003. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Collins, C., Le Bolloch, O. and Meaney, B. 2005. National Waste Report 2004. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Corinair, Technical annexes Volume 2, Default emission factors handbook. European Commission 1994.

Crowe, M. et al., 2000. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

CTC/AEA, 2005. NMVOC Inventory for Ireland (SNAP Sector 6 Solvent and Other Product Use). Report to the Environmental Protection Agency

DAF, 2003. Pesticide Usage Survey (No. 1) Grassland and Fodder Crops. Department of Agriculture and Food, Dublin, Ireland.

DAF, 2004. Pesticide Usage Survey (No. 2) Arable Crops. Department of Agriculture and Food, Dublin, Ireland.

DAFM, 2015. Explanatory Handbook for cross compliance requirements. <https://www.agriculture.gov.ie/media/migration/farmingschemesandpayments/crosscompliance/CrossComplianceHandbook130916.pdf>

Dammingen, U, Luttich, M., Haenel, H-D, Dohler, H., Eurich-Menden, B and Osterburg, B. (2007). Calculations of emissions from German agriculture – National Emission Inventory Report 2008 for 2006.

Directive 2004/42/CE of the European Parliament and of the Council of 21 April 2004 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC

DOE (Department of the Environment), 1992. The Environmental Protection Agency Act, 1992. S.I. No. 7 of 1992. Department of the Environment, Custom House, Dublin, Ireland.

DTTAS (Department of Transport, Tourism and Sport), 2010. Irish Bulletin of Vehicle and Driver Statistics 2010.

DTTAS (Department of Transport, Tourism and Sport) 2016, Irish Bulletin of Vehicle and Driver Statistics 2015, http://www.dttas.ie/publications?field_sector_tid=4

Duffy, P., Black, K., O'Brien, P., Hyde, B., Ryan, A.M., Ponzi, J. and Alam, S.. 2017. Ireland National Inventory Report 2017. Greenhouse Gas Emissions 1990-2015 Reported to the United Nations Framework Convention on Climate Change.

Dyke, P.H., Foan, C., Wenbom, M. and Coleman, P.I., 1997. A review of Dioxin releases to land and water in UK. Science of the Total Environment 127, 119-131.

EAPA, European Asphalt Pavement Association (2001). Asphalt in Figures 2001

EAPA, European Asphalt Pavement Association (2007). Asphalt in Figures 2007

EAPA, European Asphalt Pavement Association (2012). Asphalt in Figures 2012

EAPA, European Asphalt Pavement Association (2016). Asphalt in Figures 2016

EMEP/CORINAIR (2000). Atmospheric Emission Inventory Guidebook – Second edition 1999

EMEP/CORINAIR (2006). Emission Inventory Guidebook

EMEP/EEA (2009). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 9, 2009.

EMEP/EEA (2013). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 10. 2013.

EMEP/EEA (2016). Air Pollutant Emission Inventory Guidebook. EEA technical Report No. 21. 2016.

EP and CEU (European Parliament and Council of the European Union), 2001. Directive 2001/81/EC of 23 October 2001 on national emissions ceilings for certain atmospheric pollutants. O.J. L309, 27 November 2001.

EP and CEU (European Parliament and Council of the European Union), 2003. Directive 2003/87/EC establishing a scheme for greenhouse gas emission allowance trading within the Community and amending Council Directive 96/61/EC. O.J. L275, 25 October 2003.

EP and CEU (European Parliament and Council of the European Union), 2004a. Decision 280/2004/EC of the European Parliament and of the Council concerning a mechanism for monitoring Community greenhouse gas emissions and for implementing the Kyoto Protocol. O.J. L49, 19 February 2004.

EP and CEU (European Parliament and Council of the European Union), 2004b. Directive 2004/42/CE on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC. O.J. L143, 30 April 2004.

ERU Corinair 1990 Emissions Inventory for Ireland, July 1993.

Feeney, B. 2015. Craft Beer and Microbreweries in Ireland, 2016, A Report for the Independent Craft Brewers of Ireland and Bord Bia,
<http://www.bordbia.ie/industry/manufacturers/insight/publications/bbreports/RecentMarketingReports/Craft%20Beer%20and%20Microbreweries%20in%20Ireland%202016.pdf>

Finn, J. et al., 2001. Estimation of Emissions of Non-Methane Volatile Organic Compounds from SNAP Sector 6: Solvent and Other Product Use for Ireland in 1998. Prepared for the Environmental Protection Agency.

Gkatzoflias, D., Kourdis, C, Ntziachristos, L. and Samaras, Z., 2012. COPERT 4. Computer programme to calculate emissions from road transport. Laboratory of Applied Thermodynamics, Mechanical Engineering Department, Aristotle University of Thessaloniki, Greece.
<http://emisia.com/products/copert/documentation>

Hyde, B.P., Carton, O.T., O'Toole, P. and Misselbrook, T., 2003. A new inventory of ammonia emissions from Irish Agriculture. *Atmospheric Environment* 37, 55-62.

Hyde, B., Carton, O.T. and Murphy, W.E., 2008. Farm Facilities Survey – Ireland 2003. Report prepared for the Department of Agriculture by Teagasc, Johnstown Castle, Wexford, Ireland. 150 p.

IPCC (Intergovernmental Panel on Climate Change), 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories.

Irish Spirits Association (ISA) 2015, ISA homepage of the website 2017, <http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesSpirits/Home!OpenDocument>

Irish Whiskey Association 2015, Vision for Irish whiskey 2015, [http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesSpirits/Home/\\$File/Vision+for+Irish+Whiskey+May+2015.pdf](http://www.abfi.ie/Sectors/ABFI/ABFI.nsf/vPagesSpirits/Home/$File/Vision+for+Irish+Whiskey+May+2015.pdf)

Kirchmann, H., and Winter, E. 1989. Ammonia volatilisation during aerobic and anaerobic manure decomposition. *Plant and Soil* 115. Pp. 35-41.

Le Bolloch, O., Meaney, B., Cope, J. and Doyle, M., 2006. National Waste Report 2005: Data Update. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Le Bolloch, O., Cope, J., Meaney, B. and Kurz, I., 2007. National Waste Report 2006. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Le Bolloch, O., Cope, J., Kurz, I., Meaney, B. and Higgins, T., 2009. National Waste Report 2007. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Lorenz, W., Wichmann, H. and Bahadir, M., 1996. Bilanzierung der Freisetzung von polychlorierten Dibenzo-p-dioxinen und Dibenzofuranen bei Brandunfällen – ein Diskussionsbeitrag, *Gefahrstoffe – Reinhaltung der Luft* 56.

McCoole, F., Derham, J., Kurz, I. and Higgins, T. 2009. National Waste Report 2008. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

McCoole, F., Derham, J., Kurz, I. and McDonagh, M. 2011. National Waste Report 2009. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

McCoole, F., Kurz, I., McDonagh, M., Derham, J., O'Neill, D. 2012. National Waste Report 2010. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

McCoole, F., Kurz, I., McDonagh, M., Derham, J., O'Neill, D. 2013. National Waste Report 201. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Meaney, B., Collins, C., Nolan, K., Cahill, E., Delaney, J., Murray, B., Healy, J. and Carty, G. 2003. National Waste Database Report 2001, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Ministry of Health and Environmental Protection, The Netherlands. Handbook of Emission Factors, Non-Industrial Sources, 1980.

Misselbrook, T.H., Pain, B.F., Headon, D.M., 1998. Estimates of ammonia emission from dairy cow collecting yards. *Journal of Agricultural Engineering Research* 71, 127-135.

Misselbrook, T.H., Chadwick, D.R., Chambers, B.J., Smith, K., Webb, J., Demmers, T. & Sneath, R.W. (2004). Inventory of Ammonia Emissions from UK Agriculture – 2003. Annual project report to Defra for contract AM0127

Misselbrook, T.H., Chadwick, D.R., Gilhespy, S.L., Chambers, B.J., Smith, K.A., Williams, J and Dragosits, U. 2010. Inventory of ammonia emissions from UK agriculture 2009. North Wyke Research, 34 pp. (Unpublished)

Misselbrook, T.H., Gilhespy, S.L., Cardenas, L.M., Williams, J. and Dragostis, U. (2016). Inventory of Ammonia Emissions from UK Agriculture 2015. Rothamsted Research, North Wyke, 36 pp (unpublished).

Netcen/CTC, 2006. Ireland heavy Metal Inventories 1990, 1995 to 2004. A report Produced for Ireland’s Environmental Protection Agency by Netcen, an operating division of AEA Technology PLC and Clean Technology Centre, Cork. 89 pp.

O’Mara, F., 2007. Development of Emission Factors for the Irish Cattle Herd. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Office of the Revenue Commissioners, 2015 Revenue Statistics and Economic Branch. Illegal Tobacco Products Research Surveys 2015, <http://www.payeanytime.ie/en/tax/excise/tobacco/index.html>

Plejdrup, M.S., Nielsen, O.-K. & Bruun, H.G., 2017: MapElre – A high resolution spatial emission model for Ireland.

SEAI (Sustainable Energy Authority Ireland), 2016. Energy in Ireland 2016 – SEAI; http://www.seai.ie/Publications/Statistics_Publications/Energy_in_Ireland/Energy-in-Ireland-1990-2015.pdf

Thistlethwaite, G., Baggott, S.L. and Goodwin, J., 2005. National Inventory System Scoping Report for the Greenhouse Gas Emission Inventory in Ireland. AEAT/ENV/R/1990 Issue 1, NETCEN, UK.

UK NAEI, Crown 2018 copyright Defra & BEIS via naei.defra.gov.uk, licenced under the [Open Government Licence](#) (OGL).

UNEP Toolkit, 2013. Standardized Toolkit for Identification and Quantification of Dioxin and Furan Releases.

URS Dames & Moore, 2000. Inventory of Dioxin and Furan Emissions to Air, Land and Water in Ireland for 2000 and 2010. Environmental Protection Agency, Johnstown Castle Estate, Wexford, Ireland.

Xu, Y., 2008. Review and update of the HCB inventory of emissions to air, land and water for the UK. AEA. Group, Oxfordshire, UK.

Zimmerman, J., 2014. A review of crop residue burning. MODIS fire detection archive for Ireland.

Glossary

ADDF	Annual Average Daily Flow
AER	Annual Environmental Report
As	Arsenic
B[a]P	Benzo[a]pyrene
B[b]F	Benzo[b]fluoranthene
B[k]F	Benzo[k]fluoranthene
BCF	British Coatings Federations
CAP	Common Agricultural Policy
Cd	Cadmium
CEPE	European Council of Producers and Importers of Paints, Printing Inks and Artists Colours
CEPMEIP	Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance
CLEEN	Chemical Legislation European Enforcement Network
CLRTAP	Convention on Long Long-Range Transboundary Air Pollution
CO	Carbon monoxide
CORINAIR	Co-ordinated Information on the environment in the European Community-AIR. CORINAIR was one of several collaborative exercises initiated under the CORINE programme to harmonise the collection and dissemination of information on the environment in the EU
CMMS	Cattle Movement and Monitoring Scheme
Cr	Chromium
CSO	Central Statistics Office
Cu	Copper
DEHLG	Department of Environment, Heritage and Local Government
DM	Dry matter
DTTAS	Department of Transport, Tourism and Sport
DQO	Data quality objective
EAPA	European Asphalt Pavement Association
ED	Electoral Division
ELV	End-of-Life Vehicle
EMEP	European Monitoring and Evaluation Programme, a co-operative programme for monitoring and evaluation of the long-range transmissions of air pollutants in Europe
EPA	Environmental Protection Agency
E-PRTR	European Pollutant Release and Transfer Register
ESB	Electricity Supply Board
ESP	Electrostatic precipitators
ETS	Emissions Trading Scheme
EUROSTAT	Statistical Agency of the European Union
FFS	Farm Facilities Survey
Fossil Fuel	Peat, coal, oil and natural gas and associated derivatives

FUS	Fertiliser Use Survey
GHG	Greenhouse gas
Gg	Gigagram (10 ⁹ g) = kilotonne = 1,000 tonnes
GNFR	Gridded Nomenclature for Reporting Codes
HCB	Hexachlorobenzene
HFO	Heavy fuel oil
Hg	Mercury
IBEC	Irish Business and Employers' Confederation
IEA	International Energy Agency
IEF	Implied Emission Factor
IFFPG	Irish Farm Film Producers Group
IIR	Informative Inventory Report
I[123-cd]P	Indeno[1,2,3-cd]pyrene
IPC	Integrated Pollution Control
IPCC	Intergovernmental Panel on Climate Change
IPPC	Integrated Pollution Prevention and Control
KDP	Key Data Provider
ktoe	Kilotonnes of oil equivalent
LCP	Large Combustion Plant Directive
LFG	Landfill gas
LPS	Large Point Source
LTO	Landing and take-off
MoU	Memorandum of Understanding
MSW	Municipal solid waste
NAIS	National Atmospheric Inventory System
NAEI	National Atmospheric Emissions Inventory
NCT	National Car Testing
NETCEN	National Environmental Technology Centre
NEC	National Emission Ceilings
NFR	Nomenclature for Reporting Codes
NH₃	Ammonia
Ni	Nickel
NMVOC	Non-methane volatile organic compound
NO_x	Nitrogen oxides
NRA	National Roads Authority
OCLR	Office of Climate, Licensing, Research and Resource Use
OLG	Office of Licensing and Guidance
PAH	Polycyclic aromatic hydrocarbon
Pb	Lead
PCB	Polychlorinated biphenyl
PER	Pollution Emissions Register
PM	Particulate matter
PM₁₀	Particulate matter <10 µm in diameter
PM_{2.5}	Particulate matter <2.5 µm in diameter
POP	Persistent organic pollutant
PVC	Polyvinyl chloride
QA/QC	Quality assurance/quality control
S.I.	Statutory Instrument

Se	Selenium
SEAI	Sustainable Energy Authority of Ireland
SNAP	Selected Nomenclature for Air Pollution
SO₂	Sulphur dioxide
SO_x	Sulphur oxides
TAN	Total ammoniacal nitrogen
Teagasc	Irish Agriculture and Food Development Authority
TPM	Total particulate matter
TSP	Total suspended particulates
UAN	Uric acid nitrogen
UK NAEI	United Kingdom National Atmospheric Emission Inventory
UNECE	United Nations Economic Commission for Europe
VOC	Volatile organic compounds
WEEE	Waste Electrical and Electronic Equipment Regulation
Zn	Zinc

Annex A

A.1 Annex 1 Table 2016

A.2 Key Category Analysis 2016

A.3 Fuel Tourism in Road Transport and Nitrogen Oxides Emissions Based on Fuels Used

ANNEX 1: National sector emissions: Main pollutants, particulate matter, heavy metals and persistent organic pollutants

NFR 2014-2

COUNTRY:

IE

(as ISO2 code)

DATE:

15.03.2018

(as DD.MM.YYYY)

YEAR:

2016

(as YYYY, year of emissions and activity data)

Version:

v2.0

(as v1.0 for the initial submission)

XML Export for all entered years

Add a new year

[illegible]

Annex A.1 Annex 1Table (contd.)

OtherStationaryComb	1A5a	Other stationary (including military)		IE	IE	IE	NE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	NE	NA	IE	IE	IE	IE	NO	NA	TJ NCV
I,Offroad	1A5b	Other, Mobile (including military, land based and recreational boats)		IE	IE	IE	NE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	NE	NA	IE	NA	NA	NA	NO	NA	TJ NCV
D_Fugitive	1B1a	Fugitive emission from solid fuels: Coal mining and handling		NA	NO	NO	NA	0.002	NO	NO	NO	NA	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NO	NA	Coal produced [Mt]
D_Fugitive	1B1b	Fugitive emission from solid fuels: Solid fuel transformation		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NO	NA	Coal used for transformation [Mt]	
D_Fugitive	1B1c	Other fugitive emissions from solid fuels		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	Please specify	
D_Fugitive	1B2ai	Fugitive emissions oil: Exploration, production, transport		NA	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	Crude oil produced [Mt]	
D_Fugitive	1B2aiv	Fugitive emissions oil: Refining /storage		IE	2.735	IE	IE	IE	IE	IE	NE	IE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	3	Crude oil refined [Mt]	
D_Fugitive	1B2av	Distribution of oil products		NA	1.760	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1	Oil consumed [Mt]		
D_Fugitive	1B2b	Fugitive emissions from natural gas (exploration, production, processing, transmission, storage, distribution and other)		NA	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	Gas throughput [Mn3]	
D_Fugitive	1B2c	Venting and flaring (oil, gas, combined oil and gas)		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	Gas vented flared [TJ]	
D_Fugitive	1B2d	Other fugitive emissions from energy production	(a)	NE	NE	NE	NE	NO	NO	NO	NO	NE	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		
B_Industry	2A1	Cement production		IE	IE	IE	NA	IE	NE	NE	NE	IE	IE	IE	IE	IE	IE	IE	IE	1.476	IE	IE	IE	IE	IE	IE	NE	5.177	NA	NA	NA	NA	NA	3275	Clinker produced [kt]
B_Industry	2A2	Lime production		IE	IE	IE	NA	NA	NA	NA	NA	IE	NA	NA	NA	NA	NA	NA	NA	IE	NA	NA	NA	NA	NA	NA	NA	IE	NA	NA	NA	NA	229	Lime produced [kt]	
B_Industry	2A3	Glass production		IE	IE	IE	NA	NO	NO	NO	NO	NO	IE	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NE	NA	NA	NA	NA	NO	Glass produced [t]	
B_Industry	2A5a	Quarrying and mining of minerals other than coal		NA	NA	NA	NA	0.182	1.822	3.716	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	36	Material quarried [Mt]	
B_Industry	2A5b	Construction and demolition		NA	NA	NA	NA	0.046	0.463	1.543	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	4069381	Floor space constructed/demolished [M2]	
B_Industry	2A5c	Storage, handling and transport of mineral products		NA	NA	NA	NA	0.000	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	36	Amount [Mt]	
B_Industry	2A6	Other mineral products (please specify in the IIR)		NA	NA	NA	NA	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	0.016	0.010	NA	0.002	NA	0.012	NA	NA	NA	NA	NA	NO	NA	Please specify		
B_Industry	2B1	Ammonia production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	Ammonia produced [kt]	
B_Industry	2B2	Nitric acid production		NO	NA	NA	NE	NO	NO	NO	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NO	NA	Nitric acid produced [kt]		
B_Industry	2B3	Adipic acid production		NO	NA	NA	NA	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NO	NA	Adipic acid produced [kt]		
B_Industry	2B5	Carbide production		NO	NO	NO	NA	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NO	NA	Carbide produced [kt]		
B_Industry	2B6	Titanium dioxide production		NO	NO	NO	NA	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NO	NA	Titanium dioxide produced [kt]		
B_Industry	2B7	Soda ash production		NO	NO	NO	NA	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.1	NA	Soda ash produced [kt]		
B_Industry	2B10a	Chemical Industry: Other (please specify in the IIR)		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NO	NA	Please specify		
B_Industry	2B10b	Storage, handling and transport of chemical products (please specify in the IIR)		NE	NE	NE	NE	NO	NO	NO	NE	NE	NO	NO	NO	NO	NO	NO	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NO	NA	Please specify		
B_Industry	2C1	Iron and steel production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NE	NO	NA	NA	NE	Steel produced [kt]			
B_Industry	2C2	Ferroalloys production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.000	IE	IE	IE	IE	IE	IE	IE	NO	NA	NA	NA	NE	NA	Ferroalloys produced [kt]		
B_Industry	2C3	Aluminium production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NO	Aluminium produced [kt]		
B_Industry	2C4	Magnesium production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NO	Magnesium produced [kt]		
B_Industry	2C5	Lead production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	IE	NE	NE	NE	NE	NE	NE	NE	NO	IE	NA	NA	NE	Lead produced [kt]			
B_Industry	2C6	Zinc production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NO	NA	Zinc produced [kt]		
B_Industry	2C7a	Copper production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NO	NA	Copper produced [kt]		
B_Industry	2C7b	Nickel production		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NO	NA	Nickel produced [kt]		
B_Industry	2C7c	Other metal production (please specify in the IIR)		NO	NO	NO	NO	NO	NO	NO	NO	NO	0.000	NO	NO	NO	NO	NO	0.000	IE	IE	IE	IE	IE	IE	IE	NO	IE	NA	NA	NO	NA	Please specify		
B_Industry	2C7d	Storage, handling and transport of metal products (please specify in the IIR)		NE	NE	NE	NE	NE	NE	NE	NE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	NE	Amount [kt]			
E_Solvents	2D3a	Domestic solvent use including fungicides		NA	10.767	NA	NA	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	NA			
B_Industry	2D3b	Road paving with asphalt		NE	0.030	NE	NA	1.900	3.800	28.500	0.011	NE	NA	NA	NA	NA	NA	NA	NA	NE	NE	IE	NE	IE	IE	IE	IE	NE	NA	NA	NA	2	Production of warm and hot asphalt [Mt]		
B_Industry	2D3c	Asphalt roofing		NA	NE	NA	NA	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NE	NA			
E_Solvents	2D3d	Coating applications		NA	2.916	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	Paint applied [kt]			
E_Solvents	2D3e	Degreasing		NA	0.837	NA	NA	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	2	Solvents used [kt]			
E_Solvents	2D3f	Dry cleaning		NA	0.062	NA	NA	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.199	NA	Solvents used [kt]		
E_Solvents	2D3g	Chemical products		NE	1.168	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NE	NA			
E_Solvents	2D3h	Printing		NA	1.463	NA	NA	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	NA			
E_Solvents	2D3i	Other solvent use (please specify in the IIR)		NA	1.370	NA	NA	0.020	0.031	0.038	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	34	Oilseed rape crop yield [kt]			

Annex A.1 Annex 1 Table (contd.)

E_Solvents	2G	Other product use (please specify in the IIR)		0.006	0.017	0.000	0.015	0.096	0.098	0.098	0.038	0.194	0.022	0.000	0.000	0.000	0.000	0.012	0.001	NE	0.007	0.000	0.000	0.000	0.000	0.001	NE	NE		NA	NA	NA	NA	NA	3519	Tobacco products [t] see IIR for all sources of AD	
B_Industry	2H1	Pulp and paper industry		NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	NE	NE	NE	NE	NE	NA		NA	NA	NA	NA	NA	NA	Pulp production [kt]
B_Industry	2H2	Food and beverages industry		NA	21.261	NA	NA	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	Bread, Wine, Beer, Spirits production [kt]
B_Industry	2H3	Other industrial processes (please specify in the IIR)		NA	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NA	
B_Industry	2I	Wood processing		NE	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NE	NA	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	Please specify
B_Industry	2J	Production of POPs		NE	NE	NE	NE	NO	NO	NO	NO	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	NO		NA	NA	NA	NA	NO	NA
B_Industry	2K	Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)		NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NE	NO		NA	NA	NA	NA	NA	NE	NA
B_Industry	2L	Other production, consumption, storage, transportation or handling of bulk products (please specify in the IIR)		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.000	NE	NE	NE	NE	NE	NE	0.000		NA	NA	NA	NA	NA	NA	NA
K_AgriLivestock	3B1a	Manure management - Dairy cattle		0.052	9.295	NA	13.434	0.598	0.915	2.008	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	3585	Population size (1000 head)
K_AgriLivestock	3B1b	Manure management - Non-dairy cattle		0.558	26.740	NA	31.281	0.739	1.105	2.448	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	3589	Population size (1000 head)
K_AgriLivestock	3B2	Manure management - Sheep		0.049	0.356	NA	1.127	0.095	0.286	0.668	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	4770	Population size (1000 head)
K_AgriLivestock	3B3	Manure management - Swine		0.002	2.485	NA	4.719	0.008	0.183	1.251	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	1561	Population size (1000 head)
K_AgriLivestock	3B4a	Manure management - Buffalo		NO	NO	NA	NO	NO	NO	NO	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NO	Population size (1000 head)
K_AgriLivestock	3B4d	Manure management - Goats		0.001	0.002	NA	0.019	0.000	0.001	0.001	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	10	Population size (1000 head)
K_AgriLivestock	3B4e	Manure management - Horses		0.034	0.170	NA	0.788	0.013	0.020	0.044	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	92	Population size (1000 head)
K_AgriLivestock	3B4f	Manure management - Mules and asses		0.002	0.009	NA	0.054	0.001	0.001	0.003	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	9	Population size (1000 head)
K_AgriLivestock	3B4gi	Manure management - Laying hens		0.027	0.547	NA	0.841	0.010	0.133	0.630	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	3318	Population size (1000 head)
K_AgriLivestock	3B4gii	Manure management - Broilers		0.072	1.330	NA	1.494	0.025	0.246	0.493	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	12318	Population size (1000 head)
K_AgriLivestock	3B4giii	Manure management - Turkeys		0.029	0.622	NA	0.841	0.025	0.140	0.140	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	1273	Population size (1000 head)
K_AgriLivestock	3B4giv	Manure management - Other poultry		0.004	0.133	NA	0.080	0.006	0.039	0.039	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	272	Population size (1000 head)
K_AgriLivestock	3B4h	Manure management - Other animals (please specify in IIR)		0.006	0.384	NA	0.416	0.001	0.002	0.004	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	199	Population size (1000 head)
L_AgriOther	3Da1	Inorganic N-fertilizers (includes also urea application)		13.047	NA	NA	11.192	0.267	6.938	6.938	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	339104000	Use of inorganic fertilizers (kg N/yr)
L_AgriOther	3Da2a	Animal manure applied to soils		5.943	NE	NA	34.605	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Da2b	Sewage sludge applied to soils		0.010	NE	NA	0.442	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Da2c	Other organic fertilisers applied to soils (including compost)		NE	NE	NA	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Da3	Urine and dung deposited by grazing animals		11.929	NE	NA	14.194	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Da4	Crop residues applied to soils		NE	NE	NA	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Db	Indirect emissions from managed soils		NE	NE	NA	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products		NA	NA	NA	NE	0.006	0.045	0.141	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	1412	NPK fertilisers [kt]
L_AgriOther	3Dd	Off-farm storage, handling and transport of bulk agricultural products		NA	NA	NA	NE	0.009	0.058	0.231	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	2311	Wheat Oats Barley [kt]
L_AgriOther	3De	Cultivated crops	(b)	NE	3.825	NA	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3Df	Use of pesticides		NE	NE	NA	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1.151	NA		NA	NA	NA	NA	NA	NE	
L_AgriOther	3F	Field burning of agricultural residues		NE	NE	NE	NE	NE	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA		NA	NA	NA	NA	NA	NE	Area burned [k ha/yr]
L_AgriOther	3I	Agriculture other (please specify in the IIR)		NO	NO	NO	NE	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NE	NE	NE	NE	NE	NE	NE	NE	NE		NA	NA	NA	NA	NA	NA	NA

Annex A.1 Annex I Table (contd.)

J_Waste	5A	Biological treatment of waste - Solid waste disposal on land		NA	0.486	NA	NA	0.000	0.000	0.000	NE	NA	NA	NA	0.021	NA	NA	NA	NA	NA	NA	0.075	NA	NA	NA	NA	NA	NA	0.049	NA	NA	NA	NA	802	Annual deposition of MSW at the SWDS [kg]						
J_Waste	5B1	Biological treatment of waste - Composting		NA	NE	NA	0.028	NE	NE	NE	NE	0.065	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE									
J_Waste	5B2	Biological treatment of waste - Anaerobic digestion at biogas facilities		NA	NO	NA	NO	NE	NE	NE	NE	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE										
J_Waste	5C1a	Municipal waste incineration	(c)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	MSW incinerated [kt]								
J_Waste	5C1bi	Industrial waste incineration	(c)	0.007	0.055	0.000	NE	0.000	0.000	0.000	0.000	0.001	0.010	0.001	0.000	0.649	0.627	0.346	0.000	NO	NE	0.014	0.000	0.000	0.000	NE	0.000	0.004	0.006	NA	NA	11	Waste incinerated [kt]								
J_Waste	5C1bii	Hazardous waste incineration	(c)	IE	IE	IE	NE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	Waste incinerated [kt]								
J_Waste	5C1biii	Clinical waste incineration	(c)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	Waste incinerated [kt]								
J_Waste	5C1biv	Sewage sludge incineration	(c)	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	NO								
J_Waste	5C1bv	Cremation	(c)	0.004	0.000	0.001	NE	0.000	0.000	0.000	NE	0.001	0.000	0.000	0.008	0.000	0.000	0.000	0.000	NO	NO	0.000	0.000	NE	NE	0.000	0.000	0.001	NE	NA	NA	4178	Incineration of corpses [number]								
J_Waste	5C1bw	Other waste incineration (please specify in the IIR)	(c)	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA								
J_Waste	5C2	Open burning of waste		NE	NE	NE	NE	NE	NE	NE	NE	NE	NO	NO	NO	NO	NO	NO	NO	NO	0.461	0.000	0.001	0.001	NE	0.001	NE	0.783	NA	NA	2	Quantity of farm plastic burned [kt]									
J_Waste	5D1	Domestic wastewater handling		NA	NA	NA	NE	NE	NE	NE	NE	NA	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	Total organic product [Gg DC/yr]									
J_Waste	5D2	Industrial wastewater handling		NA	NA	NA	NE	NE	NE	NE	NE	NA	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	Total organic product [Gg DC/yr]									
J_Waste	5D3	Other wastewater handling		NA	NA	NA	NE	NE	NE	NE	NE	NA	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NE	Total organic product [Gg DC/yr]										
J_Waste	5E	Other waste (please specify in IIR)	(d)	NE	NE	NE	NE	0.165	0.165	0.165	NE	NE	NO	NO	NO	NO	NO	NO	NO	NO	2.252	0.013	0.016	0.006	0.001	0.036	NE	2.555	NA	NA	4	Vehicle and training material burned [kt] see IIR									
M_Other	6A	Other (included in national total for entire territory) (please specify in IIR)		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA								
	NATIONAL TOTAL	National total for the entire territory (based on fuel sold)		112.277	108.408	13.771	116.723	15.468	29.056	63.628	2.225	102.562	13.400	0.284	0.329	1.265	2.491	20.158	8.485	3.557	20.204	21.224	4.159	6.281	2.405	2.055	14.901	1.677	12.433	261465.35	88816.60	179541.20	16132.88	1025.49	NA						
	ADJUSTMENTS (Net total)	Sum of adjustments (negative value) from Annex VII		-16.065	-61.414																												NA								
	NATIONAL TOTAL FOR COMPLIANCE	National total for compliance assessment (please specify all details in the IIR)	(e)	91.234	46.839	13.766	116.700	14.941	28.214	61.851	1.829	101.272	13.129	0.280	0.351	1.265	2.417	18.306	8.449	3.552	19.344	21.045	4.150	6.268	2.394	2.046	14.858	1.677	12.388					NA							
BEMO ITEMS - NOT TO BE INCLUDED IN NATIONAL TOTALS																																									
O_AirCruise	1A3a(i)	International aviation cruise (civil)		3.216	0.096	0.724	NE	0.146	0.146	0.146	0.070	0.386	0.003	0.000	0.000	0.001	0.000	0.007	0.000	0.000	0.933	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	32156	NA	NA	NA	NA	NA	NA	TJ NCV
O_AirCruise	1A3a(ii)	Domestic aviation cruise (civil)		0.022	0.000	0.002	NE	0.001	0.001	0.001	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	70.009079	NA	NA	NA	NA	NA	NA	TJ NCV
P_InShipping	1A3d(i)	International maritime navigation		12.169	0.433	0.443	NE	0.262	0.283	0.283	0.075	1.146	0.009	0.008	0.003	0.009	0.010	0.050	0.863	0.042	0.053	0.190	0.001	0.003	0.001	0.001	0.006	NE	0.155	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	TJ NCV	
z_Memo	1A5c	Multilateral operations		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NA	NA	NA	NA	NA	NA	NA
z_Memo	1A3	Transport (fuel used)		43.977	5.264	0.256	0.761	1.574	2.021	2.024	0.764	50.066	9.220	0.044	0.048	0.005	0.649	16.454	0.416	0.056	8.017	1.356	0.067	0.093	0.078	0.065	0.303	0.008	0.359	NE	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
z_Memo	6B	Other not included in national total of the entire territory (please specify in the IIR)		NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
N_Natural	11A	Volcanoes		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	Please specify
N_Natural	11B	Forest fires		NE	NE	NE	NE	NO	NO	NO	NO	NE	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.019	0.140	0.091	0.045	0.052	0.328	NE	0.006	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	Area of forest burned [ha]
N_Natural	11C	Other natural emissions (please specify in the IIR)		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA

MEMO ITEMS - NOT TO BE INCLUDED IN NATIONAL TOTALS																																				
O_AirCruise	1A3ai(ii)	International aviation cruise (civil)		3.216	0.096	0.724	NE	0.146	0.146	0.146	0.070	0.386	0.003	0.000	0.004	0.001	0.006	0.007	0.000	0.004	0.933	NE	NE	NE	NE	NE	NE	NE	NE	32156	NA	NA	NA	NA	NA	TJ NCV
O_AirCruise	1A3ai(iii)	Domestic aviation cruise (civil)		0.022	0.000	0.002	NE	0.001	0.001	0.001	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002	NE	NE	NE	NE	NE	NE	NE	NE	70.009079	NA	NA	NA	NA	NA	TJ NCV
P_IntShipping	1A3di(i)	International maritime navigation		12.169	0.433	0.443	NE	0.262	0.283	0.283	0.075	1.146	0.009	0.008	0.003	0.009	0.010	0.050	0.663	0.042	0.053	0.190	0.001	0.003	0.001	0.001	0.006	NE	0.155	6686.4682	NA	NA	NA	NA	NA	TJ NCV
z_Memo	1A5c	Multilateral operations		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	
z_Memo	1A3	Transport (fuel used)		43.977	5.264	0.256	0.761	1.574	2.021	2.024	0.764	50.066	9.220	0.044	0.048	0.005	0.649	16.454	0.416	0.056	8.017	1.356	0.067	0.093	0.078	0.065	0.303	0.008	0.359	NE	NA	NA	NA	NA	NA	
z_Memo	6B	Other not included in national total of the entire territory (please specify in the IIR)		NO	NO	NO	NO	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
N_Natural	11A	Volcanoes		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	Please specify	
N_Natural	11B	Forest fires		NE	NE	NE	NE	NO	NO	NO	NO	NE	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.019	0.140	0.091	0.045	0.052	0.328	NE	0.006	NA	NA	NA	NA	NA	Area of forest burned [ha]	
N_Natural	11C	Other natural emissions (please specify in the IIR)		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA	NA	NA	NA	NA	

- (a) For example, fugitive emissions from the production of geothermal power could be reported here.
 (b) Does not include emissions from application of fertiliser and manure (reported under 3D). NH₃ emissions from crops should be reported here.
 (c) Excludes waste incineration for energy (this is included in 1A1) and in industry (if used as fuel).
 (d) Includes accidental fires.
 (e) The National Total for Compliance/ includes any aggregated combination of (i) adjustments to national totals; (ii) national totals based on transport fuel used; (iii) territory declared upon ratification of the relevant Protocol of the Convention.

Member States of the European Union may also use this line for reporting national totals for compliance purposes under the National Emission Ceilings Directive (NECD) if these differ from the main National Total. MS should consult the definitions of geographical coverage in the NECD to determine what should be included within the NECD National Total.

Annex A.2 Table 1: Key Category Analysis for Nitrogen Oxides

Level Assessment

NFR	NOx (kt)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	112.2769			
1A3bi	17.0176	15.16%	15.16%	x
1A3biii	13.0895	11.66%	26.81%	x
3Da1	13.0471	11.62%	38.44%	x
3Da3	11.9292	10.62%	49.06%	x
1A3bii	9.1877	8.18%	57.24%	x
1A1a	8.0003	7.13%	64.37%	x
1A2f	7.0507	6.28%	70.65%	x
1A3dii	6.5205	5.81%	76.46%	x
3Da2a	5.9433	5.29%	81.75%	x
1A4bi	4.9440	4.40%	86.15%	
1A4ai	2.5963	2.31%	88.47%	
1A4cii	2.5244	2.25%	90.71%	
1A3c	1.8475	1.65%	92.36%	
1A4ciii	1.4631	1.30%	93.66%	
1A2gviii	1.3867	1.24%	94.90%	
1A2e	1.2707	1.13%	96.03%	
1A2b	1.2125	1.08%	97.11%	
1A3ai(i)	1.1168	0.99%	98.10%	
3B1b	0.5578	0.50%	98.60%	
1A2c	0.3666	0.33%	98.93%	
1A1c	0.3174	0.28%	99.21%	
1A1b	0.3155	0.28%	99.49%	
1A3ei	0.1202	0.11%	99.60%	
3B4gii	0.0723	0.06%	99.66%	
1A4ci	0.0667	0.06%	99.72%	
3B1a	0.0522	0.05%	99.77%	
3B2	0.0486	0.04%	99.81%	
1A3biv	0.0349	0.03%	99.84%	
3B4e	0.0336	0.03%	99.87%	
3B4giii	0.0292	0.03%	99.90%	
3B4gi	0.0273	0.02%	99.92%	
1A2d	0.0216	0.02%	99.94%	
1A3aii(i)	0.0196	0.02%	99.96%	
3Da2b	0.0095	0.01%	99.97%	
5C1bi	0.0065	0.01%	99.97%	
3B4h	0.0064	0.01%	99.98%	
2G	0.0063	0.01%	99.98%	
5C1bv	0.0042	0.00%	99.99%	
3B4giv	0.0038	0.00%	99.99%	
1A2a	0.0031	0.00%	99.99%	
3B4f	0.0023	0.00%	100.00%	
3B3	0.0023	0.00%	100.00%	
3B4d	0.0014	0.00%	100.00%	

Trend Assessment

NFR	NOx, 1990 (kt)	NOx, 2016 (kt)	Trend (magnitude)	Trend %	% Cumulative	Key Category
1A1a	46.3740	8.0003	0.13477	33.97%	33.97%	x
1A3bi	37.9848	17.0176	0.04851	12.23%	46.20%	x
1A3bii	5.2845	9.1877	0.03358	8.47%	54.67%	x
1A3biii	11.4082	13.0895	0.03261	8.22%	62.89%	x
1A3dii	2.1355	6.5205	0.03017	7.61%	70.49%	x
1A2f	3.3392	7.0507	0.02858	7.21%	77.70%	x
3Da3	12.1933	11.9292	0.02267	5.71%	83.41%	x
3Da1	14.5941	13.0471	0.01985	5.00%	88.42%	
3Da2a	5.3381	5.9433	0.01419	3.58%	92.00%	
1A4cii	6.5202	2.5244	0.01067	2.69%	94.69%	
1A2gviii	1.1327	1.3867	0.00375	0.95%	95.63%	
1A4ai	3.0603	2.5963	0.00334	0.84%	96.47%	
1A3ai(i)	0.9267	1.1168	0.00297	0.75%	97.22%	
1A3c	2.1985	1.8475	0.00229	0.58%	97.80%	
1A2e	1.5126	1.2707	0.00158	0.40%	98.20%	
3B1b	0.4851	0.5578	0.00139	0.35%	98.55%	
1A1c	0.1612	0.3174	0.00124	0.31%	98.86%	
1A2a	0.2886	0.0031	0.00111	0.28%	99.14%	
1A2b	2.0331	1.2125	0.00081	0.20%	99.35%	
1A3ei	0.0542	0.1202	0.00050	0.13%	99.47%	
1A4bi	7.3261	4.9440	0.00047	0.12%	99.59%	
1A3aii(i)	0.1108	0.0196	0.00032	0.08%	99.67%	
3B4gii	0.0472	0.0723	0.00024	0.06%	99.73%	
1A4ciii	2.1497	1.4631	0.00021	0.05%	99.79%	
1A2c	0.5840	0.3666	0.00013	0.03%	99.82%	
1A3biv	0.0218	0.0349	0.00012	0.03%	99.85%	
3B1a	0.0494	0.0522	0.00011	0.03%	99.88%	
3B4e	0.0225	0.0336	0.00011	0.03%	99.91%	
3B4gi	0.0208	0.0273	0.00008	0.02%	99.93%	
5C1bi	0.0236	0.0065	0.00005	0.01%	99.94%	
1A4ci	0.0901	0.0667	0.00004	0.01%	99.95%	
3B4giii	0.0347	0.0292	0.00004	0.01%	99.96%	
1A1b	0.4668	0.3155	0.00003	0.01%	99.97%	
3B4h	0.0173	0.0064	0.00003	0.01%	99.97%	
3Da2b	0.0070	0.0095	0.00003	0.01%	99.98%	
1A2d	0.0377	0.0216	0.00002	0.01%	99.99%	
5C1bv	0.0012	0.0042	0.00002	0.00%	99.99%	
2G	0.0126	0.0063	0.00001	0.00%	99.99%	
3B3	0.0019	0.0023	0.00001	0.00%	100.00%	
3B4f	0.0021	0.0023	0.00001	0.00%	100.00%	
3B2	0.0743	0.0486	0.00000	0.00%	100.00%	
3B4giv	0.0050	0.0038	0.00000	0.00%	100.00%	
3B4d	0.0025	0.0014	0.00000	0.00%	100.00%	
NATIONAL TOTAL	169.1030	112.2769	0.00000	0.00%	100.00%	

Annex A.2 Table 2: Key Category Analysis for Sulphur Dioxide

Level Assessment

NFR	SO ₂ (kt)	% Contribution Level	% Cumulative	Key Category
National Total	13.771			
1A4bi	6.664	48.39%	48.39%	x
1A1a	3.718	27.00%	75.39%	x
1A2f	1.340	9.73%	85.12%	x
1A2e	0.990	7.19%	92.31%	
1A1c	0.245	1.78%	94.09%	
1A4ai	0.235	1.71%	95.80%	
1A2gviii	0.162	1.17%	96.98%	
1A3dii	0.116	0.84%	97.82%	
1A3ai(i)	0.092	0.67%	98.49%	
1A2c	0.079	0.58%	99.07%	
1A4cii	0.023	0.17%	99.23%	
1A1b	0.020	0.14%	99.38%	
1A3bi	0.019	0.14%	99.52%	
1A3c	0.014	0.10%	99.62%	
1A2b	0.013	0.09%	99.71%	
1A3biii	0.009	0.07%	99.78%	
1A3bii	0.008	0.06%	99.84%	
1A4ciii	0.007	0.05%	99.89%	
1A2d	0.006	0.05%	99.94%	
1A4ci	0.006	0.04%	99.98%	
1A3aii(i)	0.001	0.01%	99.99%	
5C1bv	0.001	0.00%	100.00%	
5C1bi	0.000	0.00%	100.00%	
1A3ei	0.000	0.00%	100.00%	
2G	0.000	0.00%	100.00%	
1A2a	0.000	0.00%	100.00%	
1A3biv	0.000	0.00%	100.00%	

Trend Assessment

NFR	SO ₂ , 1990 (kt)	SO ₂ , 2016 (kt)	Trend (magnitude)	Trend %	% Cumulative	Key Category
1A4bi	26.7591	6.6637	0.02543	34.94%	34.94%	x
1A1a	103.0440	3.7176	0.02213	30.41%	65.35%	x
1A2b	15.9561	0.0128	0.00651	8.94%	74.29%	x
1A2f	2.0097	1.3404	0.00651	8.94%	83.23%	x
1A4ai	11.5240	0.2353	0.00346	4.76%	87.99%	
1A2e	7.0091	0.9903	0.00253	3.48%	91.46%	
1A1c	0.1331	0.2451	0.00129	1.77%	93.23%	
1A2gviii	4.3854	0.1616	0.00092	1.27%	94.50%	
1A3bi	2.3754	0.0189	0.00088	1.20%	95.70%	
1A3biii	1.5370	0.0094	0.00058	0.80%	96.50%	
1A3bii	1.4562	0.0085	0.00055	0.76%	97.26%	
1A3ai(i)	0.0762	0.0922	0.00047	0.65%	97.91%	
1A2c	1.9811	0.0795	0.00038	0.52%	98.44%	
1A2a	0.8802	0.0000	0.00036	0.50%	98.94%	
1A4cii	1.1220	0.0231	0.00034	0.46%	99.40%	
1A3dii	1.1607	0.1163	0.00016	0.22%	99.62%	
1A2d	0.3389	0.0063	0.00011	0.14%	99.76%	
1A1b	0.4758	0.0199	0.00009	0.12%	99.88%	
1A3c	0.2515	0.0138	0.00003	0.04%	99.92%	
1A4ciii	0.1641	0.0073	0.00003	0.04%	99.96%	
1A4ci	0.1247	0.0060	0.00002	0.03%	99.98%	
1A3aii(i)	0.0084	0.0015	0.00000	0.01%	99.99%	
5C1bv	0.0002	0.0006	0.00000	0.00%	99.99%	
1A3biv	0.0045	0.0000	0.00000	0.00%	100.00%	
5C1bi	0.0013	0.0004	0.00000	0.00%	100.00%	
1A3ei	0.0001	0.0001	0.00000	0.00%	100.00%	
2G	0.0001	0.0001	0.00000	0.00%	100.00%	
National Total	182.7808	13.7706	0.00000	0.00%	100.00%	

Annex A.2 Table 3: Key Category Analysis for Non-Methane Volatile Organic Compounds

Level Assessment

NFR	NMVOC (kt)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTA	108.4076			
3B1b	26.7395	24.67%	24.67%	x
2H2	21.2606	19.61%	44.28%	x
2D3a	10.7666	9.93%	54.21%	x
3B1a	9.2951	8.57%	62.78%	x
1A4bi	8.0513	7.43%	70.21%	x
3De	3.8246	3.53%	73.74%	x
2D3d	2.9160	2.69%	76.43%	x
1B2aiv	2.7345	2.52%	78.95%	x
1A3bi	2.5589	2.36%	81.31%	x
3B3	2.4845	2.29%	83.60%	
1B2av	1.7605	1.62%	85.23%	
2D3h	1.4627	1.35%	86.58%	
1A3bv	1.3951	1.29%	87.86%	
2D3i	1.3698	1.26%	89.13%	
3B4gii	1.3304	1.23%	90.35%	
2D3g	1.1682	1.08%	91.43%	
1A2f	0.9201	0.85%	92.28%	
2D3e	0.8371	0.77%	93.05%	
1A2gviii	0.7711	0.71%	93.76%	
3B4giii	0.6224	0.57%	94.34%	
1A4ai	0.5786	0.53%	94.87%	
1A2b	0.5742	0.53%	95.40%	
1A2e	0.5586	0.52%	95.92%	
3B4gi	0.5475	0.51%	96.42%	
5A	0.4859	0.45%	96.87%	
3B4h	0.3837	0.35%	97.22%	
1A3biii	0.3736	0.34%	97.57%	
3B2	0.3559	0.33%	97.90%	
1A3bii	0.3328	0.31%	98.20%	
1A1a	0.3040	0.28%	98.48%	
1A3dii	0.2326	0.21%	98.70%	
1A4cii	0.2318	0.21%	98.91%	
1A3biv	0.2054	0.19%	99.10%	
3B4e	0.1700	0.16%	99.26%	
1A3c	0.1639	0.15%	99.41%	
1A3ai(i)	0.1517	0.14%	99.55%	
3B4giv	0.1330	0.12%	99.67%	
1A2c	0.0923	0.09%	99.76%	
2D3f	0.0622	0.06%	99.81%	
5C1bi	0.0554	0.05%	99.87%	
1A4ciii	0.0522	0.05%	99.91%	
2D3b	0.0304	0.03%	99.94%	
2G	0.0170	0.02%	99.96%	
1A1b	0.0158	0.01%	99.97%	
3B4f	0.0092	0.01%	99.98%	
1A4ci	0.0067	0.01%	99.99%	
1A2d	0.0045	0.00%	99.99%	
1A3ei	0.0040	0.00%	99.99%	
1A1c	0.0020	0.00%	100.00%	
3B4d	0.0016	0.00%	100.00%	
1A3aii(i)	0.0012	0.00%	100.00%	
1A2a	0.0010	0.00%	100.00%	
5C1bv	0.0001	0.00%	100.00%	

Trend Assessment

NFR	NMVOC, 1990 (kt)	NMVOC, 2016 (kt)	Trend (magnitude)	Trend %	% Cumulative	Key Category
1A3bi	24.8260	2.5589	0.11347	20.40%	20.40%	x
2H2	9.6172	21.2606	0.09764	17.55%	37.95%	x
1A4bi	28.5178	8.0513	0.09463	17.01%	54.97%	x
3B1b	22.5635	26.7395	0.06742	12.12%	67.09%	x
2D3a	7.9266	10.7666	0.03328	5.98%	73.07%	x
1A3bv	6.1076	1.3951	0.02257	4.06%	77.13%	x
3B1a	9.2324	9.2951	0.01609	2.89%	80.02%	x
2D3d	6.7426	2.9160	0.01530	2.75%	82.77%	
3B3	1.0167	2.4845	0.01198	2.15%	84.92%	
1B2aiv	1.5910	2.7345	0.01069	1.92%	86.84%	
2D3g	3.0231	1.1682	0.00783	1.41%	88.25%	
3De	3.8199	3.8246	0.00651	1.17%	89.42%	
2D3i	0.7012	1.3698	0.00586	1.05%	90.48%	
1A4cii	1.2898	0.2318	0.00520	0.94%	91.41%	
2D3h	2.9120	1.4627	0.00519	0.93%	92.34%	
3B4gii	0.8678	1.3304	0.00470	0.85%	93.19%	
1A2f	0.4484	0.9201	0.00406	0.73%	93.92%	
1A3bii	1.1563	0.3328	0.00379	0.68%	94.60%	
1A2gviii	0.3312	0.7711	0.00363	0.65%	95.25%	
2D3e	1.7435	0.8371	0.00338	0.61%	95.86%	
1A2b	0.1239	0.5742	0.00336	0.60%	96.46%	
1B2av	1.7850	1.7605	0.00285	0.51%	96.98%	
1A3biii	0.9086	0.3736	0.00220	0.40%	97.37%	
3B4gi	0.3083	0.5475	0.00219	0.39%	97.77%	
1A4ai	0.4736	0.5786	0.00154	0.28%	98.04%	
1A2e	0.4578	0.5586	0.00148	0.27%	98.31%	
1A3dii	0.0736	0.2326	0.00124	0.22%	98.53%	
1A1a	0.1935	0.3040	0.00110	0.20%	98.73%	
2D3f	0.2818	0.0622	0.00106	0.19%	98.92%	
5A	0.8341	0.4859	0.00102	0.18%	99.10%	
1A2a	0.1313	0.0010	0.00069	0.12%	99.23%	
5C1bi	0.2004	0.0554	0.00067	0.12%	99.35%	
3B2	0.5925	0.3559	0.00065	0.12%	99.46%	
1A3biv	0.1531	0.2054	0.00062	0.11%	99.58%	
3B4e	0.1136	0.1700	0.00059	0.11%	99.68%	
3B4h	0.4101	0.3837	0.00051	0.09%	99.77%	
3B4giii	0.7381	0.6224	0.00044	0.08%	99.85%	
1A2c	0.1740	0.0923	0.00028	0.05%	99.90%	
1A3ai(i)	0.1765	0.1517	0.00013	0.02%	99.93%	
1A3c	0.1951	0.1639	0.00011	0.02%	99.95%	
1A1b	0.0065	0.0158	0.00008	0.01%	99.96%	
2G	0.0339	0.0170	0.00006	0.01%	99.97%	
1A4ciii	0.0767	0.0522	0.00004	0.01%	99.98%	
1A3aii(i)	0.0079	0.0012	0.00003	0.01%	99.98%	
2D3b	0.0352	0.0304	0.00003	0.00%	99.99%	
3B4f	0.0083	0.0092	0.00002	0.00%	99.99%	
1A3ei	0.0018	0.0040	0.00002	0.00%	100.00%	
1A1c	0.0006	0.0020	0.00001	0.00%	100.00%	
1A2d	0.0041	0.0045	0.00001	0.00%	100.00%	
3B4d	0.0028	0.0016	0.00000	0.00%	100.00%	
1A4ci	0.0090	0.0067	0.00000	0.00%	100.00%	
3B4giv	0.1757	0.1330	0.00000	0.00%	100.00%	
5C1bv	0.0000	0.0001	0.00000	0.00%	100.00%	
NATIONAL TOTAL	143.1449	108.4076	0.00000	0.00%	100.00%	

Annex A.2 Table 4: Key Category Analysis for Ammonia and Carbon Monoxide

Level Assessment

NFR	NH3 (kt)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	116.7234			
3Da2a	34.6055	29.65%	29.65%	x
3B1b	31.2812	26.80%	56.45%	x
3Da3	14.1945	12.16%	68.61%	x
3B1a	13.4341	11.51%	80.12%	x
3Da1	11.1916	9.59%	89.71%	
3B3	4.7194	4.04%	93.75%	
3B4gii	1.4940	1.28%	95.03%	
3B2	1.1271	0.97%	95.99%	
3B4gi	0.8414	0.72%	96.71%	
3B4giii	0.8408	0.72%	97.44%	
3B4e	0.7875	0.67%	98.11%	
1A3bi	0.7420	0.64%	98.75%	
3Da2b	0.4423	0.38%	99.12%	
3B4h	0.4158	0.36%	99.48%	
1A2gviii	0.1752	0.15%	99.63%	
3B4giv	0.0799	0.07%	99.70%	
1A4bi	0.0614	0.05%	99.75%	
1A2f	0.0610	0.05%	99.80%	
3B4f	0.0536	0.05%	99.85%	
1A4ai	0.0402	0.03%	99.88%	
1A2e	0.0298	0.03%	99.91%	
5B1	0.0278	0.02%	99.93%	
1A3bii	0.0237	0.02%	99.95%	
3B4d	0.0194	0.02%	99.97%	
1A3biii	0.0179	0.02%	99.99%	
2G	0.0146	0.01%	100.00%	
1A4cii	0.0011	0.00%	100.00%	
1A3biv	0.0004	0.00%	100.00%	
1A3c	0.0002	0.00%	100.00%	

Level Assessment

NFR	CO (kt)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	102.5618			
1A3bi	42.6586	41.59%	41.59%	x
1A4bi	20.4442	19.93%	61.53%	x
1A1a	18.2510	17.80%	79.32%	x
1A2f	4.7517	4.63%	83.95%	x
1A3biii	3.0485	2.97%	86.93%	
1A2gviii	2.2396	2.18%	89.11%	
1A3bii	1.8421	1.80%	90.91%	
1A2e	1.7100	1.67%	92.57%	
1A3ai(i)	1.4970	1.46%	94.03%	
1A4ai	1.3125	1.28%	95.31%	
1A3biv	1.2946	1.26%	96.58%	
1A4cii	1.0798	1.05%	97.63%	
1A2b	0.7275	0.71%	98.34%	
1A3dii	0.6147	0.60%	98.94%	
1A3c	0.3773	0.37%	99.31%	
2G	0.1941	0.19%	99.49%	
1A2c	0.1445	0.14%	99.64%	
1A4ciii	0.1379	0.13%	99.77%	
1A1c	0.0857	0.08%	99.85%	
5B1	0.0649	0.06%	99.92%	
1A4ci	0.0267	0.03%	99.94%	
1A1b	0.0245	0.02%	99.97%	
1A3ei	0.0120	0.01%	99.98%	
1A3aii(i)	0.0116	0.01%	99.99%	
1A2d	0.0085	0.01%	100.00%	
1A2a	0.0012	0.00%	100.00%	
5C1bv	0.0007	0.00%	100.00%	
5C1bi	0.0005	0.00%	100.00%	

Annex A.2 Table 4: Key Category Analysis for Ammonia and Carbon Monoxide (continued)

Trend Assessment

NFR	NH3, 1990 (kt)	NH3, 2016 (kt)	Trend (magnitude)	Trend %	% Cumulative	Key Category
3Da1	13.6914	11.1916	0.03063	31.91%	31.91%	x
3B1b	27.4055	31.2812	0.01955	20.37%	52.28%	x
1A3bi	0.0330	0.7420	0.00644	6.71%	58.99%	x
3B2	1.7224	1.1271	0.00641	6.68%	65.67%	x
3B3	3.8461	4.7194	0.00574	5.98%	71.65%	x
3Da3	13.9244	14.1945	0.00554	5.77%	77.42%	x
3B4gii	0.9746	1.4940	0.00417	4.35%	81.77%	x
3Da2b	0.0261	0.4423	0.00378	3.93%	85.70%	
3Da2a	32.2364	34.6055	0.00306	3.19%	88.88%	
3B4e	0.5262	0.7875	0.00208	2.16%	91.05%	
3B4h	0.6034	0.4158	0.00206	2.14%	93.19%	
3B4giii	0.9972	0.8408	0.00200	2.08%	95.27%	
3B4gi	0.6415	0.8414	0.00145	1.51%	96.78%	
1A4bi	0.1479	0.0614	0.00087	0.91%	97.69%	
1A2gviii	0.0941	0.1752	0.00068	0.71%	98.40%	
3B1a	12.7005	13.4341	0.00061	0.64%	99.04%	
3B4giv	0.1056	0.0799	0.00029	0.31%	99.35%	
1A3bii	0.0039	0.0237	0.00018	0.19%	99.54%	
3B4d	0.0340	0.0194	0.00015	0.16%	99.70%	
2G	0.0290	0.0146	0.00015	0.15%	99.85%	
1A3biii	0.0045	0.0179	0.00012	0.12%	99.97%	
3B4f	0.0483	0.0536	0.00002	0.02%	99.99%	
1A4cii	0.0013	0.0011	0.00000	0.00%	100.00%	
1A3biv	0.0001	0.0004	0.00000	0.00%	100.00%	
1A3c	0.0003	0.0002	0.00000	0.00%	100.00%	
NATIONAL T	109.7978	116.7234	0.00000	0.00%	100.00%	

Annex A.2 Table 5: Key Category Analysis for Total Suspended Particulates (TSP) and Particulate Matter <10 µm in Diameter (PM₁₀)

Level Assessment					Level Assessment				
NFR	TSP (kt)	% Contribution Level	% Cumulative	Key Category	NFR	PM10 (kt)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTA	63.6280				NATIONAL TOTAL	29.05633			
2D3b	28.5000	44.79%	44.79%	x	3Da1	6.93763	23.88%	23.88%	x
1A4bi	7.5750	11.91%	56.70%	x	1A4bi	6.92137	23.82%	47.70%	x
3Da1	6.9376	10.90%	67.60%	x	2D3b	3.80000	13.08%	60.78%	x
2A5a	3.7163	5.84%	73.44%	x	2A5a	1.82173	6.27%	67.04%	x
3B1b	2.4478	3.85%	77.29%	x	3B1b	1.10537	3.80%	70.85%	x
3B1a	2.0084	3.16%	80.44%	x	1A3bvi	1.06554	3.67%	74.52%	x
2A5b	1.5429	2.42%	82.87%		3B1a	0.91521	3.15%	77.67%	x
1A3bvi	1.4186	2.23%	85.10%		1A2f	0.83727	2.88%	80.55%	x
3B3	1.2511	1.97%	87.07%		1A1a	0.67320	2.32%	82.86%	
1A3bvii	1.1632	1.83%	88.89%		1A3bvii	0.58161	2.00%	84.87%	
1A2f	0.9204	1.45%	90.34%		1A3bi	0.54781	1.89%	86.75%	
1A1a	0.8150	1.28%	91.62%		2A5b	0.46297	1.59%	88.34%	
3B2	0.6678	1.05%	92.67%		1A2gviii	0.41496	1.43%	89.77%	
3B4gi	0.6304	0.99%	93.66%		1A2e	0.31596	1.09%	90.86%	
1A3bi	0.5478	0.86%	94.52%		3B2	0.28621	0.99%	91.85%	
3B4gii	0.4927	0.77%	95.30%		1A4ai	0.26754	0.92%	92.77%	
1A2gviii	0.4482	0.70%	96.00%		1A3bii	0.26721	0.92%	93.69%	
1A2e	0.3527	0.55%	96.56%		3B4gii	0.24636	0.85%	94.53%	
1A4ai	0.3297	0.52%	97.07%		1A3biii	0.19997	0.69%	95.22%	
1A3bii	0.2672	0.42%	97.49%		3B3	0.18269	0.63%	95.85%	
3Dd	0.2311	0.36%	97.86%		5E	0.16514	0.57%	96.42%	
1A3biii	0.2000	0.31%	98.17%		3B4giii	0.14000	0.48%	96.90%	
5E	0.1651	0.26%	98.43%		3B4gi	0.13272	0.46%	97.36%	
3Dc	0.1412	0.22%	98.65%		1A3dii	0.12460	0.43%	97.79%	
3B4giii	0.1400	0.22%	98.87%		1A4cii	0.10658	0.37%	98.15%	
1A3dii	0.1246	0.20%	99.07%		2G	0.09779	0.34%	98.49%	
1A4cii	0.1066	0.17%	99.24%		1A1c	0.06830	0.24%	98.72%	
2G	0.0981	0.15%	99.39%		3Dd	0.05777	0.20%	98.92%	
1A1c	0.0709	0.11%	99.50%		1A3c	0.05077	0.17%	99.10%	
1A3c	0.0536	0.08%	99.59%		3Dc	0.04518	0.16%	99.25%	
3B4e	0.0443	0.07%	99.65%		3B4giv	0.03883	0.13%	99.39%	
3B4giv	0.0388	0.06%	99.72%		2D3i	0.03069	0.11%	99.49%	
2D3i	0.0375	0.06%	99.77%		1A4ciii	0.02796	0.10%	99.59%	
1A2c	0.0310	0.05%	99.82%		1A2c	0.02486	0.09%	99.67%	
1A4ciii	0.0280	0.04%	99.87%		1A2b	0.02220	0.08%	99.75%	
1A2b	0.0230	0.04%	99.90%		3B4e	0.02028	0.07%	99.82%	
1A3ai(i)	0.0196	0.03%	99.93%		1A3ai(i)	0.01962	0.07%	99.89%	
1A4ci	0.0183	0.03%	99.96%		1A4ci	0.01433	0.05%	99.94%	
1A1b	0.0055	0.01%	99.97%		1A1b	0.00547	0.02%	99.96%	
1A3biv	0.0049	0.01%	99.98%		1A3biv	0.00494	0.02%	99.97%	
3B4h	0.0036	0.01%	99.99%		1A2d	0.00239	0.01%	99.98%	
3B4f	0.0031	0.00%	99.99%		3B4h	0.00158	0.01%	99.99%	
1A2d	0.0030	0.00%	99.99%		3B4f	0.00147	0.01%	99.99%	
3B4d	0.0014	0.00%	100.00%		1A3aii(i)	0.00071	0.00%	99.99%	
1A3aii(i)	0.0007	0.00%	100.00%		3B4d	0.00059	0.00%	100.00%	
1A3ei	0.0005	0.00%	100.00%		1A3ei	0.00050	0.00%	100.00%	
5A	0.0004	0.00%	100.00%		5A	0.00018	0.00%	100.00%	
5C1bv	0.0002	0.00%	100.00%		5C1bv	0.00014	0.00%	100.00%	
5C1bi	0.0001	0.00%	100.00%		5C1bi	0.00005	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%		1A2a	0.00003	0.00%	100.00%	

Annex A.2 Table 6: Key Category Analysis for Particulate Matter <2.5 µm in Diameter (PM_{2.5})

Level Assessment					Trend Assessment						
NFR	PM2.5 (kt)	% Contribution Level	% Cumulative	Key Category	NFR	PM2.5 1990 (kt)	PM2.5 2015 (kt)	Trend (magnitude)	Trend %	% Cumulative	Key Category
NATIONAL TOTAL	15.4677				1A4bi	23.9144	6.8154	0.10502	41.99%	41.99%	x
1A4bi	6.8154	44.06%	44.06%	x	2D3b	2.2000	1.9000	0.02651	10.60%	52.58%	x
2D3b	1.9000	12.28%	56.35%	x	1A2f	0.4861	0.7608	0.01555	6.22%	58.80%	x
1A2f	0.7608	4.92%	61.26%	x	1A3bvi	0.2164	0.5789	0.01375	5.50%	64.30%	x
3B1b	0.7393	4.78%	66.04%	x	3B1b	0.7655	0.7393	0.01145	4.58%	68.87%	x
3B1a	0.5982	3.87%	69.91%	x	3B1a	0.5398	0.5982	0.01026	4.10%	72.97%	x
1A3bvi	0.5789	3.74%	73.65%	x	1A3bi	0.5317	0.5478	0.00893	3.57%	76.54%	x
1A3bi	0.5478	3.54%	77.20%	x	1A2gviii	0.2501	0.3940	0.00807	3.23%	79.77%	x
1A1a	0.4397	2.84%	80.04%	x	1A4cii	0.8735	0.1066	0.00788	3.15%	82.92%	x
1A2gviii	0.3940	2.55%	82.59%		1A3bvii	0.1178	0.3168	0.00753	3.01%	85.93%	
1A3bvii	0.3168	2.05%	84.63%		1A3bii	1.1464	0.2672	0.00673	2.69%	88.62%	
1A2e	0.2848	1.84%	86.47%		1A1a	0.6458	0.4397	0.00443	1.77%	90.39%	
1A3bii	0.2672	1.73%	88.20%		3Da1	0.2665	0.2668	0.00425	1.70%	92.09%	
3Da1	0.2668	1.73%	89.93%		2A5a	0.1575	0.1822	0.00321	1.28%	93.37%	
1A4ai	0.2165	1.40%	91.33%		1A2e	0.4313	0.2848	0.00271	1.08%	94.45%	
1A3biii	0.2000	1.29%	92.62%		1A3dii	0.1218	0.1163	0.00178	0.71%	95.17%	
2A5a	0.1822	1.18%	93.80%		1A2a	0.1326	0.0000	0.00166	0.66%	95.83%	
5E	0.1651	1.07%	94.87%		1A2b	0.1692	0.0216	0.00150	0.60%	96.43%	
1A3dii	0.1163	0.75%	95.62%		1A4ai	0.5621	0.2165	0.00087	0.35%	96.77%	
1A4cii	0.1066	0.69%	96.31%		1A2c	0.1120	0.0197	0.00084	0.34%	97.11%	
2G	0.0965	0.62%	96.93%		1A1c	0.0445	0.0477	0.00080	0.32%	97.43%	
3B2	0.0954	0.62%	97.55%		2A5b	0.0417	0.0463	0.00079	0.32%	97.75%	
1A3c	0.0483	0.31%	97.86%		3B2	0.1604	0.0954	0.00071	0.28%	98.03%	
1A1c	0.0477	0.31%	98.17%		1A3c	0.0575	0.0483	0.00065	0.26%	98.29%	
2A5b	0.0463	0.30%	98.47%		3B4gii	0.0161	0.0246	0.00050	0.20%	98.49%	
1A4ciii	0.0261	0.17%	98.64%		1A3biii	0.4161	0.2000	0.00048	0.19%	98.68%	
3B4giii	0.0255	0.16%	98.80%		2D3i	0.0120	0.0205	0.00043	0.17%	98.86%	
3B4gii	0.0246	0.16%	98.96%		1A3ai(i)	0.0149	0.0196	0.00037	0.15%	99.01%	
1A2b	0.0216	0.14%	99.10%		2G	0.1913	0.0965	0.00035	0.14%	99.15%	
2D3i	0.0205	0.13%	99.23%		3B4giii	0.0302	0.0255	0.00035	0.14%	99.28%	
1A2c	0.0197	0.13%	99.36%		1A4ciii	0.0383	0.0261	0.00026	0.10%	99.39%	
1A3ai(i)	0.0196	0.13%	99.48%		3B4e	0.0086	0.0129	0.00026	0.10%	99.49%	
3B4e	0.0129	0.08%	99.57%		3B4gi	0.0056	0.0100	0.00021	0.09%	99.58%	
1A4ci	0.0110	0.07%	99.64%		5E	0.3594	0.1651	0.00020	0.08%	99.66%	
3B4gi	0.0100	0.06%	99.70%		3Dd	0.0079	0.0092	0.00016	0.07%	99.73%	
3Dd	0.0092	0.06%	99.76%		3B3	0.0065	0.0081	0.00015	0.06%	99.78%	
3B3	0.0081	0.05%	99.82%		1A4ci	0.0149	0.0110	0.00013	0.05%	99.84%	
3Dc	0.0056	0.04%	99.85%		1A3biv	0.0031	0.0049	0.00010	0.04%	99.88%	
3B4giv	0.0055	0.04%	99.89%		1A1b	0.0067	0.0055	0.00007	0.03%	99.90%	
1A1b	0.0055	0.04%	99.92%		3Dc	0.0072	0.0056	0.00007	0.03%	99.93%	
1A3biv	0.0049	0.03%	99.96%		3B4giv	0.0073	0.0055	0.00007	0.03%	99.96%	
1A2d	0.0019	0.01%	99.97%		1B1a	0.0020	0.0018	0.00003	0.01%	99.97%	
1B1a	0.0018	0.01%	99.98%		1A2d	0.0060	0.0019	0.00002	0.01%	99.98%	
3B4f	0.0009	0.01%	99.98%		3B4f	0.0008	0.0009	0.00002	0.01%	99.98%	
3B4h	0.0008	0.01%	99.99%		3B4h	0.0008	0.0008	0.00001	0.00%	99.99%	
1A3aii(i)	0.0007	0.00%	99.99%		1A3aii(i)	0.0026	0.0007	0.00001	0.00%	99.99%	
1A3ei	0.0005	0.00%	100.00%		1A3ei	0.0002	0.0005	0.00001	0.00%	100.00%	
3B4d	0.0002	0.00%	100.00%		5C1bv	0.0000	0.0001	0.00000	0.00%	100.00%	
5C1bv	0.0001	0.00%	100.00%		3B4d	0.0003	0.0002	0.00000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%		5C1bi	0.0001	0.0000	0.00000	0.00%	100.00%	
5C1bi	0.0000	0.00%	100.00%		5A	0.0001	0.0000	0.00000	0.00%	100.00%	
5A	0.0000	0.00%	100.00%		2A5c	0.0000	0.0000	0.00000	0.00%	100.00%	
2A5c	0.0000	0.00%	100.00%		NATIONAL T	35.1909	15.4677	0.00000	0.00%	100.00%	

Annex A.2 Table 7: Key Category Analysis for Lead and Cadmium

Level Assessment

NFR	Pb (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	13.4001			
1A3bi	8.0127	59.80%	59.80%	x
1A4bi	2.0813	15.53%	75.33%	x
1A3bvi	1.3990	10.44%	85.77%	x
1A2f	0.6675	4.98%	90.75%	
1A1a	0.5183	3.87%	94.62%	
1A2e	0.2039	1.52%	96.14%	
1A4ai	0.1894	1.41%	97.55%	
1A2gviii	0.1694	1.26%	98.82%	
1A3biv	0.0468	0.35%	99.17%	
2G	0.0218	0.16%	99.33%	
1A2c	0.0165	0.12%	99.45%	
1A3bii	0.0144	0.11%	99.56%	
1A3dii	0.0108	0.08%	99.64%	
1A4ci	0.0107	0.08%	99.72%	
5C1bi	0.0097	0.07%	99.79%	
1A1b	0.0069	0.05%	99.84%	
1A3aii(i)	0.0064	0.05%	99.89%	
1A1c	0.0063	0.05%	99.94%	
1A4ciii	0.0024	0.02%	99.96%	
1A2b	0.0021	0.02%	99.97%	
1A2d	0.0017	0.01%	99.98%	
1A3biii	0.0015	0.01%	99.99%	
1A3ai(i)	0.0003	0.00%	100.00%	
2C7c	0.0002	0.00%	100.00%	
5C1bv	0.0002	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	

Level Assessment

NFR	Cd (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	0.2843			
1A1a	0.0653	22.98%	22.98%	x
1A2gviii	0.0623	21.93%	44.90%	x
1A4bi	0.0410	14.42%	59.32%	x
1A3bi	0.0306	10.77%	70.10%	x
1A2f	0.0304	10.71%	80.81%	x
1A4ai	0.0171	6.02%	86.83%	
1A2e	0.0133	4.67%	91.50%	
1A3bvi	0.0066	2.33%	93.83%	
1A3bii	0.0060	2.11%	95.94%	
1A3biii	0.0037	1.30%	97.24%	
1A1b	0.0028	0.97%	98.21%	
1A4cii	0.0014	0.49%	98.70%	
1A3dii	0.0008	0.29%	98.99%	
1A1c	0.0008	0.27%	99.26%	
5C1bi	0.0007	0.26%	99.53%	
1A3c	0.0004	0.12%	99.65%	
1A2c	0.0003	0.11%	99.76%	
1A4ci	0.0002	0.07%	99.83%	
1A4ciii	0.0002	0.07%	99.89%	
1A3biv	0.0001	0.05%	99.94%	
2G	0.0000	0.01%	99.96%	
1A2b	0.0000	0.01%	99.97%	
1A2d	0.0000	0.01%	99.98%	
5C1bv	0.0000	0.01%	99.99%	
1A3ai(i)	0.0000	0.01%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	

Annex A.2 Table 8: Key Category Analysis for Mercury and Arsenic

Level Assessment

NFR	Hg (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	0.3293			
1A1a	0.1117	33.93%	33.93%	x
1A4bi	0.0883	26.83%	60.75%	x
1A2f	0.0317	9.62%	70.37%	x
1A3bi	0.0236	7.18%	77.55%	x
5A	0.0209	6.36%	83.91%	x
1A2b	0.0134	4.06%	87.97%	
1A2e	0.0123	3.73%	91.70%	
1A2gviii	0.0080	2.43%	94.13%	
5C1bv	0.0075	2.29%	96.42%	
1A4ai	0.0036	1.10%	97.52%	
1A3dii	0.0025	0.76%	98.28%	
1A2c	0.0020	0.62%	98.89%	
1A1c	0.0012	0.37%	99.26%	
1A1b	0.0006	0.17%	99.43%	
1A4ciii	0.0006	0.17%	99.60%	
1A3ai(i)	0.0005	0.15%	99.75%	
5C1bi	0.0004	0.13%	99.88%	
1A3ei	0.0002	0.06%	99.94%	
1A2d	0.0001	0.03%	99.97%	
1A4ci	0.0001	0.02%	99.99%	
1A2a	0.0000	0.01%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	
2G	0.0000	0.00%	100.00%	

Level Assessment

NFR	As (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	1.2647			
5C1bi	0.6492	51.33%	51.33%	x
1A1a	0.5093	40.27%	91.60%	x
1A4bi	0.0427	3.37%	94.98%	
1A2f	0.0229	1.81%	96.79%	
1A4ai	0.0122	0.97%	97.76%	
1A2e	0.0086	0.68%	98.44%	
1A1c	0.0061	0.48%	98.92%	
1A2gviii	0.0044	0.35%	99.27%	
1A3dii	0.0033	0.26%	99.53%	
1A1b	0.0018	0.14%	99.67%	
1A2c	0.0014	0.11%	99.78%	
1A4ciii	0.0007	0.06%	99.84%	
1A4ci	0.0007	0.05%	99.90%	
1A3bi	0.0006	0.05%	99.94%	
1A3ei	0.0003	0.02%	99.96%	
1A2b	0.0001	0.01%	99.97%	
1A3ai(i)	0.0001	0.01%	99.98%	
1A2d	0.0001	0.01%	99.99%	
5C1bv	0.0001	0.01%	100.00%	
2G	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	

Annex A.2 Table 9: Key Category Analysis for Chromium and Copper

Level Assessment

NFR	Cr (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	2.4907			
5C1bi	0.6274	25.19%	25.19%	x
1A3bvi	0.5170	20.76%	45.95%	x
1A1a	0.3141	12.61%	58.56%	x
1A4bi	0.2150	8.63%	67.19%	x
1A2f	0.1853	7.44%	74.63%	x
1A4ai	0.1531	6.15%	80.77%	x
1A3bi	0.1463	5.88%	86.65%	
1A2gviii	0.1422	5.71%	92.36%	
1A2e	0.0831	3.34%	95.70%	
1A3bii	0.0308	1.23%	96.93%	
1A3biii	0.0215	0.86%	97.80%	
1A2c	0.0132	0.53%	98.33%	
1A1b	0.0106	0.42%	98.75%	
1A4ci	0.0085	0.34%	99.09%	
1A4cii	0.0069	0.28%	99.37%	
1A3dii	0.0042	0.17%	99.54%	
1A1c	0.0039	0.16%	99.69%	
1A3c	0.0018	0.07%	99.76%	
1A2b	0.0017	0.07%	99.83%	
1A2d	0.0014	0.05%	99.88%	
1A4ciii	0.0009	0.04%	99.92%	
1A3ai(i)	0.0008	0.03%	99.95%	
1A3biv	0.0006	0.02%	99.98%	
2G	0.0004	0.02%	100.00%	
5C1bv	0.0001	0.00%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	

Level Assessment

NFR	Cu (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	20.1585			
1A3bvi	11.3071	56.09%	56.09%	x
1A3bi	5.1926	25.76%	81.85%	x
1A3bii	1.0236	5.08%	86.93%	
1A3biii	0.6263	3.11%	90.03%	
1A1a	0.5431	2.69%	92.73%	
1A4bi	0.3640	1.81%	94.53%	
5C1bi	0.3462	1.72%	96.25%	
1A4cii	0.2355	1.17%	97.42%	
1A2f	0.1299	0.64%	98.06%	
1A4ai	0.0785	0.39%	98.45%	
1A3dii	0.0731	0.36%	98.82%	
1A3c	0.0599	0.30%	99.11%	
1A2e	0.0497	0.25%	99.36%	
1A2gviii	0.0471	0.23%	99.59%	
1A3biv	0.0229	0.11%	99.71%	
1A4ciii	0.0164	0.08%	99.79%	
2G	0.0124	0.06%	99.85%	
1A1b	0.0086	0.04%	99.89%	
1A2c	0.0074	0.04%	99.93%	
1A1c	0.0067	0.03%	99.96%	
1A4ci	0.0048	0.02%	99.99%	
1A2b	0.0009	0.00%	99.99%	
1A3ai(i)	0.0009	0.00%	100.00%	
1A2d	0.0008	0.00%	100.00%	
5C1bv	0.0001	0.00%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	

Annex A.2 Table 10: Key Category Analysis for Nickel and Selenium

Level Assessment

NFR	Ni (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	8.4849			
1A4ai	2.6004	30.65%	30.65%	x
1A2f	2.0183	23.79%	54.43%	x
1A2e	1.0856	12.79%	67.23%	x
1A1a	0.8947	10.55%	77.77%	x
1A2gviii	0.6827	8.05%	85.82%	x
1A2c	0.2671	3.15%	88.97%	
1A3bi	0.2149	2.53%	91.50%	
1A4bi	0.2026	2.39%	93.89%	
1A4ci	0.1733	2.04%	95.93%	
1A3dii	0.0831	0.98%	96.91%	
1A3bvi	0.0827	0.97%	97.88%	
1A3bii	0.0419	0.49%	98.38%	
1A2b	0.0335	0.40%	98.77%	
1A2d	0.0275	0.32%	99.10%	
1A3biii	0.0256	0.30%	99.40%	
1A4ciii	0.0186	0.22%	99.62%	
1A1b	0.0139	0.16%	99.78%	
1A4cii	0.0097	0.11%	99.90%	
1A1c	0.0042	0.05%	99.95%	
1A3c	0.0025	0.03%	99.98%	
1A3biv	0.0010	0.01%	99.99%	
2G	0.0008	0.01%	100.00%	
5C1bi	0.0001	0.00%	100.00%	
5C1bv	0.0001	0.00%	100.00%	
1A3ai(i)	0.0000	0.00%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	

Level Assessment

NFR	Se (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	3.5574			
1A4bi	1.8884	53.08%	53.08%	x
1A1a	1.5656	44.01%	97.09%	x
1A3bi	0.0306	0.86%	97.95%	
1A1c	0.0198	0.56%	98.51%	
1A3bvi	0.0116	0.33%	98.83%	
1A2f	0.0084	0.24%	99.07%	
1A3dii	0.0083	0.23%	99.30%	
1A3bii	0.0060	0.17%	99.47%	
1A3biii	0.0037	0.10%	99.58%	
1A2gviii	0.0032	0.09%	99.67%	
1A2e	0.0029	0.08%	99.75%	
1A4ai	0.0027	0.08%	99.82%	
1A4ciii	0.0019	0.05%	99.87%	
1A1b	0.0017	0.05%	99.92%	
1A4cii	0.0014	0.04%	99.96%	
1A3ai(i)	0.0005	0.01%	99.97%	
1A3c	0.0004	0.01%	99.98%	
1A2c	0.0003	0.01%	99.99%	
1A3biv	0.0001	0.00%	100.00%	
1A4ci	0.0001	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	
1A2d	0.0000	0.00%	100.00%	
1A2b	0.0000	0.00%	100.00%	
1A3aii(i)	0.0000	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	

Annex A.2 Table 11: Key Category Analysis for Zinc

Level Assessment

NFR	Zn (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	20.2043			
1A3bvi	4.5768	22.65%	22.65%	x
1A4bi	4.1609	20.59%	43.25%	x
1A3bi	3.0560	15.13%	58.37%	x
1A2gviii	2.4521	12.14%	70.51%	x
1A2f	1.6545	8.19%	78.70%	x
1A1a	1.4827	7.34%	86.04%	x
1A4ai	0.6506	3.22%	89.26%	
1A2e	0.6251	3.09%	92.35%	
1A3bii	0.6020	2.98%	95.33%	
1A3biii	0.3731	1.85%	97.18%	
1A4cii	0.1385	0.69%	97.86%	
1A3ai(i)	0.1189	0.59%	98.45%	
1A3dii	0.0997	0.49%	98.94%	
1A1b	0.0983	0.49%	99.43%	
1A3c	0.0353	0.17%	99.60%	
1A4ciii	0.0224	0.11%	99.72%	
1A1c	0.0163	0.08%	99.80%	
1A3biv	0.0134	0.07%	99.86%	
1A2c	0.0108	0.05%	99.92%	
2G	0.0072	0.04%	99.95%	
1A4ci	0.0053	0.03%	99.98%	
1A3aii(i)	0.0019	0.01%	99.99%	
1A2b	0.0010	0.01%	99.99%	
1A2d	0.0010	0.00%	100.00%	
2C7c	0.0003	0.00%	100.00%	
2C2	0.0001	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	
1A3ei	0.0000	0.00%	100.00%	

Annex A.2 Table 12: Key Category Analysis for Dioxins and Furans, Polychlorinated Biphenyls and Hexachlorobenzene

Level Assessment

NFR	Dioxin (g I-TEQ)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	21.2236			
1A4bi	13.3017	62.67%	62.67%	x
5E	2.2525	10.61%	73.29%	x
2A 1	1.4762	6.96%	80.24%	x
1A3bi	1.0240	4.82%	85.07%	
1A2f	0.9364	4.41%	89.48%	
1A 1a	0.5180	2.44%	91.92%	
5C2	0.4606	2.17%	94.09%	
1A3bii	0.3102	1.46%	95.55%	
1A2e	0.2666	1.26%	96.81%	
1A3biii	0.1851	0.87%	97.68%	
1A4ai	0.1790	0.84%	98.52%	
1A2gvii	0.1723	0.81%	99.34%	
5A	0.0745	0.35%	99.69%	
2A6	0.0161	0.08%	99.76%	
5C1bi	0.0143	0.07%	99.83%	
1A3dii	0.0108	0.05%	99.88%	
1A 1c	0.0086	0.04%	99.92%	
1A4ci	0.0040	0.02%	99.94%	
1A3biv	0.0036	0.02%	99.96%	
1A2c	0.0033	0.02%	99.97%	
1A4ciii	0.0024	0.01%	99.98%	
1A 1b	0.0011	0.01%	99.99%	
1A3ei	0.0011	0.01%	100.00%	
2G	0.0004	0.00%	100.00%	
1A2b	0.0003	0.00%	100.00%	
1A2d	0.0002	0.00%	100.00%	
5C1bv	0.0001	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	
2L	0.0000	0.00%	100.00%	

Level Assessment

NFR	PCB (kg)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	12.4331			
2A 1	5.1767	41.64%	41.64%	x
1A4bi	2.6739	21.51%	63.14%	x
5E	2.5549	20.55%	83.69%	x
5C2	0.7829	6.30%	89.99%	
1A2f	0.6370	5.12%	95.11%	
1A3bi	0.2932	2.36%	97.47%	
1A2e	0.1473	1.18%	98.65%	
1A3bii	0.0710	0.57%	99.23%	
5A	0.0490	0.39%	99.62%	
1A3biii	0.0354	0.29%	99.90%	
5C1bi	0.0057	0.05%	99.95%	
1A3dii	0.0032	0.03%	99.98%	
1A3biv	0.0013	0.01%	99.99%	
1A4ciii	0.0007	0.01%	99.99%	
2L	0.0004	0.00%	100.00%	
1A 1a	0.0002	0.00%	100.00%	
1A 1b	0.0002	0.00%	100.00%	
1A2gvii	0.0001	0.00%	100.00%	
1A4ai	0.0000	0.00%	100.00%	
1A 1c	0.0000	0.00%	100.00%	
1A4ci	0.0000	0.00%	100.00%	

Level Assessment

NFR	HCB (kg)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	1.6767			
3Df	1.1507	68.63%	68.63%	x
1A 1a	0.4586	27.35%	95.98%	x
1A4bi	0.0164	0.98%	96.96%	
1A2f	0.0106	0.63%	97.59%	
1A2gvii	0.0082	0.49%	98.08%	
1A4ai	0.0076	0.45%	98.53%	
1A3dii	0.0066	0.40%	98.93%	
1A 1c	0.0058	0.34%	99.27%	
1A2e	0.0046	0.27%	99.54%	
5C1bi	0.0037	0.22%	99.77%	
1A4ciii	0.0015	0.09%	99.86%	
1A3bi	0.0010	0.06%	99.92%	
5C1bv	0.0008	0.05%	99.96%	
1A3bii	0.0003	0.02%	99.98%	
1A3biii	0.0002	0.01%	99.99%	
1A4ci	0.0001	0.01%	100.00%	
1A3biv	0.0000	0.00%	100.00%	

Annex A.2 Table 13: Key Category Analysis for Benzo[a]pyrene and Benzo[b]fluoranthene

Level Assessment

NFR	B(a)P (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	4.1593			
1A4bi	3.7104	89.21%	89.21%	x
1A2f	0.2018	4.85%	94.06%	
1A2e	0.0614	1.48%	95.54%	
1A3bi	0.0556	1.34%	96.87%	
1A4ai	0.0444	1.07%	97.94%	
1A2gvii	0.0281	0.68%	98.62%	
1A3bii	0.0160	0.38%	99.00%	
5E	0.0133	0.32%	99.32%	
2A6	0.0105	0.25%	99.57%	
1A4cii	0.0055	0.13%	99.71%	
1A2c	0.0045	0.11%	99.82%	
1A3biii	0.0038	0.09%	99.91%	
1A4ci	0.0013	0.03%	99.94%	
1A3c	0.0011	0.03%	99.96%	
2G	0.0004	0.01%	99.97%	
1A2b	0.0004	0.01%	99.98%	
1A2d	0.0003	0.01%	99.99%	
5C2	0.0001	0.00%	100.00%	
1A1a	0.0001	0.00%	100.00%	
1A3biv	0.0001	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	
1A1b	0.0000	0.00%	100.00%	
5C1bi	0.0000	0.00%	100.00%	
1A1c	0.0000	0.00%	100.00%	
5C1bv	0.0000	0.00%	100.00%	

Level Assessment

NFR	B(b)F (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTA	6.2807			
1A4bi	5.2786	84.05%	84.05%	x
1A2f	0.3626	5.77%	89.82%	
1A4ai	0.2260	3.60%	93.42%	
1A2e	0.1504	2.40%	95.81%	
1A2gvii	0.0924	1.47%	97.28%	
1A3bi	0.0624	0.99%	98.28%	
1A2c	0.0257	0.41%	98.69%	
1A3biii	0.0233	0.37%	99.06%	
1A3bii	0.0179	0.29%	99.34%	
5E	0.0158	0.25%	99.59%	
1A4ci	0.0100	0.16%	99.75%	
1A4cii	0.0055	0.09%	99.84%	
1A2b	0.0026	0.04%	99.88%	
1A1a	0.0026	0.04%	99.92%	
1A2d	0.0020	0.03%	99.96%	
1A3c	0.0018	0.03%	99.98%	
5C2	0.0006	0.01%	99.99%	
2G	0.0002	0.00%	100.00%	
1A2a	0.0001	0.00%	100.00%	
1A3biv	0.0001	0.00%	100.00%	
1A1c	0.0000	0.00%	100.00%	
1A1b	0.0000	0.00%	100.00%	
5C1bi	0.0000	0.00%	100.00%	

Annex A.2 Table 14: Key Category Analysis for Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene

Level Assessment

NFR	B(k)F (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTA	2.4055			
1A4bi	2.0796	86.45%	86.45%	x
1A2f	0.1107	4.60%	91.06%	
1A3bi	0.0485	2.02%	93.07%	
1A4ai	0.0447	1.86%	94.93%	
1A2e	0.0409	1.70%	96.63%	
1A3biii	0.0260	1.08%	97.72%	
1A2gvii	0.0230	0.95%	98.67%	
1A3bii	0.0141	0.58%	99.25%	
1A2c	0.0057	0.24%	99.49%	
5E	0.0056	0.23%	99.72%	
1A1a	0.0020	0.08%	99.81%	
2A6	0.0017	0.07%	99.88%	
1A4ci	0.0011	0.05%	99.93%	
5C2	0.0006	0.03%	99.95%	
1A2b	0.0005	0.02%	99.97%	
1A2d	0.0003	0.01%	99.99%	
2G	0.0002	0.01%	99.99%	
1A2a	0.0000	0.00%	100.00%	
1A3biv	0.0000	0.00%	100.00%	
1A1c	0.0000	0.00%	100.00%	
5C1bi	0.0000	0.00%	100.00%	
1A1b	0.0000	0.00%	100.00%	

Level Assessment

NFR	I(123-cd)P (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTA	2.0553			
1A4bi	1.7880	86.99%	86.99%	x
1A2f	0.0880	4.28%	91.28%	
1A3bi	0.0535	2.60%	93.88%	
1A4ai	0.0412	2.00%	95.88%	
1A2e	0.0346	1.68%	97.57%	
1A2gvii	0.0206	1.00%	98.57%	
1A3bii	0.0149	0.72%	99.29%	
1A3biii	0.0060	0.29%	99.58%	
1A2c	0.0054	0.26%	99.85%	
1A4ci	0.0010	0.05%	99.89%	
5E	0.0009	0.04%	99.94%	
1A2b	0.0004	0.02%	99.96%	
1A2d	0.0003	0.02%	99.98%	
2G	0.0002	0.01%	99.99%	
1A1a	0.0002	0.01%	99.99%	
1A3biv	0.0001	0.00%	100.00%	
1A2a	0.0000	0.00%	100.00%	
1A1b	0.0000	0.00%	100.00%	
1A1c	0.0000	0.00%	100.00%	
5C1bv	0.0000	0.00%	100.00%	

Annex A.2 Table 15: Key Category Analysis for Polycyclic Aromatic Hydrocarbons

Level Assessment

NFR	Total PAH (t)	% Contribution Level	% Cumulative	Key Category
NATIONAL TOTAL	14.9008			
1A4bi	12.8566	86.28%	86.28%	x
1A2f	0.7632	5.12%	91.40%	
1A4ai	0.3563	2.39%	93.79%	
1A2e	0.2874	1.93%	95.72%	
1A3bi	0.2200	1.48%	97.20%	
1A2gvii	0.1640	1.10%	98.30%	
1A3bii	0.0628	0.42%	98.72%	
1A3biii	0.0592	0.40%	99.12%	
1A2c	0.0413	0.28%	99.40%	
5E	0.0357	0.24%	99.64%	
1A4ci	0.0134	0.09%	99.73%	
2A6	0.0122	0.08%	99.81%	
1A4cii	0.0111	0.07%	99.88%	
1A1a	0.0049	0.03%	99.92%	
1A2b	0.0039	0.03%	99.94%	
1A2d	0.0030	0.02%	99.96%	
1A3c	0.0028	0.02%	99.98%	
5C2	0.0014	0.01%	99.99%	
2G	0.0010	0.01%	100.00%	
1A3biv	0.0003	0.00%	100.00%	
1A2a	0.0002	0.00%	100.00%	
1A1c	0.0001	0.00%	100.00%	
1A1b	0.0000	0.00%	100.00%	
5C1bi	0.0000	0.00%	100.00%	
5C1bv	0.0000	0.00%	100.00%	

Annex A.2 Table 16: Key Category Analysis for All Pollutants

Pollutant	Key Categories									Total (%)
NO _x	1A3bi 15.16%	1A3biii 11.66%	3Da1 11.62%	3Da3 10.62%	1A3bii 8.18%	1A1a 7.13%	1A2f 6.28%	1A3dii 5.81%	3Da2a 5.29%	81.75%
CO	1A3bi 41.59%	1A4bi 19.93%	1A1a 17.80%	1A2f 4.63%						83.95%
NM VOC	3B1b 24.67%	2H2 19.61%	2D3a 9.93%	3B1a 8.57%	1A4bi 7.43%	3De 3.53%	2D3d 2.69%	1B2aiv 2.52%	1A3bi 2.36%	81.31%
SO _x	1A4bi 48.39%	1A1a 27.00%	1A2f 9.73%							85.12%
NH ₃	3Da2a 29.65%	3B1b 26.80%	3Da3 12.16%	3B1a 11.51%						80.12%
TSP	2D3b 44.79%	1A4bi 11.91%	3Da1 10.90%	2A5a 5.84%	3B1b 3.85%	3B1a 3.16%				80.44%
PM ₁₀	3Da1 23.88%	1A4bi 23.82%	2D3b 13.08%	2A5a 6.27%	3B1b 3.80%	1A3bvi 3.67%	3B1a 3.15%	1A2f 2.88%		80.55%
PM _{2.5}	1A4bi 44.06%	2D3b 12.28%	1A2f 4.92%	3B1b 4.78%	3B1a 3.87%	1A3bvi 3.74%	1A3bi 3.54%	1A1a 2.84%		80.04%
Pb	1A3bi 59.80%	1A4bi 15.53%	1A3bvi 10.44%							85.77%
Cd	1A1a 22.98%	1A2gviii 21.93%	1A4bi 14.42%	1A3bi 10.77%	1A2f 10.71%					80.81%
Hg	1A1a 33.93%	1A4bi 26.83%	1A2f 9.62%	1A3bi 7.18%	5A 6.36%					83.91%
As	5C1bi 51.33%	1A1a 40.27%								91.60%
Cr	5C1bi 25.19%	1A3bvi 20.76%	1A1a 12.61%	1A4bi 8.63%	1A2f 7.44%	1A4ai 6.15%				80.77%
Cu	1A3bvi 56.09%	1A3bi 25.76%								81.85%
Ni	1A4ai 30.65%	1A2f 23.79%	1A2e 12.79%	1A1a 10.55%	1A2gviii 8.05%					85.82%
Se	1A4bi 53.08%	1A1a 44.01%								97.09%
Zn	1A3bvi 22.65%	1A4bi 20.59%	1A3bi 15.13%	1A2gviii 12.14%	1A2f 8.19%	1A1a 7.34%				86.04%
PCDD/F	1A4bi 62.67%	5E 10.61%	2A1 6.96%							80.24%
PCBs	2A1 41.64%	1A4bi 21.51%	5E 20.55%							83.69%
HCb	3Df 68.63%	1A1a 27.35%								95.98%
PAHs	1A4bi 86.28%									86.28%
	1 Energy		2 IPPU		3 Agriculture		5 Waste			

Annex A.3

Fuel Tourism in Road Transport and Nitrogen Oxides Emissions Based on Fuels Used

Introduction

Fuel tourism is the term given to the retail purchase of petrol or diesel in one country that is subsequently used in another country. Because of the significant price differentials between the Republic of Ireland (ROI) and the United Kingdom (primarily due to higher UK Excise Tax) and the proximity of population centres in Northern Ireland, the impact of fuel tourism has been significant in the Republic of Ireland for many years. In regards to the calculation and reporting of transboundary emissions to air that arise from road transport, the reporting protocols under the Convention on Long-Range Transboundary Air Pollution (CLRTAP) and the EU National Emission Ceilings Directive provide that a Party can make adjustments to its emission estimates to account for this phenomenon. The following sections outline how the extent of fuel tourism is quantified in Ireland and provides the results for the years 1990–2016.

In the 2018 submission fuel tourism figures were provided by the Department of Communications, Climate Action and Environment (DCCAE). Petrol fuel tourism was estimated at 0.98 per cent of petrol sales in the ROI in 2016.

The customer base for diesel is broader than for petrol and it is the primary fuel consumed by the commercial sector, particularly for road freight. In recent years, diesel has also begun to attain a larger share of the private car fuel trade, particularly in light of the bias of vehicle taxation systems towards diesel cars, which generally consume less fuel in relative terms than petrol cars. The regression method used estimated diesel fuel tourism at 13.2 per cent of diesel sales in the Republic of Ireland in 2016.

The approach to estimating fuel tourism is based on log-linear OLS regression of fuel consumption in the ROI against some relevant indicator variables, including the relative price of road transport fuels between the ROI and the UK. For both petrol and diesel, after running the regression, the relative prices are re-set to zero and a new estimate for consumption is derived. Fuel tourism is then estimated as the difference between these two variables for consumption. For diesel the following variables are used.

- Relative Price
- Number of HGVs
- Dummy Variable for Year (to allow for efficiency gains over time)

The cross-price elasticity of demand for diesel between the UK and ROI is calculated as being 0.66 in this case, reflecting the greater carrying capacity of many diesel vehicles and the fact that a sizable proportion of diesel vehicles are used in a commercial rather than a domestic context. For petrol the equation is somewhat simpler (compared to the diesel estimate and includes

- Relative Price
- Number of Passenger Cars

- Dummy Variable for time set at zero for all years to 2008 and increasing sequentially thereafter

This latter variable accounts for the change to the method of applying car taxation during 2008 in ROI, where engine size was replaced by CO₂ emissions which in turn led to a sizable switch away from petrol and towards diesel cars. The cross-price elasticity in terms of petrol is smaller, at 0.21, reflecting the lower volumes of commercial traffic which use petrol as a fuel of choice

Fuel Tourism 1990–2016 Time Series

The approach outlined above has been used to estimate fuel tourism in Ireland in 1987 and for the period 1990–2016, and Figure A3.1 shows the results of the analysis. Figure A3.1 indicates that the level of fuel tourism is substantial, particularly in the case of diesel for the years 1998–2007. These results are used to produce adjusted annual emission estimates for all substances in Ireland's 2018 submission under CLRTAP, i.e. estimates of emissions based on fuels used in the country. The adjusted emissions are most relevant for Ireland in the case of nitrogen oxides (NO_x) as assessment in relation to obligations under the Sofia Protocol on NO_x emissions is undertaken with respect to emissions estimated on the basis of fuels used in Ireland. The adjusted NO_x emissions are given in Table A3.1 and are shown in Figure A3.2, which also shows Ireland's target NO_x level under the Sofia Protocol.

Fuel Tourism in Ireland 1990-2016




Figure A3.1. Percentage Fuel Tourism in Ireland 1987, 1990–2016




Figure A3.2. Emissions of Nitrogen Oxides in 1987 and 1990–2016 Based on Fuels Used in Ireland

Table A3.1. Emissions of Nitrogen Oxides in 1987 and 1990–2016 Based on Fuels Used in Ireland

	1987	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Public Electricity and Heat Production	40.142	46.374	46.188	53.065	46.944	45.100	41.391	41.864	40.192	39.384	38.769	39.720	41.145	37.621	33.812	32.333	32.384	29.874	27.330	22.022	13.320	11.516	7.995	10.303	8.569	7.575	9.530	8.000
Residential & Commercial/Institutional	7.238	10.386	10.413	9.387	9.320	9.226	8.625	8.899	8.508	8.856	8.494	8.597	8.792	8.665	8.874	8.720	9.220	8.953	8.883	9.677	9.256	9.462	8.289	8.045	8.026	7.210	7.442	7.540
Manufacturing Industries and Construction	9.207	8.928	8.728	7.388	7.786	7.777	8.050	8.127	8.975	8.980	8.988	10.529	9.413	10.883	13.580	16.009	17.313	16.592	18.417	15.733	10.338	9.935	8.455	10.407	10.530	10.711	10.853	11.312
Agriculture/Forestry/Fishing	8.703	8.760	9.368	9.844	10.385	11.764	14.310	11.881	11.957	12.346	12.557	12.913	13.075	12.447	13.164	12.874	12.758	11.590	10.528	10.269	8.570	7.428	6.560	6.195	5.552	4.842	4.274	4.054
Transport	59.604	66.072	65.575	67.938	65.093	61.953	60.614	65.222	57.752	58.438	57.282	54.132	55.223	50.583	51.970	54.780	56.832	57.840	55.187	53.973	47.421	42.886	40.958	40.006	40.593	42.410	40.879	43.977
Agriculture	0.000	32.895	32.817	32.647	33.359	34.376	35.385	35.496	34.577	36.982	36.815	34.568	32.980	32.640	33.568	32.584	31.915	31.298	30.104	29.551	29.242	30.961	28.234	28.881	31.339	30.609	30.769	31.766
Other NFR sectors	2.524	1.635	2.318	2.498	1.615	0.971	0.963	0.945	1.030	1.198	1.100	1.256	1.466	1.322	1.156	1.130	1.268	1.158	1.141	1.179	1.030	1.134	0.859	0.886	0.825	0.785	0.645	0.650
Total	127.418	175.050	175.407	182.767	174.501	171.168	169.339	172.435	162.991	166.183	164.004	161.714	162.095	154.161	156.123	158.429	161.690	157.303	151.590	142.404	119.177	113.321	101.351	104.722	105.434	104.142	104.393	107.300

Annex B

Expanded Energy Balance Sheet for 2016

Table B.1 Expanded Energy Balance Sheet 2016

2016	Units = ktoe	NACE (Rev 2)	Coal	Bituminous Coal	Anthractive + Manufactured Ovoids	Coke	Lignite \ Brown Coal Briquettes	Peat	Milled Peat	Sod Peat	Briquettes	Oil	Crude	Refinery Gas	Gasoline	Kerosene	Jet Kerosene	Fueloil	LPG	Gasoil / Diesel / DERV	Petroleum Coke	Naphta	Bitumen	White Spirit	Lubricants	Natural Gas	Renewables	Hydro	Wind	Biomass & Renewable Waste	Landfill Gas	Biogas	Liquid Biofuel	Solar	Geothermal	Non-Renewable Waste	Electricity	Heat	TOTAL			
Indigenous Production			0	0			679	551	128			0														2,493	1,026	59	529	291	39	16	24	14	55	66			4,265			
Imports			1,155	1,084	61		10	0				9,009	3,270	0	795	520	1,122	60	136	2,684	149	0	230	2	40	1,704	139			46			93				75		12,082			
Exports			9	0	8		2	5			5	1,643	0		371	13	0	1,039	17	117	0	76	3	0	8	0	0			0			0				136		1,793			
Mar. Bunkers			0				0					160						11		149						0												160				
Stock Change			228	235	-7		-0	59	68	0	-8	-34	2		-6	4	-2	-4	-0	-26	1	-2	0	0	0	54	-8			-1			-7						299			
Primary Energy Supply (incl non-energy)			1,373	1,319	47	0	8	734	619	128	-13	7,173	3,272	0	417	512	1,120	-994	119	2,392	151	-78	228	2	33	4,251	1,157	59	529	335	39	16	111	14	55	66	-61	0	14,694			
Primary Energy Requirement (excl. non-e			1,373	1,319	47	0	8	734	619	128	-13	6,911	3,272	0	417	512	1,120	-994	119	2,392	151	-78	0	0	0	4,251	1,157	59	529	335	39	16	111	14	55	66	-61	0	14,432			
Transformation Input			1,101	1,101	0	0	0	607	607	0	0	3,340	3,272	5	0	0	0	52	0	11	0	0	0	0	0	2,396	160	0	0	114	39	7	0	0	0	24	56	0	7,684			
Public Thermal Power Plants			1,101	1,101			513	513	0			63						52		11						2,068	150			111	39				24			3,920				
Combined Heat and Power Plants			0	0			9	9				5			5			0	0	0						273	10			3		7						297				
Pumped Storage Consumption																																					46		46			
Briquetting Plants			0				85	85				0														0												85				
Oil Refineries & other energy sector			0				0					3,272	3,272												54	0											10		3,336			
Transformation Output			0	0	0	0	0	81	0	0	81	3,319	0	92	590	185	0	1,046	48	1,278	0	80	0	0	0	0	58	0	0	41	14	4	0	0	0	6	2,036	0	5,500			
Public Thermal Power Plants			0				0					0													53				39	14					6	1,822			1,822			
Combined Heat and Power Plants - Electricity			0				0					0													5				1		4						189		189			
Combined Heat and Power Plants - Heat																									0													0				
Pumped Storage Generation																																						25		25		
Briquetting Plants							81				81	0														0													81			
Oil Refineries							0					3,319		92	590	185	0	1,046	48	1,278		80																	3,319			
Exchanges and transfers			12	-7	18	0	0	0	0	0	0	-15	0	0	-0	198	-198	1	0	-5	-12	0	0	0	0	0	-588	-59	-529	0	0	0	0	-0	0	0	588	0	-3			
Electricity																										-588	-59	-529							-0			588		0		
Heat																																							0			
Other			12	-7	18							-15			-0	198	-198	1		-5	-12						0												-3			
Own Use and Distribution Losses							12	12				88		87				0	0	0						51	0											254		405		
Available Final Energy Consumption			284	211	65	0	8	196	0	128	68	7,049	0	0	1,007	895	922	1	167	3,654	139	2	228	2	33	1,804	410	0	0	221	0	9	111	14	55	42	2,253	0	12,038			
Non-Energy Consumption			0	0	0	0	0	0	0	0	0	262	0	0	0	0	0	0	0	0	0				228	2	33	0	0	0	0	0	0	0	0	0	0	0	0	262		
Final non-Energy Consumption (Feedstocks)			0				0					262														0													262			
Total Final Energy Consumption			289	211	69	0	9	198	1	128	69	6,740	0	0	1,003	906	868	44	163	3,616	139	0	0	0	0	1,794	426	0	0	230	0	9	118	14	55	42	2,199	0	11,688			
Industry*			110	110	0		1	1	0	0	0	484	0	0	0	91	0	35	113	113	133	0	0	0	0	755	174	0	0	172		2	0	0	0	42	872	0	2,438			
Non-Energy Mining	05-09		0	0			0					31				3		1	0	27	0				12	0											63		105			
Food & beverages	10-11		21	21			1	1				131				53		20	32	25	0				102	22			19		2						186		463			
Textiles and textile products	13-14		0	0			0					2				1		0	0	1	0				1	0												11		15		
Wood and wood products	16		0	0			0					2				0		0	0	2	0				2	113			113								37		155			
Pulp, paper, publishing and printing	17-18		0	0			0					3				1		0	0	1	0				3	0											20		27			
Chemicals & man-made fibres	20-21		0	0			0					27				13		5	3	7	0				63	0			0								159		249			
Rubber and plastic products	22		0	0			0					9				0		0	6	3	0				4	0											38		52			
Other non-metallic mineral products	23		89	89			0					182				9		3	1	35	133				16	39			39						42	56		425				
Basic metals and fabricated metal products	24-25		0	0			0					8				0		1	5	2	0				417	0											68		493			
Machinery and equipment n.e.c.	28		0	0			0					5				0		0	2	2	0				5	0											22		33			
Electrical and optical equipment	26-27		0	0			0					41				0		0	39	1	0				120	0												108		269		
Transport equipment manufacture	29-30		0	0			0					5				0		0	4	1	0				2	0												18		25		
Other manufacturing	31-33, 12 & 15		0	0			0			0		37				10		4	20	4	0				6	0											85		128			
Transport			0	0	0	0	0	0	0	0	0	4,825	0	0	1,003	0	868	0	3	2,951	0	0	0	0	0	21	118	0	0	0	0	0	118	0	0	0	0	4	0	4,969		
Road Freight			0				0					713								713						22													735			
Road Light Goods Vehicle												309								309						9													318			
Road Private Car			0				0					2,076				831			3	1,242						65										0		2,141				
Public Passenger Services			0				0					131				13				117						4													135			
Rail			0				0					36								36						0												4		40		
Domestic Aviation			0				0					4				1										0													4			
International Aviation			0				0					866					866									0													866			
Fuel Tourism			0				0					372				10				363																			384			
Navigation			0				0					86								86																			86			
Unspecified			0				0					232				148										21	7												261			
Residential			179	101	69		9	197		128	69	1,005			0	815		0	39	145	6					563	82			32												

Annex C

Emission Factors for Energy (NFR 1)

Table C.1 Emission Factors for NFR 1A1a

Emission Factors																		
	Code	Name					Inventory Year											
NFR Source Category	1.A.1.a	Public Electricity and Heat Production					2016											
Fuel	Coal			Peat			Oil			Natural Gas			MSW Incineration			Landfill Gas		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	67.549	g/GJ	PS	92.846	g/GJ	PS	27.556	g/GJ	PS	28.354	g/GJ	PS	0.977	kg/Mg waste	PS	89.000	g/GJ	EMEP/EEA 2016
SO _x	48.203	g/GJ	PS	53.778	g/GJ	PS	108.455	g/GJ	PS	-	g/GJ	PS	0.238	kg/Mg waste	PS	-	g/GJ	EMEP/EEA 2016
NM/OC	0.900	g/GJ	EMEP/EEA 2016	0.900	g/GJ	EMEP/EEA 2016	2.30	g/GJ	EMEP/EEA 2016	2.60	g/GJ	EMEP/EEA 2016	0.006	kg/Mg waste	EMEP/EEA 2016	2.60	g/GJ	EMEP/EEA 2016
CO	313.000	g/GJ	EMEP/EEA 2016	13.000	g/GJ	EMEP/EEA 2016	15.100	g/GJ	EMEP/EEA 2016	39.000	g/GJ	EMEP/EEA 2016	0.041	kg/Mg waste	PS	39.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA			NA			NA		
TSP	8.400	g/GJ	EMEP/EEA 2016	10.200	g/GJ	EMEP/EEA 2016	35.400	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016	0.003*	kg/Mg waste	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
PM ₁₀	7.700	g/GJ	EMEP/EEA 2016	6.900	g/GJ	EMEP/EEA 2016	25.200	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016	0.003*	kg/Mg waste	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
PM _{2.5}	5.200	g/GJ	EMEP/EEA 2016	2.800	g/GJ	EMEP/EEA 2016	19.300	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016	0.003*	kg/Mg waste	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
BC	0.022	f-PM _{2.5}	EMEP/EEA 2016	NE		EMEP/EEA 2016	0.056	f-PM _{2.5}	EMEP/EEA 2016	0.025	f-PM _{2.5}	EMEP/EEA 2016	0.033	f-PM _{2.5}	EMEP/EEA 2016	0.025	f-PM _{2.5}	EMEP/EEA 2016
Pb	7.300	mg/GJ	EMEP/EEA 2016	7.300	mg/GJ	EMEP/EEA 2016	4.560	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016	0.058	g/Mg waste	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016
Cd	0.900	mg/GJ	EMEP/EEA 2016	0.900	mg/GJ	EMEP/EEA 2016	1.200	mg/GJ	EMEP/EEA 2016	0.003	mg/GJ	EMEP/EEA 2016	0.005	g/Mg waste	EMEP/EEA 2016	0.003	mg/GJ	EMEP/EEA 2016
Hg	1.400	mg/GJ	EMEP/EEA 2016	1.400	mg/GJ	EMEP/EEA 2016	0.341	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2016	0.019	g/Mg waste	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2016
As	7.100	mg/GJ	EMEP/EEA 2016	7.100	mg/GJ	EMEP/EEA 2016	3.980	mg/GJ	EMEP/EEA 2016	0.210	mg/GJ	EMEP/EEA 2016	0.006	g/Mg waste	EMEP/EEA 2016	0.210	mg/GJ	EMEP/EEA 2016
Cr	4.500	mg/GJ	EMEP/EEA 2016	4.500	mg/GJ	EMEP/EEA 2016	2.550	mg/GJ	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016	0.016	g/Mg waste	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016
Cu	7.800	mg/GJ	EMEP/EEA 2016	7.800	mg/GJ	EMEP/EEA 2016	5.310	mg/GJ	EMEP/EEA 2016	0.076	mg/GJ	EMEP/EEA 2016	0.014	g/Mg waste	EMEP/EEA 2016	0.076	mg/GJ	EMEP/EEA 2016
Ni	4.900	mg/GJ	EMEP/EEA 2016	4.900	mg/GJ	EMEP/EEA 2016	255.000	mg/GJ	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016	0.022	g/Mg waste	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016
Se	23.000	mg/GJ	EMEP/EEA 2016	23.000	mg/GJ	EMEP/EEA 2016	2.060	mg/GJ	EMEP/EEA 2016	0.011	mg/GJ	EMEP/EEA 2016	0.012	g/Mg waste	EMEP/EEA 2016	0.011	mg/GJ	EMEP/EEA 2016
Zn	19.000	mg/GJ	EMEP/EEA 2016	19.000	mg/GJ	EMEP/EEA 2016	87.800	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016	0.025	g/Mg waste	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016
PCB	3.300	ng/GJ	EMEP/EEA 2016	3.300	ng/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	0.003	µg/Mg waste	EMEP/EEA 2016	NE		EMEP/EEA 2016
PCDD/F (I-TEQ)	10.000	ng/GJ	EMEP/EEA 2016	0.525	ng/GJ	PS	2.500	ng/GJ	EMEP/EEA 2016	0.500	ng/GJ	EMEP/EEA 2016	0.0525*	µg I-TEQ/Mg waste	EMEP/EEA 2016	0.500	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	0.700	µg/GJ	EMEP/EEA 2016	0.700	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	0.560	µg/GJ	EMEP/EEA 2016	0.008	mg/Mg waste	EMEP/EEA 2016	0.560	µg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	37.000	µg/GJ	EMEP/EEA 2016	37.000	µg/GJ	EMEP/EEA 2016	4.500	µg/GJ	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016	0.018	mg/Mg waste	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016
Benzo[k]fluoranthene	29.000	µg/GJ	EMEP/EEA 2016	29.000	µg/GJ	EMEP/EEA 2016	4.500	µg/GJ	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016	0.010	mg/Mg waste	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	1.100	µg/GJ	EMEP/EEA 2016	1.100	µg/GJ	EMEP/EEA 2016	6.920	µg/GJ	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016	0.012	mg/Mg waste	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016
Total 4 PAHs	67.800	µg/GJ	EMEP/EEA 2016	67.800	µg/GJ	EMEP/EEA 2016	15.920	µg/GJ	EMEP/EEA 2016	3.080	µg/GJ	EMEP/EEA 2016	0.047	mg/Mg waste	EMEP/EEA 2016	3.080	µg/GJ	EMEP/EEA 2016
HCB	6.700	µg/GJ	EMEP/EEA 2016	6.700	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	0.045	mg/Mg waste	EMEP/EEA 2016	NE	µg/GJ	EMEP/EEA 2016

Table C.2 Emission Factors for NFR 1A1b

Emission Factors															
	Code	Name				Inventory Year									
NFR Source Category	1.A.1.b	Petroleum Refining				2016									
Fuel	Refinery Gas			Fuel Oil			Natural Gas			Gasoil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	44.616	g/GJ	PS	142.000	g/GJ	EMEP/EEA 2016	63.000	g/GJ	EMEP/EEA 2016	65.000	g/GJ	EMEP/EEA 2016	63.000	g/GJ	EMEP/EEA 2016
SO _x	5.119	g/GJ	PS	NA			0.043	g/GJ	CS	9.005	g/GJ	PS	0.281	g/GJ	EMEP/EEA 2016
NM/OC	2.580	g/GJ	EMEP/EEA 2016	2.300	g/GJ	EMEP/EEA 2016	2.580	g/GJ	EMEP/EEA 2016	0.650	g/GJ	EMEP/EEA 2016	2.580	g/GJ	EMEP/EEA 2016
CO	3.917	g/GJ	PS	NA			3.917	g/GJ	PS	16.200	g/GJ	EMEP/EEA 2016	39.300	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA			NA		
TSP	0.890	g/GJ	EMEP/EEA 2016	20.000	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016	6.470	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
PM ₁₀	0.890	g/GJ	EMEP/EEA 2016	15.000	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016	3.230	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
PM _{2.5}	0.890	g/GJ	EMEP/EEA 2016	9.000	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016	0.808	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
BC	0.184	f-PM _{2.5}	EMEP/EEA 2016	0.056	f-PM _{2.5}	EMEP/EEA 2016	0.086	f-PM _{2.5}	EMEP/EEA 2016	0.335	f-PM _{2.5}	EMEP/EEA 2016	0.086	f-PM _{2.5}	EMEP/EEA 2016
Pb	1.790	mg/GJ	EMEP/EEA 2016	4.560	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016	4.070	mg/GJ	EMEP/EEA 2016	0.0015	mg/GJ	EMEP/EEA 2016
Cd	0.712	mg/GJ	EMEP/EEA 2016	1.200	mg/GJ	EMEP/EEA 2016	0.003	mg/GJ	EMEP/EEA 2016	1.360	mg/GJ	EMEP/EEA 2016	0.0003	mg/GJ	EMEP/EEA 2016
Hg	0.086	mg/GJ	EMEP/EEA 2016	0.341	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2016	1.360	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2016
As	0.343	mg/GJ	EMEP/EEA 2016	3.980	mg/GJ	EMEP/EEA 2016	0.210	mg/GJ	EMEP/EEA 2016	1.810	mg/GJ	EMEP/EEA 2016	0.120	mg/GJ	EMEP/EEA 2016
Cr	2.740	mg/GJ	EMEP/EEA 2016	2.550	mg/GJ	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016	1.360	mg/GJ	EMEP/EEA 2016	0.00076	mg/GJ	EMEP/EEA 2016
Cu	2.220	mg/GJ	EMEP/EEA 2016	5.310	mg/GJ	EMEP/EEA 2016	0.0001	mg/GJ	EMEP/EEA 2016	2.720	mg/GJ	EMEP/EEA 2016	0.000076	mg/GJ	EMEP/EEA 2016
Ni	3.600	mg/GJ	EMEP/EEA 2016	255.000	mg/GJ	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016	1.360	mg/GJ	EMEP/EEA 2016	0.00051	mg/GJ	EMEP/EEA 2016
Se	0.420	mg/GJ	EMEP/EEA 2016	2.060	mg/GJ	EMEP/EEA 2016	0.011	mg/GJ	EMEP/EEA 2016	6.790	mg/GJ	EMEP/EEA 2016	0.0112	mg/GJ	EMEP/EEA 2016
Zn	25.500	mg/GJ	EMEP/EEA 2016	87.800	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016	1.810	mg/GJ	EMEP/EEA 2016	0.0015	mg/GJ	EMEP/EEA 2016
PCB	NE		EMEP/EEA 2016	24.251	ng/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	23.090	µg/GJ	NA EI 2006	NE		EMEP/EEA 2016
PCDD/F (I-TEQ)	NE		EMEP/EEA 2016	2.500	ng/GJ	EMEP/EEA 2016	0.500	µg/GJ	EMEP/EEA 2016	0.500	ng/GJ	EMEP/EEA 2016	0.500	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	0.669	µg/GJ	EMEP/EEA 2016	19.886	µg/GJ	EMEP/EEA 2016	0.560	µg/GJ	EMEP/EEA 2016	18.934	µg/GJ	NA EI 2006	0.560	µg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	1.140	µg/GJ	EMEP/EEA 2016	4.500	µg/GJ	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016	4.500	µg/GJ	NA EI 2006	0.840	µg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	0.631	µg/GJ	EMEP/EEA 2016	4.500	µg/GJ	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016	4.500	µg/GJ	NA EI 2006	0.840	µg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	0.631	µg/GJ	EMEP/EEA 2016	6.920	µg/GJ	EMEP/EEA 2016	0.840	µg/GJ	EMEP/EEA 2016	6.920	µg/GJ	NA EI 2006	0.840	µg/GJ	EMEP/EEA 2016
Total 4 PAHs	3.071	µg/GJ	EMEP/EEA 2016	35.806	µg/GJ	EMEP/EEA 2016	3.080	µg/GJ	EMEP/EEA 2016	34.854	µg/GJ	NA EI 2006	3.080	µg/GJ	EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016

Table C.3 Emission Factors for NFR 1A1c

Emission Factors									
	Code	Name				Inventory year			
NFR Source Category	1.A.1.c	Manufacture of Solid Fuels and Other Energy Industries				2016			
Fuel	Peat			Liquid fuels			Natural gas		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	316.662	g/GJ	PS	65.000	g/GJ	EMEP/EEA 2016	89.000	g/GJ	EMEP/EEA 2016
SO _x	285.366	g/GJ	PS	9.005	g/GJ	PS	0.043	g/GJ	PS
NM/OC	0.800	g/GJ	EMEP/EEA 2016	0.800	g/GJ	EMEP/EEA 2016	2.600	g/GJ	EMEP/EEA 2016
CO	76.580	g/GJ	PS	16.200	g/GJ	EMEP/EEA 2016	39.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA		
TSP	82.000	g/GJ	EMEP/EEA 2016	6.500	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
PM ₁₀	79.000	g/GJ	EMEP/EEA 2016	3.200	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
PM _{2.5}	55.000	g/GJ	EMEP/EEA 2016	0.800	g/GJ	EMEP/EEA 2016	0.890	g/GJ	EMEP/EEA 2016
BC	NE		EMEP/EEA 2016	0.268	g/GJ	EMEP/EEA 2016	0.022	g/GJ	EMEP/EEA 2016
Pb	7.300	mg/GJ	EMEP/EEA 2016						
Cd	0.900	mg/GJ	EMEP/EEA 2016						
Hg	1.400	mg/GJ	EMEP/EEA 2016						
As	7.100	mg/GJ	EMEP/EEA 2016						
Cr	4.500	mg/GJ	EMEP/EEA 2016						
Cu	7.800	mg/GJ	EMEP/EEA 2016						
Ni	4.900	mg/GJ	EMEP/EEA 2016						
Se	23.000	mg/GJ	EMEP/EEA 2016						
Zn	19.000	mg/GJ	EMEP/EEA 2016						
PCB	3.300	ng/GJ	EMEP/EEA 2016						
PCDD/F (I-TEQ)	10.000	µg/GJ	EMEP/EEA 2016						
Benzo[a]pyrene	0.700	µg/GJ	EMEP/EEA 2016						
Benzo[b]fluoranthene	37.000	µg/GJ	EMEP/EEA 2016						
Benzo[k]Fluoranthene	29.000	µg/GJ	EMEP/EEA 2016						
Indeno[1,2,3-cd]pyrene	11.000	µg/GJ	EMEP/EEA 2016						
Total 4 PAHs	77.700	µg/GJ	EMEP/EEA 2016						
HCB	6.700	µg/GJ	EMEP/EEA 2016						

Table C.4 Emission Factors for NFR 1A2a

Emission Factors												
	Code	Name					Inventory Year					
NFR Source Category	1.A.2.a	Combustion in Manufacturing: Iron and Steel					2016					
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	720.000	g/GJ	CS	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gas oil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	NO			74.000	g/GJ	EMEP/EEA 2016	NO		
SO _x	9.005	g/GJ	CS	NO			0.043	g/GJ	CS	NA		
NM/OC	10.000	g/GJ	EMEP/EEA 2009	NO			23.000	g/GJ	EMEP/EEA 2016	NO		
CO	40.000	g/GJ	EMEP/EEA 2009	NO			29.000	g/GJ	EMEP/EEA 2016	NO		
NH ₃	NA			NO			NA			NO		
TSP	27.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	NO		
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	NO		
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	NO		
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	NO			0.040	f-PM _{2.6}	EMEP/EEA 2016	NO		

Table C.5 Emission Factors for NFR 1A2b

Emission Factors												
	Code	Name					Inventory Year					
NFR Source Category	1.A.2.b	Combustion in Manufacturing: Non-ferrous Metals					2016					
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	720.000	g/GJ	CS	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM VOC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	47.900	g/GJ	CS	NO		
SO _x	9.005	g/GJ	CS	968.297	g/GJ	CS	0.043	g/GJ	CS	NA		
NM VOC	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016	NO		
CO	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016	NO		
NH ₃	NA			NA			NA			NO		
TSP	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	NO		
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	NO		
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	NO		
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016	NO		

Table C.6 Emission Factors for NFR 1A2c

Emission Factors												
NFR Source Category	Code	Name					Inventory Year					
1.A.2.c		Combustion in Manufacturing: Chemicals					2016					
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	720.000	g/GJ	CS	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	NO			74.000	g/GJ	EMEP/EEA 2016	91.000	g/GJ	EMEP/EEA 2016
SO _x	9.005	g/GJ	CS	NO			0.043	g/GJ	CS	11.000	g/GJ	EMEP/EEA 2016
NM/OC	10.000	g/GJ	EMEP/EEA 2009	NO			23.000	g/GJ	EMEP/EEA 2016	300.000	g/GJ	EMEP/EEA 2016
CO	40.000	g/GJ	EMEP/EEA 2009	NO			29.000	g/GJ	EMEP/EEA 2016	570.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NO			NA			37.000	g/GJ	EMEP/EEA 2016
TSP	27.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	150.000	g/GJ	EMEP/EEA 2016
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	143.000	g/GJ	EMEP/EEA 2016
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	140.000	g/GJ	EMEP/EEA 2016
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	NO			0.040	f-PM _{2.5}	EMEP/EEA 2016	0.280	f-PM _{2.5}	EMEP/EEA 2016

Table C.7 Emission Factors for NFR 1A2d

Emission Factors												
	Code	Name					Inventory Year					
NFR Source Category	1.A.2.d	Combustion in Manufacturing: Pulp, Paper and Print					2016					
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	720.000	g/GJ	CS	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	NO			74.000	g/GJ	EMEP/EEA 2016	NO		
SO _x	9.005	g/GJ	CS	NO			0.043	g/GJ	CS	NA		
NM/OC	10.000	g/GJ	EMEP/EEA 2009	NO			23.000	g/GJ	EMEP/EEA 2016	NO		
CO	40.000	g/GJ	EMEP/EEA 2009	NO			29.000	g/GJ	EMEP/EEA 2016	NO		
NH ₃	NA			NO			NA			NO		
TSP	27.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	NO		
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	NO		
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	NO			0.780	g/GJ	EMEP/EEA 2016	NO		
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	NO			0.040	f-PM _{2.5}	EMEP/EEA 2016	NO		

Table C.8 Emission Factors for NFR 1A2e

Emission Factors												
	Code	Name	Inventory Year									
NFR Source Category	1.A.2.e	Combustion in Manufacturing: Food Processing, Beverages and Tobacco	2016									
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	720.000	g/GJ	CS	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016	91.000	g/GJ	EMEP/EEA 2016
SO _x	9.005	g/GJ	CS	968.297	g/GJ	CS	0.043	g/GJ	CS	11.000	g/GJ	EMEP/EEA 2016
NM/OC	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016	300.000	g/GJ	EMEP/EEA 2016
CO	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016	570.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			37.000	g/GJ	EMEP/EEA 2016
TSP	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	150.000	g/GJ	EMEP/EEA 2016
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	143.000	g/GJ	EMEP/EEA 2016
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	140.000	g/GJ	EMEP/EEA 2016
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016	0.280	f-PM _{2.5}	EMEP/EEA 2016

Table C.9 Emission Factors for NFR 1A2f

Emission Factors												
	Code	Name				Inventory Year						
NFR Source Category	1.A.2.f	Combustion in Manufacturing: Non-metallic minerals				2016						
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	613.729	g/GJ	PS	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	158.399	g/GJ	PS	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	CS
NM VOC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gas oil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	778.633	g/GJ	PS	74.000	g/GJ	EMEP/EEA 2016	91.000	g/GJ	EMEP/EEA 2016
SO _x	9.005	g/GJ	CS	118.830	g/GJ	PS	0.043	g/GJ	CS	11.000	g/GJ	EMEP/EEA 2016
NM VOC	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016	300.000	g/GJ	EMEP/EEA 2016
CO	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016	570.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			37.000	g/GJ	EMEP/EEA 2016
TSP	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	150.000	g/GJ	EMEP/EEA 2016
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	143.000	g/GJ	EMEP/EEA 2016
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	140.000	g/GJ	EMEP/EEA 2016
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016	0.280	f-PM _{2.5}	EMEP/EEA 2016

Table C.10 Emission Factors for NFR 1A2g

Emission Factors												
	Code	Name					Inventory Year					
NFR Source Category	1.A.2.g	Combustion in Manufacturing: Other					2016					
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	720.000	g/GJ	EMEP/EEA 2016	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			NA		
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016	91.000	g/GJ	EMEP/EEA 2016
SO _x	9.005	g/GJ	CS	47.000	g/GJ	EMEP/EEA 2009	0.043	g/GJ	CS	11.000	g/GJ	EMEP/EEA 2016
NM/OC	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016	300.000	g/GJ	EMEP/EEA 2016
CO	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016	570.000	g/GJ	EMEP/EEA 2016
NH ₃	NA			NA			NA			37.000	g/GJ	EMEP/EEA 2016
TSP	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	150.000	g/GJ	EMEP/EEA 2016
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	143.000	g/GJ	EMEP/EEA 2016
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	140.000	g/GJ	EMEP/EEA 2016
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016	0.280	f-PM _{2.5}	EMEP/EEA 2016

Table C.11 Emission Factors for NFR 1A2 (Heavy Metals and Persistent Organic Pollutants)

Emission Factors (Heavy Metals and POPs)												
	Code	Name				Inventory Year						
NFR Source Category	1.A.2 (a-g)	Combustion in Manufacturing				2016						
Fuel	Coal			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	134.000	mg/GJ	EMEP/EEA 2016	16.000	mg/GJ	EMEP/EEA 2009	16.000	mg/GJ	EMEP/EEA 2009	0.011	mg/GJ	EMEP/EEA 2016
Cd	1.800	mg/GJ	EMEP/EEA 2016	0.300	mg/GJ	EMEP/EEA 2009	0.300	mg/GJ	EMEP/EEA 2009	0.0009	mg/GJ	EMEP/EEA 2016
Hg	7.900	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2009	0.540	mg/GJ	EMEP/EEA 2016
As	4.000	mg/GJ	EMEP/EEA 2016	1.000	mg/GJ	EMEP/EEA 2009	1.000	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2016
Cr	13.500	mg/GJ	EMEP/EEA 2016	12.800	mg/GJ	EMEP/EEA 2009	12.800	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016
Cu	17.500	mg/GJ	EMEP/EEA 2016	7.200	mg/GJ	EMEP/EEA 2009	7.200	mg/GJ	EMEP/EEA 2009	0.0026	mg/GJ	EMEP/EEA 2016
Ni	13.000	mg/GJ	EMEP/EEA 2016	260.000	mg/GJ	EMEP/EEA 2009	260.000	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016
Se	1.800	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2009	0.058	mg/GJ	EMEP/EEA 2016
Zn	200.000	mg/GJ	EMEP/EEA 2016	8.000	mg/GJ	EMEP/EEA 2009	8.000	mg/GJ	EMEP/EEA 2009	0.730	mg/GJ	EMEP/EEA 2016
PCB	170.000	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
PCDD/F (I-TEQ)	203.000	ng/GJ	EMEP/EEA 2016	1.400	ng/GJ	EMEP/EEA 2016	1.400	ng/GJ	EMEP/EEA 2016	0.520	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	45.500	mg/GJ	EMEP/EEA 2016	1.900	mg/GJ	EMEP/EEA 2016	1.900	mg/GJ	EMEP/EEA 2016	0.720	mg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	58.900	mg/GJ	EMEP/EEA 2016	15.000	mg/GJ	EMEP/EEA 2016	15.000	mg/GJ	EMEP/EEA 2016	2.900	mg/GJ	EMEP/EEA 2016
Benzo[k]fluoranthene	23.700	mg/GJ	EMEP/EEA 2016	1.700	mg/GJ	EMEP/EEA 2016	1.700	mg/GJ	EMEP/EEA 2016	1.100	mg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	18.500	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016	1.080	mg/GJ	EMEP/EEA 2016
Total 4 PAHs	146.600	mg/GJ	EMEP/EEA 2016	20.100	mg/GJ	EMEP/EEA 2016	20.100	mg/GJ	EMEP/EEA 2016	5.800	mg/GJ	EMEP/EEA 2016
HCB	0.620	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016

Emission Factors (Heavy Metals and POPs)												
	Code	Name				Inventory Year						
NFR Source Category	1.A.2 (a-g)	Combustion in Manufacturing				2016						
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	16.000	mg/GJ	EMEP/EEA 2009	16.000	mg/GJ	EMEP/EEA 2009	0.011	mg/GJ	EMEP/EEA 2016	27.000	mg/GJ	EMEP/EEA 2016
Cd	0.300	mg/GJ	EMEP/EEA 2009	0.300	mg/GJ	EMEP/EEA 2009	0.0009	mg/GJ	EMEP/EEA 2016	13.000	mg/GJ	EMEP/EEA 2016
Hg	0.100	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2009	0.540	mg/GJ	EMEP/EEA 2016	0.560	mg/GJ	EMEP/EEA 2016
As	1.000	mg/GJ	EMEP/EEA 2009	1.000	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2016	0.190	mg/GJ	EMEP/EEA 2016
Cr	12.800	mg/GJ	EMEP/EEA 2009	12.800	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016	23.000	mg/GJ	EMEP/EEA 2016
Cu	7.200	mg/GJ	EMEP/EEA 2009	7.200	mg/GJ	EMEP/EEA 2009	0.0026	mg/GJ	EMEP/EEA 2016	6.000	mg/GJ	EMEP/EEA 2016
Ni	260.000	mg/GJ	EMEP/EEA 2009	260.000	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016	2.000	mg/GJ	EMEP/EEA 2016
Se	0.100	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2009	0.058	mg/GJ	EMEP/EEA 2016	0.500	mg/GJ	EMEP/EEA 2016
Zn	8.000	mg/GJ	EMEP/EEA 2009	8.000	mg/GJ	EMEP/EEA 2009	0.730	mg/GJ	EMEP/EEA 2016	512.000	mg/GJ	EMEP/EEA 2016
PCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	0.060	µg/GJ	EMEP/EEA 2016
PCDD/F (I-TEQ)	1.400	ng/GJ	EMEP/EEA 2016	1.400	ng/GJ	EMEP/EEA 2016	0.520	ng/GJ	EMEP/EEA 2016	100.000	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	1.900	mg/GJ	EMEP/EEA 2016	1.900	mg/GJ	EMEP/EEA 2016	0.720	mg/GJ	EMEP/EEA 2016	10.000	mg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	15.000	mg/GJ	EMEP/EEA 2016	15.000	mg/GJ	EMEP/EEA 2016	2.900	mg/GJ	EMEP/EEA 2016	16.000	mg/GJ	EMEP/EEA 2016
Benzo[k]fluoranthene	1.700	mg/GJ	EMEP/EEA 2016	1.700	mg/GJ	EMEP/EEA 2016	1.100	mg/GJ	EMEP/EEA 2016	5.000	mg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	1.500	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016	1.080	mg/GJ	EMEP/EEA 2016	4.000	mg/GJ	EMEP/EEA 2016
Total 4 PAHs	20.100	mg/GJ	EMEP/EEA 2016	20.100	mg/GJ	EMEP/EEA 2016	5.800	mg/GJ	EMEP/EEA 2016	35.000	mg/GJ	EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	5.000	µg/GJ	EMEP/EEA 2016

Table C.12 Distances between airport pairs used to estimate fuel consumption for cruise phase

Nautical Miles		Cork	Galway	Donegal	Dublin	Knock	Kerry	Shannon	Sligo	Waterford	Other
		EICK	EICM	EIDL	EIDW	EIKN	EIKY	EINN	EISG	EIWF	
EICK	Cork		89.18	192.52	124.89	124.88	43.37	54.12	146.58	56.04	89.18
EICM	Galway	89.18		106.92	96.28	36.93	70.51	35.94	60.13	95.09	89.18
EIDL	Donegal	192.52	106.92		121.75	70.16	177.15	142.26	46.80	177.42	89.18
EIDW	Dublin	124.89	96.28	121.75		95.56	139.93	105.34	97.52	79.89	89.18
EIKN	Knock	124.88	36.93	70.16	95.56		106.99	72.70	23.53	121.02	89.18
EIKY	Kerry	43.37	70.51	177.15	139.93	106.99		38.25	130.45	89.97	89.18
EINN	Shannon	54.12	35.94	142.26	105.34	72.70	38.25		95.53	74.21	89.18
EISG	Sligo	146.58	60.13	46.80	97.52	23.53	130.45	95.53		137.05	89.18
EIWF	Waterford	56.04	95.09	177.42	79.89	121.02	89.97	74.21	137.05		89.18
	Other	89.18	89.18	89.18	89.18	89.18	89.18	89.18	89.18	89.18	

Table C.13 Number of Domestic and International LTOs

Domestic LTOs No.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	7657	6944	6885	5921	6153	7235	7742	8134	8991	10183	11018	10947	10849	11261	12143	9976	10392	10803	9611	7844	6074	3331	2190	2101	2058	1980	2118
ex Cork	2872	2604	2582	2221	2307	2713	2903	3050	3372	3819	4132	4106	4069	4223	4438	3649	3721	4608	3919	2872	1861	809	445	441	382	259	210
ex Shannon	1737	1576	1562	1343	1396	1641	1757	1845	2040	2310	2500	2484	2462	2555	2865	2809	2892	2277	1897	1349	1077	834	764	800	696	636	596
ex Galway	1425	1293	1282	1102	1145	1347	1441	1514	1674	1895	2051	2038	2019	2096	2224	1631	1615	1815	1848	1563	1746	1252	51	31	NO	11	1
ex Sligo	620	562	557	479	498	586	627	658	728	824	892	886	878	912	946	759	748	754	785	741	678	381	35	25	24	20	21
ex Donegal	581	527	523	449	467	549	588	617	682	773	836	831	824	855	717	684	747	736	754	739	697	721	733	723	732	725	732
ex Knock	445	404	400	344	358	421	450	473	523	592	641	637	631	655	753	565	557	568	481	510	454	253	79	83	67	60	62
ex Kerry	1133	1027	1019	876	910	1070	1145	1203	1330	1506	1630	1620	1605	1666	1755	1477	1515	1506	1418	1170	1048	460	781	776	775	778	782
ex Waterford	236	214	213	183	190	223	239	251	278	314	340	338	335	348	254	181	191	279	456	231	472	707	175	155	67	68	87
ex Other	347	314	312	268	279	328	350	368	407	461	499	496	491	510	539	495	518	411	476	305	282	277	241	282	191	205	152
Total	17053	15465	15334	13187	13703	16113	17242	18115	20024	22679	24538	24381	24164	25080	26634	22226	22896	23757	21645	17324	14389	9025	5494	5417	4992	4742	4761

International LTOs No.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	55852	55883	54508	53675	59073	57901	63622	68297	73636	77422	81886	84365	82626	80766	95337	80726	85400	92464	94445	79668	73356	76791	78842	82481	87460	96269	104503
ex Cork	13865	13901	10350	6557	6392	7320	8806	8593	8128	9284	10842	10893	9673	11686	13860	13456	13210	13176	13462	10840	10664	11007	11122	10459	10235	9469	10457
ex Shannon	22415	20094	17513	16318	15058	16365	17806	15141	17258	19383	20308	18090	13898	14191	14831	15811	16504	17172	16075	14037	10493	10785	9601	9328	10870	9561	10074
ex Galway	1625	1586	1453	1350	1421	1439	1592	1624	1747	1872	1994	2000	1873	1881	1915	2139	1989	2937	2174	1627	1353	793	69	87	NO	49	0
ex Sligo	62	61	56	52	54	55	61	62	67	72	76	77	72	72	27	34	36	164	204	36	54	45	21	23	12	8	13
ex Donegal	203	198	181	168	177	179	198	202	218	233	248	249	233	234	121	147	268	320	410	218	260	291	264	299	330	210	222
ex Knock	2061	2011	1843	1713	1802	1825	2019	2059	2216	2374	2529	2536	2376	2386	2203	2381	2571	2378	2726	2728	2593	2930	3275	3179	3187	2667	3135
ex Kerry	1352	1319	1209	1124	1182	1198	1325	1351	1454	1557	1659	1664	1559	1566	1846	1626	1675	1622	1751	1429	1658	1538	1246	1378	1238	1200	1254
ex Waterford	1218	1188	1089	1012	1064	1078	1193	1217	1309	1402	1494	1498	1404	1410	1161	1122	1351	1800	2152	1502	1404	1468	1452	536	496	740	483
ex Other	615	600	550	511	538	545	603	615	661	708	755	757	709	712	943	695	854	849	862	419	712	747	702	745	659	621	662
Total	99269	96840	88753	82480	86761	87906	97225	99161	106692	114309	121792	122128	114423	114905	132244	118137	123858	132882	134261	112504	102547	106395	106594	108515	114487	120794	130803

Table C.14 Domestic LTO EFs by aircraft type

Aircraft Type	kg of fuel per LTO	NO_x kg/ LTO	HC kg/ LTO	CO kg/ LTO
A30B	1540.55	23.20	5.54	25.84
A310	1540.55	23.20	5.54	25.84
A320	802.33	10.83	1.92	17.59
A321	802.33	10.83	1.92	17.59
A332	2231.52	36.13	2.11	21.50
A333	2231.52	36.13	2.11	21.50
A343	2231.52	36.13	2.11	21.50
AT43	115.20	1.02	0.00	0.86
AT72	137.00	1.35	0.00	0.86
ATP	569.51	4.19	1.01	9.69
B462	569.51	4.19	1.01	9.69
B463	569.51	4.19	1.01	9.69
B733	825.39	8.25	0.67	11.83
B734	825.39	8.25	0.67	11.83
B737	784.12	7.84	0.63	11.24
B738	763.48	7.64	0.62	10.94
B752	1253.00	19.73	1.23	12.55
B762	1617.09	26.03	0.88	6.08
B763	1617.09	26.03	0.88	6.08
B764	1617.09	26.03	0.88	6.08
BE20	51.80	0.24	0.13	0.76
BE40	58.30	0.24	0.23	1.87
CL30	569.51	4.19	1.01	9.69
CL60	569.51	4.19	1.01	9.69
DC10	2381.18	41.71	22.83	61.62
GLF2	569.51	4.19	1.01	9.69
GLF4	569.51	4.19	1.01	9.69
GLF5	569.51	4.19	1.01	9.69
H25B	569.51	4.19	1.01	9.69
LJ31	569.51	4.19	1.01	9.69
LJ45	569.51	4.19	1.01	9.69
LJ60	569.51	4.19	1.01	9.69
MD11	1003.06	12.34	1.92	6.52
MD82	1003.06	12.34	1.92	6.52
MD83	1003.06	12.34	1.92	6.52
T154	2190.00	14.00	75.90	116.81
Other	49.57	0.29	0.20	0.75

Table C.15 International LTO EFs by aircraft type

Aircraft Type	kg of fuel per LTO	NO _x kg/ LTO	HC kg/ LTO	CO kg/ LTO	Aircraft Type	kg of fuel per LTO	NO _x kg/ LTO	HC kg/ LTO	CO kg/ LTO	Aircraft Type	kg of fuel per LTO	NO _x kg/ LTO	HC kg/ LTO	CO kg/ LTO
A30B	1540.55	23.20	5.54	25.84	C10T	30.40	0.17	0.03	0.29	G150	569.51	4.19	1.01	9.69
A109	51.80	0.24	0.13	0.76	C25A	30.40	0.17	0.03	0.29	GALX	3413.87	55.94	37.25	78.23
A124	3413.87	55.94	37.25	78.23	C25B	30.40	0.17	0.03	0.29	GLEX	569.51	4.19	1.01	9.69
A306	1540.55	23.20	5.54	25.84	C56X	30.40	0.17	0.03	0.29	GLF2	569.51	4.19	1.01	9.69
A310	1540.55	23.20	5.54	25.84	C130	306.50	2.22	0.90	1.95	GLF3	569.51	4.19	1.01	9.69
A318	802.33	10.83	1.92	17.59	C160	306.50	2.22	0.90	1.95	GLF4	569.51	4.19	1.01	9.69
A319	802.33	10.83	1.92	17.59	C172	30.40	0.17	0.03	0.29	GLF5	569.51	4.19	1.01	9.69
A320	802.33	10.83	1.92	17.59	C182	30.40	0.17	0.03	0.29	H25A	569.51	4.19	1.01	9.69
A321	802.33	10.83	1.92	17.59	C208	30.40	0.17	0.03	0.29	H25B	569.51	4.19	1.01	9.69
A332	2231.52	36.13	2.11	21.50	C210	30.40	0.17	0.03	0.29	JS31	45.60	0.38	0.05	0.52
A333	2231.52	36.13	2.11	21.50	C340	50.53	0.29	0.20	0.75	JS41	62.70	0.47	0.09	0.82
A748	569.51	4.19	1.01	9.69	C441	50.53	0.29	0.20	0.75	L101	1412.83	12.57	65.40	26.37
AC90	50.53	0.29	0.20	0.75	C500	50.53	0.29	0.20	0.75	L188	260.50	1.80	0.84	1.81
AJET	569.51	4.19	1.01	9.69	C510	50.53	0.29	0.20	0.75	L410	127.70	1.30	0.00	0.73
AN72	569.51	0.29	1.01	9.69	C525	666.07	5.19	32.86	32.72	LJ31	569.51	4.19	1.01	9.69
AT43	115.20	1.02	0.00	0.86	C550	666.07	5.19	32.86	32.72	LJ35	569.51	4.19	1.01	9.69
AT45	115.20	1.02	0.00	0.86	C551	666.07	5.19	32.86	32.72	LJ45	569.51	4.19	1.01	9.69
AT72	137.00	1.35	0.00	0.86	C560	666.07	5.19	32.86	32.72	LJ60	569.51	4.19	1.01	9.69
ATP	569.51	4.19	1.01	9.69	C650	666.07	5.19	32.86	32.72	MD11	1003.06	12.34	1.92	6.52
B190	61.10	0.26	0.63	2.23	C680	666.07	5.19	32.86	32.72	MD82	1003.06	12.34	1.92	6.52
B350	59.40	0.25	0.23	1.89	C750	666.07	5.19	32.86	32.72	MD83	1003.06	12.34	1.92	6.52
B461	569.51	4.19	1.01	9.69	CL30	569.51	4.19	1.01	9.69	MD87	1003.06	12.34	1.92	6.52
B462	569.51	4.19	1.01	9.69	CL60	569.51	4.19	1.01	9.69	MD90	1003.06	12.34	1.92	6.52
B463	569.51	4.19	1.01	9.69	CN35	666.07	5.19	32.86	32.72	MU2	126.70	1.23	0.00	0.71
B732	919.70	7.97	0.58	4.82	CRJ2	666.07	5.19	32.86	32.72	P46T	51.18	0.29	0.20	0.75
B733	825.39	8.25	0.67	11.83	D228	126.70	1.23	0.00	0.71	P68	51.18	0.29	0.20	0.75
B734	825.39	8.25	0.67	11.83	D328	126.70	1.23	0.00	0.71	PA31	51.18	0.29	0.20	0.75
B735	825.39	8.25	0.67	11.83	DC10	2381.18	41.71	22.83	61.62	PA32	51.18	0.29	0.20	0.75
B736	825.39	8.25	0.67	11.83	DC86	876.10	7.26	0.77	5.35	PA34	51.18	0.29	0.20	0.75
B737	784.12	7.84	0.63	11.24	DC87	876.10	7.26	0.77	5.35	PA44	51.18	0.29	0.20	0.75
B738	763.48	7.64	0.62	10.94	DC93	876.10	7.26	0.77	5.35	PA46	51.18	0.29	0.20	0.75
B742	3413.87	55.94	37.25	78.23	DH8C	189.00	1.89	0.64	1.57	PC9	51.18	0.29	0.20	0.75
B744	3402.16	56.64	1.85	19.50	DH8D	215.40	2.49	0.00	1.15	PC12	51.18	0.29	0.20	0.75
B752	1253.00	19.73	1.23	12.55	E135	569.51	4.19	1.01	9.69	PRM1	569.51	4.19	1.01	9.69
B753	1253.00	19.73	1.23	12.55	E145	569.51	4.19	1.01	9.69	RJ1H	569.51	4.19	1.01	9.69
B762	1617.09	26.03	0.88	6.08	E170	569.51	4.19	1.01	9.69	RJ70	569.51	4.19	1.01	9.69
B763	1617.09	26.03	0.88	6.08	E190	569.51	4.19	1.01	9.69	RJ85	569.51	4.19	1.01	9.69
B764	1617.09	26.03	0.88	6.08	F2TH	569.51	4.19	1.01	9.69	SB20	155.40	1.13	0.04	0.85
B772	2562.84	53.64	22.77	61.38	F27	173.60	0.39	1.73	7.53	SF34	75.20	0.50	0.22	0.43
BE9L	51.80	0.24	0.13	0.76	F50	126.70	1.28	0.00	0.73	SH36	86.00	0.42	0.68	3.21
BE9T	51.80	0.24	0.13	0.76	F70	744.38	5.79	1.42	13.68	SR22	51.18	0.29	0.20	0.75
BE20	51.80	0.24	0.13	0.76	F100	744.38	5.79	1.42	13.68	SW4	46.80	0.39	0.04	0.51
BE40	58.30	0.24	0.23	1.87	F900	569.51	4.19	1.01	9.69	T154	2190.00	14.00	75.90	116.81
BE58	58.30	0.24	0.23	1.87	FA7X	569.51	4.19	1.01	9.69	TBM7	51.18	0.29	0.20	0.75
BE76	58.30	0.24	0.23	1.87	FA50	569.51	4.19	1.01	9.69	TRIN	51.18	0.29	0.20	0.75

Table C.16 Domestic LTO IEFs by airport

IEF kg fuel/LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	539.4	484.8	509.3	615.3	545.3	555.0	559.5	506.5	420.2	398.6	351.2	337.7	382.2	331.8
ex Cork	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	391.7	216.7	238.1	412.6	407.9	443.1	505.2	518.3	415.1	180.8	181.2	189.8	239.8	234.5
ex Shannon	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1361.6	1432.0	1491.0	1476.9	1351.3	1377.9	1287.7	1114.3	816.3	858.9	704.2	740.5	914.9	871.6
ex Galway	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	132.2	117.7	118.4	122.9	146.2	174.4	124.3	121.3	113.6	97.4	83.2	NA	49.6	49.6
ex Sligo	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	112.8	111.2	110.5	113.2	112.5	112.3	114.9	115.0	93.4	53.4	50.3	49.6	49.6	50.4
ex Donegal	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	113.5	112.9	111.4	113.6	113.9	113.2	114.9	114.7	97.7	49.6	49.7	50.3	108.2	112.4
ex Knock	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	198.8	182.9	186.2	231.3	231.1	217.9	180.9	161.5	151.2	233.5	226.5	354.5	290.5	262.3
ex Kerry	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	339.6	137.1	151.2	150.1	141.9	370.3	714.9	712.0	469.0	122.1	118.0	117.7	118.7	133.5
ex Waterford	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	141.4	78.9	83.2	79.2	205.3	293.1	109.4	140.8	115.7	74.1	50.0	57.3	49.6	49.6
ex Other	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	132.8	147.4	142.4	132.3	154.4	118.7	116.8	117.8	142.3	113.9	99.4	120.3	120.6	80.4
Weighted average	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	498.3	446.0	490.7	568.2	494.5	508.2	515.5	453.8	336.8	328.9	289.3	293.0	341.3	313.4

IEF kg NO _x /LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	7.2	6.9	7.3	8.7	7.3	7.2	6.8	6.1	5.2	5.6	4.8	4.6	5.1	4.4
ex Cork	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	3.9	2.1	2.4	4.1	4.0	4.4	5.1	5.3	4.2	1.8	1.8	1.8	2.5	2.4
ex Shannon	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	20.9	21.8	23.0	23.0	20.5	21.1	19.8	16.9	11.9	12.7	10.1	10.8	13.5	13.0
ex Galway	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.0	1.1	1.1	1.3	1.5	1.2	1.1	1.0	0.8	0.5	NA	0.3	0.3
ex Sligo	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0.8	0.3	0.3	0.3	0.3	0.3
ex Donegal	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0.8	0.3	0.3	0.3	0.9	1.0
ex Knock	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	1.8	1.9	2.5	2.4	2.2	1.8	1.5	1.4	2.3	2.2	3.5	2.9	2.5
ex Kerry	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	1.3	1.4	1.4	1.4	3.6	7.1	7.1	4.6	1.1	1.0	1.0	1.1	1.3
ex Waterford	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	0.6	0.6	0.6	1.5	2.2	1.0	1.3	1.1	0.6	0.3	0.4	0.3	0.3
ex Other	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	1.0	1.0	0.9	1.1	0.8	0.8	0.8	1.0	0.8	0.7	0.8	1.2	0.5
Weighted average	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.2	6.9	7.9	6.4	6.4	6.2	5.4	4.0	4.4	3.8	3.8	4.5	4.1

Table C.16 Domestic LTO IEFs by airport (continued)

IEF g HC/LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	486.4	498.7	545.8	518.1	464.7	519.9	461.3	396.1	355.6	362.0	327.7	328.9	258.0	257.7
ex Cork	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	367.2	229.1	259.4	416.3	372.8	378.7	425.7	488.3	410.7	221.1	236.1	283.3	447.7	512.7
ex Shannon	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1467.9	1844.2	1844.1	1435.3	1338.5	1571.7	1205.1	1036.1	692.2	700.8	596.5	529.8	766.8	708.0
ex Galway	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	36.4	12.4	16.7	14.8	65.2	126.1	9.8	10.0	33.9	166.0	251.2	NA	201.2	201.2
ex Sligo	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	18.9	16.9	9.8	9.0	16.8	7.1	5.6	73.9	195.4	201.2	201.2	201.2	201.2
ex Donegal	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	9.3	15.1	8.5	4.7	10.6	4.0	4.1	60.6	201.2	200.6	202.3	22.2	9.1
ex Knock	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	248.7	173.4	334.4	379.7	296.5	250.5	150.8	155.8	250.4	429.9	415.8	587.0	493.5	418.2
ex Kerry	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	240.3	36.6	62.1	57.9	44.1	292.5	595.0	593.8	397.9	30.2	32.3	52.6	36.3	26.1
ex Waterford	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	232.8	169.6	158.5	180.2	429.1	505.6	98.8	87.8	36.9	127.3	199.4	213.3	201.2	201.2
ex Other	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	321.4	339.7	335.2	315.1	349.2	299.7	303.5	307.2	342.6	295.7	278.7	311.7	283.4	248.3
Weighted average	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	465.7	481.7	544.3	507.4	438.3	490.9	425.9	366.7	285.4	316.8	294.8	292.6	267.1	249.6

IEF g CO/LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	5616.1	4772.5	5011.0	5754.5	5419.5	6206.2	6401.4	5747.6	4239.6	3199.9	2796.3	2666.0	2816.8	2478.7
ex Cork	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	5389.0	2618.6	3142.8	5724.7	5545.8	6066.5	7108.6	7515.8	5941.7	2468.9	2536.4	2824.7	4120.1	4268.9
ex Shannon	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	13360.8	15486.7	15727.4	13951.0	12686.2	14136.5	11681.8	9855.9	6080.6	6106.7	5075.6	4591.1	6688.1	5860.2
ex Galway	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	1092.6	902.4	894.4	890.9	1344.6	1870.4	877.2	868.4	859.1	1137.8	1326.3	NA	749.0	749.0
ex Sligo	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	865.7	881.5	865.4	869.4	857.2	866.6	859.3	860.1	821.3	752.3	749.0	749.1	749.1	749.0
ex Donegal	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	865.9	870.3	867.8	871.8	860.3	869.3	861.0	860.9	842.0	749.0	749.4	761.3	850.7	858.1
ex Knock	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2495.3	2246.9	2756.5	2939.7	3218.3	2719.8	1941.7	1644.1	1976.2	3753.1	3646.7	5401.0	4527.9	4009.0
ex Kerry	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	4291.6	1109.5	1305.4	1217.9	1068.6	4846.7	10266.2	10226.5	6569.8	1026.8	994.2	1024.5	929.9	982.3
ex Waterford	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	1891.3	986.0	977.6	1004.6	3381.6	4637.5	979.9	1271.5	868.9	797.3	749.8	882.5	749.0	749.0
ex Other	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2170.1	2426.0	2340.1	2166.0	2504.7	1933.5	1897.7	1922.5	2336.7	1838.1	1605.3	1966.5	1618.6	1278.7
Weighted average	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	5363.0	4603.3	5091.6	5633.5	5140.3	5832.6	5936.2	5199.7	3410.5	2746.0	2455.0	2389.2	2723.7	2428.0

Table C.17 International LTO IEFs by airport

IEF kg fuel/LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	781.5	790.1	786.4	770.7	772.2	789.7	791.8	770.0	771.8	766.0	745.1	735.0	734.4	730.7
ex Cork	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	631.9	580.4	623.4	640.7	651.3	653.6	652.1	622.2	600.8	597.4	588.2	555.0	539.9	532.7
ex Shannon	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	959.3	997.9	939.4	955.7	1003.3	948.8	940.1	929.6	868.5	775.3	766.4	724.6	653.6	806.2
ex Galway	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	139.5	121.4	172.5	134.5	142.2	143.5	135.2	127.5	128.3	256.4	191.7	NA	167.4	NA
ex Sligo	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	67.5	49.6	49.6	49.6	109.3	115.1	50.0	49.6	88.0	49.2	128.9	49.5	122.2	50.0
ex Donegal	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	95.9	100.6	107.6	90.1	88.6	85.1	105.2	94.5	96.6	75.1	87.2	86.7	106.0	109.6
ex Knock	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	732.0	619.6	732.5	761.0	760.2	751.8	751.7	747.3	724.0	633.8	627.5	661.9	701.0	630.0
ex Kerry	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	578.8	629.1	588.2	574.5	560.5	566.2	581.8	551.1	555.2	595.0	614.8	604.7	590.7	574.7
ex Waterford	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	162.2	164.4	168.2	182.6	173.6	177.5	137.1	131.9	120.0	123.6	157.4	170.2	133.2	117.1
ex Other	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	245.7	249.0	213.2	279.7	236.8	299.3	233.5	208.1	214.2	198.9	206.2	204.9	209.6	171.3
Weighted average	773.7	769.9	772.1	777.7	774.3	775.1	773.5	768.6	771.8	772.5	770.8	767.4	763.4	761.2	766.5	763.1	756.9	759.4	764.8	771.3	743.6	739.6	728.5	717.7	707.0	702.8	710.5

IEF kg NO _x /LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	8.6	9.1	9.2	9.2	9.5	9.6	9.4	9.4	9.4	9.2	9.0	8.9	8.8
ex Cork	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	7.4	6.4	7.2	7.5	7.6	7.7	8.0	7.6	7.3	7.2	6.9	6.5	6.3	6.1
ex Shannon	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	12.4	13.0	12.1	12.4	13.0	12.1	12.0	12.5	11.8	10.0	9.8	9.1	7.7	10.4
ex Galway	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.1	1.5	1.3	1.4	1.4	1.3	1.2	1.2	1.9	1.4	NA	1.2	NA
ex Sligo	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.3	0.3	1.0	1.1	0.3	0.3	0.7	0.3	0.9	0.3	0.9	0.3
ex Donegal	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.9	0.9	0.8	0.7	0.7	0.9	0.8	0.8	0.5	0.6	0.6	0.9	1.0
ex Knock	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	6.7	7.8	8.2	8.1	7.7	7.8	7.8	7.6	6.7	6.6	6.9	7.4	6.6
ex Kerry	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	6.0	5.6	5.5	5.4	5.5	5.7	5.4	5.4	5.9	6.0	5.9	5.8	5.6
ex Waterford	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.6	1.5	1.5	1.3	1.2	1.1	1.1	1.6	1.8	1.3	1.0
ex Other	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.5	2.1	1.7	2.2	1.7	1.5	1.6	1.4	1.5	1.5	1.5	1.2
Weighted average	9.3	9.3	9.3	9.4	9.3	9.3	9.3	9.2	9.2	9.3	9.2	9.2	9.1	9.1	8.6	8.9	9.1	9.1	9.2	9.4	9.1	9.1	8.9	8.8	8.6	8.4	8.5

Table C.17 International LTO IEFs by airport (continued)

IEF g HC/LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1319.4	1273.9	1417.0	1391.7	1314.4	1250.9	1276.9	1311.2	1370.5	1276.5	1242.2	1184.5	1134.2	1130.4
ex Cork	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1245.8	1023.4	1118.4	1244.1	1234.3	1326.7	1400.3	1373.6	1286.5	1233.3	1218.5	1177.1	1094.8	1121.4
ex Shannon	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	3278.0	4805.3	2983.4	2614.7	3808.3	2871.8	2790.7	3071.6	2857.9	1800.2	1719.1	1558.0	1518.8	1599.8
ex Galway	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	183.4	398.3	306.8	297.1	36.4	208.2	18.9	18.2	157.0	3254.3	4031.9	NA	1647.1	NA
ex Sligo	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	161.4	201.2	201.2	201.2	38.0	86.0	201.2	201.2	168.9	178.9	3025.0	183.1	4239.9	178.8
ex Donegal	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	44.9	23.3	53.3	51.9	60.4	27.6	44.2	300.2	221.6	312.1	193.1	37.9	16.3
ex Knock	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	810.9	916.7	882.6	870.8	747.2	713.3	767.2	778.8	735.3	701.5	691.8	772.9	772.7	733.3
ex Kerry	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	3984.7	6263.7	5648.8	4898.5	4953.0	3729.6	1293.6	1105.4	1205.8	1054.0	1418.5	1418.0	1309.3	1069.2
ex Waterford	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1316.0	1504.4	2115.8	2816.1	1755.4	709.7	278.6	31.8	165.4	178.9	501.3	678.5	876.5	359.0
ex Other	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	5006.8	2756.7	3075.0	5958.6	5696.2	7627.5	6377.7	3556.0	3756.7	2865.2	2727.0	2693.6	3649.3	2799.2
Weighted average	1778.0	1741.9	1723.9	1727.7	1680.6	1704.5	1698.1	1638.7	1657.2	1672.2	1666.1	1629.6	1577.7	1580.5	1705.9	1634.3	1601.7	1664.1	1483.9	1450.2	1458.2	1480.3	1292.8	1273.6	1213.9	1165.1	1159.4

IEF g CO/LTO	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	12251.0	10630.9	11609.4	12621.3	12572.8	12710.2	12941.2	12671.4	12734.7	12669.3	12335.8	12060.4	11860.7	11773.6
ex Cork	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	11361.6	9498.5	10716.1	11365.1	11664.4	11886.4	12436.9	11963.6	11325.4	11267.2	10915.1	10368.0	10112.6	9949.1
ex Shannon	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	14309.6	13639.4	12646.8	12761.6	15346.7	15086.6	15232.3	15453.7	14151.2	11564.6	11593.1	10929.8	10561.0	11736.4
ex Galway	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1185.6	1157.2	1936.6	1132.0	1004.8	1207.3	935.3	925.9	1113.4	5986.6	5594.4	NA	3617.6	NA
ex Sligo	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	780.3	749.0	749.0	749.0	841.7	875.1	749.0	749.0	985.8	759.2	3927.8	806.3	4631.2	802.3
ex Donegal	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	836.8	837.8	850.1	836.3	837.9	826.6	835.2	834.0	1062.7	446.7	645.0	548.6	813.4	851.0
ex Knock	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	11012.3	9888.0	10655.4	11771.8	10851.1	10958.1	11455.4	11506.5	10954.4	9499.5	9453.9	10027.6	10744.2	9442.0
ex Kerry	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	10738.7	12959.6	12094.1	11259.5	11243.2	10368.5	8933.1	8313.1	8452.7	8941.9	9482.0	9375.3	9094.9	8724.1
ex Waterford	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2470.9	2826.7	3126.3	3832.4	2987.0	2319.7	1232.1	972.4	1060.9	1056.7	1476.8	1758.7	1619.3	1302.0
ex Other	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	7153.7	5645.7	5344.5	8370.4	7484.4	9785.0	7871.5	5574.7	5784.7	4858.8	4968.9	4920.8	5637.8	4471.3
Weighted average	12182.0	12140.9	12143.9	12178.0	12133.2	12150.6	12137.9	12078.7	12106.7	12118.3	12105.5	12067.0	12016.3	12005.2	10617.5	11324.8	12131.4	12333.4	12420.2	12715.3	12381.8	12288.2	12036.0	11870.7	11596.1	11450.9	11444.5

Table C.18 Domestic Cruise IEFs by airport

IEF kg fuel/Cruise	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	442.3	459.8	537.6	496.4	510.5	526.2	490.8	409.1	390.2	359.3	348.4	394.5	349.1
ex Cork	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	245.3	266.9	399.9	410.7	439.8	494.8	501.9	394.4	173.2	168.7	188.6	256.9	254.9
ex Shannon	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	1010.3	1055.9	1059.3	978.1	990.0	938.6	826.5	625.6	655.3	549.0	573.7	710.4	679.6
ex Galway	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	159.4	160.3	158.5	176.9	196.1	160.5	159.2	147.9	124.7	102.8	NA	89.6	36.5
ex Sligo	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	164.2	164.1	166.9	165.6	163.6	167.6	168.0	143.7	108.2	97.6	92.5	91.6	93.6
ex Donegal	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	212.1	210.6	213.8	215.2	212.4	216.0	215.6	191.0	126.4	127.0	127.4	208.0	213.5
ex Knock	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	192.6	202.7	242.8	244.8	230.1	201.6	186.8	176.0	240.2	232.3	323.4	276.9	251.6
ex Kerry	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	246.7	247.8	247.0	242.4	452.1	757.9	753.8	533.6	242.7	240.6	229.9	229.6	239.0
ex Waterford	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	104.8	109.8	105.5	210.1	287.0	130.4	164.8	151.2	101.9	72.7	88.4	87.5	83.3
ex Other	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	160.2	157.6	148.5	165.4	139.9	140.7	141.8	159.7	138.0	128.0	143.8	136.8	113.5
Weighted average	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	402.0	433.5	492.9	451.1	467.0	485.6	440.6	334.9	326.7	298.9	303.9	354.8	332.5

IEF kg NO _x /Cruise	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	9.2	10.6	8.9	9.0	8.3	7.2	5.9	6.2	5.3	5.0	5.7	5.0
ex Cork	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	2.9	3.2	5.1	4.9	5.3	6.0	6.2	5.0	2.2	2.1	2.5	3.7	3.6
ex Shannon	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	23.8	25.2	26.6	26.5	23.3	24.6	22.2	18.5	11.7	12.4	9.3	9.9	12.9	12.4
ex Galway	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.5	1.5	1.6	1.8	2.1	1.6	1.5	1.4	1.2	0.9	NA	0.7	0.3
ex Sligo	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.4	1.4	1.5	1.4	1.5	1.5	1.5	1.3	0.8	0.7	0.7	0.7	0.7
ex Donegal	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.8	1.8	1.9	1.9	1.9	1.9	1.9	1.7	0.9	1.0	1.0	1.8	1.9
ex Knock	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.2	2.5	3.2	3.2	2.8	2.3	2.0	2.0	3.3	3.1	4.5	4.0	3.2
ex Kerry	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	2.6	2.8	2.8	2.7	5.4	9.2	9.2	6.3	2.2	2.2	2.2	2.2	2.5
ex Waterford	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	0.9	1.0	1.0	2.4	3.5	1.4	1.8	1.5	0.8	0.5	0.7	0.7	0.6
ex Other	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.7	1.6	1.5	1.8	1.4	1.4	1.4	1.7	1.3	1.2	1.4	1.6	1.0
Weighted average	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.9	7.7	8.5	9.5	7.8	7.9	7.5	6.4	4.7	5.0	4.2	4.3	5.1	4.7

Table C.18 Domestic Cruise IEFs by airport (continued)

IEF g HC/Cruise	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	226.6	248.8	250.3	286.7	229.5	233.4	187.5	150.2	137.3	178.0	142.6	148.1	80.3	117.4
ex Cork	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	78.3	38.4	49.5	85.9	80.5	85.8	95.1	112.9	104.7	63.2	76.6	87.4	125.2	170.6
ex Shannon	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	790.9	842.0	888.3	911.3	789.6	845.9	720.7	538.7	245.7	255.8	136.1	153.7	212.0	243.8
ex Galway	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	7.8	11.1	10.2	21.8	36.4	6.5	6.5	25.1	96.9	144.3	NA	145.0	59.1
ex Sligo	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	10.6	14.9	14.6	10.4	10.3	13.9	5.8	4.4	58.7	164.0	158.0	149.7	148.3	151.6
ex Donegal	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	8.8	10.7	16.3	10.4	6.4	9.5	4.1	4.2	60.4	204.6	204.5	205.2	22.5	8.3
ex Knock	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	71.5	39.4	104.1	113.6	59.9	72.0	47.9	63.7	110.0	157.0	166.6	186.6	198.2	164.4
ex Kerry	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	71.9	19.4	23.0	25.9	22.3	94.8	158.8	159.1	120.8	14.5	21.7	42.3	30.8	13.2
ex Waterford	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	110.3	108.1	105.6	116.5	172.0	169.9	60.4	39.3	23.0	74.3	116.7	135.1	141.6	134.8
ex Other	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	170.8	178.0	173.6	175.8	179.6	168.2	159.8	160.9	165.4	158.0	156.9	160.9	157.1	153.7
Weighted average	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	209.0	218.9	243.4	270.0	205.8	212.3	173.5	137.6	109.7	154.5	127.7	137.0	89.6	103.9

IEF g CO/Cruise	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ex Dublin	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1970.0	1934.1	1918.7	1968.0	1960.0	1936.5	2037.0	2035.7	1831.1	1901.7	1869.0	1786.3	2065.4	1883.5
ex Cork	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1706.3	1521.9	1626.5	1563.0	1742.6	1758.8	1894.4	1836.8	1476.1	990.8	964.5	985.2	1069.5	1055.1
ex Shannon	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2335.6	2276.0	2381.3	2546.1	2400.7	2341.4	2305.5	2098.6	1778.2	1860.3	1696.9	1761.5	1916.1	1932.6
ex Galway	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1370.2	1501.4	1453.5	1334.4	1332.7	1265.4	1326.2	1377.5	1266.7	928.3	831.1	NA	857.1	349.2
ex Sligo	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1624.9	1625.9	1628.8	1650.2	1643.8	1582.6	1615.1	1628.1	1357.5	1014.7	933.8	884.8	876.3	895.8
ex Donegal	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2109.3	2116.3	2084.2	2115.6	2131.8	2087.3	2114.0	2116.1	1842.1	1209.3	1215.1	1212.0	2054.1	2114.3
ex Knock	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1350.8	1358.6	1225.5	1278.9	1450.2	1398.1	1397.3	1347.3	1085.9	1042.9	1065.7	1115.8	1102.3	1114.6
ex Kerry	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1927.2	1914.6	1646.1	1632.7	1623.5	1889.4	2400.6	2383.5	2163.3	2240.6	2263.8	2071.6	1941.3	1773.5
ex Waterford	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	951.1	931.2	936.0	877.4	903.9	1006.4	953.2	1049.3	1231.4	996.4	700.0	791.9	836.9	797.0
ex Other	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	936.9	967.8	948.4	968.3	958.8	934.5	890.3	890.3	901.2	890.9	883.2	888.3	911.2	881.0
Weighted average	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1850.8	1813.4	1824.9	1854.3	1844.2	1818.3	1926.5	1868.3	1617.5	1677.7	1631.9	1620.6	1881.4	1804.0

Table C.19 Other Aviation EFs (SO₂, PM, Heavy Metals and POPs)

Emission Factors						
	Code	Name				Inventory Year
NFR Source Category	1.A.3.a	Civil Aviation & International Aviation				2016
	Gasoline			Kerosene		
Pollutant	Value	Unit	Reference	Value	Unit	Reference
SO _x	0.800	kg/LTO	Guidebook	0.970	kg/LTO	Guidebook
TSP	0.150	kg/LTO	CEPMEIP	0.150	kg/LTO	CEPMEIP
PM ₁₀	0.150	kg/LTO	CEPMEIP	0.150	kg/LTO	CEPMEIP
PM _{2.5}	0.150	kg/LTO	CEPMEIP	0.150	kg/LTO	CEPMEIP
Pb	186.391	mg/Mg	CS	0.080	mg/Mg	EMEP/EEA 2016
Cd	0.045	mg/Mg	EMEP/EEA 2016	0.006	mg/Mg	EMEP/EEA 2016
Hg	0.195	mg/Mg	EMEP/EEA 2016	0.120	mg/Mg	EMEP/EEA 2016
As	0.007	mg/Mg	EMEP/EEA 2016	0.030	mg/Mg	EMEP/EEA 2016
Cr	0.141	mg/Mg	EMEP/EEA 2016	0.200	mg/Mg	EMEP/EEA 2016
Cu	0.101	mg/Mg	EMEP/EEA 2016	0.220	mg/Mg	EMEP/EEA 2016
Ni	0.052	mg/Mg	EMEP/EEA 2016	0.008	mg/Mg	EMEP/EEA 2016
Se	0.004	mg/Mg	EMEP/EEA 2016	0.110	mg/Mg	EMEP/EEA 2016
Zn	0.740	mg/Mg	EMEP/EEA 2016	29.000	mg/Mg	EMEP/EEA 2016
PCDD/F (I-TEQ)	NE			NE		
Benzo[a]pyrene	NE			NE		
Benzo[b]fluoranthene	NE			NE		
Benzo[k]Fluoranthene	NE			NE		
Indeno[1,2,3-cd]pyrene	NE			NE		
Total 4 PAHs	NE			NE		
HCB	NA			NA		

Table C.20 Emission Factors for NFR 1A3. (c and d)

Emission Factors				Emission Factors						
	Code	Name	Inventory year		Code	Name	Inventory year			
NFR Source Category	1.A.3.c	Railways	2016	NFR Source Category	1.A.3.d	National Navigation	2016			
Fuel	Gas oil			Fuel	Gas oil			Fuel Oil		
Pollutant	Value	Unit	Reference	Pollutant	Value	Unit	Reference	Value	Unit	Reference
NO _x	1209.931	g/GJ	EMEP/EEA 2016	NO _x	1812.587	g/GJ	EMEP/EEA 2016	1923.087	g/GJ	EMEP/EEA 2016
SO _x	9.005	g/GJ	CS	SO _x	32.325	g/GJ	CS	329.777	g/GJ	CS
NM/OC	107.370	g/GJ	EMEP/EEA 2016	NM/OC	64.653	g/GJ	EMEP/EEA 2016	65.477	g/GJ	EMEP/EEA 2016
CO	247.066	g/GJ	EMEP/EEA 2016	CO	170.868	g/GJ	EMEP/EEA 2016	179.456	g/GJ	EMEP/EEA 2016
NH ₃	0.162	g/GJ	EMEP/EEA 2016	NH ₃	NA		EMEP/EEA 2016	NA		EMEP/EEA 2016
TSP	35.097	g/GJ	EMEP/EEA 2016	TSP	34.635	g/GJ	EMEP/EEA 2016	150.355	g/GJ	EMEP/EEA 2016
PM ₁₀	33.250	g/GJ	EMEP/EEA 2016	PM ₁₀	34.635	g/GJ	EMEP/EEA 2016	150.355	g/GJ	EMEP/EEA 2016
PM _{2.5}	31.634	g/GJ	EMEP/EEA 2016	PM _{2.5}	32.326	g/GJ	EMEP/EEA 2016	135.804	g/GJ	EMEP/EEA 2016
BC	0.650	f-TSP	EMEP/EEA 2016	BC	0.310	f-TSP	EMEP/EEA 2016	0.120	f-TSP	EMEP/EEA 2016
Pb	NE		EMEP/EEA 2016	Pb	0.130	g/t	EMEP/EEA 2016	0.180	g/t	EMEP/EEA 2016
Cd	0.010	g/t	EMEP/EEA 2016	Cd	0.010	g/t	EMEP/EEA 2016	0.020	g/t	EMEP/EEA 2016
Hg	NE		EMEP/EEA 2016	Hg	0.030	g/t	EMEP/EEA 2016	0.020	g/t	EMEP/EEA 2016
As	NE		EMEP/EEA 2016	As	0.040	g/t	EMEP/EEA 2016	0.680	g/t	EMEP/EEA 2016
Cr	0.050	g/t	EMEP/EEA 2016	Cr	0.050	g/t	EMEP/EEA 2016	0.720	g/t	EMEP/EEA 2016
Cu	1.700	g/t	EMEP/EEA 2016	Cu	0.880	g/t	EMEP/EEA 2016	1.250	g/t	EMEP/EEA 2016
Ni	0.070	g/t	EMEP/EEA 2016	Ni	1.000	g/t	EMEP/EEA 2016	32.000	g/t	EMEP/EEA 2016
Se	0.010	g/t	EMEP/EEA 2016	Se	0.100	g/t	EMEP/EEA 2016	0.210	g/t	EMEP/EEA 2016
Zn	1.000	g/t	EMEP/EEA 2016	Zn	1.200	g/t	EMEP/EEA 2016	1.200	g/t	EMEP/EEA 2016
PCB	NE		EMEP/EEA 2016	PCB	0.038	mg/t	EMEP/EEA 2016	0.570	mg/t	EMEP/EEA 2016
PCDD/F (I-TEQ)	NE		EMEP/EEA 2016	PCDD/F (I-TEQ)	0.130	µg/t	EMEP/EEA 2016	0.470	µg/t	EMEP/EEA 2016
Benzo[a]pyrene	0.030	g/t	EMEP/EEA 2016	Benzo[a]pyrene	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
Benzo[b]fluoranthene	0.050	g/t	EMEP/EEA 2016	Benzo[b]fluoranthene	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
Benzo[k]fluoranthene	NE		EMEP/EEA 2016	Benzo[k]fluoranthene	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	NE		EMEP/EEA 2016	Indeno[1,2,3-cd]pyrene	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
Total 4 PAHs	0.080	g/t	EMEP/EEA 2016	Total 4 PAHs	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016	HCB	0.080	mg/t	EMEP/EEA 2016	0.140	mg/t	EMEP/EEA 2016

Table C.21 Emission Factors for NFR 1A3e

Emission Factors			
	Code	Name	Inventory year
NFR Source Category	1.A.3.e	Pipeline compressors	2016
Fuel	Gasoil		
Pollutant	Value	Unit	Reference
NO _x	48.000	g/GJ	EMEP/EEA 2016
SO _x	0.043	g/GJ	CS
NMVOC	1.600	g/GJ	EMEP/EEA 2016
CO	4.800	g/GJ	EMEP/EEA 2016
NH ₃	NA		
TSP	0.200	g/GJ	EMEP/EEA 2016
PM ₁₀	0.200	g/GJ	EMEP/EEA 2016
PM _{2.5}	0.200	g/GJ	EMEP/EEA 2016
BC	0.025	f-PM _{2.5}	EMEP/EEA 2016
Pb	0.0015	mg/GJ	EMEP/EEA 2016
Cd	0.00025	mg/GJ	EMEP/EEA 2016
Hg	0.100	mg/GJ	EMEP/EEA 2016
As	0.120	mg/GJ	EMEP/EEA 2016
Cr	0.00076	mg/GJ	EMEP/EEA 2016
Cu	0.000076	mg/GJ	EMEP/EEA 2016
Ni	0.00051	mg/GJ	EMEP/EEA 2016
Se	0.011	mg/GJ	EMEP/EEA 2016
Zn	0.0015	mg/GJ	EMEP/EEA 2016
PCB	NE		EMEP/EEA 2016
PCDD/F (I-TEQ)	0.500	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	0.560	µg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	0.840	µg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	0.840	µg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	0.840	µg/GJ	EMEP/EEA 2016
Total 4 PAHs	3.080	µg/GJ	EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016

Table C.22 Emission Factors for NFR 1A4a

Emission Factors												
	Code	Name				Inventory Year						
NFR Source Category	1.A.4.a	Commercial/Institutional				2016						
Fuel	Coal			Anthracite & Ovoids			Lignite			Sod Peat		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	173.000	g/GJ	EMEP/EEA 2016	173.000	g/GJ	EMEP/EEA 2016	173.000	g/GJ	EMEP/EEA 2016
SO _x	301.407	g/GJ	CS	574.110	g/GJ	CS	840.000	g/GJ	EMEP/EEA 2016	840.000	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	88.800	g/GJ	EMEP/EEA 2016	88.800	g/GJ	EMEP/EEA 2016	88.800	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	931.000	g/GJ	EMEP/EEA 2016	931.000	g/GJ	EMEP/EEA 2016	931.000	g/GJ	EMEP/EEA 2016
NH ₃	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
TSP	124.000	g/GJ	EMEP/EEA 2016	124.000	g/GJ	EMEP/EEA 2016	124.000	g/GJ	EMEP/EEA 2016	124.000	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	117.000	g/GJ	EMEP/EEA 2016	117.000	g/GJ	EMEP/EEA 2016	117.000	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	108.000	g/GJ	EMEP/EEA 2016	108.000	g/GJ	EMEP/EEA 2016	108.000	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.064	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Peat Briquettes			Kerosene			Fuel Oil			LPG		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	173.000	g/GJ	EMEP/EEA 2016	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016
SO _x	840.000	g/GJ	EMEP/EEA 2016	21.538	g/GJ	CS	329.777	g/GJ	CS	0.670	g/GJ	EMEP/EEA 2016
NM/OC	88.800	g/GJ	EMEP/EEA 2016	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016
CO	931.000	g/GJ	EMEP/EEA 2016	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016
NH ₃	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
TSP	124.000	g/GJ	EMEP/EEA 2016	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM ₁₀	117.000	g/GJ	EMEP/EEA 2016	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
PM _{2.5}	108.000	g/GJ	EMEP/EEA 2016	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016
Fuel	Gasoil			Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	100.000	g/GJ	EMEP/EEA 2009	74.000	g/GJ	EMEP/EEA 2016	91.000	g/GJ	EMEP/EEA 2016
SO _x	9.005	g/GJ	CS	968.297	g/GJ	CS	0.043	g/GJ	CS	11.000	g/GJ	EMEP/EEA 2016
NM/OC	10.000	g/GJ	EMEP/EEA 2009	10.000	g/GJ	EMEP/EEA 2009	23.000	g/GJ	EMEP/EEA 2016	300.000	g/GJ	EMEP/EEA 2016
CO	40.000	g/GJ	EMEP/EEA 2009	40.000	g/GJ	EMEP/EEA 2009	29.000	g/GJ	EMEP/EEA 2016	570.000	g/GJ	EMEP/EEA 2016
NH ₃	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	37.000	g/GJ	EMEP/EEA 2016
TSP	27.500	g/GJ	EMEP/EEA 2009	27.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	150.000	g/GJ	EMEP/EEA 2016
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	21.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	143.000	g/GJ	EMEP/EEA 2016
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	16.500	g/GJ	EMEP/EEA 2009	0.780	g/GJ	EMEP/EEA 2016	140.000	g/GJ	EMEP/EEA 2016
BC	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.560	f-PM _{2.5}	EMEP/EEA 2016	0.040	f-PM _{2.5}	EMEP/EEA 2016	0.280	f-PM _{2.5}	EMEP/EEA 2016

Table C.23 (a) Emission Factors for NFR 1A4a (Heavy Metals and Persistent Organic Pollutants)

Emission Factors (Heavy Metals and POPs)												
	Code	Name					Inventory Year					
NFR Source Category	1.A.4.a	Commercial/Institutional					2016					
Fuel	Coal			Anthracite & Ovoids			Lignite			Sod Peat		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	134.00	mg/GJ	EMEP/EEA 2016	134.00	mg/GJ	EMEP/EEA 2016	134.00	mg/GJ	EMEP/EEA 2016	134.00	mg/GJ	EMEP/EEA 2016
Cd	1.80	mg/GJ	EMEP/EEA 2016	1.80	mg/GJ	EMEP/EEA 2016	1.80	mg/GJ	EMEP/EEA 2016	1.80	mg/GJ	EMEP/EEA 2016
Hg	7.90	mg/GJ	EMEP/EEA 2016	7.90	mg/GJ	EMEP/EEA 2016	7.90	mg/GJ	EMEP/EEA 2016	7.90	mg/GJ	EMEP/EEA 2016
As	4.00	mg/GJ	EMEP/EEA 2016	4.00	mg/GJ	EMEP/EEA 2016	4.00	mg/GJ	EMEP/EEA 2016	4.00	mg/GJ	EMEP/EEA 2016
Cr	13.50	mg/GJ	EMEP/EEA 2016	13.50	mg/GJ	EMEP/EEA 2016	13.50	mg/GJ	EMEP/EEA 2016	13.50	mg/GJ	EMEP/EEA 2016
Cu	17.50	mg/GJ	EMEP/EEA 2016	17.50	mg/GJ	EMEP/EEA 2016	17.50	mg/GJ	EMEP/EEA 2016	17.50	mg/GJ	EMEP/EEA 2016
Ni	13.00	mg/GJ	EMEP/EEA 2016	13.00	mg/GJ	EMEP/EEA 2016	13.00	mg/GJ	EMEP/EEA 2016	13.00	mg/GJ	EMEP/EEA 2016
Se	1.80	mg/GJ	EMEP/EEA 2016	1.80	mg/GJ	EMEP/EEA 2016	1.80	mg/GJ	EMEP/EEA 2016	1.80	mg/GJ	EMEP/EEA 2016
Zn	200.00	mg/GJ	EMEP/EEA 2016	200.00	mg/GJ	EMEP/EEA 2016	200.00	mg/GJ	EMEP/EEA 2016	200.00	mg/GJ	EMEP/EEA 2016
PCB	170.00	µg/GJ	EMEP/EEA 2016	170.00	µg/GJ	EMEP/EEA 2016	170.00	µg/GJ	EMEP/EEA 2016	170.00	µg/GJ	EMEP/EEA 2016
PCDD/F (I-TEQ)	203.00	ng/GJ	EMEP/EEA 2016	203.00	ng/GJ	EMEP/EEA 2016	203.00	ng/GJ	EMEP/EEA 2016	203.00	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	45.50	mg/GJ	EMEP/EEA 2016	45.50	mg/GJ	EMEP/EEA 2016	45.50	mg/GJ	EMEP/EEA 2016	45.50	mg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	58.90	mg/GJ	EMEP/EEA 2016	58.90	mg/GJ	EMEP/EEA 2016	58.90	mg/GJ	EMEP/EEA 2016	58.90	mg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	23.70	mg/GJ	EMEP/EEA 2016	23.70	mg/GJ	EMEP/EEA 2016	23.70	mg/GJ	EMEP/EEA 2016	23.70	mg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	18.50	mg/GJ	EMEP/EEA 2016	18.50	mg/GJ	EMEP/EEA 2016	18.50	mg/GJ	EMEP/EEA 2016	18.50	mg/GJ	EMEP/EEA 2016
Total 4 PAHs	146.60	mg/GJ	EMEP/EEA 2016	146.60	mg/GJ	EMEP/EEA 2016	146.60	mg/GJ	EMEP/EEA 2016	146.60	mg/GJ	EMEP/EEA 2016
HCB	0.62	µg/GJ	EMEP/EEA 2016	0.62	µg/GJ	EMEP/EEA 2016	0.62	µg/GJ	EMEP/EEA 2016	0.62	µg/GJ	EMEP/EEA 2016

Emission Factors (Heavy Metals and POPs)												
	Code	Name					Inventory Year					
NFR Source Category	1.A.4.a	Commercial/Institutional					2016					
Fuel	Peat Briquettes			Fuel Oil			LPG			Gas oil		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	134.00	mg/GJ	EMEP/EEA 2016	16.00	mg/GJ	EMEP/EEA 2009	0.011	mg/GJ	EMEP/EEA 2016	16.00	mg/GJ	EMEP/EEA 2009
Cd	1.80	mg/GJ	EMEP/EEA 2016	0.300	mg/GJ	EMEP/EEA 2009	0.0009	mg/GJ	EMEP/EEA 2016	0.30	mg/GJ	EMEP/EEA 2009
Hg	7.90	mg/GJ	EMEP/EEA 2016	0.10	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2016	0.10	mg/GJ	EMEP/EEA 2009
As	4.00	mg/GJ	EMEP/EEA 2016	1.00	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2016	1.00	mg/GJ	EMEP/EEA 2009
Cr	13.50	mg/GJ	EMEP/EEA 2016	12.80	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016	12.80	mg/GJ	EMEP/EEA 2009
Cu	17.50	mg/GJ	EMEP/EEA 2016	7.20	mg/GJ	EMEP/EEA 2009	0.0026	mg/GJ	EMEP/EEA 2016	7.20	mg/GJ	EMEP/EEA 2009
Ni	13.00	mg/GJ	EMEP/EEA 2016	260.000	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016	260.000	mg/GJ	EMEP/EEA 2009
Se	1.80	mg/GJ	EMEP/EEA 2016	0.10	mg/GJ	EMEP/EEA 2009	0.058	mg/GJ	EMEP/EEA 2016	0.10	mg/GJ	EMEP/EEA 2009
Zn	200.00	mg/GJ	EMEP/EEA 2016	8.00	mg/GJ	EMEP/EEA 2009	0.730	mg/GJ	EMEP/EEA 2016	8.00	mg/GJ	EMEP/EEA 2009
PCB	170.00	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
PCDD/F (I-TEQ)	203.00	ng/GJ	EMEP/EEA 2016	6.000	ng/GJ	EMEP/EEA 2016	0.52	ng/GJ	EMEP/EEA 2016	6.00	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	45.50	mg/GJ	EMEP/EEA 2016	1.900	mg/GJ	EMEP/EEA 2016	0.72	mg/GJ	EMEP/EEA 2016	1.90	mg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	58.90	mg/GJ	EMEP/EEA 2016	15.000	mg/GJ	EMEP/EEA 2016	2.90	mg/GJ	EMEP/EEA 2016	15.00	mg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	23.70	mg/GJ	EMEP/EEA 2016	1.700	mg/GJ	EMEP/EEA 2016	1.10	mg/GJ	EMEP/EEA 2016	1.70	mg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	18.50	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016	1.08	mg/GJ	EMEP/EEA 2016	1.50	mg/GJ	EMEP/EEA 2016
Total 4 PAHs	146.60	mg/GJ	EMEP/EEA 2016	20.100	mg/GJ	EMEP/EEA 2016	5.80	mg/GJ	EMEP/EEA 2016	20.10	mg/GJ	EMEP/EEA 2016
HCB	0.62	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016

Table C.23 (b) Emission Factors for NFR 1A4a (Heavy Metals and Persistent Organic Pollutants)

Emission Factors (Heavy Metals and POPs)									
	Code	Name				Inventory Year			
NFR Source Category	1.A.4.a	Commercial/Institutional				2016			
Fuel	Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	16.00	mg/GJ	EMEP/EEA 2009	0.011	mg/GJ	EMEP/EEA 2016	27.000	mg/GJ	EMEP/EEA 2016
Cd	0.300	mg/GJ	EMEP/EEA 2009	0.0009	mg/GJ	EMEP/EEA 2016	13.000	mg/GJ	EMEP/EEA 2016
Hg	0.10	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2016	0.560	mg/GJ	EMEP/EEA 2016
As	1.00	mg/GJ	EMEP/EEA 2009	0.100	mg/GJ	EMEP/EEA 2016	0.190	mg/GJ	EMEP/EEA 2016
Cr	12.80	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016	23.000	mg/GJ	EMEP/EEA 2016
Cu	7.20	mg/GJ	EMEP/EEA 2009	0.0026	mg/GJ	EMEP/EEA 2016	6.000	mg/GJ	EMEP/EEA 2016
Ni	260.00	mg/GJ	EMEP/EEA 2009	0.013	mg/GJ	EMEP/EEA 2016	2.000	mg/GJ	EMEP/EEA 2016
Se	0.10	mg/GJ	EMEP/EEA 2009	0.058	mg/GJ	EMEP/EEA 2016	0.500	mg/GJ	EMEP/EEA 2016
Zn	8.00	mg/GJ	EMEP/EEA 2009	0.730	mg/GJ	EMEP/EEA 2016	512.000	mg/GJ	EMEP/EEA 2016
PCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	0.060	µg/GJ	EMEP/EEA 2016
PCDD/F (I-TEQ)	6.000	ng/GJ	EMEP/EEA 2016	0.520	ng/GJ	EMEP/EEA 2016	100.000	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	1.900	mg/GJ	EMEP/EEA 2016	0.720	mg/GJ	EMEP/EEA 2016	10.000	mg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	15.000	mg/GJ	EMEP/EEA 2016	2.900	mg/GJ	EMEP/EEA 2016	16.000	mg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	1.700	mg/GJ	EMEP/EEA 2016	1.100	mg/GJ	EMEP/EEA 2016	5.000	mg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	1.500	mg/GJ	EMEP/EEA 2016	1.080	mg/GJ	EMEP/EEA 2016	4.000	mg/GJ	EMEP/EEA 2016
Total 4 PAHs	20.100	mg/GJ	EMEP/EEA 2016	5.800	mg/GJ	EMEP/EEA 2016	35.000	mg/GJ	EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	5.000	µg/GJ	EMEP/EEA 2016

Table C.24 Emission Factors for NFR 1A4b

Emission Factors																
	Code	Name	Inventory Year													
NFR Source Category	1.A.4.b	Residential	2016													
Fuel	Coal				Anthracite & Ovoids				Lignite				Sod Peat			
Pollutant	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier
NO _x	110.000	g/GJ	EMEP/EEA 2016	1	110.000	g/GJ	EMEP/EEA 2016		110.000	g/GJ	EMEP/EEA 2016		110.000	g/GJ	EMEP/EEA 2016	
SO _x	301.407	g/GJ	CS		574.110	g/GJ	CS		756.000	g/GJ	CS		300.000	g/GJ	EMEP/EEA 2016	
NM/OC	484.000	g/GJ	EMEP/EEA 2016		484.000	g/GJ	EMEP/EEA 2016		484.000	g/GJ	EMEP/EEA 2016		484.000	g/GJ	EMEP/EEA 2016	
CO	931.000	g/GJ	EMEP/EEA 2016		931.000	g/GJ	EMEP/EEA 2016		931.000	g/GJ	EMEP/EEA 2016		931.000	g/GJ	EMEP/EEA 2016	
NH ₃	0.300	g/GJ	EMEP/EEA 2016		0.300	g/GJ	EMEP/EEA 2016		0.300	g/GJ	EMEP/EEA 2016		0.300	g/GJ	EMEP/EEA 2016	
TSP	444.000	g/GJ	EMEP/EEA 2016		444.000	g/GJ	EMEP/EEA 2016		444.000	g/GJ	EMEP/EEA 2016		444.000	g/GJ	EMEP/EEA 2016	
PM ₁₀	404.000	g/GJ	EMEP/EEA 2016		404.000	g/GJ	EMEP/EEA 2016		404.000	g/GJ	EMEP/EEA 2016		404.000	g/GJ	EMEP/EEA 2016	
PM _{2.5}	398.000	g/GJ	EMEP/EEA 2016		398.000	g/GJ	EMEP/EEA 2016		398.000	g/GJ	EMEP/EEA 2016		398.000	g/GJ	EMEP/EEA 2016	
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016		0.064	f-PM _{2.5}	EMEP/EEA 2016		0.064	f-PM _{2.5}	EMEP/EEA 2016		0.064	f-PM _{2.5}	EMEP/EEA 2016	
Fuel	Peat Briquettes				Kerosene				LPG				Gasoil			
Pollutant	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier
NO _x	110.000	g/GJ	EMEP/EEA 2016		51.000	g/GJ	EMEP/EEA 2016		42.000	g/GJ	EMEP/EEA 2016		51.000	g/GJ	EMEP/EEA 2016	
SO _x	280.000	g/GJ	EMEP/EEA 2016		21.538	g/GJ	CS		0.300	g/GJ	EMEP/EEA 2016		9.005	g/GJ	CS	
NM/OC	484.000	g/GJ	EMEP/EEA 2016		0.690	g/GJ	EMEP/EEA 2016		1.800	g/GJ	EMEP/EEA 2016		0.690	g/GJ	EMEP/EEA 2016	
CO	931.000	g/GJ	EMEP/EEA 2016		57.000	g/GJ	EMEP/EEA 2016		22.000	g/GJ	EMEP/EEA 2016		57.000	g/GJ	EMEP/EEA 2016	
NH ₃	0.300	g/GJ	EMEP/EEA 2016		NE		EMEP/EEA 2016		NA		EMEP/EEA 2016		NE		EMEP/EEA 2016	
TSP	444.000	g/GJ	EMEP/EEA 2016		1.500	g/GJ	EMEP/EEA 2016		0.200	g/GJ	EMEP/EEA 2016		1.500	g/GJ	EMEP/EEA 2016	
PM ₁₀	404.000	g/GJ	EMEP/EEA 2016		1.500	g/GJ	EMEP/EEA 2016		0.200	g/GJ	EMEP/EEA 2016		1.500	g/GJ	EMEP/EEA 2016	
PM _{2.5}	398.000	g/GJ	EMEP/EEA 2016		1.500	g/GJ	EMEP/EEA 2016		0.200	g/GJ	EMEP/EEA 2016		1.500	g/GJ	EMEP/EEA 2016	
BC	0.064	f-PM _{2.5}	EMEP/EEA 2016		0.085	f-PM _{2.5}	EMEP/EEA 2016		0.054	f-PM _{2.5}	EMEP/EEA 2016		0.085	f-PM _{2.5}	EMEP/EEA 2016	
Fuel	Petroleum Coke				Natural Gas				Biomass							
Pollutant	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier	Value	Unit	Reference	Tier
NO _x	51.000	g/GJ	EMEP/EEA 2016		42.000	g/GJ	EMEP/EEA 2016		50.000	g/GJ	EMEP/EEA 2016					
SO _x	968.297	g/GJ	CS		0.043	g/GJ	CS		11.000	g/GJ	EMEP/EEA 2016					
NM/OC	0.690	g/GJ	EMEP/EEA 2016		1.800	g/GJ	EMEP/EEA 2016		600.000	g/GJ	EMEP/EEA 2016					
CO	57.000	g/GJ	EMEP/EEA 2016		22.000	g/GJ	EMEP/EEA 2016		4000.000	g/GJ	EMEP/EEA 2016					
NH ₃	NA		EMEP/EEA 2016		NA		EMEP/EEA 2016		70.000	g/GJ	EMEP/EEA 2016					
TSP	1.500	g/GJ	EMEP/EEA 2016		0.200	g/GJ	EMEP/EEA 2016		800.000	g/GJ	EMEP/EEA 2016					
PM ₁₀	1.500	g/GJ	EMEP/EEA 2016		0.200	g/GJ	EMEP/EEA 2016		760.000	g/GJ	EMEP/EEA 2016					
PM _{2.5}	1.500	g/GJ	EMEP/EEA 2016		0.200	g/GJ	EMEP/EEA 2016		740.000	g/GJ	EMEP/EEA 2016					
BC	0.085	f-PM _{2.5}	EMEP/EEA 2016		0.054	f-PM _{2.5}	EMEP/EEA 2016		0.100	f-PM _{2.5}	EMEP/EEA 2016					

Table C.25 (a) Emission Factors for NFR 1A4b (Heavy Metals and Persistent Organic Pollutants)

Emission Factors (Heavy Metals and POPs)												
	Code	Name					Inventory Year					
NFR Source Category	1.A.4.b	Residential					2016					
Fuel	Coal			Anthracite & Ovoids			Lignite			Sod Peat		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	130.000	mg/GJ	EMEP/EEA 2016	130.000	mg/GJ	EMEP/EEA 2016	130.000	mg/GJ	EMEP/EEA 2016	130.000	mg/GJ	EMEP/EEA 2016
Cd	1.500	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016	1.500	mg/GJ	EMEP/EEA 2016
Hg	5.100	mg/GJ	EMEP/EEA 2016	5.100	mg/GJ	EMEP/EEA 2016	5.100	mg/GJ	EMEP/EEA 2016	5.100	mg/GJ	EMEP/EEA 2016
As	2.500	mg/GJ	EMEP/EEA 2016	2.500	mg/GJ	EMEP/EEA 2016	2.500	mg/GJ	EMEP/EEA 2016	2.500	mg/GJ	EMEP/EEA 2016
Cr	11.200	mg/GJ	EMEP/EEA 2016	11.200	mg/GJ	EMEP/EEA 2016	11.200	mg/GJ	EMEP/EEA 2016	11.200	mg/GJ	EMEP/EEA 2016
Cu	22.300	mg/GJ	EMEP/EEA 2016	22.300	mg/GJ	EMEP/EEA 2016	22.300	mg/GJ	EMEP/EEA 2016	22.300	mg/GJ	EMEP/EEA 2016
Ni	12.700	mg/GJ	EMEP/EEA 2016	12.700	mg/GJ	EMEP/EEA 2016	12.700	mg/GJ	EMEP/EEA 2016	12.700	mg/GJ	EMEP/EEA 2016
Se	120.000	mg/GJ	EMEP/EEA 2016	120.000	mg/GJ	EMEP/EEA 2016	120.000	mg/GJ	EMEP/EEA 2016	120.000	mg/GJ	EMEP/EEA 2016
Zn	220.000	mg/GJ	EMEP/EEA 2016	220.000	mg/GJ	EMEP/EEA 2016	220.000	mg/GJ	EMEP/EEA 2016	220.000	mg/GJ	EMEP/EEA 2016
PCB	170.000	µg/GJ	EMEP/EEA 2016	170.000	µg/GJ	EMEP/EEA 2016	170.000	µg/GJ	EMEP/EEA 2016	170.000	µg/GJ	EMEP/EEA 2016
PCDD/F (I-TEQ)	800.000	ng/GJ	EMEP/EEA 2016	800.000	ng/GJ	EMEP/EEA 2016	800.000	ng/GJ	EMEP/EEA 2016	800.000	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	0.230	µg/GJ	EMEP/EEA 2016	0.230	µg/GJ	EMEP/EEA 2016	0.230	µg/GJ	EMEP/EEA 2016	0.230	µg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	0.330	µg/GJ	EMEP/EEA 2016	0.330	µg/GJ	EMEP/EEA 2016	0.330	µg/GJ	EMEP/EEA 2016	0.330	µg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	0.130	µg/GJ	EMEP/EEA 2016	0.130	µg/GJ	EMEP/EEA 2016	0.130	µg/GJ	EMEP/EEA 2016	0.130	µg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	0.110	µg/GJ	EMEP/EEA 2016	0.110	µg/GJ	EMEP/EEA 2016	0.110	µg/GJ	EMEP/EEA 2016	0.110	µg/GJ	EMEP/EEA 2016
Total 4 PAHs	0.800	µg/GJ	EMEP/EEA 2016	0.800	µg/GJ	EMEP/EEA 2016	0.800	µg/GJ	EMEP/EEA 2016	0.800	µg/GJ	EMEP/EEA 2016
HCB	0.620	µg/GJ	EMEP/EEA 2016	0.620	µg/GJ	EMEP/EEA 2016	0.620	µg/GJ	EMEP/EEA 2016	0.620	µg/GJ	EMEP/EEA 2016

Emission Factors (Heavy Metals and POPs)												
	Code	Name					Inventory Year					
NFR Source Category	1.A.4.b	Residential					2016					
Fuel	Peat Briquettes			Kerosene			LPG			Gasoil		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	130.000	mg/GJ	EMEP/EEA 2016	0.012	mg/GJ	EMEP/EEA 2016	0.0015	mg/GJ	EMEP/EEA 2016	0.012	mg/GJ	EMEP/EEA 2016
Cd	1.500	mg/GJ	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016	0.00025	mg/GJ	EMEP/EEA 2016	0.001	mg/GJ	EMEP/EEA 2016
Hg	5.100	mg/GJ	EMEP/EEA 2016	0.120	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2016	0.120	mg/GJ	EMEP/EEA 2016
As	2.500	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016	0.120	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016
Cr	11.200	mg/GJ	EMEP/EEA 2016	0.200	mg/GJ	EMEP/EEA 2016	0.00076	mg/GJ	EMEP/EEA 2016	0.200	mg/GJ	EMEP/EEA 2016
Cu	22.300	mg/GJ	EMEP/EEA 2016	0.130	mg/GJ	EMEP/EEA 2016	0.000076	mg/GJ	EMEP/EEA 2016	0.130	mg/GJ	EMEP/EEA 2016
Ni	12.700	mg/GJ	EMEP/EEA 2016	0.005	mg/GJ	EMEP/EEA 2016	0.00051	mg/GJ	EMEP/EEA 2016	0.005	mg/GJ	EMEP/EEA 2016
Se	120.000	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016	0.011	mg/GJ	EMEP/EEA 2016	0.002	mg/GJ	EMEP/EEA 2016
Zn	220.000	mg/GJ	EMEP/EEA 2016	0.420	mg/GJ	EMEP/EEA 2016	0.0015	mg/GJ	EMEP/EEA 2016	0.420	mg/GJ	EMEP/EEA 2016
PCB	170.0	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
PCDD/F (I-TEQ)	800.0	ng/GJ	EMEP/EEA 2016	5.90	ng/GJ	EMEP/EEA 2016	1.500	ng/GJ	EMEP/EEA 2016	5.90	ug/TJ	EMEP/EEA 2016
Benzo[a]pyrene	230.0	mg/GJ	EMEP/EEA 2016	0.08	µg/GJ	EMEP/EEA 2016	0.001	µg/GJ	EMEP/EEA 2016	0.08	g/TJ	EMEP/EEA 2016
Benzo[b]fluoranthene	330.0	mg/GJ	EMEP/EEA 2016	0.04	µg/GJ	EMEP/EEA 2016	0.001	µg/GJ	EMEP/EEA 2016	0.04	g/TJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	130.0	mg/GJ	EMEP/EEA 2016	0.07	µg/GJ	EMEP/EEA 2016	0.001	µg/GJ	EMEP/EEA 2016	0.07	g/TJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	110.0	mg/GJ	EMEP/EEA 2016	0.16	µg/GJ	EMEP/EEA 2016	0.001	µg/GJ	EMEP/EEA 2016	0.16	g/TJ	EMEP/EEA 2016
Total 4 PAHs	800.0	mg/GJ	EMEP/EEA 2016	0.35	µg/GJ	EMEP/EEA 2016	0.003	µg/GJ	EMEP/EEA 2016	0.35	g/TJ	EMEP/EEA 2016
HCB	0.620	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016

Table C.25 (b) Emission Factors for NFR 1A4b (Heavy Metals and Persistent Organic Pollutants)

Emission Factors (Heavy Metals and POPs)									
	Code	Name				Inventory Year			
NFR Source Category	1.A.4.b	Residential				2016			
Fuel	Petroleum Coke			Natural Gas			Biomass		
Pollutant	Value	Unit	Reference	Value	Unit	Reference	Value	Unit	Reference
Pb	0.012	mg/GJ	EMEP/EEA 2016	0.0015	mg/GJ	EMEP/EEA 2016	27.000	mg/GJ	EMEP/EEA 2016
Cd	0.001	mg/GJ	EMEP/EEA 2016	0.00025	mg/GJ	EMEP/EEA 2016	13.000	mg/GJ	EMEP/EEA 2016
Hg	0.120	mg/GJ	EMEP/EEA 2016	0.100	mg/GJ	EMEP/EEA 2016	0.560	mg/GJ	EMEP/EEA 2016
As	0.002	mg/GJ	EMEP/EEA 2016	0.120	mg/GJ	EMEP/EEA 2016	0.190	mg/GJ	EMEP/EEA 2016
Cr	0.200	mg/GJ	EMEP/EEA 2016	0.00076	mg/GJ	EMEP/EEA 2016	23.000	mg/GJ	EMEP/EEA 2016
Cu	0.130	mg/GJ	EMEP/EEA 2016	0.000076	mg/GJ	EMEP/EEA 2016	6.000	mg/GJ	EMEP/EEA 2016
Ni	0.005	mg/GJ	EMEP/EEA 2016	0.00051	mg/GJ	EMEP/EEA 2016	2.000	mg/GJ	EMEP/EEA 2016
Se	0.002	mg/GJ	EMEP/EEA 2016	0.011	mg/GJ	EMEP/EEA 2016	0.500	mg/GJ	EMEP/EEA 2016
Zn	0.420	mg/GJ	EMEP/EEA 2016	0.0015	mg/GJ	EMEP/EEA 2016	512.000	mg/GJ	EMEP/EEA 2016
PCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	0.060	µg/GJ	EMEP/EEA 2016
PCDD/F (I-TEQ)	5.900	ng/GJ	EMEP/EEA 2016	1.500	ug/TJ	EMEP/EEA 2016	800.000	ng/GJ	EMEP/EEA 2016
Benzo[a]pyrene	0.080	µg/GJ	EMEP/EEA 2016	0.001	g/TJ	EMEP/EEA 2016	121.000	mg/GJ	EMEP/EEA 2016
Benzo[b]fluoranthene	0.040	µg/GJ	EMEP/EEA 2016	0.001	g/TJ	EMEP/EEA 2016	111.000	mg/GJ	EMEP/EEA 2016
Benzo[k]Fluoranthene	0.070	µg/GJ	EMEP/EEA 2016	0.001	g/TJ	EMEP/EEA 2016	42.000	mg/GJ	EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	0.160	µg/GJ	EMEP/EEA 2016	0.001	g/TJ	EMEP/EEA 2016	71.000	mg/GJ	EMEP/EEA 2016
Total 4 PAHs	0.350	µg/GJ	EMEP/EEA 2016	0.003	g/TJ	EMEP/EEA 2016	345.000	mg/GJ	EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016	5.000	µg/GJ	EMEP/EEA 2016

Table C.26 Emission Factors for NFR 1A4c

Emission Factors						
	Code	Name	Inventory year			
NFR Source Category	1.A.4.c	Agriculture/Forestry/Fishing	2016			
Fuel	Gasoil			Gasoil		
	Stationary (1 A 4 c i)			Mobile (1 A 4 c ii)		
Pollutant	Value	Unit	Reference	Value	Unit	Reference
NO _x	100.000	g/GJ	EMEP/EEA 2009	420.742	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
SO _x	9.005	g/GJ	CS	3.848	g/GJ	CS
NM/VO	10.000	g/GJ	EMEP/EEA 2009	38.640	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
CO	40.000	g/GJ	EMEP/EEA 2009	179.962	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
NH ₃	NE		EMEP/EEA 2016	0.184	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
TSP	27.500	g/GJ	EMEP/EEA 2009	17.766	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
PM ₁₀	21.500	g/GJ	EMEP/EEA 2009	17.763	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
PM _{2.5}	16.500	g/GJ	EMEP/EEA 2009	17.763	g/GJ	Danish Inventory (Winther and Nielsen, 2006)
Pb	16.000	mg/GJ	EMEP/EEA 2009	NE		EMEP/EEA 2016
Cd	0.300	mg/GJ	EMEP/EEA 2009	0.010	mg/kg	EMEP/EEA 2016
Hg	0.100	mg/GJ	EMEP/EEA 2009	NE		EMEP/EEA 2016
As	1.000	mg/GJ	EMEP/EEA 2009	NE		EMEP/EEA 2016
Cr	12.800	mg/GJ	EMEP/EEA 2009	0.050	mg/kg	EMEP/EEA 2016
Cu	7.200	mg/GJ	EMEP/EEA 2009	1.700	mg/kg	EMEP/EEA 2016
Ni	260.000	mg/GJ	EMEP/EEA 2009	0.070	mg/kg	EMEP/EEA 2016
Se	0.100	mg/GJ	EMEP/EEA 2016	0.010	mg/kg	EMEP/EEA 2016
Zn	8.000	mg/GJ	EMEP/EEA 2009	1.000	mg/kg	EMEP/EEA 2016
PCDD/F (I-TEQ)	6.000	ng/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016
Benzo[a]pyrene	1.900	µg/GJ	EMEP/EEA 2016	40.000	µg/kg	EMEP/EEA 2016
Benzo[b]fluoranthene	15.000	µg/GJ	EMEP/EEA 2016	40.000	µg/kg	EMEP/EEA 2016
Benzo[k]Fluoranthene	1.700	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	1.500	µg/GJ	EMEP/EEA 2016	NE		EMEP/EEA 2016
Total 4 PAHs	20.100	µg/GJ	EMEP/EEA 2016	80.000	µg/kg	EMEP/EEA 2016
HCB	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016

Table C.27 Tier 2 Implied Emission Factors for NFR 1A4cii

Implied Emission Factors																												
Pollutant	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
NOx	kg/TJ	804.24	825.70	852.26	877.05	900.05	921.29	940.76	959.04	976.10	985.43	993.48	993.25	967.08	929.99	893.12	859.35	819.05	775.16	730.86	690.09	654.27	615.01	576.46	540.28	499.74	458.09	420.74
NMVOC	kg/TJ	159.09	155.49	150.99	146.78	142.89	139.29	136.00	132.91	130.03	126.28	122.73	118.22	110.76	103.24	96.22	89.98	84.12	78.52	72.84	67.81	63.24	58.61	53.84	49.44	45.39	41.81	38.64
CO	kg/TJ	423.24	416.92	409.08	401.75	394.96	388.69	382.95	377.55	372.52	365.07	357.95	348.18	330.90	313.91	298.10	284.06	270.75	257.99	244.86	233.47	223.07	213.78	205.45	197.88	191.01	185.04	179.96
PM ₁₀	kg/TJ	107.74	103.91	98.78	94.00	89.59	85.54	81.85	78.38	75.13	71.74	68.56	65.20	60.84	56.41	52.29	48.61	45.17	41.83	38.53	35.49	32.72	30.05	27.18	24.50	22.03	19.80	17.76
PM _{2.5}	kg/TJ	107.74	103.91	98.78	94.00	89.59	85.54	81.85	78.38	75.13	71.74	68.56	65.20	60.84	56.41	52.29	48.61	45.17	41.83	38.53	35.49	32.72	30.05	27.18	24.50	22.03	19.80	17.76
TSP	kg/TJ	107.74	103.91	98.78	94.00	89.59	85.54	81.85	78.38	75.13	71.74	68.56	65.20	60.84	56.41	52.29	48.62	45.18	41.84	38.53	35.50	32.72	30.05	27.19	24.51	22.03	19.80	17.77

Table C.28 Emission Factors for NFR 1A4ciii

Emission Factors			
	Code	Name	Inventory year
NFR Source Category	1.A.4.c iii	Fishing	2016
Fuel	Gasoil		
Pollutant	Value	Unit	Reference
NO _x	1812.59	g/GJ	EMEP/EEA 2016
SO _x	9.00	g/GJ	CS
NM/OC	64.65	g/GJ	EMEP/EEA 2016
CO	170.87	g/GJ	EMEP/EEA 2016
NH ₃	NA		EMEP/EEA 2016
TSP	34.64	g/GJ	EMEP/EEA 2016
PM ₁₀	34.64	g/GJ	EMEP/EEA 2016
PM _{2.5}	32.33	g/GJ	EMEP/EEA 2016
BC	0.31	f-TSP	EMEP/EEA 2016
Pb	130	mg/t	EMEP/EEA 2016
Cd	10	mg/t	EMEP/EEA 2016
Hg	30	mg/t	EMEP/EEA 2016
As	40	mg/t	EMEP/EEA 2016
Cr	50	mg/t	EMEP/EEA 2016
Cu	880	mg/t	EMEP/EEA 2016
Ni	1000	mg/t	EMEP/EEA 2016
Se	100	mg/t	EMEP/EEA 2016
Zn	1200	mg/t	EMEP/EEA 2016
PCB	0.038	mg/t	EMEP/EEA 2016
PCDD/F (I-TEQ)	0.130	µg/t	EMEP/EEA 2016
Benzo[a]pyrene	NE		EMEP/EEA 2016
Benzo[b]fluoranthene	NE		EMEP/EEA 2016
Benzo[k]Fluoranthene	NE		EMEP/EEA 2016
Indeno[1,2,3-cd]pyrene	NE		EMEP/EEA 2016
Total 4 PAHs	NE		EMEP/EEA 2016
HCB	0.080	mg/t	EMEP/EEA 2016

Table C.29 Emission Factors for NFR 1A3di

Emission Factors						
	Code	Name	Inventory year			
NFR Source Category	1.A.3.d.(i)	International Navigation	2016			
Fuel	Fuel Oil			Gas oil		
Pollutant	Value	Unit	Reference	Value	Unit	Reference
NO _x	79.300	kg/t	EMEP/EEA 2016	78.500	kg/t	EMEP/EEA 2016
SO _x	329.777	g/GJ	CS	32.325	g/GJ	CS
NM/OC	2.700	kg/t	EMEP/EEA 2016	2.800	kg/t	EMEP/EEA 2016
CO	7.400	kg/t	EMEP/EEA 2016	7.400	kg/t	EMEP/EEA 2016
NH ₃	NE		EMEP/EEA 2016	NE		EMEP/EEA 2016
TSP	6.200	kg/t	EMEP/EEA 2016	1.500	kg/t	EMEP/EEA 2016
PM ₁₀	6.200	kg/t	EMEP/EEA 2016	1.500	kg/t	EMEP/EEA 2016
PM _{2.5}	5.600	kg/t	EMEP/EEA 2016	1.400	kg/t	EMEP/EEA 2016
Pb	292.483	mg/t	NAEI 2008	40.965	mg/t	NAEI 2008
Cd	515.924	mg/t	NAEI 2008	17.647	mg/t	NAEI 2008
Hg	23.453	mg/t	NAEI 2008	19.936	mg/t	NAEI 2008
As	292.483	mg/t	NAEI 2008	40.965	mg/t	NAEI 2008
Cr	566.868	mg/t	NAEI 2008	24.314	mg/t	NAEI 2008
Cu	693.432	mg/t	NAEI 2008	297.516	mg/t	NAEI 2008
Ni	31959.485	mg/t	NAEI 2008	2217.647	mg/t	NAEI 2008
Se	361.125	mg/t	NAEI 2008	266.280	mg/t	NAEI 2008
Zn	1205.584	mg/t	NAEI 2008	276.492	mg/t	NAEI 2008
PCDD/F (I-TEQ)	104.278	ug/TJ	NAEI 2006	23.090	ug/TJ	NAEI 2006
Benzo[a]pyrene	113.978	mg/TJ	NAEI 2006	108.525	mg/TJ	NAEI 2006
Benzo[b]fluoranthene	492.288	mg/TJ	NAEI 2006	468.736	mg/TJ	NAEI 2006
Benzo[k]Fluoranthene	97.003	mg/TJ	NAEI 2006	92.362	mg/TJ	NAEI 2006
Indeno[1,2,3-cd]pyrene	184.305	mg/TJ	NAEI 2006	175.487	mg/TJ	NAEI 2006
Total 4 PAHs	887.574	mg/TJ	NAEI 2006	845.109	mg/TJ	NAEI 2006
HCB	NE			NE		




Figure C.1. Emission Factor Trends for Nitrogen Oxides in NFR 1A3b




Figure C.2. Emission Factor Trends for Sulphur Dioxide in NFR 1A3b




Figure C.3. Emission Factor Trends for Ammonia in NFR 1A3b




Figure C.4. Emission Factor Trends for Non-Methane Volatile Organic Compounds in NFR 1A3b




Figure C.5. Emission Factor Trends for PM_{2.5} in NFR 1A3b

Annex D

Emission Factors for Industrial Processes (NFR 2)

Table D.1 Emission Factors for NFR 2A

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	2A	Mineral Products	2016
NFR Source Category	Cement Production (2A1)		
Pollutant	Value	Unit	Reference
PCDD/F (I-TEQ)	0.05 to 5	µg I-TEQ / t	UNEP Toolkit (2013)
PCBs	1.000	µg / t	EMEP/CORINAIR Guidebook (2006)
NFR Source Category	Glass Production (2A3)		
Pollutant	Value	Unit	Reference
PCDD/F (I-TEQ)	0.015 to 0.2	µg I-TEQ / t	UNEP Toolkit (2013)
NFR Source Category	Glass Production (2A3) - Lead crystal		
Pollutant	Value	Unit	Reference
TSP	10	g/t	EEA/EMEP Guidebook 2016
PM ₁₀	9	g/t	EEA/EMEP Guidebook 2016
PM _{2.5}	8	g/t	EEA/EMEP Guidebook 2016
BC	0.062	% of PM _{2.5}	EEA/EMEP Guidebook 2016
Pb	10.000	g/t	EEA/EMEP Guidebook 2016
Ni	0.175	g/t	NAEI, 2003
Zn	1.014	g / t	NAEI, 2003
NFR Source Category	Glass Production (2A3) - Container		
Pollutant	Value	Unit	Reference
TSP	280	g/t	EEA/EMEP Guidebook 2016
PM ₁₀	250	g/t	EEA/EMEP Guidebook 2016
PM _{2.5}	220	g/t	EEA/EMEP Guidebook 2016
BC	0.062	% of PM _{2.5}	EEA/EMEP Guidebook 2016
As	0.290	g/t	EEA/EMEP Guidebook 2016
Cd	0.120	g/t	EEA/EMEP Guidebook 2016
Cr	0.370	g/t	EEA/EMEP Guidebook 2016
Pb	2.900	g/t	EEA/EMEP Guidebook 2016
Ni	0.240	g/t	EEA/EMEP Guidebook 2016
Se	1.500	g/t	EEA/EMEP Guidebook 2016

Emission Factors			
NFR Source Category	Glass Production (2A3) - Glass wool		
Pollutant	Value	Unit	Reference
TSP	670	g/t	EEA/EMEP Guidebook 2016
PM ₁₀	590	g/t	EEA/EMEP Guidebook 2016
PM _{2.5}	520	g/t	EEA/EMEP Guidebook 2016
BC	2	% of PM _{2.5}	EEA/EMEP Guidebook 2016
As	0.020	g/t	NAEI, 2003
Cd	0.028	g/t	NAEI, 2003
Cr	0.202	g/t	NAEI, 2003
Cu	0.202	g/t	NAEI, 2003
Pb	0.210	g/t	NAEI, 2003
Hg	0.069	g/t	NAEI, 2003
Ni	0.202	g/t	NAEI, 2003
Se	0.013	g/t	NAEI, 2003
Zn	2.355	g/t	NAEI, 2003
NFR Source Category	Bricks and Ceramic Production (2A6)		
Pollutant	Value	Unit	Reference
PCDD/F (I-TEQ)	0.110	µg I-TEQ / t	URS Dames and Moore (2000) & UNEP Toolkit (2013)
NFR Source Category	Asphalt Production (2A6)		
Pollutant	Value	Unit	Reference
PCDD/F (I-TEQ)	0.007 to 0.07	µg I-TEQ / t	UNEP Toolkit (2013)
NFR Source Category	Quarrying and mining of minerals other than coal (2A5a)		
Pollutant	Value	Unit	Reference
TSP	102	g/t	EEA/EMEP Guidebook 2016
PM ₁₀	50	g/t	EEA/EMEP Guidebook 2016
PM _{2.5}	5	g/t	EEA/EMEP Guidebook 2016
NFR Source Category	Storage handling and transport of mineral products (2A5c)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.6	g/t	EEA/EMEP Guidebook 2016

Table D.1 Emission Factors for NFR 2A (continued)

Emission Factors					
NFR Source Category	Construction and demolition (2A5b)				
Pollutant	Value			Unit	Reference
	PM2.5	PM10	TSP		
Bungalow	0.009	0.086	0.29	(kg m2/year)	EEA/EMEP Guidebook 2016
House Detached	0.009	0.086	0.29	(kg m2/year)	EEA/EMEP Guidebook 2016
House Semi-detached	0.009	0.086	0.29	(kg m2/year)	EEA/EMEP Guidebook 2016
House terraced	0.009	0.086	0.29	(kg m2/year)	EEA/EMEP Guidebook 2016
Flat/Apartment	0.030	0.3	1	(kg m2/year)	EEA/EMEP Guidebook 2016
Non-Residential	0.100	1	3.3	(kg m2/year)	EEA/EMEP Guidebook 2016
Roads	0.230	2.3	7.7	(kg m2/year)	EEA/EMEP Guidebook 2016

Table D.2 Emission Factors for NFR 2B

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	2B	Chemical Industry	2016
NFR Source Category	Nitric Acid Production (2B2)		
Pollutant	Value	Unit	Reference
NO _x	7.25 to 10.08	kg / t	Plant specific

Table D.3 Emission Factors for NFR 2C

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	2C	Metal Production	2016
NFR Source Category	Iron and Steel Production (2C1)		
Pollutant	Value	Unit	Reference
Cd	0.663	g / t	Plant specific
Cr	4.381	g / t	Plant specific
Pb	5.378	g / t	Plant specific
Ni	8.264	g / t	Plant specific
Zn	85.111	g / t	Plant specific
As	0.4	g / t	EEA/EMEP Guidebook 2016
Cu	0.070	g / t	EEA/EMEP Guidebook 2016
Hg	0.100	g / t	EEA/EMEP Guidebook 2016
Se	0.020	g / t	EEA/EMEP Guidebook 2016
PCDD/F (I-TEQ)	0.003	g I-TEQ / kt	EEA/EMEP Guidebook 2016
NFR Source Category	Ferroalloys production (2C2)		
Pollutant	Value	Unit	Reference
HCB	5	g / t	UNECE Guidebook
NFR Source Category	Aluminium Production (2C3)		
Pollutant	Value	Unit	Reference
Zn	2.725	g / t	UK NAEI

Table D.4 Emission Factors for NFR 2D

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	2D	Solvent and product use	2016
NFR Source Category	Domestic Solvent use including fungicides (2D3a) NMVOC		
Category	Value	Unit	Reference
Household products (aerosol)	200.00	g/person	EEA/EMEP Guidebook 2016
Household (cleaning) Products -Non aerosol	252.00	g/person	EEA/EMEP Guidebook 2016
Car Care Products- Aerosol	161.00	g/person	EEA/EMEP Guidebook 2016
Car Care Products-Non Aerosol	303.00	g/person	EEA/EMEP Guidebook 2016
Cosmetics and toiletries-Aerosol	355.00	g/person	EEA/EMEP Guidebook 2016
Cosmetics and toiletries-Non Aerosol	494.00	g/person	EEA/EMEP Guidebook 2016
DIY/Buildings-Adhesives	76.00	g/person	EEA/EMEP Guidebook 2016
DIY/Buildings-Paint thinner	205.00	g/person	EEA/EMEP Guidebook 2016
DIY/Buildings-Paint and varnish remover	68.00	g/person	EEA/EMEP Guidebook 2016
DIY/Buildings-Sealants & Filling Agents	23.00	g/person	EEA/EMEP Guidebook 2016
Pharmaceutical Products	48.00	g/person	EEA/EMEP Guidebook 2016
Pesticides	76.00	g/person	EEA/EMEP Guidebook 2016
NMVOC		g / t	EEA/EMEP Guidebook 2016
NFR Source Category	Road paving with Asphalt (2D3b)		
Pollutant	Value	Unit	Reference
NMVOC	16	g / t	EEA/EMEP Guidebook 2016
TSP	15000	g / t	EEA/EMEP Guidebook 2016
PM10	2000	g / t	EEA/EMEP Guidebook 2016
PM2.5	1000	g / t	EEA/EMEP Guidebook 2016
NFR Source Category	Paint application; vehicle refinishing (2D3d)		
Category	Value	Unit	Reference
Thinner /Prepartory and Cleaning	835	g/L	CS
Body filler	175	g/L	CS
Top Coat	420	g/L	CS
Primer	540	g/L	CS
NFR Source Category	Paint application; Decorative paints Solvent & Water based (2D3d) (NMVOC)		
Category	Value	Unit	Reference
Interior matt walls and ceilings (water based)	23	g/L	CS
Interior glossy walls and ceilings (water based)	50	g/L	CS
Exterior walls of mineral substrate (water based)	9	g/L	CS

Emission Factors			
Interior/exterior trim and cladding paints for wood and metal (water based)	43	g/L	CS
Primers (water based)	45	g/L	CS
Interior matt walls and ceilings (solvent based)	30	g/L	CS
Interior glossy walls and ceilings (solvent based)	76	g/L	CS
Exterior walls of mineral substrate (solvent based)	126	g/L	CS
Interior/exterior trim and cladding paints for wood and metal (solvent based)	324	g/L	CS
Primers (solvent based)	201	g/L	CS
NFR Source Category	Paint application; Marine (2D3d) (NMVOC)		
Category	Value	Unit	Reference
Anti Fouling Emission Factor g/L	440	g/L	CS
Top Coat Emission Factor & Primer Emission Factor g/L	400.0	g/L	CS
NFR Source Category	Degreasing (2D3e)		
Pollutant	Value	Unit	Reference
NMVOC	460.0	g/kg	EEA/EMEP Guidebook 2016
NFR Source Category	Dry Cleaning (2D3e)		
Pollutant	Value	Unit	Reference
NMVOC	0.63	g/kg	CS
NFR Source Category	Preservation of Wood (2D3i)		
Pollutant	Value	Unit	Reference
Benzo(a)pyrene	1.050	mg/kg	EEA/EMEP Guidebook 2016
Benzo(b)fluoranthene	0.530	mg/kg	EEA/EMEP Guidebook 2016
Benzo(k)fluoranthene	0.530	mg/kg	EEA/EMEP Guidebook 2016
Indeno(1,2,3-cd)pyrene	0.530	mg/kg	EEA/EMEP Guidebook 2016
NFR Source Category	Fat Edible and non edible oil extraction (2D3i)		
Pollutant	Value	Unit	Reference
NMVOC	1.570	g/kg	EEA/EMEP Guidebook 2016
TSP	1.1	g/kg	EEA/EMEP Guidebook 2016
PM10	0.9	g/kg	EEA/EMEP Guidebook 2016
PM2.5	0.6	g/kg	EEA/EMEP Guidebook 2016
NFR Source Category	Industrial Adhesive usage (2D3i)		
Pollutant	Value	Unit	Reference
NMVOC	522.000	g/kg	EEA/EMEP Guidebook 2016

Table D.5 Emission Factors for NFR 2G-2L

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	2	Industrial Processes	2016
NFR Source Category	Fireworks (2G)		
Pollutant	Value	Unit	Reference
NOx	260	g/t	EEA/EMEP Guidebook 2016
SO2	3020	g/t	EEA/EMEP Guidebook 2016
CO	7150	g/t	EEA/EMEP Guidebook 2016
TSP	109830	g/t	EEA/EMEP Guidebook 2016
PM10	99920	g/t	EEA/EMEP Guidebook 2016
PM2.5	51940	g/t	EEA/EMEP Guidebook 2016
Cd	1.48	g/t	EEA/EMEP Guidebook 2016
Ni	30	g/t	EEA/EMEP Guidebook 2016
Zn	260	g/t	EEA/EMEP Guidebook 2016
Cu	444	g/t	EEA/EMEP Guidebook 2016
Cr	15.60	g/t	EEA/EMEP Guidebook 2016
Hg	0.06	g/t	EEA/EMEP Guidebook 2016
Pb	784	g/t	EEA/EMEP Guidebook 2016
As	1.33	g/t	EEA/EMEP Guidebook 2016
NFR Source Category	Tobacco (2G)		
Pollutant	Value	Unit	Reference
NOx	1.8	g/t	EEA/EMEP Guidebook 2016
NH3	4.15	g/t	EEA/EMEP Guidebook 2016
CO	55.1	g/t	EEA/EMEP Guidebook 2016
NMVOC	4.8	g/kg	EEA/EMEP Guidebook 2016
TSP	27	g/t	EEA/EMEP Guidebook 2016
PM10	27	g/t	EEA/EMEP Guidebook 2016
PM2.5	27	g/t	EEA/EMEP Guidebook 2016
BC	40	%of PM2.5	EEA/EMEP Guidebook 2016
Cd	5	g/t	EEA/EMEP Guidebook 2016
Ni	3	g/t	EEA/EMEP Guidebook 2016
Zn	3	g/t	EEA/EMEP Guidebook 2016
Cu	5	g/t	EEA/EMEP Guidebook 2016
PCDD/F	0.10	µg I-TEQ/Mg	EEA/EMEP Guidebook 2016
Benzo(a)pyrene	0.11	g/t	EEA/EMEP Guidebook 2016

Emission Factors			
Benzo(b)fluoranthene	0.05	g/t	EEA/EMEP Guidebook 2016
Benzo(k)fluoranthene	0.05	g/t	EEA/EMEP Guidebook 2016
Indeno(1,2,3-cd)pyrene	0.05	g/t	EEA/EMEP Guidebook 2016
NFR Source Category	Bread Baking (2H2) (NMVOC)		
Category	Value	Unit	Reference
Bread	4.50	kg/t	EEA/EMEP Guidebook 2016
Baking Goods	1.00	kg/t	EEA/EMEP Guidebook 2016
NFR Source Category	Alcohol production (2H2) (NMVOC)		
Category	Value	Unit	Reference
Production of Beer	0.04	kg/hL	EEA/EMEP Guidebook 2016
Production of Spirits	15.00	kg/hL	EEA/EMEP Guidebook 2016
NFR Source Category	Meat/fish frying/curing (2H2) (NMVOC)		
Pollutant	Value	Unit	Reference
NMVOC	0.33	kg/t	EEA/EMEP Guidebook 2016
NFR Source Category	Coffee roasting (2H2)		
Pollutant	Value	Unit	Reference
NMVOC	0.55	kg/t	EEA/EMEP Guidebook 2016
NFR Source Category	Animal feedstock (2H2)		
Pollutant	Value	Unit	Reference
NMVOC	1.00	kg/t	EEA/EMEP Guidebook 2016
NFR Source Category	Leakage from Electrical Equipment (2L)		
Pollutant	Value	Unit	Reference
PCBs	0.060	g/kgfluid	UK NAEI (2006)
PCDD/F (I-TEQ)	83.500	µg I-TEQ / kg PCB	Dyke (1997)
NFR Source Category	Fragmentisers & Shredders (2L)		
Pollutant	Value	Unit	Reference
PCBs	0.004	g/capita/year	UNECE Guidebook (2000)

Annex E

Agricultural Activity Data and Emission Factors (NFR 3)

Table E.1 Activity Data for Agriculture

Population (1000s)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Total Cattle	6822	6921	6973	6959	6965	7009	7282	7491	7592	7348	7012	7022	6961	6971	6973	6951	6925	6827	6828	6801	6555	6428	6691	6829	6840	6926	7173
Dairy Cows	1341	1309	1262	1256	1247	1239	1241	1227	1216	1187	1165	1165	1146	1146	1139	1025	1054	1054	1060	1060	1039	1076	1101	1123	1177	1268	1347
All Other Cattle	5481	5612	5711	5703	5718	5770	6041	6264	6376	6160	5847	5857	5814	5826	5834	5926	5872	5773	5768	5742	5516	5352	5590	5706	5663	5658	5827
Other Cows	730	801	903	958	990	1022	1098	1183	1222	1192	1171	1178	1153	1166	1179	1121	1171	1185	1198	1169	1125	1103	1138	1118	1085	1065	1073
Dairy Heifers	172	156	187	191	207	230	238	244	226	212	205	202	223	221	234	214	204	197	195	196	234	252	268	271	317	331	325
Other Heifers	80	71	106	116	104	123	134	149	128	121	133	140	142	139	142	191	193	212	180	156	170	202	181	149	174	188	184
Cattle < 1 yrs	1716	1765	1695	1738	1736	1746	1852	1938	1965	1821	1752	1824	1799	1761	1771	1962	1953	1941	1959	1889	1761	1846	2036	1969	1878	2042	2126
Cattle < 1 yrs - male	903	919	889	914	904	915	974	1023	1055	965	919	955	953	922	930	958	951	947	969	918	827	892	1023	959	902	994	1046
Cattle < 1 yrs - female	813	846	806	824	832	831	878	915	910	856	833	869	846	839	842	1005	1002	994	990	971	935	954	1013	1009	977	1048	1080
Cattle 1 - 2 yrs	1663	1692	1638	1587	1586	1586	1639	1717	1783	1706	1517	1515	1593	1577	1535	1642	1506	1466	1496	1542	1408	1270	1376	1551	1469	1373	1517
Cattle 1 - 2 yrs - male	986	981	982	958	952	964	996	1055	1086	1039	912	913	992	983	950	972	845	818	832	851	760	673	770	873	821	790	873
Cattle 1 - 2 yrs - female	677	711	656	630	634	622	643	662	697	667	605	602	601	594	585	670	661	648	664	690	647	597	606	678	648	583	644
Cattle > 2 yrs	1093	1099	1152	1078	1058	1023	1036	986	1002	1058	1016	941	845	902	911	734	782	715	687	738	772	640	554	609	701	628	579
Cattle > 2 yrs - male	826	798	830	773	740	712	732	690	708	737	722	642	560	599	605	537	565	510	476	501	506	426	361	388	456	424	391
Cattle > 2 yrs - female	266	301	322	305	318	311	304	296	294	321	295	299	284	303	305	197	217	206	211	237	265	214	193	221	245	204	188
Bulls	27	29	32	36	38	40	44	48	50	51	53	56	59	60	63	61	63	57	54	52	47	38	37	39	38	33	22
Total Sheep	8021	8484	8736	8977	8559	8364	8329	8051	8572	8547	7957	7455	6682	6481	6703	6431	6187	5656	5105	4727	4328	4429	4843	4918	5019	4870	4770
Ewes Lowland	2397	2543	2622	2576	2511	2427	2369	2390	3056	2936	2814	2704	2637	2552	2464	2627	2414	2207	2057	1928	1920	1954	2036	2016	1978	1960	1964
Ewes Upland	1961	2080	2145	2108	2055	1986	1938	1955	1310	1258	1206	1159	1130	1094	1056	657	604	552	514	482	480	489	509	504	494	490	491
Rams Lowland	64	67	70	69	67	66	62	64	81	79	77	75	73	72	70	77	74	69	63	58	59	59	61	62	60	60	61
Rams Upland	53	55	57	56	55	54	51	52	35	34	33	32	31	31	30	19	19	17	16	14	15	15	15	15	15	15	15
Other Sheep>1 - Lowland	164	96	89	99	107	113	106	118	172	153	143	128	129	144	140	124	122	109	112	103	96	101	116	112	97	110	70
Other Sheep>1 - Upland	134	79	73	81	88	92	86	97	74	66	61	55	55	62	60	31	31	27	28	26	24	25	29	28	24	27	17
Lambs - Lowland	1787	1960	2024	2194	2022	1994	2044	1856	2692	2815	2535	2312	1838	1768	2019	2317	2339	2140	1853	1693	1387	1429	1661	1745	1880	1766	1722
Lambs - Upland	1462	1604	1656	1795	1654	1632	1672	1519	1154	1206	1086	991	788	758	865	579	585	535	463	423	347	357	415	436	470	442	430
Total Pigs	1222	1325	1404	1504	1514	1546	1643	1708	1810	1775	1727	1760	1791	1729	1704	1679	1632	1544	1486	1444	1508	1551	1532	1511	1530	1506	1561
Gilts in Pig	21	22	25	23	22	24	25	27	26	25	21	23	20	20	22	20	22	21	21	20	19	19	20	19	20	20	19
Gilts not yet Served	12	14	15	14	15	18	17	18	19	16	18	19	20	18	19	20	19	16	16	17	15	15	15	15	15	15	16
Sows in Pig	83	90	96	101	99	100	103	108	109	109	110	107	110	104	102	100	96	96	91	89	92	90	84	82	83	82	82
Other Sows for Breeding	31	31	33	33	30	31	36	37	38	38	32	37	33	32	30	34	31	28	25	27	29	27	25	29	30	27	30
Boars	6	7	7	6	6	5	5	5	5	4	4	4	3	3	3	2	2	2	2	2	2	1	1	1	1	1	1
Pigs 20 Kg +	749	803	837	905	918	952	1016	1064	1144	1094	1038	1036	1062	1043	1028	1010	1034	939	932	911	953	965	960	926	941	934	977
Pigs Under 20 Kg	319	358	392	422	425	417	442	450	469	489	504	535	543	508	500	494	429	443	400	378	400	434	426	438	440	427	436
Total Poultry	11772	12698	13272	13072	14034	14438	15375	15548	15686	15490	15680	16024	15545	16152	17190	16573	15934	13324	13258	15277	15212	15032	15631	14989	16504	16993	17181
Layer	1868	1800	2231	1832	1730	1371	1701	1580	1559	1537	1572	1676	1613	1907	1906	1950	1970	1813	1813	2145	2145	2145	2600	2828	2917	3268	3318
Broiler	8035	8905	9067	9522	10393	11092	11730	12096	12287	12200	12426	12629	12322	12672	13375	12818	12360	9696	9696	11904	11904	11520	11520	10764	12127	12223	12318
Turkey	1509	1633	1615	1358	1552	1616	1585	1513	1482	1393	1322	1358	1248	1209	1461	1274	1097	1330	1330	874	874	1078	1222	1125	1189	1231	1273
Ducks	347	347	347	347	347	347	347	347	347	347	347	347	348	349	350	435	520	497	475	409	344	279	279	265	265	265	265
Geese	12	12	12	12	12	12	12	12	12	12	12	13	13	13	12	11	10	10	10	10	10	10	10	7	7	7	7
Horses	62	63	65	66	67	68	70	72	73	76	70	71	73	70	73	80	87	89	96	98	106	106	111	102	95	93	92
Mules	8	7	8	9	8	7	8	7	8	7	5	5	5	6	6	6	7	7	9	9	8	9	10	8	8	9	9
Goats	17	17	18	18	16	16	15	15	15	14	8	8	8	8	8	7	7	7	9	10	11	11	10	9	9	11	10
Farmed Deer	12	12	13	15	15	16	16	18	17	16	12	12	12	11	11	10	9	10	10	9	5	3	2	2	2	1	1
Mink	185	143	130	124	124	124	133	143	143	143	146	146	146	146	146	149	149	149	149	190	183	183	198	198	198	198	198
Fox	26	22	13	9	7	7	8	8	9	9	4	4	4	4	4	2	2	3	1	0	0	0	NO	NO	NO	NO	NO
Fertiliser (1000's tonnes/N)	379.3	370.1	358.3	378.0	404.8	428.8	416.9	380.4	432.0	442.9	407.6	368.7	363.5	388.1	362.5	352.2	342.1	321.6	309.0	306.8	362.4	295.8	296.5	353.0	331.8	331.0	339.1

Table E.2.1 Input Data on Manure Management Practices - Cattle

Cattle	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Number of days housed																											
Dairy Cows	118	118	118	118	118	118	118	118	118	118	118	118	118	118	118	118	118	117	117	117	117	117	117	117	117	117	117
Suckler Cows	141	141	141	141	141	141	141	141	141	141	141	141	141	141	141	142	141	141	141	141	141	141	142	141	141	142	141
Dairy Heifer	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128
Other Heifer	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139
Under1yr	221	221	221	221	221	221	221	221	221	221	221	221	221	221	221	222	222	222	223	223	227	224	224	223	222	221	221
Oneto2yrs	156	156	156	156	156	156	156	156	156	156	156	156	156	156	154	155	154	154	154	156	157	154	153	153	156	155	153
Over2yrs	23	23	23	23	23	23	23	23	23	23	23	23	23	23	25	26	24	26	23	26	26	23	21	22	22	28	26
Bulls	156	156	156	156	156	156	156	156	156	156	156	156	156	156	154	155	154	154	154	156	157	154	153	153	156	155	153
Number of days grazing																											
Dairy Cows	247	247	247	247	247	247	247	247	247	247	247	247	247	247	247	247	247	248	248	248	248	248	248	248	248	248	248
Suckler Cows	224	224	224	224	224	224	224	224	224	224	224	224	224	224	224	223	224	224	224	224	224	224	223	224	224	223	224
Dairy Heifer	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237
Other Heifer	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226
Under1yr	144	144	144	144	144	144	144	144	144	144	144	144	144	144	144	143	143	143	142	142	138	141	141	142	143	144	144
Oneto2yrs	209	209	209	209	209	209	209	209	209	209	209	209	209	209	211	210	211	211	211	209	208	211	212	212	209	210	212
Over2yrs	342	342	342	342	342	342	342	342	342	342	342	342	342	342	340	339	341	339	342	339	339	342	344	343	343	337	339
Bulls	209	209	209	209	209	209	209	209	209	209	209	209	209	209	211	210	211	211	211	209	208	211	212	212	209	210	212
Proportion to each AWMS																											
Liquid																											
Dairy Cows	0.32	0.32	0.32	0.32	0.31	0.31	0.31	0.31	0.31	0.30	0.30	0.30	0.30	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.28	0.28
Suckler Cows	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.27	0.26	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27
Dairy Heifer	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
Other Heifer	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38
Under1yr	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.42	0.41	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.41	0.41
Oneto2yrs	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.33	0.34	0.34	0.34	0.34	0.33	0.33	0.33	0.34	0.34	0.33
Over2yrs	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.04	0.04	0.04
Bulls	0.42	0.42	0.42	0.41	0.41	0.41	0.40	0.40	0.40	0.39	0.39	0.39	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38
Solid																											
Dairy Cows	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Suckler Cows	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Dairy Heifer	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Other Heifer	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Under1yr	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.18	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.19	0.19	0.19
Oneto2yrs	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09
Over2yrs	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.03	0.03	0.03
Bulls	0.14	0.14	0.14	0.14	0.14	0.14	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13
Pasture																											
Dairy Cows	0.66	0.66	0.66	0.66	0.67	0.67	0.67	0.67	0.68	0.68	0.68	0.68	0.69	0.69	0.69	0.69	0.69	0.69	0.69	0.69	0.70	0.70	0.69	0.69	0.69	0.70	0.70
Suckler Cows	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63
Dairy Heifer	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65
Other Heifer	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62
Under1yr	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.38	0.39	0.39	0.39	0.39	0.39	0.39
Oneto2yrs	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.58	0.58	0.58	0.58	0.58	0.57	0.57	0.58	0.58	0.58	0.57	0.58	0.58
Over2yrs	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.93	0.93	0.93	0.93	0.94	0.93	0.93	0.94	0.94	0.94	0.94	0.92	0.93
Bulls	0.44	0.44	0.45	0.45	0.45	0.46	0.46	0.47	0.47	0.48	0.48	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49

Table E.2.2 Input Data on Manure Management Practices - Other Livestock

Animal Category	Days housed	% housed	% outwintered	Housing Type		Proportion to each AWMS		
				% Slurry based	% Straw based	Liquid	Solid	Pasture
Sheep								
Lowland Ewes	84.00	47.07	52.93	0.00	100.00	NA	0.23	0.77
Upland Ewes	85.00	44.34	55.66	0.00	100.00	NA	0.23	0.77
Rams	85.00	22.34	77.66	0.00	100.00	NA	0.15	0.85
Lambs	28.00	100.00	0.00	0.00	100.00	NA	0.03	0.97
Other sheep	67.00	7.00	93.00	0.00	100.00	NA	0.08	0.92
Pigs								
Gilts in pig	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Gilts not yet served	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Sows in pig	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Other sows for breeding	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Boars	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Pigs < 20 kg	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Pigs > 20 kg	365.00	100.00	0.00	100.00	0.00	1.00	0.00	0.00
Poultry								
Layers	365.00	88.00	12.00	84.20	15.80	0.00	0.88	0.12
Broilers	365.00	100.00	0.00	0.00	100.00	0.00	1.00	0.00
Turkeys	365.00	100.00	0.00	0.00	100.00	0.00	1.00	0.00
Horses	180.00	100.00	0.00	0.00	100.00	0.00	0.49	0.51
Mules and Asses	180.00	100.00	0.00	0.00	100.00	0.00	0.49	0.51
Goats	84.00	100.00	0.00	0.00	100.00	0.00	0.23	0.77

Table E.3 Nitrogen excretion (kg/head/year)

[illegible]

Table E.4 Fertiliser Compound Statistics and Emission Estimates 1990-2016

Tonnes N/year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Ammonium sulphate	6,522	5,219	4,419	3,755	2,408	2,245	1,680	1,297	1,288	1,259	1,047	1,203	1,173	1,348	1,912	802	2,003	2,744	2,239	2,266	2,097	2,845	3,687	4,044	2,973	2,511	2,078
CAN	139,145	134,804	124,731	138,893	151,775	166,035	160,721	145,864	165,330	167,928	153,646	134,413	127,219	137,384	115,148	126,011	116,632	118,789	130,704	141,810	149,171	123,280	136,006	148,901	137,119	128,666	124,866
NK mixtures	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	3,338	3,015	2,835	2,439	2,691	1,880	2,009	3,336	3,906	2,842	2,839	2,249	2,379	2,861	2,782	2,252	2,392
NPK mixtures	169,725	164,430	152,144	169,418	185,131	202,525	196,043	177,921	201,665	204,834	186,583	172,215	176,005	192,486	196,145	179,398	178,847	158,834	125,966	100,661	143,487	122,404	124,653	166,446	153,436	156,411	157,748
NP mixtures	2,588	2,507	2,320	2,583	2,823	3,088	2,989	2,713	3,075	3,123	2,631	2,165	1,948	1,556	1,185	976	892	768	1,484	1,007	368	376	586	1,254	1,265	1,504	1,619
Other straight N compo	688	666	616	686	750	821	794	721	817	830	2,741	3,195	3,647	3,636	3,506	5,056	1,458	5,269	2,714	2,411	2,846	570	151	NO	NO	NO	NO
Urea	60,643	62,494	74,072	62,649	61,924	54,113	54,690	51,834	59,824	64,944	57,611	52,463	50,687	49,230	41,938	38,042	40,296	31,848	41,947	55,809	61,587	44,076	29,075	29,538	34,209	38,598	48,817
Protected urea	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	1,197	1,584
Total	379,311	370,121	358,302	377,985	404,811	428,826	416,918	380,350	431,999	442,918	407,597	368,669	363,514	388,080	362,524	352,166	342,137	321,588	308,960	306,805	362,395	295,800	296,537	353,044	331,784	331,139	339,104
EF (g per kg)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Ammonium sulphate	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90	90
CAN	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
NK mixtures	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
NPK mixtures	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
NP mixtures	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
Other straight N compo	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Urea	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155	155
Protected urea	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
Tonnes NH ₃ /year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Ammonium sulphate	587	470	398	338	217	202	151	117	116	113	94	108	106	121	172	72	180	247	202	204	189	256	332	364	268	226	187
CAN	1,113	1,078	998	1,111	1,214	1,328	1,286	1,167	1,323	1,343	1,229	1,075	1,018	1,099	921	1,008	933	950	1,046	1,134	1,193	986	1,088	1,191	1,097	1,029	999
NK mixtures	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	50	45	43	37	40	28	30	50	59	43	43	34	36	43	42	34	36
NPK mixtures	2,546	2,466	2,282	2,541	2,777	3,038	2,941	2,669	3,025	3,073	2,799	2,583	2,640	2,887	2,942	2,691	2,683	2,383	1,889	1,510	2,152	1,836	1,870	2,497	2,302	2,346	2,366
NP mixtures	39	38	35	39	42	46	45	41	46	47	39	32	29	23	18	15	13	12	22	15	6	6	9	19	19	23	24
Other straight N compo	7	7	6	7	8	8	8	7	8	8	27	32	36	36	35	51	15	53	27	24	28	6	2	NO	NO	NO	NO
Urea	9,400	9,687	11,481	9,711	9,598	8,388	8,477	8,034	9,273	10,066	8,930	8,132	7,856	7,631	6,500	5,897	6,246	4,936	6,502	8,650	9,546	6,832	4,507	4,578	5,302	5,983	7,567
Protected urea	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	10	13
Other	13,691	13,745	15,200	13,747	13,856	13,010	12,907	12,035	13,791	14,651	13,169	12,008	11,728	11,835	10,629	9,761	10,100	8,630	9,746	11,581	13,157	9,955	7,842	8,692	9,029	9,650	11,192

Table E.5 Timing of Slurry Spreading – Cattle

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Spring	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.52	0.52	0.52	0.52	0.52	0.52	0.52
Summer	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.36	0.36	0.36	0.36	0.36	0.36	0.36
Autumn	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.12	0.12	0.12	0.12	0.12	0.12	0.12
Winter	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table E.6.1 NMVOC Emission Factors for NFR 3B

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	3	Agriculture	2016
NFR Source Category	Manure management - cattle (3B1 a&b)		
Pollutant	Value	Unit	Reference
NMVOC	0.0002002	kgNMVOC/kg MJ feed intake	Inventory Guidebook (2016) (silage feeding)
NMVOC	0.00005005	kgNMVOC/kg MJ feed intake	Inventory Guidebook (2016) (silage store)
NMVOC	0.0000353	kgNMVOC/kg MJ feed intake	Inventory Guidebook (2016) (house)
NFR Source Category	Manure management - sheep (3B2)		
Pollutant	Value	Unit	Reference
NMVOC	0.001614	kgNMVOC/kgVS	Inventory Guidebook (2016) (Housing)
NMVOC	1.2619	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure store ratio)
NMVOC	0.787977	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure application ratio)
NMVOC	0.000023	kgNMVOC/kgVS	Inventory Guidebook (2016) (Grazing)
NFR Source Category	Manure management - pigs (3B3)		
Pollutant	Value	Unit	Reference
NMVOC	0.001703	kgNMVOC/kgVS	Inventory Guidebook (2016) (Housing, gilts)
NMVOC	0.007042	kgNMVOC/kgVS	Inventory Guidebook (2016) (Housing, sow s)
NMVOC	0.001703	kgNMVOC/kgVS	Inventory Guidebook (2016) (Housing, other pigs)
NFR Source Category	Manure management - goats (3B4d)		
Pollutant	Value	Unit	Reference
NMVOC	0.001614	kgNMVOC/kgVS	Inventory Guidebook (2016) (House)
NMVOC	3.101	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure store ratio)
NMVOC	3.887	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure application ratio)
NMVOC	0.0000235	kgNMVOC/kgVS	Inventory Guidebook (2016) (Grazing)
NFR Source Category	Manure management - horses (3B4e)		
Pollutant	Value	Unit	Reference
NMVOC	0.001614	kgNMVOC/kgVS	Inventory Guidebook (2016) (House)
NMVOC	1.2326	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure store ratio)
NMVOC	1.014	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure application ratio)
NMVOC	0.0000235	kgNMVOC/kgVS	Inventory Guidebook (2016) (Grazing)
NFR Source Category	Manure management - mules & asses (3B4f)		
Pollutant	Value	Unit	Reference
NMVOC	0.001614	kgNMVOC/kgVS	Inventory Guidebook (2016) (House)
NMVOC	1.2326	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure store ratio)
NMVOC	1.014	kgNMVOC/kgVS	Inventory Guidebook (2016) (Manure application ratio)
NMVOC	0.0000235	kgNMVOC/kgVS	Inventory Guidebook (2016) (Grazing)
NFR Source Category	Manure management - poultry (3B4g)		
Pollutant	Value	Unit	Reference
NMVOC	0.165	kg/AAP/yr	Inventory Guidebook (2016) (Layers)
NMVOC	0.108	kg/AAP/yr	Inventory Guidebook (2016) (Broilers)
NMVOC	0.489	kg/AAP/yr	Inventory Guidebook (2016) (Other poultry)
NFR Source Category	Manure management - other animals (3B4h)		
Pollutant	Value	Unit	Reference
NMVOC	0.045	kg/AAP/yr	Inventory Guidebook (2016) (Farmed Deer)
NMVOC	1.941	kg/AAP/yr	Inventory Guidebook (2016) (Mink & Fox)

Table E.6.2 Feed intake values for silage and grass consumed by cattle

Silage Feed intake (MJ/yr)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Dairy Cows	21078	21031	20984	20937	20890	20843	20796	20750	20703	20656	20609	20562	20515	20468	20504	20813	20822	20840	20629	20515	20923	20936	20650	20809	20769	21067	20953
Other Cows	19612	19594	19575	19556	19537	19519	19500	19481	19462	19444	19425	19406	19387	19369	19252	18825	19212	19525	18838	19733	19709	19315	18749	19576	19390	19008	19202
Dairy Heifers	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844
Other Heifers	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844
Cattle <1 yrs -male	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647	9647
Cattle <1 yrs -female	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704	8704
Cattle 1-2 yrs - male	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748	15748
Cattle 1-2 yrs - female	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844	20844
Cattle >2yrs - male	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310	1310
Cattle >2yrs - female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bulls	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344	18344
Grass Feed intake (MJ/yr)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Dairy Cows	51624	51878	52132	52385	52639	52893	53147	53401	53655	53909	54162	54416	54670	54924	54908	57047	57113	57139	56106	55040	58213	58385	56573	57191	56991	58944	58179
Other Cows	40899	40897	40894	40892	40889	40887	40884	40881	40879	40876	40874	40871	40869	40866	40841	40958	40940	40790	40956	40879	40863	41007	41176	40908	41077	41087	41067
Dairy Heifers	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170
Other Heifers	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170
Cattle <1 yrs -male	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034	10034
Cattle <1 yrs -female	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531	8531
Cattle 1-2 yrs - male	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167	28167
Cattle 1-2 yrs - female	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170	26170
Cattle >2yrs - male	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544	14544
Cattle >2yrs - female	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013	3013
Bulls	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127	34127

Table E.7 TSP Emission Factors for NFR 3B

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	3	Agriculture	2016
NFR Source Category	Manure management - cattle (3B1 a&b)		
Pollutant	Value	Unit	Reference
TSP	1.380	kg/AAP/yr	Inventory Guidebook (2016) (Dairy cow s - slurry)
TSP	1.380	kg/AAP/yr	Inventory Guidebook (2016) (Dairy cow s - solid)
TSP	0.590	kg/AAP/yr	Inventory Guidebook (2016) (Other cattle - slurry)
TSP	0.590	kg/AAP/yr	Inventory Guidebook (2016) (Other cattle - solid)
TSP	0.340	kg/AAP/yr	Inventory Guidebook (2016) (Calves - slurry)
TSP	0.340	kg/AAP/yr	Inventory Guidebook (2016) (Calves - solid)
NFR Source Category	Manure management - sheep (3B2)		
Pollutant	Value	Unit	Reference
TSP	0.140	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - pigs (3B3)		
Pollutant	Value	Unit	Reference
TSP	1.05	kg/AAP/yr	Inventory Guidebook (2016) (fattening pigs)
TSP	0.27	kg/AAP/yr	Inventory Guidebook (2016) (w eaners)
TSP	0.62	kg/AAP/yr	Inventory Guidebook (2016) (sow s)
NFR Source Category	Manure management - goats (3B4d)		
Pollutant	Value	Unit	Reference
TSP	0.140	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - horses (3B4e)		
Pollutant	Value	Unit	Reference
TSP	0.48	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - mules & asses (3B4f)		
Pollutant	Value	Unit	Reference
TSP	0.34	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - poultry (3B4g)		
Pollutant	Value	Unit	Reference
TSP	0.190	kg/AAP/yr	Inventory Guidebook (2016) (Layers)
TSP	0.040	kg/AAP/yr	Inventory Guidebook (2016) (Broilers)
TSP	0.110	kg/AAP/yr	Inventory Guidebook (2016) (Turkeys)
TSP	0.240	kg/AAP/yr	Inventory Guidebook (2016) (Geese)
TSP	0.140	kg/AAP/yr	Inventory Guidebook (2016) (Ducks)
NFR Source Category	Manure management - other animals (3B4h)		
Pollutant	Value	Unit	Reference
TSP	0.018	kg/AAP/yr	Inventory Guidebook (2016) (Mink & Fox)

Table E.8 PM₁₀ Emission Factors for NFR 3B

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	3	Agriculture	2016
NFR Source Category	Manure management - cattle (3B1 a&b)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.630	kg/AAP/yr	Inventory Guidebook (2016) (Dairy cows - slurry)
PM ₁₀	0.630	kg/AAP/yr	Inventory Guidebook (2016) (Dairy cows - solid)
PM ₁₀	0.270	kg/AAP/yr	Inventory Guidebook (2016) (Other cattle - slurry)
PM ₁₀	0.240	kg/AAP/yr	Inventory Guidebook (2016) (Other cattle - solid)
PM ₁₀	0.160	kg/AAP/yr	Inventory Guidebook (2016) (Calves - slurry)
PM ₁₀	0.160	kg/AAP/yr	Inventory Guidebook (2016) (Calves - solid)
NFR Source Category	Manure management - sheep (3B2)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.0600	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - pigs (3B3)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.14	kg/AAP/yr	Inventory Guidebook (2016) (fattening pigs)
PM ₁₀	0.05	kg/AAP/yr	Inventory Guidebook (2016) (weaners)
PM ₁₀	0.17	kg/AAP/yr	Inventory Guidebook (2016) (sows)
NFR Source Category	Manure management - goats (3B4d)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.0600	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - horses (3B4e)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.22	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - mules & asses (3B4f)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.16	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - poultry (3B4g)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.040	kg/AAP/yr	Inventory Guidebook (2016) (Layers)
PM ₁₀	0.020	kg/AAP/yr	Inventory Guidebook (2016) (Broilers)
PM ₁₀	0.110	kg/AAP/yr	Inventory Guidebook (2016) (Turkeys)
PM ₁₀	0.240	kg/AAP/yr	Inventory Guidebook (2016) (Geese)
PM ₁₀	0.140	kg/AAP/yr	Inventory Guidebook (2016) (Ducks)
NFR Source Category	Manure management - other animals (3B4h)		
Pollutant	Value	Unit	Reference
PM ₁₀	0.0080	kg/AAP/yr	Inventory Guidebook (2016) (Mink & Fox)

Table E.9 PM_{2.5} Emission Factors for NFR 3B

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	3	Agriculture	2016
NFR Source Category	Manure management - cattle (3B1 a&b)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.410	kg/AAP/yr	Inventory Guidebook (2016) (Dairy cow s - slurry)
PM _{2.5}	0.410	kg/AAP/yr	Inventory Guidebook (2016) (Dairy cow s - solid)
PM _{2.5}	0.180	kg/AAP/yr	Inventory Guidebook (2016) (Other cattle - slurry)
PM _{2.5}	0.180	kg/AAP/yr	Inventory Guidebook (2016) (Other cattle - solid)
PM _{2.5}	0.100	kg/AAP/yr	Inventory Guidebook (2016) (Calves - slurry)
PM _{2.5}	0.100	kg/AAP/yr	Inventory Guidebook (2016) (Calves - solid)
NFR Source Category	Manure management - sheep (3B2)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.0200	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - pigs (3B3)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.006	kg/AAP/yr	Inventory Guidebook (2016) (fattening pigs)
PM _{2.5}	0.002	kg/AAP/yr	Inventory Guidebook (2016) (w eaners)
PM _{2.5}	0.010	kg/AAP/yr	Inventory Guidebook (2016) (sow s)
NFR Source Category	Manure management - goats (3B4d)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.0200	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - horses (3B4e)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.14	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - mules & asses (3B4f)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.10	kg/AAP/yr	Inventory Guidebook (2016)
NFR Source Category	Manure management - poultry (3B4g)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.003	kg/AAP/yr	Inventory Guidebook (2016) (Layers)
PM _{2.5}	0.002	kg/AAP/yr	Inventory Guidebook (2016) (Broilers)
PM _{2.5}	0.020	kg/AAP/yr	Inventory Guidebook (2016) (Turkeys)
PM _{2.5}	0.030	kg/AAP/yr	Inventory Guidebook (2016) (Geese)
PM _{2.5}	0.020	kg/AAP/yr	Inventory Guidebook (2016) (Ducks)
NFR Source Category	Manure management - other animals (3B4h)		
Pollutant	Value	Unit	Reference
PM _{2.5}	0.0040	kg/AAP/yr	Inventory Guidebook (2016) (Mink & Fox)

Table E.10 Emission Factors for NFR 3D

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	3	Agriculture	2016
NFR Source Category	Inorganic N-fertilizers (includes also urea application) (3Da1)		
Pollutant	Value	Unit	Reference
TSP	1.56	kg/ha	Inventory Guidebook (2016)
PM ₁₀	1.56	kg/ha	Inventory Guidebook (2016)
PM _{2.5}	0.06	kg/ha	Inventory Guidebook (2016)
NFR Source Category	Farm-level agricultural operations including storage, handling and transport of agricultural products (3Dc)		
Pollutant	Value	Unit	Reference
PM ₁₀	32.00	kg/ha	Inventory Guidebook (2016)
PM _{2.5}	4.00	kg/ha	Inventory Guidebook (2016)
TSP	100.00	kg/ha	Inventory Guidebook (2016) PM ₁₀ value
NFR Source Category	Off-farm storage, handling and transport of bulk agricultural products (3Dd)		
Pollutant	Value	Unit	Reference
TSP	100.00	g / t	CORINAIR
PM ₁₀	25.00	g / t	CORINAIR
PM _{2.5}	4.00	g / t	CORINAIR
NFR Source Category	Cultivated crops (3De)		
Pollutant	Value	Unit	Reference
NM/OC	0.86	kg/ha	Inventory Guidebook (2016)

Table E.11 Activity data associated with NMVOC and PM emission calculations for NFR 3D

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Utilised agricultural area ('000 ha)	4442	4442	4413	4404	4391	4389	4341	4432	4415	4418	4443	4410	4372	4370	4305	4302	4261	4276	4629	4594	4569	4556	4533	4478	4466	4430	4447
Wheat, Oats & Barley (kt)	1965	1965	2018	1626	1610	1796	2142	1943	1865	2011	2174	2165	1964	2147	2501	1940	2083	1997	2461	2063	2040	2509	2125	2401	2598	2634	2311
Total fertilizer (kt)	1793	1745	1646	1767	1820	1921	1896	1699	1830	1850	1730	1546	1523	1628	1538	1479	1396	1330	1242	1172	1424	1204	1231	1487	1403	1395	1412

Annex F

Emission Factors for Waste (NFR 5)

Table F.1 Emission Factors for NFR 5A and NFR 5C1

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	5	Waste	2016
NFR Source Category	Solid Waste Disposal on Land (5A)		
Pollutant	Value	Unit	Reference
NMVOC	5.65	g NMVOC/m ³ landfill gas	2016 EMEP/EEA Guidebook
TSP	0.463	g/Mg	2016 EMEP/EEA Guidebook
PM ₁₀	0.219	g/Mg	2016 EMEP/EEA Guidebook
PM _{2.5}	0.033	g/Mg	2016 EMEP/EEA Guidebook
PCDD/F (I-TEQ)	0.953	µg I-TEQ/tonne of escaping gas	UK NAEI
PCDD/F (I-TEQ)	0.614	µg I-TEQ/tonne of flared gas	UK NAEI
PCBs	0.0008	kg/ktonne of escaping gas	UK NAEI
NFR Source Category	Clinical Waste Incineration (5C1biii)		
Pollutant	Value	Unit	Reference
NO _x	2.3	kg/Mg waste	2016 EMEP/EEA Guidebook
CO	0.19	kg/Mg waste	2016 EMEP/EEA Guidebook
NMVOC	0.70	kg/Mg waste	2016 EMEP/EEA Guidebook
SO _x	0.54	kg/Mg waste	2016 EMEP/EEA Guidebook
TSP	17.0	kg/Mg waste	2016 EMEP/EEA Guidebook
PM ₁₀	11.9	kg/Mg waste	2016 EMEP/EEA Guidebook
PM _{2.5}	6.8	kg/Mg waste	2016 EMEP/EEA Guidebook
BC	2.3	% of TSP	2016 EMEP/EEA Guidebook
As	0.10	g/tonne	2016 EMEP/EEA Guidebook
Cd	3.0	g/tonne	2016 EMEP/EEA Guidebook

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	5	Waste	2016
NFR Source Category	Industrial (incl. Hazardous & Sludge) Waste Incineration (5C1bii,iv)		
Pollutant	Value	Unit	Reference
NO _x	0.87	kg/Mg waste	2016 EMEP/EEA Guidebook
CO	0.07	kg/Mg waste	2016 EMEP/EEA Guidebook
NMVOC	7.4	kg/Mg waste	2016 EMEP/EEA Guidebook
SO _x	0.047	kg/Mg waste	2016 EMEP/EEA Guidebook
TSP	0.010	kg/Mg waste	2016 EMEP/EEA Guidebook
PM ₁₀	0.007	kg/Mg waste	2016 EMEP/EEA Guidebook
PM _{2.5}	0.004	kg/Mg waste	2016 EMEP/EEA Guidebook
BC	3.5	% of PM _{2.5}	2016 EMEP/EEA Guidebook
As	0.016	g/tonne	2016 EMEP/EEA Guidebook
Cd	0.100	g/tonne	2016 EMEP/EEA Guidebook
Cr	0.121	g/tonne	UK NAEI
Cu	0.085	g/tonne	UK NAEI
Pb	1.300	g/tonne	2016 EMEP/EEA Guidebook
Hg	0.056	g/tonne	2016 EMEP/EEA Guidebook
Ni	0.140	g/tonne	2016 EMEP/EEA Guidebook
Zn	NE		2016 EMEP/EEA Guidebook
PCBs	1.223 - 4.83	kg/Mt burned	UK NAEI
HCB	0.5 - 1.975	kg/Mt burned	UK NAEI
Benzo[a]pyrene	0.15 - 0.592	kg/Mt burned	UK NAEI

Cr	0.4	g/tonne	2016 EMEP/EEA Guidebook
Cu	0.6	g/tonne	2016 EMEP/EEA Guidebook
Pb	364	g/tonne	2016 EMEP/EEA Guidebook
Hg	54	g/tonne	2016 EMEP/EEA Guidebook
Ni	0.3	g/tonne	2016 EMEP/EEA Guidebook
Zn	0.019	g/tonne	UK NAEI
PCDD/F	372.1	g-I-TEQ/mt burnt	UK NAEI
PCBs	0.77 - 3.15	kg/Mt burned	UK NAEI
HCB	0.5 - 2.053	kg/Mt burned	UK NAEI
Benzo[a]pyrene	0.7 - 2.875	kg/Mt burned	UK NAEI
Benzo[b]fluoranthene	3.15 - 12.937	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	3.15 - 12.937	kg/Mt burned	UK NAEI

Benzo[b]fluoranthene	0.65 - 2.567	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	0.65 - 2.567	kg/Mt burned	UK NAEI
NFR Source Category	Crematoria (5C1bv)		
Pollutant	Value	Unit	Reference
NOx	0.825	kg/body	2016 EMEP/EEA Guidebook
CO	0.140	kg/body	2016 EMEP/EEA Guidebook
NMVOC	0.013	kg/body	2016 EMEP/EEA Guidebook
SOx	0.113	kg/body	2016 EMEP/EEA Guidebook
TSP	0.039	kg/body	2016 EMEP/EEA Guidebook
PM10	0.027	kg/body	2016 EMEP/EEA Guidebook
PM2.5	0.015	kg/body	2016 EMEP/EEA Guidebook
As	13.610	mg/cremation	2016 EMEP/EEA Guidebook
Cd	5.030	mg/cremation	2016 EMEP/EEA Guidebook
Pb	30.030	mg/cremation	2016 EMEP/EEA Guidebook
Cr	13.560	mg/cremation	2016 EMEP/EEA Guidebook
Ni	17.330	mg/cremation	2016 EMEP/EEA Guidebook
Hg	1.490	g/cremation	2016 EMEP/EEA Guidebook
Cu	12.430	mg/cremation	2016 EMEP/EEA Guidebook
Se	19.780	mg/cremation	2016 EMEP/EEA Guidebook
Zn	160.120	mg/cremation	2016 EMEP/EEA Guidebook
PCDD/F	0.0270	ug - ITEQ/ cremation	2016 EMEP/EEA Guidebook
HCB	0.150	mg/cremation	2016 EMEP/EEA Guidebook
Benzo[a]pyrene	0.0132	mg/cremation	2016 EMEP/EEA Guidebook
Benzo[b]fluoranthene	0.0072	mg/cremation	2016 EMEP/EEA Guidebook
Benzo[k]Fluoranthene	0.0064	mg/cremation	2016 EMEP/EEA Guidebook
Indeno[1,2,3-cd]pyrene	0.0070	mg/cremation	2016 EMEP/EEA Guidebook

Table F.2 Emission Factors for NFR 5C2 and NFR 5E

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	5	Waste	2016
NFR Source Category	Open Burning of Agricultural Wastes - Farm Plastics (5C2)		
Pollutant	Value	Unit	Reference
PCDD/F	300	g-I-TEQ/Mt burned	UNEP Toolkit (2013)
PCBs	510	kg/Mt burned	UK NAEI
Benzo[a]pyrene	89.5	kg/Mt burned	UK NAEI
Benzo[b]fluoranthene	405	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	405	kg/Mt burned	UK NAEI
NFR Source Category	Accidental Vehicle Fires (5E)		
Pollutant	Value	Unit	Reference
PCDD/F	100	ug I-TEQ/vehicle	UNEP Toolkit (2013)
PCBs	25.5	mg / vehicle fire	UK NAEI
Benzo[a]pyrene	0.060	mg / vehicle fire	UK NAEI
Benzo[b]fluoranthene	0.095	mg / vehicle fire	UK NAEI
Benzo[k]Fluoranthene	0.034	mg / vehicle fire	UK NAEI
Indeno[1,2,3-cd]pyrene	0.065	mg / vehicle fire	UK NAEI
NFR Source Category	Accidental Buidling Fires (5E)		
Pollutant	Value	Unit	Reference
PCDD/F	400	ug I-TEQ/t	UNEP Toolkit (2013)
PCBs	510	kg/Mt burned	UK NAEI
Benzo[a]pyrene	1.2	kg/Mt burned	UK NAEI
Benzo[b]fluoranthene	1.9	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	0.670	kg/Mt burned	UK NAEI
Indeno[1,2,3-cd]pyrene	1.30	kg/Mt burned	UK NAEI

Emission Factors			
	Code	Name	Inventory Year
NFR Source Category	5	Waste	2016
NFR Source Category	Domestic Bonfires (5E)		
Pollutant	Value	Unit	Reference
PCDD/F	60	g-I-TEQ/Mt burned	UNEP Toolkit (2013)
PCBs	1.14	kg/Mt burned	UK NAEI
Benzo[a]pyrene	1300	kg/Mt burned	UK NAEI
Benzo[b]fluoranthene	1500	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	500	kg/Mt burned	UK NAEI
Indeno[1,2,3-cd]pyrene	90	kg/Mt burned	UK NAEI
NFR Source Category	Domestic Burning of Household Waste (5E)		
Pollutant	Value	Unit	Reference
PCDD/F	40	g-I-TEQ/Mt burned	UNEP Toolkit (2013)
PCBs	510	kg/Mt burned	UK NAEI
Benzo[a]pyrene	89.5	kg/Mt burned	UK NAEI
Benzo[b]fluoranthene	405	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	405	kg/Mt burned	UK NAEI
NFR Source Category	Wood Burining on Construction Sites (5E)		
Pollutant	Value	Unit	Reference
PCDD/F	60	g-I-TEQ/Mt burned	UNEP Toolkit (2013)
PCBs	1.99	kg/Mt burned	UK NAEI
Benzo[a]pyrene	1300	kg/Mt burned	UK NAEI
Benzo[b]fluoranthene	1500	kg/Mt burned	UK NAEI
Benzo[k]Fluoranthene	500	kg/Mt burned	UK NAEI
Indeno[1,2,3-cd]pyrene	90	kg/Mt burned	UK NAEI

Annex G

Uncertainty Analysis

A Tier 1 uncertainty analysis was conducted for NO_x, SO₂, NMVOC, NH₃, CO and PM_{2.5}. The method used was that specified in the IPCC Good Practice Guidance. This allows the calculation of an absolute uncertainty for the emissions in 2016, and the uncertainty in the trend between 1990 and 2016.

Uncertainty Associated with the Activity Data

The uncertainty values assigned to activity data in the combustion sectors (i.e. quantities of fuel) were taken from the Tier 1 uncertainty analysis of Greenhouse Gas emissions in Ireland's National Inventory Report. This was possible because the pollutants included here use the same activity data for combustion emissions. Uncertainties assigned to the activity data for sources not included in the carbon uncertainty analysis, i.e. process emissions, were determined by considering the variability in the trend of the activity data with time, and the extent to which recalculations had been made on data in previous years. In this way, it is possible to obtain a general assessment of how uncertain the data appears to be. This, combined with expert opinion, allowed a numerical value for the uncertainty in the activity data to be made.

Uncertainty Associated with Emission Factors

The EMEP/EEA Emissions Inventory Guidebook doesn't provide uncertainty information with each of the reported emission factors, but it does provide a guide as to the levels of uncertainty that are typically found by sector across the pollutants. Uncertainty ranges are used to give a general indication of how well emission factors are characterised.

This information was used as context when uncertainties were assigned to the emission factors. Expert judgement and knowledge of the inventory methodologies were also used, so that input datasets considered being higher or lower in uncertainty than the typical default levels could be reflected in the uncertainty value assigned to the relevant emission factor. Trends in the emission factor uncertainty across the sectors or fuel types were also incorporated to provide consistency in the approach across the pollutants. For example, in terms of the expected levels of uncertainty in emission factors the following relationship was assumed for the majority of cases:

Electricity Generation < Industrial Combustion < Commercial Combustion < Residential Combustion

Similar trends across the fuel were also generally used:

Gaseous Fuels < Petroleum Fuels < Coal < Peat

Uncertainty values were generally restricted to 10%, 20%, 50%, 100%, 300% and 500% (although other values were used on occasion).

This overall approach proves a good framework for populating the emission factor uncertainties in a consistent and transparent way.

The following tables show the contributions to the overall analysis for each pollutant by source sector.

Table G.1 Tier 1 Emissions Uncertainty Analysis NOx

IPCC Source Category	Gas	Emissions in 1990	Emissions in 2016	Activity Data (AD) Uncertainty	Emission Factor (EF) Uncertainty	Combined Uncertainty	Combined Uncertainty as % of Emissions in 2016	Combined Emissions Uncertainty Squared	Type A Sensitivity	Type B Sensitivity	Uncertainty in Trend in Total Emissions due to AD	Uncertainty in Trend in Total Emissions due to EF	Combined Uncertainty in Trend in Total Emissions	Combined Trend Uncertainty Squared
		tonnes	tonnes	%	%	%	%	%	%	%	%	%	%	%
1A1 Energy-Liquid	NOx	2766.3	74.2	1.0	10.0	10.05	0.01	0.00	-0.01	0.00	0.00	-0.10	0.10	0.01
1A1 Energy-Solid	NOx	34882.2	5380.3	1.0	10.0	10.05	0.48	0.23	-0.10	0.03	0.04	-1.05	1.05	1.10
1A1 Energy-Gas	NOx	9353.5	2815.0	1.0	10.0	10.05	0.25	0.06	-0.02	0.02	0.02	-0.20	0.20	0.04
1A1 Energy-Biomass & renewable waste	NOx	0.0	111.6	1.0	50.0	50.01	0.05	0.00	0.00	0.00	0.00	0.03	0.03	0.00
1A1 Energy-non-renewable waste	NOx	0.0	107.3	1.0	50.0	50.01	0.05	0.00	0.00	0.00	0.00	0.03	0.03	0.00
1A1 Energy-Landfill Gas	NOx	0.0	144.8	1.0	300.0	300.00	0.39	0.15	0.00	0.00	0.00	0.26	0.26	0.07
1A2 Industry-Liquid exc Pet Coke	NOx	3611.4	1347.0	10.0	20.0	22.36	0.27	0.07	-0.01	0.01	0.11	-0.12	0.17	0.03
1A2 Industry-Coal + Biomass	NOx	2723.7	3109.4	2.0	50.0	50.04	1.39	1.92	0.01	0.02	0.05	0.38	0.39	0.15
1A2 Industry-Pet Coke	NOx	1409.9	4348.6	5.0	20.0	20.62	0.80	0.64	0.02	0.03	0.18	0.40	0.44	0.20
1A2 Industry-Gas	NOx	1182.9	2506.7	7.0	20.0	21.19	0.47	0.22	0.01	0.01	0.15	0.20	0.25	0.06
1A3a Transport-Oil-Aviation	NOx	1037.5	1136.4	1.0	7.5	7.57	0.08	0.01	0.00	0.01	0.01	0.02	0.02	0.00
1A3b Transport-Oil-Road	NOx	54699.2	39329.6	1.0	10.0	10.05	3.52	12.39	0.02	0.23	0.33	0.18	0.37	0.14
1A3c Transport-Oil-Rail	NOx	2198.5	1847.5	1.0	70.0	70.01	1.15	1.33	0.00	0.01	0.02	0.16	0.16	0.03
1A3d Transport-Oil-Navigation	NOx	2135.5	6520.5	1.0	40.0	40.01	2.32	5.40	0.03	0.04	0.05	1.21	1.21	1.46
1A3e Transport-Gas	NOx	54.2	120.2	1.0	50.0	50.01	0.05	0.00	0.00	0.00	0.00	0.02	0.02	0.00
1A4 Comm-Liquid	NOx	2523.1	1025.4	10.0	20.0	22.36	0.20	0.04	0.00	0.01	0.09	-0.08	0.12	0.01
1A4 Comm-Coal + Biomass	NOx	4.7	98.8	5.0	50.0	50.25	0.04	0.00	0.00	0.00	0.00	0.03	0.03	0.00
1A4 Comm-Peat	NOx	231.5	0.0	10.0	100.0	100.50	0.00	0.00	0.00	0.00	0.00	-0.09	0.09	0.01
1A4 Comm-Gas + Biogas	NOx	301.0	1472.1	2.5	20.0	20.16	0.26	0.07	0.01	0.01	0.03	0.15	0.15	0.02
1A4 Res-Liquid (excl Pet Coke)	NOx	764.2	2119.5	10.0	50.0	50.99	0.96	0.93	0.01	0.01	0.18	0.48	0.51	0.26
1A4 Res-Coal + Biomass	NOx	2975.6	916.0	5.0	100.0	100.12	0.82	0.67	-0.01	0.01	0.04	-0.63	0.63	0.39
1A4 Res-Petcoke	NOx	41.2	11.7	5.0	50.0	50.25	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Res-Peat	NOx	3338.8	906.8	10.0	100.0	100.50	0.81	0.66	-0.01	0.01	0.08	-0.77	0.78	0.61
1A4 Res-Gas	NOx	206.2	990.0	2.5	20.0	20.16	0.18	0.03	0.01	0.01	0.02	0.10	0.10	0.01
1A4 Agric/Forestry/Fishing Liquid	NOx	8760.0	4054.2	10.0	100.0	100.50	3.63	13.17	-0.01	0.02	0.34	-1.04	1.10	1.20
2B2 Nitric Acid Production	NOx	960.0	0.0	1.0	100.0	100.00	0.00	0.00	0.00	0.00	0.00	-0.38	0.38	0.14
2G Other Product Use	NOx	12.6	6.3	100.0	30.0	104.40	0.01	0.00	0.00	0.00	0.01	0.00	0.01	0.00
3B1a Dairy cattle	NOx	49.4	52.2	1.0	50.0	50.01	0.02	0.00	0.00	0.00	0.00	0.01	0.01	0.00
3B1b Non-dairy cattle	NOx	485.1	557.8	1.0	50.0	50.01	0.25	0.06	0.00	0.00	0.00	0.07	0.07	0.00
3B2 Sheep	NOx	74.3	48.6	1.0	100.0	100.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B3 Swine	NOx	1.9	2.3	1.0	100.0	100.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4d Goats	NOx	2.5	1.4	1.0	100.0	100.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Horses	NOx	22.5	33.6	1.0	100.0	100.00	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00
3B4f Mules and asses	NOx	2.1	2.3	1.0	100.0	100.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4g Poultry	NOx	107.7	132.7	1.0	100.0	100.00	0.12	0.01	0.00	0.00	0.00	0.04	0.04	0.00
3B4h Other animals	NOx	17.3	6.4	1.0	100.0	100.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3D Synthetic Fertilizer	NOx	14594.1	13047.1	11.2	200.0	200.31	23.28	541.84	0.02	0.08	1.22	3.97	4.15	17.23
3D Organic fertiliser	NOx	17538.4	17882.1	11.2	200.0	200.31	31.90	1017.83	0.04	0.11	1.67	7.37	7.56	57.11
5C1 Incineration	NOx	34.0	10.7	25.0	50.0	55.90	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total NOx		169103.0	112276.9				39.97	1,597.74					2.67	7.12

Table G.2 Tier 1 Emissions Uncertainty Analysis SO₂

IPCC Source Category	Gas	Emissions in 1990	Emissions in 2016	Activity Data (AD) Uncertainty	Emission Factor (EF) Uncertainty	Combined Uncertainty	Combined Uncertainty as % of Emissions in 2016	Combined Emissions Uncertainty Squared	Type A Sensitivity	Type B Sensitivity	Uncertainty in Trend in Total Emissions due to AD	Uncertainty in Trend in Total Emissions due to EF	Combined Uncertainty in Trend in Total Emissions	Combined Trend Uncertainty Squared
		tonnes	tonnes	%	%	%	%	%	%	%	%	%	%	%
1A1 Energy-Liquid	SO ₂	19611.0	287.0	1.0	5.0	5.10	0.11	0.01	-0.01	0.00	0.00	-0.03	0.03	0.00
1A1 Energy-Solid	SO ₂	83692.1	3622.4	1.0	10.0	10.05	2.64	6.99	-0.01	0.02	0.03	-0.15	0.15	0.02
1A1 Energy-Gas	SO ₂	349.8	19.8	1.0	5.0	5.10	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Biomass & renewable waste	SO ₂	0.0	27.2	1.0	50.0	50.01	0.10	0.01	0.00	0.00	0.00	0.01	0.01	0.00
1A1 Energy-non-renewable waste	SO ₂	0.0	26.1	1.0	50.0	50.01	0.09	0.01	0.00	0.00	0.00	0.01	0.01	0.00
1A1 Energy-Landfill Gas	SO ₂	0.0	0.0	1.0	300.0	300.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A2 Industry-Liquid exc Pet Coke	SO ₂	27831.5	604.0	10.0	5.0	11.18	0.49	0.24	-0.01	0.00	0.05	-0.04	0.06	0.00
1A2 Industry-Coal + Biomass	SO ₂	4599.8	1321.4	2.0	20.0	20.10	1.93	3.72	0.01	0.01	0.02	0.11	0.11	0.01
1A2 Industry-Pet Coke	SO ₂	128.2	663.7	5.0	5.0	7.07	0.34	0.12	0.00	0.00	0.03	0.02	0.03	0.00
1A2 Industry-Gas	SO ₂	0.9	1.9	7.0	5.0	8.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3a Transport-Oil-Aviation	SO ₂	84.6	93.7	1.0	7.5	7.57	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3b Transport-Oil-Road	SO ₂	5373.1	36.8	1.0	5.0	5.10	0.01	0.00	0.00	0.00	0.00	-0.01	0.01	0.00
1A3c Transport-Oil-Rail	SO ₂	251.5	13.8	1.0	5.0	5.10	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3d Transport-Oil-Navigation	SO ₂	1160.7	116.3	1.0	30.0	30.02	0.25	0.06	0.00	0.00	0.00	0.00	0.00	0.00
1A3e Transport-Gas	SO ₂	0.1	0.1	1.0	5.0	5.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Comm-Liquid	SO ₂	10384.2	222.5	10.0	10.0	14.14	0.23	0.05	0.00	0.00	0.02	-0.03	0.04	0.00
1A4 Comm-Coal + Biomass	SO ₂	15.5	11.9	5.0	20.0	20.62	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Comm-Peat	SO ₂	1124.1	0.0	10.0	50.0	50.99	0.00	0.00	0.00	0.00	0.00	-0.02	0.02	0.00
1A4 Comm-Gas + Biogas	SO ₂	0.2	0.8	2.5	5.0	5.59	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Res-Liquid (excl Pet Coke)	SO ₂	1337.4	790.4	10.0	10.0	14.14	0.81	0.66	0.00	0.00	0.06	0.04	0.07	0.01
1A4 Res-Coal + Biomass	SO ₂	15199.7	3234.3	5.0	20.0	20.62	4.84	23.44	0.01	0.02	0.13	0.23	0.26	0.07
1A4 Res-Petcoke	SO ₂	1245.6	223.0	5.0	10.0	11.18	0.18	0.03	0.00	0.00	0.01	0.01	0.01	0.00
1A4 Res-Peat	SO ₂	8976.2	2415.0	10.0	50.0	50.99	8.94	79.97	0.01	0.01	0.19	0.48	0.51	0.26
1A4 Res-Gas	SO ₂	0.3	1.0	2.5	5.0	5.59	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Agric/Forestry/Fishing Liquid	SO ₂	1410.8	36.4	10.0	10.0	14.14	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2G Other product use	SO ₂	0.1	0.1	100.0	30.0	104.40	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Incineration	SO ₂	3.6	0.9	25.0	25.0	35.36	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total SO₂		182780.8	13770.6				10.74	115.32					0.61	0.38

Table G.3 Tier 1 Emissions Uncertainty Analysis NMVOC

IPCC Source Category	Gas	Emissions in 1990 tonnes	Emissions in 2016 tonnes	Activity Data (AD) Uncertainty %	Emission Factor (EF) Uncertainty %	Combined Uncertainty %	Combined Uncertainty as % of Emissions in 2016 %	Combined Emissions Uncertainty Squared %	Type A Sensitivity %	Type B Sensitivity %	Uncertainty in Trend in Total Emissions due to AD % %	Uncertainty in Trend in Total Emissions due to EF % %	Combined Uncertainty in Trend in Total Emissions % %	Combined Trend Uncertainty Squared %
1A1 Energy-Liquid	NMVOC	34.6	6.1	1.0	10.0	10.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Solid	NMVOC	72.2	63.7	1.0	10.0	10.05	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Gas	NMVOC	93.8	242.2	1.0	10.0	10.05	0.02	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A1 Energy-Biomass & renewable waste	NMVOC	0.0	4.2	1.0	50.0	50.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-non-renewable waste	NMVOC	0.0	1.3	1.0	50.0	50.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Landfill Gas	NMVOC	0.0	4.2	1.0	300.0	300.00	0.01	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A2 Industry-Liquid exc Pet Coke	NMVOC	305.0	208.7	10.0	20.0	22.36	0.04	0.00	0.00	0.00	0.02	0.00	0.02	0.00
1A2 Industry-Coal + Biomass	NMVOC	978.2	1677.8	2.0	20.0	20.10	0.31	0.10	0.01	0.01	0.03	0.13	0.14	0.02
1A2 Industry-Pet Coke	NMVOC	19.7	55.9	5.0	20.0	20.62	0.01	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A2 Industry-Gas	NMVOC	367.7	979.3	7.0	20.0	21.19	0.19	0.04	0.00	0.01	0.07	0.10	0.12	0.01
1A3a Transport-Oil-Aviation	NMVOC	184.4	152.8	1.0	7.5	7.57	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3b Transport-Oil-Road	NMVOC	33151.7	4865.8	1.0	15.0	15.03	0.67	0.46	-0.14	0.03	0.05	-2.12	2.12	4.48
1A3c Transport-Oil-Rail	NMVOC	195.1	163.9	1.0	64.0	64.01	0.14	0.02	0.00	0.00	0.00	0.04	0.04	0.00
1A3d Transport-Oil-Navigation	NMVOC	73.6	232.6	1.0	50.0	50.01	0.08	0.01	0.00	0.00	0.00	0.04	0.04	0.00
1A3e Transport-Gas	NMVOC	1.8	4.0	1.0	10.0	10.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Comm-Liquid	NMVOC	258.8	108.1	10.0	20.0	22.36	0.02	0.00	0.00	0.00	0.01	-0.01	0.02	0.00
1A4 Comm-Coal + Biomass	NMVOC	2.4	13.0	5.0	20.0	20.62	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Comm-Peat	NMVOC	118.8	0.0	10.0	50.0	50.99	0.00	0.00	0.00	0.00	0.00	-0.03	0.03	0.00
1A4 Comm-Gas + Biogas	NMVOC	93.6	457.5	2.5	20.0	20.16	0.09	0.01	0.00	0.00	0.01	0.05	0.06	0.00
1A4 Res-Liquid (excl Pet Coke)	NMVOC	13.9	30.7	10.0	20.0	22.36	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Res-Coal + Biomass	NMVOC	13803.6	3988.3	5.0	50.0	50.25	1.85	3.42	-0.05	0.03	0.20	-2.26	2.26	5.13
1A4 Res-Petcoke	NMVOC	0.6	0.2	5.0	20.0	20.62	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Res-Peat	NMVOC	14690.9	3989.7	10.0	50.0	50.99	1.88	3.52	-0.05	0.03	0.39	-2.49	2.52	6.36
1A4 Res-Gas	NMVOC	8.8	42.4	2.5	20.0	20.16	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00
1A4 Agric/Forestry/Fishing Liquid	NMVOC	1375.4	290.7	10.0	50.0	50.99	0.14	0.02	-0.01	0.00	0.03	-0.26	0.26	0.07
1B1a Fugitive emission from solid fuels: Co	NMVOC	20.0	0.0	5.0	300.0	300.04	0.00	0.00	0.00	0.00	0.00	-0.03	0.03	0.00
1B2 Fuel Extraction and Distribution	NMVOC	3376.1	4495.0	5.0	300.0	300.04	12.44	154.78	0.01	0.03	0.22	4.06	4.07	16.54
2D3a Domestic solvent use including fungic	NMVOC	7926.6	10766.6	100.0	30.0	104.40	10.37	107.51	0.03	0.08	10.64	1.00	10.68	114.14
2D3b Road paving with asphalt	NMVOC	35.2	30.4	33.0	30.0	44.60	0.01	0.00	0.00	0.00	0.01	0.00	0.01	0.00
2D3d Coating applications; paint applicatio	NMVOC	6742.6	2916.0	33.0	30.0	44.60	1.20	1.44	-0.02	0.02	0.95	-0.46	1.06	1.11
2D3e Degreasing	NMVOC	1743.5	837.1	50.0	30.0	58.31	0.45	0.20	0.00	0.01	0.41	-0.10	0.43	0.18
2D3f Dry cleaning	NMVOC	281.8	62.2	50.0	30.0	58.31	0.03	0.00	0.00	0.00	0.03	-0.03	0.04	0.00
2D3g Chemical products	NMVOC	3023.1	1168.2	100.0	30.0	104.40	1.13	1.27	-0.01	0.01	1.15	-0.23	1.18	1.39
2D3h Printing	NMVOC	2912.0	1462.7	100.0	30.0	104.40	1.41	1.98	-0.01	0.01	1.45	-0.16	1.45	2.11
2D3i Other solvent use	NMVOC	701.2	1369.8	100.0	30.0	104.40	1.32	1.74	0.01	0.01	1.35	0.18	1.36	1.86
2G4 Other product use	NMVOC	33.9	17.0	100.0	30.0	104.40	0.02	0.00	0.00	0.00	0.02	0.00	0.02	0.00
2H2 Food and beverage industry	NMVOC	9617.2	21260.6	33.0	30.0	44.60	8.75	76.50	0.10	0.15	6.93	2.93	7.52	56.62
3B1a Dairy cattle	NMVOC	9232.4	9295.1	1.0	300.0	300.00	25.72	661.66	0.02	0.06	0.09	4.82	4.82	23.28
3B1b Other cattle	NMVOC	22563.5	26739.5	1.0	300.0	300.00	74.00	5475.64	0.07	0.19	0.26	20.20	20.20	407.93
3B2 Sheep	NMVOC	592.5	355.9	1.0	300.0	300.00	0.98	0.97	0.00	0.00	0.00	-0.19	0.19	0.04
3B3 Swine	NMVOC	1016.7	2484.5	1.0	300.0	300.00	6.88	47.27	0.01	0.02	0.02	3.59	3.59	12.91
3B4d Goats	NMVOC	2.8	1.6	1.0	300.0	300.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Horses	NMVOC	113.6	170.0	1.0	300.0	300.00	0.47	0.22	0.00	0.00	0.00	0.18	0.18	0.03
3B4f Mules and asses	NMVOC	8.3	9.2	1.0	300.0	300.00	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00
3B4g Poultry	NMVOC	2089.9	2633.2	1.0	300.0	300.00	7.29	53.10	0.01	0.02	0.03	2.20	2.20	4.85
3B4h Other animals	NMVOC	410.1	383.7	1.0	300.0	300.00	1.06	1.13	0.00	0.00	0.00	0.15	0.15	0.02
3De Cultivated crops	NMVOC	3819.9	3824.6	20.0	300.0	300.67	10.61	112.52	0.01	0.03	0.76	1.95	2.09	4.38
5A Waste disposal to Land	NMVOC	834.1	485.9	200.0	300.0	360.56	1.62	2.61	0.00	0.00	0.96	-0.31	1.01	1.01
5C Incineration	NMVOC	203.2	55.5	25.0	25.0	35.36	0.02	0.00	0.00	0.00	0.01	-0.02	0.02	0.00
Total NMVOC		143144.9	108407.6				81.90	6708.13					25.78	664.49

Table G.4 Tier 1 Emissions Uncertainty Analysis NH₃

IPCC Source Category	Gas	Emissions in 1990	Emissions in 2016	Activity Data (AD) Uncertainty	Emission Factor (EF) Uncertainty	Combined Uncertainty	Combined Uncertainty as % of Emissions in 2016	Combined Emissions Uncertainty Squared	Type A Sensitivity	Type B Sensitivity	Uncertainty in Trend in Total Emissions due to AD %	Uncertainty in Trend in Total Emissions due to EF %	Combined Uncertainty in Trend in Total Emissions %	Combined Trend Uncertainty Squared
		tonnes	tonnes	%	%	%	%	%	%	%	%	%	%	%
1A2 Industry-Biomass	NH ₃	94.1	266.0	1.0	200.0	200.00	0.46	0.21	0.00	0.00	0.00	0.30	0.30	0.09
1A3b Transport-Oil-Road	NH ₃	41.5	783.9	1.0	20.0	20.02	0.13	0.02	0.01	0.01	0.01	0.13	0.14	0.02
1A3c Transport-Oil-Railways	NH ₃	0.3	0.2	1.0	57.0	57.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Coal, Peat, Biomass	NH ₃	147.9	101.6	10.0	200.0	200.25	0.17	0.03	0.00	0.00	0.01	-0.10	0.10	0.01
1A4 Agric/Forestry/Fishing Liquid	NH ₃	1.3	1.1	10.0	100.0	100.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2G Other product use	NH ₃	29.0	14.6	100.0	6.0	100.18	0.01	0.00	0.00	0.00	0.02	0.00	0.02	0.00
3B1a Dairy cattle	NH ₃	12700.5	13434.1	1.0	50.0	50.01	5.76	33.13	0.00	0.12	0.17	-0.03	0.18	0.03
3B1b Non-dairy cattle	NH ₃	27405.5	31281.2	1.0	50.0	50.01	13.40	179.62	0.02	0.28	0.40	0.98	1.06	1.11
3B2 Sheep	NH ₃	1722.4	1127.1	1.0	100.0	100.00	0.97	0.93	-0.01	0.01	0.01	-0.64	0.64	0.41
3B3 Swine	NH ₃	3846.1	4719.4	1.0	100.0	100.00	4.04	16.35	0.01	0.04	0.06	0.57	0.58	0.33
3B4d Goats	NH ₃	34.0	19.4	1.0	100.0	100.00	0.02	0.00	0.00	0.00	0.00	-0.02	0.02	0.00
3B4e Horses	NH ₃	526.2	787.5	1.0	100.0	100.00	0.67	0.46	0.00	0.01	0.01	0.21	0.21	0.04
3B4f Mules and asses	NH ₃	48.3	53.6	1.0	100.0	100.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4g Poultry	NH ₃	2718.8	3256.1	1.0	100.0	100.00	2.79	7.78	0.00	0.03	0.04	0.33	0.34	0.11
3B4h Other animals	NH ₃	603.4	415.8	1.0	100.0	100.00	0.36	0.13	0.00	0.00	0.01	-0.21	0.21	0.04
3D Synthetic Fertilizer	NH ₃	13691.4	11191.6	11.2	200.0	200.31	19.21	368.89	-0.03	0.10	1.61	-6.12	6.33	40.05
3D Organic fertiliser	NH ₃	46186.9	49242.3	11.2	200.0	200.31	84.51	7141.35	0.00	0.45	7.10	0.26	7.11	50.53
5B1 Biological treatment of waste - Comp	NH ₃	0.0	27.8	10.0	300.0	300.17	0.07	0.01	0.00	0.00	0.00	0.08	0.08	0.01
Total NH₃		109797.8	116723.4				88.03	7,748.69					9.63	92.70

Table G.5 Tier 1 Emissions Uncertainty Analysis CO

IPCC Source Category	Gas	Emissions in 1990	Emissions in 2016	Activity Data (AD) Uncertainty	Emission Factor (EF) Uncertainty	Combined Uncertainty	Combined Uncertainty as % of Emissions in 2016	Combined Emissions Uncertainty Squared	Type A Sensitivity	Type B Sensitivity	Uncertainty in Trend in Total Emissions due to AD %	Uncertainty in Trend in Total Emissions due to EF %	Combined Uncertainty in Trend in Total Emissions %	Combined Trend Uncertainty Squared
		tonnes	tonnes	%	%	%	%	%	%	%	%	%	%	%
1A1 Energy-Liquid	CO	227.2	40.5	1.0	10.0	10.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Solid	CO	16759.9	14776.1	1.0	10.0	10.05	1.45	2.10	0.03	0.04	0.06	0.28	0.29	0.08
1A1 Energy-Gas	CO	1354.9	3421.1	1.0	10.0	10.05	0.34	0.11	0.01	0.01	0.01	0.09	0.09	0.01
1A1 Energy-Biomass & renewable waste	CO	0.0	51.0	1.0	50.0	50.01	0.02	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A1 Energy-non-renewable waste	CO	0.0	8.9	1.0	50.0	50.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Landfill Gas	CO	0.0	63.4	1.0	50.0	50.01	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A2 Industry-Liquid exc Pet Coke	CO	653.5	524.0	10.0	20.0	22.36	0.11	0.01	0.00	0.00	0.02	0.02	0.03	0.00
1A2 Industry-Coal + Biomass	CO	9459.3	7589.1	2.0	50.0	50.04	3.70	13.71	0.01	0.02	0.06	0.69	0.69	0.48
1A2 Industry-Pet Coke	CO	482.3	235.1	5.0	20.0	20.62	0.05	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A2 Industry-Gas	CO	463.6	1234.8	7.0	20.0	21.19	0.26	0.07	0.00	0.00	0.04	0.06	0.07	0.01
1A3a Transport-Oil-Aviation	CO	1300.8	1508.5	1.0	7.5	7.57	0.11	0.01	0.00	0.00	0.01	0.02	0.03	0.00
1A3b Transport-Oil-Road	CO	248617.0	48843.8	1.0	15.0	15.03	7.16	51.26	-0.07	0.14	0.20	-1.05	1.07	1.15
1A3c Transport-Oil-Rail	CO	448.9	377.3	1.0	65.0	65.01	0.24	0.06	0.00	0.00	0.00	0.05	0.05	0.00
1A3d Transport-Oil-Navigation	CO	199.8	614.7	1.0	65.0	65.01	0.39	0.15	0.00	0.00	0.00	0.10	0.10	0.01
1A3e Transport-Gas	CO	5.4	12.0	1.0	50.0	50.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Comm-Liquid	CO	1009.0	409.9	10.0	20.0	22.36	0.09	0.01	0.00	0.00	0.02	0.01	0.02	0.00
1A4 Comm-Coal + Biomass	CO	25.2	325.7	5.0	50.0	50.25	0.16	0.03	0.00	0.00	0.01	0.05	0.05	0.00
1A4 Comm-Peat	CO	1245.9	0.0	10.0	100.0	100.50	0.00	0.00	0.00	0.00	0.00	-0.11	0.11	0.01
1A4 Comm-Gas + Biogas	CO	118.0	576.9	2.5	20.0	20.16	0.11	0.01	0.00	0.00	0.01	0.03	0.03	0.00
1A4 Res-Liquid (excl Pet Coke)	CO	782.0	2328.0	10.0	50.0	50.99	1.16	1.34	0.01	0.01	0.09	0.30	0.32	0.10
1A4 Res-Coal + Biomass	CO	31875.5	9910.1	5.0	100.0	100.12	9.67	93.60	0.00	0.03	0.20	0.14	0.25	0.06
1A4 Res-Petcoke	CO	46.1	13.1	5.0	50.0	50.25	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Res-Peat	CO	28258.8	7674.4	10.0	100.0	100.50	7.52	56.55	0.00	0.02	0.31	-0.19	0.37	0.13
1A4 Res-Gas	CO	108.0	518.6	2.5	20.0	20.16	0.10	0.01	0.00	0.00	0.01	0.03	0.03	0.00
1A4 Agric/Forestry/Fishing Liquid	CO	3670.0	1244.3	10.0	100.0	100.50	1.22	1.49	0.00	0.00	0.05	0.05	0.07	0.00
2G Other Product use	CO	386.0	194.1	100.0	30.0	104.40	0.20	0.04	0.00	0.00	0.08	0.01	0.08	0.01
5B1 Biological treatment of waste - Comp	CO	0.0	64.9	10.0	300.0	300.17	0.19	0.04	0.00	0.00	0.00	0.06	0.06	0.00
5C Incineration	CO	2.9	1.2	25.0	50.0	55.90	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total CO		347499.9	102561.8				14.85	220.59					1.44	2.06

Table G.6 Tier 1 Emissions Uncertainty Analysis PM2.5

IPCC Source Category	Gas	Emissions in 1990	Emissions in 2016	Activity Data (AD) Uncertainty	Emission Factor (EF) Uncertainty	Combined Uncertainty	Combined Uncertainty as % of Emissions in 2016	Combined Emissions Uncertainty Squared	Type A Sensitivity	Type B Sensitivity	Uncertainty in Trend in Total Emissions due to AD	Uncertainty in Trend in Total Emissions due to EF	Combined Uncertainty in Trend in Total Emissions	Combined Trend Uncertainty Squared
		tonnes	tonnes	%	%	%	%	%	%	%	%	%	%	%
1A1 Energy-Liquid	PM _{2.5}	280.7	51.1	1.0	300.0	300.00	0.99	0.98	0.00	0.00	0.00	-0.62	0.62	0.38
1A1 Energy-Solid	PM _{2.5}	384.2	347.1	1.0	100.0	100.00	2.24	5.04	0.01	0.01	0.01	0.51	0.51	0.26
1A1 Energy-Gas	PM _{2.5}	32.1	83.0	1.0	100.0	100.00	0.54	0.29	0.00	0.00	0.00	0.20	0.20	0.04
1A1 Energy-Biomass & renewable waste	PM _{2.5}	0.0	9.9	1.0	50.0	50.01	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A1 Energy-non-renewable waste	PM _{2.5}	0.0	0.3	1.0	50.0	50.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A1 Energy-Landfill Gas	PM _{2.5}	0.0	1.4	1.0	50.0	50.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A2 Industry-Liquid exc Pet Coke	PM _{2.5}	407.1	167.9	10.0	50.0	50.99	0.55	0.31	0.00	0.00	0.07	-0.02	0.07	0.00
1A2 Industry-Coal + Biomass	PM _{2.5}	1135.2	1189.6	2.0	50.0	50.04	3.85	14.81	0.02	0.03	0.10	0.98	0.99	0.97
1A2 Industry-Pet Coke	PM _{2.5}	32.5	92.2	5.0	50.0	50.25	0.30	0.09	0.00	0.00	0.02	0.11	0.11	0.01
1A2 Industry-Gas	PM _{2.5}	12.4	33.1	7.0	50.0	50.49	0.11	0.01	0.00	0.00	0.01	0.04	0.04	0.00
1A3a Transport-Oil-Aviation	PM _{2.5}	17.4	20.3	1.0	7.5	7.57	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A3b Transport-Oil-Road	PM _{2.5}	2097.3	1019.9	1.0	10.0	10.05	0.66	0.44	0.00	0.03	0.04	0.03	0.05	0.00
1A3b Transport-Tyre and brake wear	PM _{2.5}	216.4	578.9	1.0	50.0	50.01	1.87	3.50	0.01	0.02	0.02	0.69	0.69	0.47
1A3b Transport-Road Abrasion	PM _{2.5}	117.8	316.8	1.0	50.0	50.01	1.02	1.05	0.01	0.01	0.01	0.38	0.38	0.14
1A3c Transport-Oil-Rail	PM _{2.5}	57.5	48.3	1.0	169.0	169.00	0.53	0.28	0.00	0.00	0.00	0.11	0.11	0.01
1A3d Transport-Oil-Navigation	PM _{2.5}	121.8	116.3	1.0	50.0	50.01	0.38	0.14	0.00	0.00	0.00	0.09	0.09	0.01
1A3e Transport-Gas	PM _{2.5}	0.2	0.5	1.0	50.0	50.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Comm-Liquid	PM _{2.5}	411.5	165.1	10.0	50.0	50.99	0.54	0.30	0.00	0.00	0.07	-0.02	0.07	0.00
1A4 Comm-Coal + Biomass	PM _{2.5}	2.9	35.8	5.0	50.0	50.25	0.12	0.01	0.00	0.00	0.01	0.05	0.05	0.00
1A4 Comm-Peat	PM _{2.5}	144.5	0.0	10.0	50.0	50.99	0.00	0.00	0.00	0.00	0.00	-0.09	0.09	0.01
1A4 Comm-Gas + Biogas	PM _{2.5}	3.2	15.5	2.5	50.0	50.06	0.05	0.00	0.00	0.00	0.00	0.02	0.02	0.00
1A4 Res-Liquid (excl Pet Coke)	PM _{2.5}	19.5	60.6	10.0	100.0	100.50	0.39	0.16	0.00	0.00	0.02	0.15	0.15	0.02
1A4 Res-Coal + Biomass	PM _{2.5}	11812.2	3468.9	5.0	200.0	200.06	44.87	2013.07	-0.05	0.10	0.70	-9.76	9.78	95.74
1A4 Res-Petcoke	PM _{2.5}	1.2	0.3	5.0	100.0	100.12	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1A4 Res-Peat	PM _{2.5}	12080.6	3280.8	10.0	300.0	300.17	63.67	4053.53	-0.06	0.09	1.32	-17.24	17.29	298.90
1A4 Res-Gas	PM _{2.5}	1.0	4.7	2.5	100.0	100.03	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00
1A4 Agric/Forestry/Fishing Liquid	PM _{2.5}	926.7	143.7	10.0	100.0	100.50	0.93	0.87	-0.01	0.00	0.06	-0.75	0.75	0.56
1B1 Fugitive emission from solid fuels	PM _{2.5}	2.0	1.8	5.0	500.0	500.02	0.06	0.00	0.00	0.00	0.00	0.01	0.01	0.00
2A3 Glass production	PM _{2.5}	17.7	0.0	5.0	300.0	300.04	0.00	0.00	0.00	0.00	0.00	-0.07	0.07	0.00
2A5a Quarrying and mining excl coal	PM _{2.5}	157.5	182.2	10.0	150.0	150.33	1.77	3.13	0.00	0.01	0.07	0.48	0.49	0.24
2A5b Construction and demolition	PM _{2.5}	41.7	46.3	10.0	300.0	300.17	0.90	0.81	0.00	0.00	0.02	0.24	0.24	0.06
2A5c Storage and handling - mineral produ	PM _{2.5}	0.0	0.0	10.0	300.0	300.17	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2C2 Ferroalloys production	PM _{2.5}	42.3	0.0	10.0	500.0	500.10	0.00	0.00	0.00	0.00	0.00	-0.26	0.26	0.07
2D3b Road paving with asphalt	PM _{2.5}	2200.0	1900.0	10.0	500.0	500.10	61.43	3773.71	0.03	0.05	0.76	13.25	13.27	176.10
2D3i Other solvent use	PM _{2.5}	12.0	20.5	100.0	30.0	104.40	0.14	0.02	0.00	0.00	0.08	0.01	0.08	0.01
2G Other product use	PM _{2.5}	191.3	96.5	100.0	30.0	104.40	0.65	0.42	0.00	0.00	0.39	0.01	0.39	0.15
3B1a Dairy cattle	PM _{2.5}	539.8	598.2	1.0	50.0	50.01	1.93	3.74	0.01	0.02	0.02	0.51	0.51	0.26
3B1b Other cattle	PM _{2.5}	765.5	739.3	1.0	50.0	50.01	2.39	5.71	0.01	0.02	0.03	0.57	0.57	0.33
3B2 Sheep	PM _{2.5}	160.4	95.4	1.0	100.0	100.00	0.62	0.38	0.00	0.00	0.00	0.07	0.07	0.01
3B3 Swine	PM _{2.5}	6.5	8.1	1.0	100.0	100.00	0.05	0.00	0.00	0.00	0.00	0.01	0.01	0.00
3B4d Goats	PM _{2.5}	0.3	0.2	1.0	100.0	100.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4e Horses	PM _{2.5}	8.6	12.9	1.0	100.0	100.00	0.08	0.01	0.00	0.00	0.00	0.03	0.03	0.00
3B4f Mules and asses	PM _{2.5}	0.8	0.9	1.0	100.0	100.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3B4g Poultry	PM _{2.5}	59.2	65.6	1.0	100.0	100.00	0.42	0.18	0.00	0.00	0.00	0.11	0.11	0.01
3B4h Other animals	PM _{2.5}	0.8	0.8	1.0	100.0	100.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3Da1 Inorganic fertilizers	PM _{2.5}	266.5	266.8	10.0	200.0	200.25	3.45	11.93	0.00	0.01	0.11	0.85	0.86	0.74
3Dc Handling of Farm Products - Farm lev	PM _{2.5}	7.2	5.6	10.0	500.0	500.10	0.18	0.03	0.00	0.00	0.00	0.04	0.04	0.00
3Dd Handling of Farm Products - Off farm	PM _{2.5}	7.9	9.2	10.0	500.0	500.10	0.30	0.09	0.00	0.00	0.00	0.08	0.08	0.01
5A Solid waste disposal on land	PM _{2.5}	0.1	0.0	34.6	500.0	501.20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5C Incineration	PM _{2.5}	27.3	0.1	25.0	100.0	103.08	0.00	0.00	0.00	0.00	0.00	-0.03	0.03	0.00
5E Other waste	PM _{2.5}	359.4	165.1	25.0	100.0	103.08	1.10	1.21	0.00	0.00	0.00	0.00	0.00	0.00
Total PM_{2.5}		35190.88	15467.7				99.48	9,896.26					23.99	575.52

AN GHNÍOMHAIREACHT UM CHAOMHNÚ COMHSHAOIL

Tá an Gníomhaireacht um Chaomhnú Comhshaoil (GCC) freagrach as an gcomhshaoil a chaomhnú agus a fheabhsú mar shócmhainn luachmhar do mhuintir na hÉireann. Táimid tiomanta do dhaoine agus don chomhshaoil a chosaint ó éifeachtaí díobhálacha na radaíochta agus an truaillithe.

Is féidir obair na Gníomhaireachta a roinnt ina trí phríomhréimse:

Rialú: Déanaimid córais éifeachtacha rialaithe agus comhlíonta comhshaoil a chur i bhfeidhm chun torthaí maíthe comhshaoil a sholáthar agus chun díriú orthu siúd nach gcloíonn leis na córais sin.

Eolas: Soláthraimid sonraí, faisnéis agus measúnú comhshaoil atá ar ardchaighdeán, spriocdhírthe agus tráthúil chun bonn eolais a chur faoin gcinnteoireacht ar gach leibhéal.

Tacaíocht: Bímid ag saothrú i gcomhar le grúpaí eile chun tacú le comhshaoil atá glan, táirgiúil agus cosanta go maith, agus le hiompar a chuirfidh le comhshaoil inbhuanaithe.

Ár bhFreagrachtaí

Ceadúnú

Déanaimid na gníomhaíochtaí seo a leanas a rialú ionas nach ndéanann siad dochar do shláinte an phobail ná don chomhshaoil:

- saoráidí dramhaíola (*m.sh. láithreáin líonta talún, loisceoirí, stáisiúin aistrithe dramhaíola*);
- gníomhaíochtaí tionsclaíocha ar scála mór (*m.sh. déantúsaíocht cógaisíochta, déantúsaíocht stroighne, stáisiúin chumhachta*);
- an diantalmhaíocht (*m.sh. muca, éanlaith*);
- úsáid shrianta agus scaoileadh rialaithe Orgánach Géinmhodhnaithe (OGM);
- foinsí radaíochta ianúcháin (*m.sh. trealamh x-gha agus radaiteiripe, foinsí tionsclaíocha*);
- áiseanna móra stórála peitril;
- scardadh dramhuisce;
- gníomhaíochtaí dumpála ar farraige.

Forfheidhmiú Náisiúnta i leith Cúrsaí Comhshaoil

- Clár náisiúnta iniúchtaí agus cigireachtaí a dhéanamh gach bliain ar shaoráidí a bhfuil ceadúnas ón nGníomhaireacht acu.
- Maoirseacht a dhéanamh ar fhreagrachtaí cosanta comhshaoil na n-údarás áitiúil.
- Caighdeán an uisce óil, arna sholáthar ag soláthraithe uisce phoiblí, a mhaoirsiú.
- Obair le húdaráis áitiúla agus le gníomhaireachtaí eile chun dul i ngleic le coireanna comhshaoil trí chomhordú a dhéanamh ar líonra forfheidhmiúcháin náisiúnta, trí dhíríú ar chiontóirí, agus trí mhaoirsiú a dhéanamh ar leasúchán.
- Cur i bhfeidhm rialachán ar nós na Rialachán um Dhramhthrealamh Leictreach agus Leictreonach (DTLL), um Shrian ar Shubstaintí Guaiseacha agus na Rialachán um rialú ar shubstaintí a idíonn an ciseal ózóin.
- An dlí a chur orthu siúd a bhriseann dlí an chomhshaoil agus a dhéanann dochar don chomhshaoil.

Bainistíocht Uisce

- Monatóireacht agus tuairisciú a dhéanamh ar cháilíocht aibhneacha, lochanna, uiscí idirchriosacha agus cósta na hÉireann, agus screamhuisc; leibhéil uisce agus sruthanna aibhneacha a thomhas.
- Comhordú náisiúnta agus maoirsiú a dhéanamh ar an gCreat-Treoir Uisce.
- Monatóireacht agus tuairisciú a dhéanamh ar Cháilíocht an Uisce Snámha.

Monatóireacht, Anailís agus Tuairisciú ar an gComhshaoil

- Monatóireacht a dhéanamh ar cháilíocht an aeir agus Treoir an AE maidir le hAer Glan don Eoraip (CAFÉ) a chur chun feidhme.
- Tuairisciú neamhspleách le cabhrú le cinnteoireacht an rialtais náisiúnta agus na n-údarás áitiúil (*m.sh. tuairisciú tréimhsiúil ar staid Chomhshaoil na hÉireann agus Tuarascálacha ar Tháscairí*).

Rialú Astaíochtaí na nGás Ceaptha Teasa in Éirinn

- Fardail agus réamh-mheastacháin na hÉireann maidir le gáis cheaptha teasa a ullmhú.
- An Treoir maidir le Trádáil Astaíochtaí a chur chun feidhme i gcomhair breis agus 100 de na táirgeoirí dé-ocsaíde carbóin is mó in Éirinn.

Taighde agus Forbairt Comhshaoil

- Taighde comhshaoil a chistiú chun brúnna a shainaitheint, bonn eolais a chur faoi bheartais, agus réitigh a sholáthar i réimsí na haeráide, an uisce agus na hinbhuanaitheachta.

Measúnacht Straitéiseach Timpeallachta

- Measúnacht a dhéanamh ar thionchar pleananna agus clár beartaithe ar an gcomhshaoil in Éirinn (*m.sh. mórphleananna forbartha*).

Cosaint Raideolaíoch

- Monatóireacht a dhéanamh ar leibhéil radaíochta, measúnacht a dhéanamh ar nochtadh mhuintir na hÉireann don radaíocht ianúcháin.
- Cabhrú le pleananna náisiúnta a fhorbairt le haghaidh éigeandálaí ag eascairt as taismí núicléacha.
- Monatóireacht a dhéanamh ar fhorbairtí thar lear a bhaineann le saoráidí núicléacha agus leis an tsábháilteacht raideolaíochta.
- Sainseirbhísí cosanta ar an radaíocht a sholáthar, nó maoirsiú a dhéanamh ar sholáthar na seirbhísí sin.

Treoir, Faisnéis Inrochtana agus Oideachas

- Comhairle agus treoir a chur ar fáil d'earnáil na tionsclaíochta agus don phobal maidir le hábhair a bhaineann le caomhnú an chomhshaoil agus leis an gcosaint raideolaíoch.
- Faisnéis thráthúil ar an gcomhshaoil ar a bhfuil fáil éasca a chur ar fáil chun rannpháirtíocht an phobail a spreagadh sa chinnteoireacht i ndáil leis an gcomhshaoil (*m.sh. Timpeall an Tí, léarscáileanna radóin*).
- Comhairle a chur ar fáil don Rialtas maidir le hábhair a bhaineann leis an tsábháilteacht raideolaíoch agus le cúrsaí práinnfhreagartha.
- Plean Náisiúnta Bainistíochta Dramhaíola Guaisí a fhorbairt chun dramhaíl ghuaiseach a chosaint agus a bhainistiú.

Múscailt Feasachta agus Athrú Iompraíochta

- Feasacht comhshaoil níos fearr a ghiniúint agus dul i bhfeidhm ar athrú iompraíochta dearfach trí thacú le gnóthais, le pobail agus le teaghlacha a bheith níos éifeachtúla ar acmhainní.
- Tástáil le haghaidh radóin a chur chun cinn i dtithe agus in ionaid oibre, agus gníomhartha leasúcháin a spreagadh nuair is gá.

Bainistíocht agus struchtúr na Gníomhaireachta um Chaomhnú Comhshaoil

Tá an ghníomhaíocht á bainistiú ag Bord lánaimseartha, ar a bhfuil Ard-Stiúrthóir agus cúigear Stiúrthóirí. Déantar an obair ar fud cúig cinn d'Oifigí:

- An Oifig um Inmharthanacht Comhshaoil
- An Oifig Forfheidhmithe i leith cúrsaí Comhshaoil
- An Oifig um Fianaise is Measúnú
- Oifig um Chosaint Radaíochta agus Monatóireachta Comhshaoil
- An Oifig Cumarsáide agus Seirbhísí Corparáideacha

Tá Coiste Comhairleach ag an nGníomhaireacht le cabhrú léi. Tá dáréag comhaltaí air agus tagann siad le chéile go rialta le plé a dhéanamh ar ábhair inní agus le comhairle a chur ar an mBord.



Headquarters
PO Box 3000, Johnstown Castle Estate
County Wexford, Y35 W821, Ireland
Bosca Poist 3000, Eastát Chaisleán Bhaile Sheáin
Contae Loch Garman, Y35 W821, Éire

T: +353 53 9160600
F: +353 53 9160699
E: info@epa.ie
W: www.epa.ie
Lo Call: 1890 33 55 99

EPA Regional Inspectorate Dublin
McCumiskey House
Richview
Clonskeagh Road
Dublin 14
D14 YR62
Tel: 01-268 0100
Fax: 01-268 0199

EPA Regional Inspectorate Cork
Inniscarra
Co. Cork
P31 VX59
Tel: 021-4875540
Fax: 021-4875545

EPA Regional Inspectorate Castlebar
John Moore Road
Castlebar
Co. Mayo
F23 KT91
Tel: 094-9048400
Fax: 094-9021934

EPA Regional Inspectorate Kilkenny
Seville Lodge
Callan Road
Kilkenny
R95 ED28
Tel: 056-7796700
Fax: 056-7796798

EPA Regional Inspectorate Monaghan
The Glen
Monaghan
H18 YT02
Tel: 047-77600
Fax: 047-84987

E: info@epa.ie
W: www.epa.ie
LoCall: 1890 33 55 99



Printed on an environmentally friendly paper stock